## The Failure of the Ergodic Assumption

M. Ignaccolo<sup>1</sup>, M. Latka<sup>2</sup> and B.J. West<sup>1,3</sup>

The well established procedure of constructing phenomenological ensemble from a single long time series is investigated. It is determined that a time series generated by a simple Uhlenbeck-Ornstein Langevin equation is mean ergodic. However the probability ensemble average yields a variance that is different from that determined using the phenomenological ensemble (time average). We conclude that the latter ensemble is often neither stationary nor ergodic and consequently the probability ensemble averages can misrepresent the underlying dynamic process.

PACS numbers:

The ergodic hypothesis, which states the equivalence of time averages and phase space averages, began with Boltzmann [\[1\]](#page-3-0), who conjectured that a single trajectory can densely cover a surface of constant energy in phase space. This 'proof 'of the hypothesis as well as many subsequent proofs were subsequently shown to be fatally flawed. It was not until metric decomposability was introduced by Birkoff [\[2](#page-3-1)] that a rigorous mathematical theory of ergodicity began to take shape. Kinchin [\[3](#page-3-2)], who wrote a seminal work on the mathematical foundations of statistical mechanics, offered a second meaning for the ergodic hypothesis, that being, to assume the truth of the hypothesis and judge the theory constructed on this basis by its practical success or failure. This latter perspective is the one adopted by the vast majority of physicists, with the subsequent replacement of phase space averages with averages over an ensemble probability density. We call this latter use of ergodicity the 'ergodic assumption'.

The ergodic assumption was quite evident in quantum measurements, which historically employed single particle ensembles to evaluate averages. However the advances in instrumentation and measurement techniques over the past decade or so enabled the measurement of single molecule time series and these relatively recent experiments forced the ergodic assumption out of the shadows into the foreground. Margolin and Barkai [\[4](#page-3-3)] have argued that a wide variety of complex phenomena whose dynamics consist of random switching between two states, such as the fluorescent intermittency of single molecules [\[5](#page-3-4)] and nanocrystals [\[6\]](#page-3-5) may display non-ergodicity, which is to say the ergodic assumption breaks down. The break down occurs because the probability density for sojourning within one of the two states is given by the inverse power law  $\psi(t) \propto t^{-\mu}$  so that when  $\mu < 2$  the average sojourn time within a state diverges and the time series is manifestly non-ergodic . A similar breakdown is obseved in neuroscience where the cognitive activity associated with perceptual judgement is determined to span the interval  $1 < \mu < 3$  [\[8](#page-3-6)] and consequently to manifest non-ergoic behavior in a variety of experiments [\[9,](#page-3-7) [10\]](#page-3-8).

In this Letter we show that the ergodic assumption is even more delicate than determined in these experiments, which is to say it is less applicable to complex networks in general, or physical networks in particular, than was previously believed. To establish the extent of the failure

of the ergodic assumption we begin with a review of its introduction into the physics literature.

A stochastic physical process is denoted by a dynamic variable  $X(t)$  that can be the velocity of a particle undergoing Brownian motion, the erratic voltage measured in an electroencephalogram or the time series from any of a vast number of other complex physical or physiological phenomena. In any event in the absence of direct evidence to the contrary it is assumed that the underlying process is stationary in time implying that this observable can be determined experimentally from a single historical record  $X(t)$  measured over a sufficiently long time. As stated in a classic paper by Wang and Uhlenbeck (WU) in 1945 [\[11\]](#page-3-9):

..One can then cut the record in pieces of length T (where T is long compared to all "periods" occurring in the process), and one may consider the different pieces as the different records of an ensemble of observations. In computing average values one has in general to distinguish between an ensemble average and a time average. However, for a stationary process these two ways of averaging will always give the same result...

This technique for generating empirical realizations of physical ensembles subsequently proved to be very useful and has been the form of the ergodic assumption adopted to determine average quantities from time series for three generations of scientists. The stationarity assumption implies that single variable distributions are independent of time, so that their ensemble averages and time averages are the same. Moreover, the joint distribution functions for pairs of dynamic variables depends only on the difference in time between the measurements of the two variables. Formally, this implies that the difference variable  $Z(t, \tau) = X(t + \tau) - X(t)$  is only a function of the time difference  $\tau$ , that is,  $Z(t, \tau) = Z(\tau)$ . However a stationary time series need not have stationary increments and the difference variable can depend on t and  $\tau$ separately. Since the introduction of phenomenological ensembles over a half century ago the phenomena of interest have become more complex and consequently the WU approach to evaluating averages is often not adequate. To establish this inadequacy we go back to the prequel of WU, another classic, and consider a process first analyzed in detail by Uhlenbeck and Ornstein (UO) [\[12\]](#page-3-10) in 1930.

Let us consider the UO stochastic process about which we know everything so as to test the ergodic assumption. Consider the UO Langevin equation

<span id="page-1-0"></span>
$$
\frac{dX(t)}{dt} = -\lambda X(t) + F(t)
$$
\n(1)

where in a physics context  $X(t)$  is the one-dimensional velocity, the particle mass is set to one,  $\lambda$  is the dissipation parameter and  $F(t)$  is a random force. In other contexts the dynamic variable has been taken to be the voltage measured in an EEG record [\[13](#page-3-11)], or any of a number of other interesting observables. The solution to the UO Langevin equation, as we show below, actually violates the stationarity assumption made by Wang and Uhlenbeck in 1945. To establish this violation we calculate the variance using the phenomenological ensemble and find that it is different from that obtained using the multi-trajectory ensemble (MTE), that is, the ensemble obtained by solving the stochastic differential equation [\(1\)](#page-1-0) using an ensemble of realizations of the random force. The significance of this result can not be overestimated since it directly contradicts a half century of analyses made assuming the ergodicity of the phenomenological ensembles constructed from the time series.

The formal solution of the OU Langevin equation is, with the initial condition  $X(0)$ ,

<span id="page-1-1"></span>
$$
X(t) = e^{-\lambda t} \left[ X(0) + \int_{0}^{t} F(t') e^{\lambda t'} dt' \right].
$$
 (2)

A proper interpretation of this solution requires the specification of the statistics of the random force. If we assume the random force is a zero-centered delta correlated Gaussian process, as is often done, then the solution [\(2\)](#page-1-1) is generally accepted as a stationary ergodic process.

Recall that an ergodic process is one for which the time average of an analytic function of the dynamic variable

<span id="page-1-2"></span>
$$
\langle g(X) \rangle_T \equiv \lim_{T \to \infty} \frac{1}{T} \int_0^T g(X(t')) dt' \tag{3}
$$

and the MTE average

$$
\langle g(X) \rangle_{MTE} \equiv \int g(x)P(x)dx \tag{4}
$$

are equal

<span id="page-1-3"></span>
$$
\langle g(X) \rangle_T = \langle g(X) \rangle_{MTE} \,. \tag{5}
$$

If  $g(x) = x$ , then [\(3\)](#page-1-2) is a generalization of the law of large numbers and in the limit of long time lags the time average of the dynamic variable exists in proba-bility [\[14\]](#page-3-12). The process in this case is *mean ergodic*, or first-order ergodic, which requires that the autocorrelation function have the appropriate asymptotic behavior, that being, for  $C_X(\tau) = \langle X(t)X(t + \tau)\rangle_T$  we must have  $\frac{1}{2T} \int_{0}^{T}$  $\int\limits_0^{\infty} C_X(\tau) d\tau \to 0$  as  $T \to \infty$  in which case [\(5\)](#page-1-3) is true [\[15](#page-3-13)]. If we consider the quadratic variable  $Z(t) = X(t)X(t + \tau)$  then  $Z(t)$  is mean ergodic if  $rac{1}{2T}$  $\int$  $\int_{0}^{1} C_{Z}(\tau) d\tau \to 0$  as  $T \to \infty$ , which implies that  $X(t)$  is second-order ergodic. The question remains whether the phenomenological ensemble averages constructed from a single historical trajectory of long but finite length T satisfies [\(5\)](#page-1-3) for any  $g(x)$  other than linear as explicitly assumed in WU. It is probably obvious but it should be said that first-order ergodicity is important in statistical physics because it guarantees that the microcanonical ensemble of Gibbs produces the correct statistical average in phase space for Hamiltonian systems [\[14](#page-3-12)].

To answer the question regarding the validity of [\(5\)](#page-1-3) for quantities higher than first-order consider the solution to the UO Langevin equation, which since the dynamic equation is linear has the same statistics as the random force. Assuming the random force is a zero-centered delta correlated Gaussian process, we know that the solution [\(2\)](#page-1-1) is completely determined by the mean and variance. The solution with  $X(0) = 0$  yields the average values for the dynamic variable  $\langle X \rangle_T = \langle X \rangle_{MTE} = 0$ . The autocorrelation function is  $C_X(\tau) = \frac{\sigma_F^2}{2\lambda} e^{-\lambda \tau}$  so that the condition  $\frac{1}{2T} \int_{0}^{T}$  $\int\limits_0^{\cdot} C_X(\tau) d\tau \to 0$  as  $T \to \infty$  is satisfied and consequently the solution to the UO Lagevin equation is first-order ergodic.

Now let us consider the variance using the MTE averages consisting of an infinite number of trajectories all starting from  $X(0) = 0$  (without loss of generality) yielding

<span id="page-1-4"></span>
$$
\sigma^{2}(t) \equiv \left\langle X^{2}(t) \right\rangle_{MTE} - \left\langle X(t) \right\rangle_{MTE}^{2} = \frac{\sigma_{F}^{2}}{2\lambda} \left[ 1 - e^{-2\lambda t} \right]. \tag{6}
$$

Note that [\(6\)](#page-1-4) is also the variance obtained using the Gaussian solution to the Fokker-Planck equation for the ensemble probability distribution.

Now let us examine the single trajectory case. To create an ensemble of trajectories consider different portions of the single trajectory. Let  $X(t)$   $(X(0)=0)$  be a single trajectory of maximum finite length  $T$  and consider the set of trajectories for  $t \in [0, T - \tau]$ 

<span id="page-1-5"></span>
$$
Z(t,\tau) = X(t+\tau) - X(t). \tag{7}
$$

Using [\(2\)](#page-1-1), we can write  $X(t + \tau)$  as

<span id="page-2-0"></span>
$$
X(t+\tau) = e^{-\lambda \tau} \left[ X(t) + \int_{0}^{\tau} F(t+t')e^{\lambda t'}dt' \right]
$$
 (8)

where the initial condition for the trajectory at  $t + \tau$  is the time series at time  $t$ . Inserting  $(8)$  into  $(7)$ , we obtain for  $t \in [0, T - \tau]$ 

<span id="page-2-1"></span>
$$
Z(t,\tau) = e^{-\lambda\tau} \int_{0}^{\tau} F(t+t')e^{\lambda t'}dt' + X(t) \left[e^{-\lambda\tau} - 1\right]
$$
\n(9)

Note that the first term of the rhs of [\(9\)](#page-2-1) is the solution of the OU Langevin equation [\(1\)](#page-1-0) with  $t=\tau$  and  $X(0)=0$ , apart from the fact that in the integrand of [\(9\)](#page-2-1) only the time in the range  $[t, t+\tau]$  are considered for the function F (instead of the range  $(0, \tau)$  as in [\(2\)](#page-1-1)). We now define the solution starting from zero to be

<span id="page-2-2"></span>
$$
X_0(t,\tau) = e^{-\lambda \tau} \int_0^{\tau} F(t+t')e^{\lambda t'}dt' \qquad (10)
$$

so that the term  $X(t)$  in [\(9\)](#page-2-1) is equivalent to  $X_0(0, t)$ , and therefore with  $t \in [0, T - \tau]$  [\(9\)](#page-2-1) reduces to

$$
Z(t,\tau) = X_0(t,\tau) + X_0(0,t) \left[ e^{-\lambda \tau} - 1 \right]
$$
 (11)

This equation reveals that the statistics of the variable  $Z(t, \tau)$  depends on the particular value of t through the second term on the rhs. In fact the term  $X_0(0, t)$  has zero mean and variance that depends on t according to [\(6\)](#page-1-4). Also notice that the first term of the above equation depends on  $t$  but the "dependence" is a time translation of the variable  $F$  in the integrand of [\(10\)](#page-2-2). But the random force  $F$  is stationary by assumption and thus  $X_0(t, \tau)$  is statistically equivalent to  $X_0(0, \tau)$ . The function  $Z(t, \tau)$  is the sum of two statistically independent Gaussian variables and therefore is itself a Gaussian variable with  $\langle Z(t, \tau) \rangle_T = \langle Z(t, \tau) \rangle_{MTE} = 0$  and secondmoment

<span id="page-2-3"></span>
$$
\langle Z^2(t,\tau) \rangle_{MTE} = \sigma^2(\tau) + \left[ e^{-\lambda \tau} - 1 \right]^2 \sigma^2(t). \tag{12}
$$

We use this variance to demonstrate that the phenomenological ensemble created is not equivalent to MTE because the trajectories  $Z(t, \tau)$  are not statistically equivalent to those of  $X(t)$ . We show that  $\sigma_{straj}^2$ , the variance calculated with the ensemble  $Z(t, \tau)$ , is different from that calculated using the MTE [\(6\)](#page-1-4).

To calculate the variance  $\sigma_{straj}^2(\tau)$  using the phenomenological ensemble, we simply need to calculate the mean value (among the trajectories) using [\(12\)](#page-2-3). Here the variance is calculated using the time average since only the time series is assumed to be available to us

$$
\sigma_{straj}^2(\tau) = \frac{1}{T - \tau} \int_0^{T - \tau} dt \left\langle Z^2(t, \tau) \right\rangle_{MTE}.
$$
 (13)



FIG. 1: The variance calculated for the OU process using the single long time series and the phenomenological WU ensemble. Note the difference in the dissipation rates for the two ways of calculating the variance.

Inserting the mean of the phenomenological second moment into the integrand yields

$$
\sigma_{straj}^2(\tau) = \frac{1}{T - \tau} \int\limits_0^{T - \tau} dt \left\{ \sigma^2(\tau) + \left[ e^{-\lambda \tau} - 1 \right]^2 \sigma^2(t) \right\}
$$
\n(14)

and using [\(6\)](#page-1-4) allows to write after integrating over time

$$
\sigma_{straj}^2(\tau) = \sigma^2(\tau) + \left[e^{-\lambda \tau} - 1\right]^2
$$

$$
\times \left[\frac{\sigma_F^2}{2\lambda}\right] \left[1 + \frac{1}{T - \tau} \frac{e^{-2\lambda(T-\tau)} - 1}{2\lambda}\right] (15)
$$

Since  $\lambda T \gg 1$  (because T is the length of the record) we can write

$$
\sigma_{straj}^2(\tau) \approx \sigma^2(\tau) + \left[e^{-\lambda \tau} - 1\right]^2 \left[\frac{\sigma_F^2}{2\lambda}\right] \tag{16}
$$

and substitution from [\(6\)](#page-1-4) yields

$$
\sigma_{straj}^2(\tau) \approx \frac{\sigma_F^2}{\lambda} (1 - e^{-\lambda \tau}). \tag{17}
$$

The last expression shows that the phenomenological ensemble trajectories  $Z(t, \tau)$  behave as a MTE with an effective dissipation that is half the dissipation rate in the Langevin equation generating the process, that is,

$$
\lambda_{eff} = \lambda/2. \tag{18}
$$

Consequently, it takes half as long for the MTE variance to decay as it does for the phenomenological variance of the time series.

Consider what has been established in this Letter. First a long time series is generated numerically from the U) Langevin equation. This long series is separated into a large number of equal length time series as prescribed by Wang and Uhlenbeck [\[11](#page-3-9)] to form a phenomenological ensemble of realizations of the UO process. This phenomenological ensemble of trajectories is shown to be mean ergodic. However when they are used to calculate the variance of the process anticipating that the same result will be obtained as that for a MTE average, the probability ensemble average, of the variance. The expected result turns out to be wrong. The dissipation parameter determined by the MTE variance is twice that obtained

- <span id="page-3-0"></span>[1] L. Boltzmann, Lectures on Gas Theory, first published in 1895, translated by S.G. Brush, Dover, New York (1995).
- <span id="page-3-1"></span>[2] G.D. Birkoff, "Proof of the Ergodic Theorem", PNAS 17, 656-660 (1931).
- <span id="page-3-2"></span>[3] A.I. Kinchin, Mathematical Foundaitons of Statistical Mechanics, translated by G. Gamow, Dover, New York (1949).
- <span id="page-3-3"></span>[4] G. Margolin and E. Barkai, "Nonergodicity of blinking nanocrystals and other Lévy-walk processes", Phys. Rev. Lett. **94**, 080601 (2005).
- <span id="page-3-4"></span>[5] M. Haase, c.G. Hübner, E. Reuther, A. Herrmann, K. Müllen and Th. Basché, J. Phys. Chem. B 108, 10445 (2004).
- <span id="page-3-5"></span>[6] M. Nirmal, B.O. Dabbousi, M.G. Bawendi, J.J. Macklin, J.K. Trautman, T.D. Harris, and L.E. Brus, Nature 383, 802 (1996); X. Brokmann, J. P. Hermier, G. Messin, P. Desbiolles, J. P. Bouchaud and M. Dahan, Phys. Rev. Lett. **90**, 120601 (2003).
- [7] B.J. West, E.L. Geneston and P. Grigolini, "Maximizing information exchange between complex networks", Phys.

using the phenomenological ensemble. The inescapable conclusion is that the phenomenological prescription for calculating the average of any nonlinear function of the dynamic variable using time series is not equivalent to that using a MTE. The fact that [\(5\)](#page-1-3) is not satisfied for  $q(x) \neq x$  implies that the ergodic assumption is violated by any finite time phenomenological ensemble. Given the simplicity of the dynamic model discussed here; it is mean ergodic; combined with the recent findings concerning the non-ergodicity of experimental data mentioned earlier, it is not unreasonable to conclude that the application of the ergodic assumption in general is probably unwarranted.

Rept. 468, 1-99 (2008).

- <span id="page-3-6"></span>[8] P. Grigolini, G. Aquino, M. Bologna, M. Lukovic and B.J. West, "A theory of 1/f noise in human cognition", Physica A 388, 4192 (2009).
- <span id="page-3-7"></span>[9] D.L. Gilden, T. Thornton, and M.W. Mallon, "1/f noise in Human Cognition", Science 267, 1837 (1995).
- <span id="page-3-8"></span>[10] J. correll, "1/f noise and effort on implicit measures of bias", J. Personality and Social Psych. 94, 48 (2008).
- <span id="page-3-9"></span>[11] M.C. Wang and G.E. Uhlenbeck, "On the theory of the Brownian motion II", Rev. Mod. Phys. 17, 323 (1945).
- <span id="page-3-10"></span>[12] G.E. Uhlenbeck and L.S. Ornstein, "On the theory of the Brownian motion", Phys. Rev. 36, 823 (1930).
- <span id="page-3-11"></span>[13] M. Ignaccolo, M. Latka, W. Jernajczyk, P. Grigolini and B.J. West, "The dynamics of EEG entropy", J. Biol. Phys. (2009).
- <span id="page-3-12"></span>[14] J.L. McCauley, "Time vs. ensemble averages for nonstationary time series", Physica a 387 , 5518-5522 (2008).
- <span id="page-3-13"></span>[15] B.V. Gnedeko, *Theory of Probability* (trans. from Russian by B.D. Seckler) Chelsea, NY (1968).