# Solution of boundary value and eigenvalue problems for second order elliptic operators in the plane using pseudoanalytic formal powers

Raúl Castillo Pérez<sup>1</sup>, Vladislav V. Kravchenko<sup>2</sup> and  $Rabindranath Reséndiz Vázquez<sup>1</sup>$ <sup>1</sup>SEPI, ESIME Zacatenco, Instituto Politécnico Nacional, Av. IPN S/N, C.P. 07738, D.F. MEXICO

 $^2$ Department of Mathematics, CINVESTAV del IPN, Unidad Queretaro, Libramiento Norponiente No. 2000, Fracc. Real de Juriquilla, Queretaro, Qro. C.P. 76230 MEXICO e-mail: vkravchenko@qro.cinvestav.mx<sup>∗</sup>

October 30, 2018

#### Abstract

We propose a method for solving boundary value and eigenvalue problems for the elliptic operator  $D = \text{div } p \text{ grad } +q$  in the plane using pseudoanalytic function theory and in particular pseudoanalytic formal powers. Under certain conditions on the coefficients  $p$  and  $q$  with the aid of pseudoanalytic function theory a complete system of null solutions of the operator can be constructed following a simple algorithm consisting in recursive integration. This system of solutions is used for solving boundary value and spectral problems for the operator  $D$  in bounded simply connected domains. We study theoretical and numerical aspects of the method.

## 1 Introduction

The main numerical techniques for solving problems related to elliptic linear partial differential equations with variable coefficients in one way or another involve a discretization of a domain and solution of systems of thousands of algebraic equations. Seldom the method of separation of variables is applied due to its natural limitations related to the requirements of a complete agreement between the geometry of the domain and the symmetry of the coefficients. Moreover, the method of separation of variables implies solution of Sturm-Liouville spectral problems which is not an easy task itself.

In the present paper we propose a different method based on some old and new results from pseudoanalytic function theory [\[3\]](#page-19-0), [\[15\]](#page-20-0). Its applicability is not so universal as the

<sup>∗</sup>Research was supported by CONACYT, Mexico via the research project 50424. The first named author wishes to thank support from the CONACYT and the National Polytechnic Institute for the possibility of a postdoctoral stay in the Department of Mathematics of the CINVESTAV in Queretaro, as well as from the SIBE program of the National Polytechnic Institute, Mexico.

applicability of the finite difference method or the finite element method. First of all, it is applicable to problems in bounded domains in the plane and up to now only for the operator  $div p grad + q$ . Moreover, at present we can apply the method only when the equation

<span id="page-1-0"></span>
$$
(\operatorname{div} p \operatorname{grad} + q) u(x, y) = 0 \tag{1}
$$

possesses a particular solution  $u_0$  such that the function  $f = p^{1/2}u_0$  is sufficiently smooth, nonvanishing in the domain of interest and representable in the form  $f = S(s)T(t)$  where s and t represent an orthogonal coordinate system. This separable form of f may cause associations with the method of separation of variables. Nevertheless this is a completely different technique, based on different ideas and free of the mentioned above limitations of the method of separation of variables.

The heart of the method is the construction of a complete system of solutions for [\(1\)](#page-1-0) in the domain of interest, complete in the sense explained below (see [\[7,](#page-19-1) Sect. 1.3] for related ideas and additional details). The system of solutions is used for approximating the solution of a boundary value problem. Due to the linearity of equation [\(1\)](#page-1-0) after the construction of the system of solutions the problem reduces to approximation of boundary conditions, for which a variety of methods can be used. Here we apply the collocation method.

The complete system of solutions is constructed in the following way. The knowledge of a particular solution of [\(1\)](#page-1-0) allows us to propose a corresponding Vekua equation [\[19\]](#page-20-1) closely related to [\(1\)](#page-1-0) in the sense that the real part of any of its solutions has the form  $p^{1/2}u$ where u is a solution of  $(1)$ , and vice versa given u one can easily construct a corresponding solution of the Vekua equation [\[12\]](#page-20-2), [\[15\]](#page-20-0). The relation between [\(1\)](#page-1-0) and the Vekua equation is similar to the relation between the Laplace equation and the Cauchy-Riemann system. L. Bers developed [\[3\]](#page-19-0), [\[4\]](#page-19-2) a theory of so-called pseudoanalytic formal powers. They are generalizations of the analytic powers  $(z - z_0)^n$  in the sense that they are solutions of the corresponding Vekua equation and behave locally like the analytic powers. The theory of Bers includes generalizations of Taylor series, Runge's theorem and other basic facts from analytic function theory. Thus, under certain quite natural conditions the system of pseudoanalytic formal powers is complete in the space of all pseudoanalytic functions (solutions of the Vekua equation) in the same sense as the system of powers  $(z - z_0)^n$  is complete in the space of analytic functions. To construct the pseudoanalytic formal powers the knowledge of a corresponding generating sequence is required. Recently [\[14\]](#page-20-3), [\[15\]](#page-20-0) an algorithm for construction of generating sequences under additional conditions on the coefficients in the Vekua equation was proposed. This implies that when  $f = p^{1/2}u_0$  is representable in a separable form the complete system of formal powers for the Vekua equation associated with [\(1\)](#page-1-0) can be constructed explicitly following Bers' recursive procedure.

We investigate the efficiency of the proposed method which we call MPFP, the Method of Pseudoanalytic Formal Powers. We show its fast convergence and compare its accuracy with that of the finite element method. In general, we show that in problems addmitting the explicit construction of formal powers and hence the application of the MPFP its use is advantageous compared to other computational techniques based on discretization of the problem.

It is worth mentioning that the MPFP is a direct generalization of the method of harmonic polynomials for solving boundary value problems for the Laplace equation which has been considered in dozens of works (see, e.g., [\[5\]](#page-19-3), [\[10\]](#page-19-4), [\[11\]](#page-20-4), [\[18\]](#page-20-5)). Indeed, in a special case when  $p \equiv 1, q \equiv 0$  and  $u_0 \equiv 1$  the corresponding complete system of solutions constructed by means of the MPFP coincides with the system of harmonic polynomials {Re(z - z<sub>0</sub>)<sup>n</sup>, Im(z - z<sub>0</sub>)<sup>n</sup>} $_{n=0}^{\infty}$ .

The knowledge of a complete system of solutions for an equation corresponding to any value of a spectral parameter allows one to use it for solving eigenvalue problems. We consider this possibility in section [6.](#page-18-0) The numerical results are highly promising, and it is clear that in the case of eigenvalue problems as well as for boundary value problems further work should be done in investigation of optimal ways of application of the MPFP. For example, for solving eigenvalue problems by means of the MPFP we used the simplest possible idea reducing the problem to calculation of zeros of a certain determinant obtained by evaluating the first N solutions from the constructed complete system in  $N$  points on the boundary of the domain under consideration. Meanwhile, in principle, this natural approach works there exist other techniques offering different ways of using the available exact solution systems (see [\[2,](#page-19-5) Sect. 1.13], where similar questions are discussed).

## <span id="page-2-2"></span>2 Factorization of the operator div  $p$  grad  $+q$ .

Let  $\Omega$  be a domain in  $\mathbb{R}^2$ . Throughout the whole paper we suppose that  $\Omega$  is a simply connected domain. Denote  $\partial_{\overline{z}} = \frac{1}{2}$  $\frac{1}{2} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$  and  $\partial_z = \frac{1}{2}$  $\frac{1}{2} \left( \frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right)$ . By C we denote the operator of complex conjugation.

Note that the operator  $\partial_{\overline{z}}$  applied to a real valued function  $\varphi$  can be regarded as a kind of gradient, and if we know that  $\partial_{\overline{z}}\varphi = \Phi$  in a whole complex plane or in a convex domain, where  $\Phi = \Phi_1 + i \Phi_2$  is a given complex valued function such that its real part  $\Phi_1$  and imaginary part  $\Phi_2$  satisfy the equation

<span id="page-2-1"></span>
$$
\partial_y \Phi_1 - \partial_x \Phi_2 = 0,\tag{2}
$$

then we can reconstruct  $\varphi$  up to an arbitrary real constant c in the following way

$$
\varphi(x,y) = 2\left(\int_{x_0}^x \Phi_1(\eta, y)d\eta + \int_{y_0}^y \Phi_2(x_0, \xi)d\xi\right) + c
$$

where  $(x_0, y_0)$  is an arbitrary fixed point in the domain of interest. Note that this formula can be easily extended to any simply connected domain by considering the integral along an arbitrary rectifiable curve Γ leading from  $(x_0, y_0)$  to  $(x, y)$ 

<span id="page-2-0"></span>
$$
\varphi(x,y) = 2\left(\int_{\Gamma} \Phi_1 dx + \Phi_2 dy\right) + c.
$$
\n(3)

By  $\overline{A}$  we denote the integral operator in [\(3\)](#page-2-0):

$$
\overline{A}[\Phi](x,y) = 2\left(\int_{x_0}^x \Phi_1(\eta,y)d\eta + \int_{y_0}^y \Phi_2(x_0,\xi)d\xi\right) + c.
$$

Thus if  $\Phi$  satisfies [\(2\)](#page-2-1), there exists a family of real valued functions  $\varphi$  such that  $\partial_{\overline{z}}\varphi = \Phi$ , given by the formula  $\varphi = \overline{A}[\Phi]$ .

The following result is in the core of the method proposed in the present work.

**Theorem 1** [\[13\]](#page-20-6) Let p and q be real valued functions,  $p \in C^2(\Omega)$  and  $p \neq 0$  in  $\Omega$ ,  $u_0$  be a positive particular solution of the equation

<span id="page-3-2"></span>
$$
(\text{div } p \,\text{grad } + q)u = 0 \qquad \text{in } \Omega. \tag{4}
$$

Then for any real valued continuously twice differentiable function  $\varphi$  the following equality holds

<span id="page-3-0"></span>
$$
\frac{1}{4}(\text{div } p \text{ grad} + q)\varphi = p^{1/2} \left(\partial_z + \frac{f_{\overline{z}}}{f}C\right) \left(\partial_{\overline{z}} - \frac{f_{\overline{z}}}{f}C\right) p^{1/2}\varphi, \tag{5}
$$

where

$$
f = p^{1/2}u_0.
$$
 (6)

**Remark 2** Let  $q \equiv 0$ . Then  $u_0$  can be chosen as  $u_0 \equiv 1$ . Hence [\(5\)](#page-3-0) gives us the equality

$$
\frac{1}{4}\operatorname{div}(p\operatorname{grad}\varphi) = p^{1/2}\left(\partial_z + \frac{\partial_{\overline{z}}p^{1/2}}{p^{1/2}}C\right)\left(\partial_{\overline{z}} - \frac{\partial_{\overline{z}}p^{1/2}}{p^{1/2}}C\right)(p^{1/2}\varphi).
$$

Let  $f$  be a real function of  $x$  and  $y$ . Consider the Vekua equation

<span id="page-3-1"></span>
$$
W_{\overline{z}} = \frac{f_{\overline{z}}}{f} \overline{W} \qquad \text{in } \Omega. \tag{7}
$$

This equation plays a crucial role in all that follows and hence we will call it the main Vekua equation. We notice that the operator of this equation is precisely the second factor in [\(5\)](#page-3-0).

Denote  $W_1 = \text{Re } W$  and  $W_2 = \text{Im } W$ .

<span id="page-3-3"></span>**Theorem 3** [\[13\]](#page-20-6) Let  $W = W_1 + iW_2$  be a solution of [\(7\)](#page-3-1). Assume that  $f = p^{1/2}u_0$ , where  $u_0$  is a positive solution of [\(4\)](#page-3-2) in  $\Omega$ . Then  $u = p^{-1/2}W_1$  is a solution of (4) in  $\Omega$ , and  $v = p^{1/2}W_2$  is a solution of the equation

<span id="page-3-4"></span>
$$
(\operatorname{div}\frac{1}{p}\operatorname{grad}+q_1)v=0 \qquad \text{in } \Omega,
$$
\n(8)

where

<span id="page-3-5"></span>
$$
q_1 = -\frac{1}{p} \left( \frac{q}{p} + 2 \left\langle \frac{\nabla p}{p}, \frac{\nabla u_0}{u_0} \right\rangle + 2 \left( \frac{\nabla u_0}{u_0} \right)^2 \right). \tag{9}
$$

<span id="page-3-6"></span>Theorem [3](#page-3-3) shows us that as much as real and imaginary parts of a complex analytic function are harmonic functions, the real and imaginary parts of a solution of the main Vekua equation [\(7\)](#page-3-1) multiplied by  $p^{-1/2}$  and  $p^{1/2}$  respectively are solutions of the associated elliptic equations [\(4\)](#page-3-2) and [\(8\)](#page-3-4). The following natural question arises then. We know that given an arbitrary real valued harmonic function in a simply connected domain, a conjugate harmonic function can be constructed explicitly such that the obtained couple of harmonic functions represent the real and imaginary parts of a complex analytic function. What is the corresponding more general fact for solutions of associated elliptic equations [\(4\)](#page-3-2) and [\(8\)](#page-3-4) (which we slightly generalizing the definition of I. N. Vekua call metaharmonic functions). The precise result is given in the following theorem.

**Theorem 4** [\[13\]](#page-20-6) Let  $f = p^{1/2}u_0$ , where  $u_0$  is a positive solution of [\(4\)](#page-3-2) in a simply connected domain  $\Omega$  and u be a solution of [\(4\)](#page-3-2). Then a solution v of [\(8\)](#page-3-4) with  $q_1$  defined by [\(9\)](#page-3-5) such that  $W = p^{1/2}u + i p^{-1/2}v$  is a solution of [\(7\)](#page-3-1), is constructed according to the formula

<span id="page-4-0"></span>
$$
v = u_0^{-1} \overline{A}(ipu_0^2 \partial_{\overline{z}}(u_0^{-1}u)).
$$
\n
$$
(10)
$$

Let v be a solution of [\(8\)](#page-3-4), then the corresponding solution u of [\(4\)](#page-3-2) such that  $W = p^{1/2}u +$  $ip^{-1/2}v$  is a solution of [\(7\)](#page-3-1), is constructed according to the formula

<span id="page-4-1"></span>
$$
u = -u_0 \overline{A}(ip^{-1}u_0^{-2}\partial_{\overline{z}}(u_0v)).
$$
\n(11)

**Remark 5** When  $p \equiv 1$ ,  $q \equiv 0$  and  $u_0 \equiv 1$ , equalities [\(10\)](#page-4-0) and [\(11\)](#page-4-1) turn into the well known formulas in complex analysis for constructing conjugate harmonic functions.

## 3 Formal powers

Briefly speaking formal powers are solutions of a Vekua equation

<span id="page-4-2"></span>
$$
W_{\overline{z}} = aW + b\overline{W} \tag{12}
$$

(with a and b being complex valued functions) generalizing the usual analytic powers  $\{(z-z_0)^n\}_{n=1}^{\infty}$  $n=0$ in the sense that locally when  $z \to z_0$  they behave asymptotically like the usual powers and under some additional conditions on the coefficients a and b they form a complete system in the space of all solutions of the Vekua equation in the same sense as the analytic powers  $\{(z-z_0)^n\}_{n=0}^{\infty}$  form a complete system in the space of analytic functions. Generalizations of the extension theorem, the Runge theorem and of other important results about the convergence of corresponding series are valid. The construction of formal powers is one of the main problems of pseudoanalytic function theory. Recently it was solved [\[14\]](#page-20-3), [\[15\]](#page-20-0) for a wide class of Vekua equations of the form [\(7\)](#page-3-1) which as was shown in the preceding section are of main interest for studying problems for second order equations of the form [\(4\)](#page-3-2).

The main ingredient for obtaining the explicit form of formal powers for a certain Vekua equation is the generating sequence, a concept introduced by Bers. If one knows a generating sequence for a given Vekua equation then the construction of formal powers reduces to a simple algorithm. Here we briefly explain the main ideas and steps refering the reader to [\[3\]](#page-19-0) and [\[15\]](#page-20-0) for further details.

#### 3.1 Generating pair and generating sequence

**Definition 6** A pair of solutions F and G of a Vekua equation [\(12\)](#page-4-2) in  $\Omega$  possessing partial derivatives with respect to the real variables x and y is said to be a generating pair if it satisfies the inequality

<span id="page-4-3"></span>
$$
\operatorname{Im}(FG) > 0 \qquad \text{in } \Omega. \tag{13}
$$

Condition [\(13\)](#page-4-3) implies that every complex function W defined in a subdomain of  $\Omega$ admits the unique representation  $W = \phi F + \psi G$  where the functions  $\phi$  and  $\psi$  are real valued. Thus, the pair  $(F, G)$  generalizes the pair  $(1, i)$  which corresponds to usual complex analytic function theory. The following expressions are known as characteristic coefficients of the pair  $(F, G)$ 

$$
a_{(F,G)} = -\frac{\overline{F}G_{\overline{z}} - F_{\overline{z}}\overline{G}}{F\overline{G} - \overline{F}G}, \qquad b_{(F,G)} = \frac{FG_{\overline{z}} - F_{\overline{z}}G}{F\overline{G} - \overline{F}G},
$$

$$
A_{(F,G)} = -\frac{\overline{F}G_{z} - F_{z}\overline{G}}{F\overline{G} - \overline{F}G}, \qquad B_{(F,G)} = \frac{FG_{z} - F_{z}G}{F\overline{G} - \overline{F}G}.
$$

If  $(F, G)$  is a generating pair of a Vekua equation [\(12\)](#page-4-2) then  $a_{(F,G)} = a$  and  $b_{(F,G)} = b$ . The other two characteristic coefficients are related to the concept of a derivative [\[3\]](#page-19-0). The  $(F, G)$ -derivative  $W = \frac{d_{(F, G)}W}{dz}$  of a continuously differentiable function W exists and has the  $\mathop{\mathrm{form}}$   $\blacksquare$ 

<span id="page-5-1"></span>
$$
W = W_z - A_{(F,G)}W - B_{(F,G)}\overline{W}
$$
\n(14)

if and only if

$$
W_{\overline{z}} = a_{(F,G)}W + b_{(F,G)}\overline{W}.
$$

Solutions of this equation are called  $(F, G)$ -pseudoanalytic functions.

**Definition 7** Let  $(F, G)$  and  $(F_1, G_1)$  - be two generating pairs in  $\Omega$ .  $(F_1, G_1)$  is called successor of  $(F, G)$  and  $(F, G)$  is called predecessor of  $(F_1, G_1)$  if

<span id="page-5-0"></span>
$$
a_{(F_1,G_1)} = a_{(F,G)} \qquad and \qquad b_{(F_1,G_1)} = -B_{(F,G)}.\tag{15}
$$

This definition arises naturally in relation to the notion of the  $(F, G)$ -derivative due to the following fact.

<span id="page-5-2"></span>**Theorem 8** Let W be an  $(F, G)$ -pseudoanalytic function and let  $(F_1, G_1)$  be a successor of  $(F, G)$ . Then W is an  $(F_1, G_1)$ -pseudoanalytic function.

Thus, to the difference of analytic functions whose derivatives are again analytic, the  $(F, G)$ -derivatives of pseudoanalytic functions are in general solutions of another Vekua equation with the coefficients given by [\(15\)](#page-5-0). Obviously this process of construction of new Vekua equations associated with the previous ones via relations [\(15\)](#page-5-0) can be continued and we arrive at the following definition.

**Definition 9** A sequence of generating pairs  $\{(F_m, G_m)\}\$ ,  $m = 0, \pm 1, \pm 2, \ldots$ , is called a generating sequence if  $(F_{m+1}, G_{m+1})$  is a successor of  $(F_m, G_m)$ . If  $(F_0, G_0) = (F, G)$ , we say that  $(F, G)$  is embedded in  $\{(F_m, G_m)\}.$ 

**Definition 10** A generating sequence  $\{(F_m, G_m)\}\)$  is said to have period  $\mu > 0$  if  $(F_{m+\mu}, G_{m+\mu})$ is equivalent to  $(F_m, G_m)$  that is their characteristic coefficients coincide.

We will need the following notation introduced by Bers. The  $(F, G)$ -integral is defined as follows

$$
\int_{\Gamma} W d_{(F,G)} z = F(z_1) \operatorname{Re} \int_{\Gamma} \frac{2 \overline{G}}{F \overline{G} - \overline{F} G} W dz - G(z_1) \operatorname{Re} \int_{\Gamma} \frac{2 \overline{F}}{F \overline{G} - \overline{F} G} W dz
$$

where  $\Gamma$  is a rectifiable curve leading from  $z_0$  to  $z_1$ .

Let W be an  $(F, G)$ -pseudoanalytic function. Using a generating sequence in which  $(F, G)$ is embedded we can define the higher derivatives of  $W$  by the recursion formula

$$
W^{[0]} = W; \qquad W^{[m+1]} = \frac{d_{(F_m, G_m)} W^{[m]}}{dz}, \quad m = 0, 1, \dots.
$$

A generating sequence defines an infinite sequence of Vekua equations. If for a given (original) Vekua equation we know not only a corresponding generating pair but the whole generating sequence, that is a couple of exact and independent solutions for each of the Vekua equations from the infinite sequence of equations corresponding to the original one, we are able to construct an infinite system of solutions of the original Vekua equation as is shown in the next definition.

<span id="page-6-1"></span>**Definition 11** The formal power  $Z_m^{(0)}(a, z_0; z)$  with center at  $z_0 \in \Omega$ , coefficient a and exponent 0 is defined as the linear combination of the generators  $F_m$ ,  $G_m$  with real constant coefficients  $\lambda$ ,  $\mu$  chosen so that  $\lambda F_m(z_0) + \mu G_m(z_0) = a$ . The formal powers with exponents  $n = 1, 2, \ldots$  are defined by the recursion formula

<span id="page-6-2"></span>
$$
Z_m^{(n)}(a, z_0; z) = n \int_{z_0}^z Z_{m+1}^{(n-1)}(a, z_0; \zeta) d_{(F_m, G_m)} \zeta.
$$
 (16)

This definition implies the following properties.

- 1.  $Z_m^{(n)}(a, z_0; z)$  is an  $(F_m, G_m)$ -pseudoanalytic function of z.
- 2. If a' and a'' are real constants, then  $Z_m^{(n)}(a'+ia'', z_0; z) = a'Z_m^{(n)}(1, z_0; z) + a''Z_m^{(n)}(i, z_0; z)$ .
- 3. The formal powers satisfy the differential relations

$$
\frac{d_{(F_m,G_m)}Z_m^{(n)}(a,z_0;z)}{dz} = nZ_{m+1}^{(n-1)}(a,z_0;z).
$$

4. The asymptotic formulas

$$
Z_m^{(n)}(a, z_0; z) \sim a(z - z_0)^n, \quad z \to z_0 \tag{17}
$$

hold.

Assume now that

<span id="page-6-0"></span>
$$
W(z) = \sum_{n=0}^{\infty} Z^{(n)}(a_n, z_0; z)
$$
 (18)

where the absence of the subindex  $m$  means that all the formal powers correspond to the same generating pair  $(F, G)$ , and the series converges uniformly in some neighborhood of  $z_0$ . It can be shown that the uniform limit of pseudoanalytic functions is pseudoanalytic, and that a uniformly convergent series of  $(F, G)$ -pseudoanalytic functions can be  $(F, G)$ -differentiated term by term. Hence the function W in [\(18\)](#page-6-0) is  $(F, G)$ -pseudoanalytic and its rth derivative admits the expansion

$$
W^{[r]}(z) = \sum_{n=r}^{\infty} n(n-1)\cdots(n-r+1)Z_r^{(n-r)}(a_n, z_0; z).
$$

From this the Taylor formulas for the coefficients are obtained

<span id="page-7-0"></span>
$$
a_n = \frac{W^{[n]}(z_0)}{n!}.
$$
\n(19)

**Definition 12** Let  $W(z)$  be a given  $(F, G)$ -pseudoanalytic function defined for small values of  $|z-z_0|$ . The series

<span id="page-7-1"></span>
$$
\sum_{n=0}^{\infty} Z^{(n)}(a_n, z_0; z)
$$
 (20)

with the coefficients given by  $(19)$  is called the Taylor series of W at  $z_0$ , formed with formal powers.

The Taylor series always represents the function asymptotically:

$$
W(z) - \sum_{n=0}^{N} Z^{(n)}(a_n, z_0; z) = O\left(|z - z_0|^{N+1}\right), \quad z \to z_0,
$$
\n(21)

for all N. This implies (since a pseudoanalytic function can not have a zero of arbitrarily high order without vanishing identically) that the sequence of derivatives  $\{W^{[n]}(z_0)\}\$  determines the function W uniquely.

If the series [\(20\)](#page-7-1) converges uniformly in a neighborhood of  $z_0$ , it converges to the function W.

#### <span id="page-7-5"></span>3.2 Convergence theorems

S. Agmon and L. Bers [\[1\]](#page-19-6) and L. Bers developed a theory of expansions in pseudoanalytic formal powers which in its generality is presented in [\[3\]](#page-19-0), [\[4\]](#page-19-2). We do need here the general results concerning a general Vekua equation [\(12\)](#page-4-2). Fortunately the situation with the main Vekua equation [\(7\)](#page-3-1) in a bounded simply connected domain under quite natural conditions on the function f is much easier than in the general case, and we have the following expansion theorem and Runge theorem [\[14\]](#page-20-3), [\[15\]](#page-20-0).

**Theorem 13** Let D be a disk of a finite radius R and center  $z_0$ , and  $f \in C^1(\overline{D})$  be positive in  $\overline{D}$ . Then any solution W of [\(7\)](#page-3-1) in D admits a unique normally convergent expansion<sup>[1](#page-7-2)</sup> of the form  $W(z) = \sum_{n=0}^{\infty} Z^{(n)}(a_n, z_0; z)$ .

<span id="page-7-3"></span>**Theorem 14** any solution W of  $(7)$  defined in a simply connected domain can be expanded into a normally convergent series of formal polynomials (linear combinations of formal powers with positive exponents).

Remark 15 This theorem admits a direct generalization onto the case of a multiply connected domain (see  $\vert 4 \vert$ ).

We mention here another important result obtained by Menke in [\[17\]](#page-20-7) which gives a useful estimate for the rate of convergence of the series from the preceding theorem in the case when  $W$  is a Hölder continuous function up to the boundary of the domain of interest.

<span id="page-7-4"></span><span id="page-7-2"></span><sup>&</sup>lt;sup>1</sup>Following [\[3\]](#page-19-0), [\[9\]](#page-19-7) we shall say that a sequence of functions  $W_n$  converges normally in a domain  $\Omega$  if it converges uniformly on every bounded closed subdomain of Ω.

**Theorem 16** Let W be a pseudoanalytic function in a domain  $\Omega$  bounded by a Jordan curve and satisfy the Hölder condition on  $\partial\Omega$  with the exponent  $\alpha$  ( $0 < \alpha \leq 1$ ). Then for any  $\varepsilon > 0$ and any natural n there exists a pseudopolynomial of order n satisfying the inequality

$$
|W(z) - P_n(z)| \le \frac{\text{Const}}{n^{\alpha - \varepsilon}} \qquad \text{for any } z \in \overline{\Omega}
$$

where the constant does not depend on n, but only on  $\varepsilon$ .

The following statements are direct corollaries of the relations established in section [2](#page-2-2) between pseudoanalytic functions (solutions of [\(7\)](#page-3-1)) and solutions of second-order elliptic equations, and of the convergence theorems formulated above. Here we assume the existence of a positive solution  $u_0$  of [\(4\)](#page-3-2) in the domain  $\Omega$  and the function f in [\(7\)](#page-3-1) to be defined by  $f = p^{1/2}u_0$  and belong to  $C^1(\overline{\Omega})$ .

**Definition 17** Let  $u(z)$  be a given solution of the equation [\(4\)](#page-3-2) defined for small values of  $|z-z_0|$ , and let  $W(z)$  be a solution of [\(7\)](#page-3-1) constructed according to theorem [4,](#page-3-6) such that  $\mathrm{Re}\,W=p^{1/2}u.$  The series

<span id="page-8-2"></span>
$$
p^{-1/2}(z) \sum_{n=0}^{\infty} \text{Re } Z^{(n)}(a_n, z_0; z)
$$
 (22)

with the coefficients given by  $(19)$  is called the Taylor series of u at  $z_0$ , formed with formal powers.

<span id="page-8-3"></span>**Theorem 18** [\[13\]](#page-20-6), [\[15\]](#page-20-0) Let  $u(z)$  be a solution of [\(4\)](#page-3-2) defined for  $|z - z_0| < R$ . Then it admits a unique expansion of the form

$$
u(z) = p^{-1/2}(z) \sum_{n=0}^{\infty} \text{Re } Z^{(n)}(a_n, z_0; z)
$$

which converges normally for  $|z - z_0| < R$ .

<span id="page-8-0"></span>**Theorem 19** An arbitrary solution of  $(4)$  defined in a simply connected domain where there exists a positive particular solution  $u_0$  such that  $f = p^{1/2}u_0 \in C^1(\overline{\Omega})$  can be expanded into a normally convergent series of formal polynomials multiplied by  $p^{-1/2}$ .

More precisely the last theorem has the following meaning. Due to Property 2 of formal powers we have that  $Z^{(n)}(a, z_0; z)$  for any Taylor coefficient a can be expressed through  $Z^{(n)}(1, z_0; z)$  and  $Z^{(n)}(i, z_0; z)$ . Then due to theorem [14](#page-7-3) any solution W of [\(7\)](#page-3-1) can be expanded into a normally convergent series of linear combinations of  $Z^{(n)}(1, z_0; z)$  and  $Z^{(n)}(i, z_0; z)$ . Consequently, any solution of [\(4\)](#page-3-2) can be expanded into a normally convergent series of linear combinations of real parts of  $Z^{(n)}(1, z_0; z)$  and  $Z^{(n)}(i, z_0; z)$  multiplied by  $p^{-1/2}$ .

Obviously, for solutions of [\(4\)](#page-3-2) the results on the interpolation and on the degree of approximation like, e.g., theorem [16](#page-7-4) are also valid.

Let us stress that theorem [19](#page-8-0) gives us the following result. The functions

<span id="page-8-1"></span>
$$
\left\{ p^{-1/2}(z) \operatorname{Re} Z^{(n)}(1, z_0; z), \quad p^{-1/2}(z) \operatorname{Re} Z^{(n)}(i, z_0; z) \right\}_{n=0}^{\infty} \tag{23}
$$

represent a complete system of solutions of [\(4\)](#page-3-2) in the sense that any solution of [\(4\)](#page-3-2) can be represented by a normally convergent series formed by functions [\(23\)](#page-8-1) in any simply connected domain  $\Omega$  where a positive solution of [\(4\)](#page-3-2) exists, and the rate of convergence of the series can be estimated with the aid of theorem [16.](#page-7-4)

#### 3.3 Explicit construction of generating sequences and formal powers

The results of section [2](#page-2-2) show us that the theory of the elliptic equation

$$
(\operatorname{div} p \operatorname{grad} + q)u = 0
$$

is closely related to equation [\(7\)](#page-3-1):

<span id="page-9-0"></span>
$$
W_{\overline{z}} = \frac{f_{\overline{z}}}{f} \overline{W}.
$$
\n(24)

It is interesting that for this equation we always know a generating pair. Namely, it is easy to see that the functions  $F = f$  and  $G = \frac{i}{f}$  $\frac{i}{f}$  satisfy [\(24\)](#page-9-0) together with the condition [\(13\)](#page-4-3). Then the corresponding characteristic coefficients  $A_{(F,G)}$  and  $B_{(F,G)}$  have the form

$$
A_{(F,G)} = 0,
$$
  $B_{(F,G)} = \frac{f_z}{f},$ 

and the  $(F, G)$ -derivative according to  $(14)$  is defined as follows

$$
W = W_z - \frac{f_z}{f} \overline{W} = \left(\partial_z - \frac{f_z}{f} C\right) W.
$$

Due to Theorem [8](#page-5-2) we obtain the following statement.

**Proposition 20** Let W be a solution of [\(24\)](#page-9-0). Then its  $(F, G)$ -derivative, the function  $w =$ W is a solution of the equation  $\left(\partial_{\overline{z}} + \frac{f_z}{f}C\right)w = 0.$ 

In spite of having given a generating pair for [\(24\)](#page-9-0) in general it is not known how to construct a corresponding generating sequence necessary for calculating the system of formal powers. Nevertheless a recent result from [\[14\]](#page-20-3), [\[15\]](#page-20-0) which we formulate in the following statement gives an answer to this question in a quite general situation.

<span id="page-9-1"></span>**Theorem 21** Let  $F = S(s)T(t)$  and  $G = \frac{i}{S(s)^{s}}$  $\frac{i}{S(s)T(t)}$  where S and T are arbitrary differentiable nonvanishing real valued functions,  $\Phi = s+it$  is an analytic function of the variable  $z = x+iy$ in  $\Omega$  such that  $\Phi_z$  is bounded and has no zeros in  $\Omega$ . Then the generating pair  $(F, G)$  is embedded in the generating sequence  $(F_m, G_m)$ ,  $m = 0, \pm 1, \pm 2, \ldots$  in  $\Omega$  defined as follows

$$
F_m = (\Phi_z)^m F
$$
 and  $G_m = (\Phi_z)^m G$  for even m

and

$$
F_m = \frac{(\Phi_z)^m}{S^2} F \quad and \quad G_m = (\Phi_z)^m S^2 G \quad \text{for odd } m.
$$

In order to appreciate the generality of this construction let us remind that orthogonal coordinate systems in a plane are obtained (see [\[16\]](#page-20-8)) from Cartesian coordinates  $x, y$  by means of the relation

$$
s + it = \Phi(x + iy)
$$

where  $\Phi$  is an arbitrary analytic function. Quite often a transition to more general coordinates is useful

$$
\xi = \xi(s), \quad \eta = \eta(t).
$$

 $\xi$  and  $\eta$  preserve the property of orthogonality. To illustrate the point, besides the obvious example of Cartesian coordinates which are generated by the analytic function  $z$  we give some other examples taken from [\[16\]](#page-20-8).

### Example 22 Polar coordinates

$$
s + it = \ln(x + iy),
$$
  
\n
$$
s = \ln \sqrt{x^2 + y^2}, \quad t = \arctan \frac{y}{x}.
$$
\n(25)

Usually the following new coordinates are introduced

$$
r = e^s = \sqrt{x^2 + y^2}
$$
,  $\varphi = t = \arctan \frac{y}{x}$ .

#### Example 23 Parabolic coordinates

$$
\frac{s+it}{\sqrt{2}} = \sqrt{x+iy},
$$
  

$$
s = \sqrt{r+x}, \quad t = \sqrt{r-x}.
$$

More frequently the parabolic coordinates are introduced as follows

$$
\xi = s^2, \quad \eta = t^2.
$$

#### Example 24 Elliptic coordinates

$$
s + it = \arcsin \frac{x + iy}{\alpha},
$$
  
\n
$$
\sin s = \frac{s_1 - s_2}{2\alpha}, \quad \cosh t = \frac{s_1 + s_2}{2\alpha}
$$
  
\nwhere  $s_1 = \sqrt{(x + \alpha)^2 + y^2}, s_2 = \sqrt{(x - \alpha)^2 + y^2}.$  The substitution

 $\xi = \sin s$ ,  $\eta = \cosh t$ 

is frequently used.

Example 25 Bipolar coordinates

$$
s + it = \ln \frac{\alpha + x + iy}{\alpha - x - iy},
$$
  

$$
\tanh s = \frac{2\alpha x}{\alpha^2 + x^2 + y^2}, \quad \tan t = \frac{2\alpha y}{\alpha^2 - x^2 - y^2}
$$

The following substitution is frequently used

$$
\xi = e^{-s}, \quad \eta = \pi - t.
$$

The last theorem opens the way for explicit construction of formal powers corresponding to the main Vekua equation  $(24)$  in the case when f has the form

<span id="page-11-0"></span>
$$
f = S(s)T(t) \tag{26}
$$

.

and hence for explicit construction of complete systems of solutions for corresponding secondorder elliptic equations admitting a particular solution of this form.

## <span id="page-11-3"></span>4 Description of the method

We consider boundary value problems of Dirichlet, Neumann or mixed type for the elliptic equation of the form [\(4\)](#page-3-2) in a bounded, simply connected domain  $\Omega \subset \mathbb{R}^2$ . The main assumption required for the applicability of the method of pseudoanalytic formal powers (MPFP) is the existence in  $\overline{\Omega}$  of a positive solution  $u_0$  such that the function  $f = p^{1/2}u_0 \in C^1(\overline{\Omega})$ be representable in a separable form [\(26\)](#page-11-0) in an orthogonal coordinate system. Let us stress that very often such a particular solution  $u_0$  is readily available. The simplest example of such situation is when  $q \equiv 0$  and p is of the form [\(26\)](#page-11-0). For example, the cases when  $p(x,y) = X(x)Y(y)$  or  $p = p(\sqrt{x^2 + y^2})$  frequently occur in practice [\[8\]](#page-19-8).

When the equation of the form

<span id="page-11-1"></span>
$$
(-\Delta + q(y))u(x, y) + \lambda^2 u(x, y) = 0 \tag{27}
$$

is considered, it is sufficient to obtain a particular solution for the ordinary differential equation

<span id="page-11-4"></span>
$$
(-\frac{d^2}{dy^2} + q(y))h(y) = 0.
$$
\n(28)

Then a particular solution of [\(27\)](#page-11-1) can be constructed as follows

<span id="page-11-2"></span>
$$
u_0(x, y) = e^{\lambda x} h(y).
$$
\n(29)

It has a convenient separable form. Notice that in this example we come to an important open problem. It is related to the requirement that  $u_0$  should be different from zero in the domain of interest. Meanwhile in many practically significant situations it is easy to guarantee that  $h(y) \neq 0$  when  $(x, y) \in \overline{\Omega}$ , sometimes this condition becomes a considerable obstacle. Moreover, when  $\lambda$  in [\(27\)](#page-11-1) is purely imaginary, the solution [\(29\)](#page-11-2) is not acceptable because it is no longer real valued. In this case one should take instead of  $e^{ikx}$ , where  $\lambda = ik$ , a solution in the form of  $\sin kx$  or  $\cos kx$  but then for a big domain or large k one cannot avoid the appearance of zeros of the resulting particular solution  $u_0$  and in this case the proposed scheme in general does not work.

One possibility to overcome this problem is to include under consideration a complex valued particular solution  $u_0$  but then we would need to consider a corresponding bicomplex main Vekua equation (see [\[13\]](#page-20-6) and [\[15\]](#page-20-0)). Then the whole algorithm for the construction of generating sequences and formal powers would go through with no modification compared to the complex case, but up to now there is no proof of the completeness of the system of formal powers for a bicomplex Vekua equation. As a consequence there is no guarantee that the infinite system of exact solutions obtained similar to [\(23\)](#page-8-1) will be complete in the space of solutions of [\(4\)](#page-3-2) in  $\Omega$ . Our conjecture is that at least in the case when  $u_0(x, y) = q(x)h(y)$ where  $q$  and  $h$  are complex valued nonvanishing functions the system of formal powers for the corresponding bicomplex main Vekua equation is complete in the same sense as was established earlier for the complex case. We continue this discussion in section [6](#page-18-0) where we use complex valued particular solutions of the form [\(29\)](#page-11-2) for solving eigenvalue problems for operators of the form  $-\Delta + q(y)$ .

Turning back to equation [\(4\)](#page-3-2) we assume that it admits a positive solution  $u_0$  in the domain  $\Omega$  such that  $f = p^{1/2} u_0 \in C^1(\overline{\Omega})$  is representable in a separable form [\(26\)](#page-11-0) and that  $\Phi = s+it$  is an analytic function of the variable  $z = x+iy$  in  $\Omega$  such that  $\Phi_z$  is bounded and has no zeros in  $\Omega$ . Then applying theorem [21](#page-9-1) one can construct a corresponding generating sequence. Construction of formal powers  $\left\{Z^{(n)}(1, z_0; z), Z^{(n)}(i, z_0; z)\right\}_{n=0}^{\infty}$  reduces then to the recursive algorithm described in Definition [11,](#page-6-1) and in this way one obtains the complete system of solutions for [\(4\)](#page-3-2) in  $\Omega$  given by [\(23\)](#page-8-1). By construction Re  $Z^{(0)}(i, z_0; z) \equiv 0$ . Taking this into account we introduce the notations

$$
u_1(z) = p^{-1/2}(z) \operatorname{Re} Z^{(1)}(1, z_0; z), \quad u_2(z) = p^{-1/2}(z) \operatorname{Re} Z^{(1)}(i, z_0; z),
$$
  

$$
u_3(z) = p^{-1/2}(z) \operatorname{Re} Z^{(2)}(1, z_0; z), \quad u_4(z) = p^{-1/2}(z) \operatorname{Re} Z^{(2)}(i, z_0; z), \dots
$$

and obtain the complete system of solutions for [\(4\)](#page-3-2) given by  $\{u_0, u_1, u_2, \ldots\}$ . We look for an approximate solution of a boundary value problem for [\(4\)](#page-3-2) in the form

<span id="page-12-0"></span>
$$
u^N = \sum_{k=0}^N b_k u_k \tag{30}
$$

where  $b_k$  are real coefficients which should be found from boundary conditions. To obtain  $N+1$  equations for finding  ${b_k}_{k=0}^N$  one can use, e.g., the collocation method. Chosing  $N+1$ points  $\zeta_i \in \partial \Omega$  we obtain  $N+1$  equations

$$
\sum_{k=0}^{N} b_k B[u_k](\zeta_j) = v(\zeta_j), \quad j = \overline{0, N}
$$

where  $v$  is a given function and  $B$  is the linear operator of the boundary condition. For the Dirichlet condition one has  $B[u] = u$  and for the Neumann condition,  $B[u] = \frac{\partial u}{\partial \vec{n}}$  – the normal derivative of u. Finding  ${b_k}_{k=0}^N$  we have the approximate solution  $u^N$ .

Thus, the proposed here MPFP belongs to the class of boundary methods because due to the linearity of the problem the function [\(30\)](#page-12-0) is an exact solution of [\(4\)](#page-3-2) in  $\Omega$  and only

boundary conditions should be approximated. An estimate for the rate of convergence of the method is given in theorem [16.](#page-7-4) In the next section we discuss the numerical realization of MPFP and results of numerical tests.

## 5 Approximate solution of boundary value problems

As a first example we considered the Dirichlet problem for the equation

<span id="page-13-0"></span>
$$
\left(-\Delta + c^2\right)u = 0\tag{31}
$$

where  $c$  is a real constant. The interest in this relatively simple equation is not due to its numerical simplicity. In fact this is not the case,- numerical solution of this equation is not less difficult than that of an equation with c being a reasonably good function with a range of values comparable with  $c$ . The attractiveness of this example consists in the possibility to calculate a large number of the functions  $u_k$  (see the preceding section) symbolically, using an appropriate software for symbolic calculations like Mathematica (Wolfram), Maple or Matlab. In this work we used Matlab 2006 and a PC of 2 GB in RAM and a processor of 1.73 GHz. Implementation of the symbolically calculated base functions  $u_k$  gives us the possibility to estimate the accuracy of the MPFP itself without considering the precision of recursive numerical integrations. We also compare the results obtained using symbolically calculated  $u_k$  with the results obtained purely numerically.

For equation [\(31\)](#page-13-0) it is easy to propose a positive particular solution. It can be chosen, e.g., as  $f = e^{cy}$ . Then the first functions  $u_k$  constructed as described in the preceding section taking as a center of the formal powers the origin will have the form [\[15\]](#page-20-0)

$$
u_0(x, y) = e^{cy}, \t u_1(x, y) = xe^{cy}, \t u_2(x, y) = -\frac{\sinh(cy)}{c},
$$
  

$$
u_3(x, y) = \left(x^2 - \frac{y}{c}\right)e^{cy} + \frac{\sinh(cy)}{c^2}, \t u_4(x, y) = -\frac{2x\sinh(cy)}{c}, \dots
$$
 (32)

It is interesting to mention that using Matlab we obtained the first 101 functions of this system calculated symbolically. According to theorem [19](#page-8-0) this system of solutions is complete in any bounded simply connected domain containing the origin. First we show results obtained with the help of the system of functions  $u_k$  calculated symbolically.

## 5.1 Numerical results obtained with symbolically calculated base functions

We begin with the unitary disk  $D$  with center in the origin. As a test exact solution we take the function

<span id="page-13-2"></span><span id="page-13-1"></span>
$$
u = e^{cx}.\tag{33}
$$

Thus, the problem we consider is to solve [\(31\)](#page-13-0) in D with the boundary condition  $u|_{\partial D} = e^{cx}$ . We look for an approximate solution  $u^N$  in the form [\(30\)](#page-12-0) with the base functions [\(32\)](#page-13-1). We use the collocation method for satisfying the boundary condition, the collocation points are distributed uniformly on  $\partial D$ . Their number is equal to the number of solutions  $u_k$ .

According to theory from subsection [3.2](#page-7-5) the coefficients  $b_k$  in [\(30\)](#page-12-0) are obtained in the case under consideration from the Taylor coefficients which appear in [\(22\)](#page-8-2). More precisely we have that according to theorem [18](#page-8-3) the solution  $u$  can be represented as follows

$$
u(z) = \sum_{n=0}^{\infty} \text{Re } Z^{(n)}(a_n, 0; z) = \sum_{n=0}^{\infty} \left( a'_n \text{ Re } Z^{(n)}(1, 0; z) + a''_n \text{ Re } Z^{(n)}(i, 0; z) \right)
$$

where  $a_n = a'_n + i a''_n$  are the Taylor coefficients given by [\(19\)](#page-7-0). In the case of the exact solution [\(33\)](#page-13-2) the Taylor coefficients have the form [\[15,](#page-20-0) Sect. 7.3]

$$
a_n = \frac{c^n}{n!} (1+i).
$$

Thus, the exact values for the coefficients  $b_k$  from [\(30\)](#page-12-0) in our example are as follows

$$
b_0 = 1
$$
,  $b_1 = b_2 = c$ ,  $b_3 = b_4 = \frac{c^2}{2}$ ,....

Having compared the numerically calculated constants  $b_k$  which we denote by  $b_k$  for  $N = 34$  with their exact values in the case  $c = 1$  we obtained their coincidence up to  $10^{-14}$ for every  $k = 0, \ldots, 34$ . For smaller values of c the situation is the same. The difference between  $b_k$  and  $b_k$  tends to become larger for larger values of c. In Table [5.1](#page-13-2) we show results for  $c = 5$  and  $N = 34$ .

Table 1. Comparison of the values of $b_k$ and $b_k$ as k increases				
$\boldsymbol{k}$	The values of $b_k$	The values of $b_k$	$ b_k-b_k $	
5	20.83333333333382	20.833333333333333	0.00000000000048	
8	26.04166666666448	26.04166666666667	0.00000000000219	
13	15.50099206509864	15.50099206349206	0.00000000160657	
17	5.38228885848900	5.38228891093474	0.00000005244574	
25	0.19601580023149	0.19603324996120	0.00001744972971	
31	0.00743227875004	0.00729290364439	0.00013937510565	
34	0.00172611010091	0.00214497166011	0.00041886155920	

The maximum number of functions  $u_k$  that we used here is limited not by the possibility of obtaining them symbolically but rather by the time required for numerical calculations involving the corresponding quite long symbolic expressions.

In the following two tables, the convergence of MPFP is shown by comparison of the maximum absolute error obtained for different values of N, for the case  $c = 1$  and  $c = 5$ .

Tables 2 and 3 Maximum absolute error depending on N for  $c = 1$  and  $c = 5$ 

N	Maximum absolute error	N	Maximum absolute error
8	0.00698626935341	6	$3.59578971016677 \times 10^{2}$
14	$2.534633673767495\times10^{-5}$	14	22.38029523897584
22	$1.432881036045330 \times 10^{-9}$	22	0.73431266884919
28	$4.276579090856103 \times 10^{-13}$	32	0.00194275813006
32	$1.776356839400251 \times 10^{-15}$	44	$0.59167057031573\times10^{-7}$
36	$8.881784197001252\times 10^{-16}$	54	$0.72795103278622\times 10^{-11}$
38	$1.110223024625157 \times 10^{-15}$	60	$8.7818\overline{64124784988\times 10^{-14}}$

In Table 4 for a fixed number  $(N = 34)$  of the base functions  $u_k$  we show the dependence of the maximum absolute error on the parameter  $c$ . Here we also indicate the maximum absolute error obtained for the same problem using the standard PDE tool of Matlab.

absolute error for increasing values of c as $\alpha = 34$			
$\mathcal{C}$	Maximum absolute error of MPFP   PDE tool (2129 nodes)		
0.1	$0.89 \times 10^{-15}$	$1.5 \times 10^{-6}$	
0.5	$0.26 \times 10^{-14}$	$4.5 \times 10^{-6}$	
	$0.12 \times 10^{-14}$	$1.6 \times 10^{-4}$	
$\mathcal{D}_{\mathcal{L}}$	$0.14 \times \overline{10^{-10}}$	$1.4 \times 10^{-3}$	
5	$0.29 \times 10^{-3}$	$3.0 \times 10^{-2}$	
10	$4.06 \times \overline{10^2}$	8.0	

Table 4. Performance of MPFP compared to Matlab's PDE tool in terms of the maximum absolute error for increasing values of  $c$  as  $N = 34$ 

As it can be observed in the last table the result of application of MPFP in the case of  $c = 10$  is less satisfactory as that of PDE tool. This is due to the fact that for larger values of c one should consider a bigger  $N$ . In Table 5 we show the absolute error of MPFP for  $c = 10$  and  $N \geq 42$ .

Table 5. Improvement in the maximum absolute error due to MPFP as the number of functions  $u_k$  keeps increasing

	rancelons $u_k$ access increasing
N	Maximum absolute error of MPFP
42	3.89
44	3.25
46	1.81
48	0.81
50	0.41
52	0.10

Thus, one can see that for  $N \geq 42$  the result obtained with the aid of MPFP is more accurate than that given by Matlab. We stress that in the case of using MPFP a system of  $N+1$  linear algebraic equations is solved which means solution of dozens of equations instead of thousands required by the finite element method implemented in the PDE tool.

We experimented also with the shape of the domain. We considered the elliptic form as well as a unitary disk with a triangle shaped deformation. In the first case it is possible to see how the maximum absolute error increases with the excentricity e, Table 6. Here in all cases the area of the considered ellipses was kept constant, equal to  $\pi$ , while the excentricity was being increased.

Table 6. Maximum absolute error for different values of the excentricity of the elliptic domain with the area of the domain being equal to  $\pi$ . The case  $e = 0$  corresponds to the unitary disc.

allicaty aloc.					
	$\boxed{30}$ $\boxed{2.2 \times 10^{-14}$ $\boxed{0.4 \times 10^{-13}}$ $\boxed{0.5 \times 10^{-13}}$ $\boxed{0.3 \times 10^{-12}}$ $\boxed{0.2 \times 10^{-11}}$ $\boxed{1 \times 10^{-10}}$				

For the case of the domain with a triangular deformation (see Fig. 1), the errors were



Figure 1: The unitary disk with a triangular deformation.

tested for different heights of the peak and different values of  $c$  and  $N$  with satisfactory results. In Table 7 we present the maximum absolute error of the approximate solution of the boundary value problem in dependence on the height of the triangular peak over the unitary circunference.

Height of the peak over the unitary disk   Maximum absolute error	
0.5	$0.92 \times 10^{-12}$
	$0.62 \times 10^{-11}$
	$0.72 \times 10^{-10}$

Table 7. Maximum absolute error for  $N = 31$ ,  $c = 1$  and different heights of the peak

#### 5.2 Results obtained with numerically calculated base functions

The use of the numerically calculated base functions which we denote by  $\tilde{u}_k$  poses the natural question about the accuracy of their calculation. Consideration of equation [\(31\)](#page-13-0) gives us the possibility to compare  $\tilde{u}_k$  with the symbolically calculated exact solutions [\(32\)](#page-13-1). In the following table we give the difference between  $u_k$  and  $\tilde{u}_k$  for  $c = 1$ .

Table 8. Maximum absolute error of calculation of the base functions for  $c = 1$ 

$k_1$	$ u_k - \widetilde{u}_k $
1	$\overline{0.00000667646050} \times 10^{-4}$
.5	$0.00022726096338 \times 10^{-4}$
11	$0.01492959925020 \times 10^{-4}$
16	$\sqrt{0.07644765777970 \times 10^{-4}}$
20	$0.22545767650040\times10^{-4}$

Regarding the results given in the last table it is important to note that in fact the weight of  $u_k$  in the expansion of a solution decreases as  $\left(\frac{k+1}{2}\right)$  $\frac{+1}{2}$ )! or  $\left(\frac{k}{2}\right)$  $(\frac{k}{2})!$  for odd or even k respectively. This is due to the factor  $1/n!$  in the definition of the Taylor coefficients [\(19\)](#page-7-0). That is in fact the real accuracy of calculation of the base functions would be given by  $|u_k - \tilde{u}_k|/K!$  where  $K = \begin{cases} \frac{k+1}{2} \end{cases}$  $\frac{+1}{2}$ )! for k odd  $\frac{k}{2}$  $\frac{k}{2}$ )! for k even  $\frac{k}{2}$ . It is easy to see that in the case of the results given in Table 7 one then obtains  $|u_{20} - \tilde{u}_{20}|/10! \simeq 6.213 \times 10^{-12}$  which is a remarkably good agreement.

For the numerical computation of  $\tilde{u}_k$  we implemented the following procedure. Before integrating on each new step according to [\(16\)](#page-6-2) along segments joining the center of the formal powers with points on the boundary of the domain the integrand was represented as a cubic spline which then was integrated using the standard Matlab routine for integration of splines. This procedure is simple but clearly not optimal. Nevertheless the approximate results presented in this work show that even such integration procedure gives satisfactory agreement between the exact base functions and those calculated numerically.

The accuracy of the approximate solution obtained with the aid of  $\tilde{u}_k$  in our numerical tests did not differ significantly from that of the solution obtained using the exactly calculated  $u_k$ . The order of the maximum absolute error for a given N and c coincided in both cases. Hence here we present results corresponding to another test problem for which we did not have the exactly calculated base functions.

Consider the equation

<span id="page-17-0"></span>
$$
\left(-\Delta + \frac{e^y}{4}\right)u(x, y) = 0.
$$
\n(34)

An exact solution for this equation can be found using the fact that the change of variables  $\xi = e^{\frac{y}{2}} \cos \frac{x}{2}$ ,  $\eta = e^{\frac{y}{2}} \sin \frac{x}{2}$  leads to [\(31\)](#page-13-0) in the new variables. Thus, e.g., the function  $u(x,y) = \exp(e^{\frac{y}{2}} \cos \frac{x}{2})$  is an exact solution of [\(34\)](#page-17-0). Consequently, as a test problem we can consider the problem of finding a solution of  $(34)$  in  $\Omega$ , satisfying the boundary condition

<span id="page-17-1"></span>
$$
u(x,y) = \exp(e^{\frac{y}{2}} \cos \frac{x}{2}), \quad (x,y) \in \partial \Omega.
$$
 (35)

In order to construct a particular solution  $u_0$  in a separable form we solve numerically the ordinary differential equation

$$
\left(-\frac{d^2}{dy^2} + \frac{e^y}{4}\right)u_0(y) = 0.
$$

The obtained solution we then use for constructing the system of functions  $\{u_k\}$ . Some results on the accuracy of the approximate solution are given in the following table.

Table 9. Maximum absolute error of the approximate solution of the test problem [\(34\)](#page-17-0), [\(35\)](#page-17-1) considered in a unitary disk in dependence on N



## <span id="page-18-0"></span>6 Approximate solution of eigenvalue problems

In this section we consider the application of MPFP to solution of eigenvalue problems for operators of the form  $-\Delta+q(y)$ . For simplicity we keep working with the Dirichlet boundary conditions and suppose that q is continuous and  $q(y) \geq 0$  in  $\overline{\Omega}$ . Then the spectrum of the operator is discrete and positive. As was explained in section [4](#page-11-3) for the equation

<span id="page-18-1"></span>
$$
(-\Delta + q(y))u_0(x, y) = \lambda^2 u_0(x, y)
$$
\n(36)

it is easy to propose a particular solution in a separable form for any value of  $\lambda$ . We are interested here in positive values, and hence a natural choice of a nonvanishing solution would be  $u_0(x, y) = e^{i\lambda x}h(y)$  where  $h(y)$  is a positive solution of [\(28\)](#page-11-4). As was observed in section [4](#page-11-3) the completeness of the system of solutions  ${u_k}_{k=0}^{\infty}$  obtained in this case is up to now an open problem due to the fact that  $u_0$  is complex valued and one should consider bicomplex pseudoanalytic formal powers for which the whole theory is still underdeveloped. Nevertheless we used the constructed system of exact solutions  ${u_k}_{k=0}^{\infty}$  for finding the eigenvalues  $\lambda^2$  in the following way. Assuming that  $\{u_k\}_{k=0}^{\infty}$  is complete in the same sense as was proved in the case of the real-valued particular solution  $u_0$  (subsection [3.2\)](#page-7-5) we have then that if a nontrivial solution u of [\(36\)](#page-18-1) exists satisfying the boundary condition  $u|_{\partial\Omega} = 0$  then  $u \simeq \sum_{k=0}^{N} b_k u_k$  and the coefficients  $b_k$  are such that the trivial boundary condition is approximately fulfilled. This means that one can require that  $\sum_{k=0}^{N} b_k u_k(z_j) = 0$  for  $z_j \in \partial \Omega$  and  $j = \overline{0, N}$ . This is possible iff the determinant of the matrix  $U = (u_{jk})_{j,k=0}^N$  vanishes where  $u_{ik} = u_k(z_i)$ . The determinant of U for a fixed N is a function of  $\lambda$ . Thus, the problem of finding eigenvalues reduces to the problem of finding zeros of the function det  $U(\lambda)$ .

As a test problem we considered the problem of calculating the eigenvalues of the Dirichlet problem for the Helmholtz equation  $(\Delta + \lambda^2) u = 0$ . For every  $\lambda$  a system of exact solutions  ${u_k}_{k=0}^{\infty}$  can be constructed using [\(32\)](#page-13-1) where c should be replaced by i $\lambda$ . Then following the described scheme we looked for zeros of det  $U(\lambda)$ . As it is well known (see, e.g., [\[6\]](#page-19-9)) the eigenvalues of the Dirichlet problem for the Helmholtz equation in a unitary disk are squares of zeros of Bessel functions  $J_n(x)$ . Our numerical experiments showed that a relatively small value of  $N = 21$  was needed for computing the first five eigenvalues with the accuracy of four decimals. With  $N = 23$  we obtained six first eigenvalues with the same accuracy. Thus,

indeed, the MPFP is clearly competitive in solving eigenvalue problems for elliptic operators. In applying the MPFP we detected a similar problem to that described by Alexidze in [\[2,](#page-19-5) Sect. 1.13] where the method of fundamental solutions (or auxiliary sources) was applied to eigenvalue problems. The considered determinant shows a very fast decrement (in spite of this the method gives good numerical results). [\[2\]](#page-19-5) contains references to other publications where different ways of using the knowledge of a system of exact solutions for numerical solution of eigenvalue problems were studied. In this direction further research is needed.

## 7 Conclusions

A new approach for solving boundary value and eigenvalue problems for elliptic operators in bounded planar domains is proposed. It is based on some classical and some new results from pseudoanalytic function theory which allow one to construct complete systems of solutions of the elliptic equations. We showed the practical applicability of the numerical method based on this construction, studied the rate of its convergence, accuracy and other parameters of its performance.

## References

- <span id="page-19-6"></span>[1] Agmon S and Bers L 1952 The expansion theorem for pseudo-analytic functions. Proc. Amer. Math. Soc. 3 757-764.
- <span id="page-19-5"></span>[2] Alexidze M A 1991 Fundamental Functions in Approximate Solutions of Boundary Value Problems (in Russian). Moscow: Nauka.
- <span id="page-19-0"></span>[3] Bers L 1952 Theory of pseudo-analytic functions. New York University.
- <span id="page-19-2"></span>[4] Bers L 1956 Formal powers and power series. Communications on Pure and Applied Mathematics 9, 693-711.
- <span id="page-19-3"></span>[5] Cannon J R 1964 The numerical solution of the Dirichlet problem for Laplace's equation by linear programming. J. Soc. Indust. Appl. Math. 12, 233–237.
- <span id="page-19-9"></span>[6] Coleman M P 2005 An Introduction to Partial Differential Equations with MATLAB. Boca Raton: Chapman & Hall/CRC.
- <span id="page-19-1"></span>[7] Colton D L 1976 Solution of Boundary Value Problems by the Method of Integral Operators. London: Pitman Publ.
- <span id="page-19-8"></span>[8] Demidenko Eu. 2006 Separable Laplace equation, magic Toeplitz matrix, and generalized Ohm's law. Applied Mathematics and Computation 181, 1313-1327.
- <span id="page-19-7"></span>[9] Dettman J W 1984 Applied complex variables. NY: Dover Publications, Inc.
- <span id="page-19-4"></span>[10] Genev V N 1984 Solution of the Neumann problem by the method of harmonic polynomials in a simply connected domain bounded by a smooth contour. (Russian) Godishnik Vissh. Uchebn. Zaved. Prilozhna Mat. 19, no. 2, 95–106.
- <span id="page-20-4"></span>[11] Hozejowski L., Hozejowska S and Piasecka M 2003 Application of harmonic polynomials as complete solutions of Laplace equation in an inverse heat conduction problem. PAMM Proc. Appl. Math. Mech. 2, 362–363.
- <span id="page-20-2"></span>[12] Kravchenko V V 2005 On a relation of pseudoanalytic function theory to the twodimensional stationary Schrödinger equation and Taylor series in formal powers for its solutions. J. of Phys. A , 38, No. 18, 3947-3964.
- <span id="page-20-6"></span>[13] Kravchenko V V 2006 On a factorization of second order elliptic operators and applications. Journal of Physics A: Mathematical and General, 39, No. 40, 12407-12425.
- <span id="page-20-3"></span>[14] Kravchenko V V 2008 Recent developments in applied pseudoanalytic function theory. In "Some topics on value distribution and differentiability in complex and p-adic analysis", eds. A. Escassut, W. Tutschke and C. C. Yang, Science Press 293-328.
- <span id="page-20-0"></span>[15] Kravchenko V V 2009 Applied Pseudoanalytic Function Theory. Basel: Birkhäuser, Series: Frontiers in Mathematics.
- <span id="page-20-8"></span>[16] Madelung E 1957 Die Mathematischen Hilfsmittel des Physikers. Berlin: Springer-Verlag.
- <span id="page-20-7"></span>[17] Menke K 1974 Zur Approximation pseudoanalytischer Funktionen durch Pseudopolynome. Manuscripta Math. 11 111-125.
- <span id="page-20-5"></span>[18] Suetin P K 1988 Orthogonal polynomials in two variables (in Russian). Moscow: Nauka.
- <span id="page-20-1"></span>[19] Vekua I N 1959 Generalized analytic functions. Moscow: Nauka (in Russian); English translation Oxford: Pergamon Press 1962.