

Banach spaces without approximation properties of type p

Oleg Reinov* and Qaisar Latif

ABSTRACT. The main purpose of this note is to show that the question posed in the paper of Sinha D.P. and Karn A.K. ("Compact operators which factor through subspaces of l_p Math. Nachr. 281, 2008, 412-423; see the very end of that paper) has a negative answer, and that the answer could be obtained, essentially, in 1985 after the papers [2], [3] by Reinov O.I. have been appeared in 1982 and in 1985 respectively.

The main purpose of this note is to show that the question posed in [1] (see the very end of that paper) has a negative answer, and that the answer could be obtained, essentially, in 1985 after the papers [2], [3] have been appeared in 1982 and in 1985 respectively.

For the sake of completeness (and since the article [3] is now quite difficult to get) we will reconstruct some of the facts from [3], maybe, in a somewhat changed form (a translation of [3] with some remarks can be found in arXiv:1002.3902v1 [math.FA]).

All the notation and terminology, we use, are from [2], [4], or [5] and more or less standard. For the general theory of absolutely p -summing, p -nuclear and other operator ideals, we refer to [6]. We will keep, in particular, the following standard notations. If A is a bounded subset of a Banach space X , then $\overline{\Gamma(A)}$ is the closed absolutely convex hull of A ; X_A is the Banach space with "the unit ball" $\overline{\Gamma(A)}$; $\Phi_A : X_A \rightarrow X$ is the canonical embedding. For $B \subset X$, it is denoted by \overline{B}^τ and $\overline{B}^{\|\cdot\|}$ the closures of the set B in the topology τ and in the norm $\|\cdot\|$ respectively. When it is necessary, we denote by $\|\cdot\|_X$ the norm in X . Other notations: Π_p , QN_p , N_p , I_p are the ideals of absolutely p -summing, quasi- p -nuclear, p -nuclear, strictly p -integral operators, respectively; $X^* \widehat{\otimes}_p Y$ is the complete tensor product associated with $\text{N}_p(X, Y)$; $X^* \widehat{\otimes}_p Y = \overline{X^* \otimes Y}^{\pi_p}$ (i.e. the closure of the set of finite dimensional operators in $\Pi_p(X, Y)$). Finally, if $p \in [1, +\infty]$ then p' is the adjoint exponent.

*The research was supported by the Higher Education Commission of Pakistan.

AMS Subject Classification 2000: 46B28. Spaces of operators; tensor products; approximation properties .

Key words: p -nuclear operators, approximation properties, compact approximation, nuclear tensor norms, tensor products.

Now, we recall the main definition from [3] of the topology τ_p of the π_p -compact convergence, which is just the same as the λ_p -topology in [1]¹, where it was proved, essentially, that that $T \in K_p^d(X, Y)$ if and only if $T \in QN_p(X, Y)$.

For Banach spaces X, Y , the topology τ_p of π_p -compact convergence in the space $\Pi_p(Y, X)$ is the topology, a local base (in zero) of which is defined by sets of type

$$\omega_{K, \varepsilon} = \{U \in \Pi_p(Y, X) : \pi_p(U\Phi_K) < \varepsilon\},$$

where $\varepsilon > 0$, $K = \overline{\Gamma(K)}$ — a compact subset of Y .

Proposition 1 [3]. *Let R be a linear subspace in $\Pi_p(Y, X)$, containing $Y^* \otimes X$. Then $(R, \tau_p)'$ is isomorphic to a factor space of the space $X^* \widehat{\otimes}_{p'} Y$. More precisely, if $\varphi \in (R, \tau_p)'$, then there exists an element $z = \sum_1^\infty x'_n \otimes y_n \in X^* \widehat{\otimes}_{p'} Y$ such that*

$$\varphi(U) = \text{trace } U \circ z, \quad U \in R. \quad (*)$$

On the other hand, for every $z \in X^* \widehat{\otimes}_{p'} Y$ the relation $(*)$ defines a linear continuous functional on (R, τ_p) .

Proof. Let φ be a linear continuous functional on (R, τ_p) . Then one can find a neighborhood of zero $\omega_K = \omega_{K, \varepsilon}$, such that φ is bounded on it: $\forall U \in \omega_K, |\varphi(U)| \leq 1$. We may assume that $\varepsilon = 1$. Consider the operator $U\Phi_K : Y_K \xrightarrow{\Phi_K} Y \xrightarrow{U} X$. Since the mapping Φ_K is compact, $U\Phi_K \in QN_p(Y_K, X)$. Put $\varphi_K(U\Phi_K) = \varphi(U)$ for $U \in R$. On the linear subspace $R_K = \{V \in QN_p(Y_K, X) : V = U\Phi_K\}$ of the space $QN_p(Y_K, X)$, the linear functional φ_K is bounded: if $V = U\Phi_K \in R_K$ and $\pi_p(V) \leq 1$, then $|\varphi_K(V)| = |\varphi(U)| \leq 1$. Therefore, φ_K can be extended to a linear continuous functional $\tilde{\varphi}$ on the whole $QN_p(Y_K, X)$; moreover, because of the injectivity of the ideal QN_p , considering X as a subspace of some space $C(K_0)$, we may assume that $\tilde{\varphi} \in QN_p(Y_K, C(K_0))^*$. Let us mention that

$$\tilde{\varphi}(jU\Phi_K) = \varphi_K(U\Phi_K) = \varphi(U) \quad (1)$$

(here j is an isometric embedding of X into $C(K_0)$).

Furthermore, since $QN_p(Y_K, C(K_0))^* = I_{p'}(C(K_0), (Y_K)^{**})$, we can find an operator $\Psi : C(K_0) \rightarrow (Y_K)^{**}$, for which

$$\tilde{\varphi}(A) = \text{trace } \Psi A, \quad A \in (Y_K)^* \otimes C(K_0).$$

Let $A_n \in (Y_K)^* \otimes C(K_0)$, $\pi_p(A_n - jU\Phi_K) \rightarrow 0$. Then

$$\tilde{\varphi}(jU\Phi_K) = \lim \text{trace } \Psi A_n. \quad (2)$$

Consider the operator $\Phi_K^{**}\Psi : C(K_0) \xrightarrow{\Psi} (Y_K)^{**} \xrightarrow{\Phi_K^{**}} Y$. Since $\Psi \in I_{p'}$, and Φ_K is compact, we have $\Phi_K^{**}\Psi \in N_{p'}(C(K_0), Y) = C(K_0)^* \widehat{\otimes}_{p'} Y$. Let $\sum_1^\infty \mu_n \otimes y_n \in C(K_0)^* \widehat{\otimes}_{p'} Y$ be a representation of the operator $\Phi_K^{**}\Psi$. Put $z = \sum j^*(\mu_n) \otimes y_n$. The element z generates an operator $\Phi_K^{**}\Psi j$ from X to Y . We will show now that

¹ Definition 4.3 from [1]: "Given a compact subset $K \subset X$ we define a seminorm $\|\cdot\|_K$ on $\Pi_p(X, Y)$ given by $\|T\|_K = \inf\{\kappa_p^d(Ti_Z) : i_Z : Z \rightarrow X \text{ as above}\}$. The family of seminorms $\{\|\cdot\|_K : K \subset X \text{ is compact}\}$ determines a locally convex topology λ_p on $\Pi_p(X, Y)$."; — for details, and for the definition of maps i_Z , see [1], Lemma 4.2.

trace $U \circ z = \tilde{\varphi}(jU\Phi_K)$ (note that $U \circ z$ is an element of the space $X^* \widehat{\otimes} X$, so the trace is well defined). We have:

$$\begin{aligned} \text{trace } U \circ z &= \text{trace} \left(\sum j^*(\mu_n) \otimes Uy_n \right) = \sum \langle j^*(\mu_n), Uy_n \rangle = \\ &= \sum \langle \mu_n, jUy_n \rangle = \text{trace } jU\Phi_K^{**}\Psi = \text{trace} (jU\Phi_K)^{**}\Psi, \end{aligned} \quad (3)$$

where $(jU\Phi_K)^{**}\Psi : C(K_0) \xrightarrow{\Psi} (Y_K)^{**} \xrightarrow{\Phi_K^{**}} Y \xrightarrow{U} X \xrightarrow{j} C(K_0)$. Since $\pi_p(A_n - jU\Phi_K) \rightarrow 0$, then $\pi_p(A_n^{**} - (jU\Phi_K)^{**}) \rightarrow 0$. Moreover, if $A := A_n = \sum_1^N w_m \otimes f_m \in (Y_K)^* \otimes C(K_0)$, then

$$\text{trace } A_n^{**}\Psi = \sum_m \langle \Psi^* w_m, f_m \rangle = \sum_m \langle w_m, \Psi f_m \rangle = \text{trace } \Psi A.$$

Hence, $\text{trace} (jU\Phi_K)^{**}\Psi = \lim \text{trace } A_n^{**}\Psi = \lim \text{trace } \Psi A_n$. Now, it follows from (3) and (2) that $\tilde{\varphi}(jU\Phi_K) = \text{trace } U \circ z$. Finally, we get from (1): $\varphi(U) = \text{trace } U \circ z$. Thus, the functional φ is defined by an element of $X^* \widehat{\otimes}_{p'} Y$.

Inversely, if $z \in X^* \widehat{\otimes}_{p'} Y$, put $\varphi(U) = \text{trace } U \circ z$ for $U \in \mathbb{R}$ (the trace is defined since $U \circ z \in X^* \widehat{\otimes} X$). We have to show that the linear functional φ is bounded on a neighborhood $\omega_{K,\varepsilon}$ of zero in τ_p . For this, we need the following fact, which proof is rather standard (see [3], Lemma 1.2): *If $z \in X^* \widehat{\otimes}_q Y$, then $z \in X^* \widehat{\otimes}_q Y_K$, where $K = \overline{\Gamma(K)}$ is a compact in Y .* Now, let K be a compact subset of Y , for which $z \in X^* \widehat{\otimes}_{p'} Y_K$. If $U \in \omega_{K,1}$, then $\pi_p(U\Phi_K) < 1$ and $|\text{trace } U\Phi_K \circ z| \leq \|z\|_{X^* \widehat{\otimes}_{p'} Y_K} \cdot \pi_p(U\Phi_K) \leq C$.

Corollary 1. $(\mathbb{R}, \tau_p)' = (\mathbb{R}, \sigma)'$, where $\sigma = \sigma(\mathbb{R}, X^* \widehat{\otimes}_{p'} Y)$. Thus, the closures of convex subsets of the space $\Pi_p(Y, X)$ in τ_p and in σ are the same.

Proposition 2 [3]. *If the canonical mapping $j : X^* \widehat{\otimes}_{p'} Y \rightarrow N_{p'}(X, Y)$ is one-to-one then $\Pi_p(Y, X) = \overline{Y^* \otimes X}^{\tau_p}$.*

Proof If the map j is one-to-one then the annihilator $j^{-1}(0)^\perp$ of its kernel in the space, dual to $X^* \widehat{\otimes}_{p'} Y$, coincides with $\Pi_p(Y, X^{**})$. On the other hand, in any case $j^{-1}(0)^\perp = \overline{Y^* \otimes X}^*$ (the closure in *-weak topology of the space $\Pi_p(Y, X^{**})$); by Corollary from Proposition 1,

$$\Pi_p(Y, X) \cap \overline{Y^* \otimes X}^* = \overline{Y^* \otimes X}^{\tau_p}.$$

Therefore, $\Pi_p(Y, X) = \overline{Y^* \otimes X}^{\tau_p}$.

For a reflexive space X , the dual space to $X^* \widehat{\otimes}_{p'} Y$ is equal to $\Pi_p(Y, X)$. Consequently, it follows from the last two statements

Corollary 2. *For a reflexive space X the canonical mapping $j : X^* \widehat{\otimes}_{p'} Y \rightarrow N_{p'}(X, Y)$ is one-to-one iff the set of finite rank operators is dense in the space $\Pi_p(Y, X)$ in the topology τ_p of π_p -compact convergence.*

Recall a definition of the *approximation property* AP_p of order $p, p \in (0, \infty]$ (see, e.g., [2] and [7]): a Banach space X has the AP_p if for every Banach space Y (equivalently, for every reflexive Banach space Y , see [2] or [5]) one has the equality $Y^* \widehat{\otimes}_p X = N_p(Y, X)$. It follows now from Proposition 1 and Corollary 2:

Corollary 3. For $p \in [1, \infty]$ and for every Banach space X the following are equivalent:

- 1) X has the AP_p ;
- 2) for each Banach space Y $\overline{X^* \otimes_{p'} Y}^{\tau_{p'}} = \Pi_{p'}(X, Y)$;
- 3) for each reflexive Banach space Y $\overline{X^* \otimes_{p'} Y}^{\tau_{p'}} = \Pi_{p'}(X, Y)$;
- 3') for each reflexive Banach space Y $\overline{X^* \otimes_{p'} Y}^{\lambda_{p'}} = \Pi_{p'}(X, Y)$.

As shown in [2], [4] and [5], for each $p, p \in [1, \infty], p \neq 2$, there exist reflexive Banach spaces without the AP_p . Thus, we get from the last consequence

Corollary 4. For $1 \leq p \leq \infty, p \neq 2$, there are two reflexive Banach spaces X and Y such that the natural map $Y^* \widehat{\otimes}_p X \rightarrow N_p(Y, X)$ is not one-to-one and $\overline{X^* \otimes_{p'} Y}^{\tau_{p'}} \neq \Pi_{p'}(X, Y)$.

Since, as we said, τ_q equals to the topology λ_q from [1], we get firstly Theorem 4.11 [1] (the case $p > 2$ below) and secondly the negative answer to the question in the end of the paper [1] (whether for $p \in [1, 2)$ every Banach space has the approximation property of type p of [1]²). Or, generally, we have:

Theorem. Let $1 \leq p \neq 2 \leq \infty$. Then there is a (reflexive) Banach space that fails to have the approximation property of type p of [1].

References

- [1] Sinha D.P., Karn A.K.: *Compact operators which factor through subspaces of l_p* , Math. Nachr. 281(2008), 412-423.
- [2] O.I. Reinov: *Approximation properties of order p and the existence of non- p -nuclear operators with p -nuclear second adjoints*, Math. Nachr. 109(1982), 125-134.
- [3] O.I. Reinov: *Approximation of operators in Banach spaces*, Application of functional analysis in the approximation theory (KGU, Kalinin) (1985), 128-142.
- [4] O.I. Reinov: *Disappearance of tensor elements in the scale of p -nuclear operators*, Theory of operators and theory of functions (LGU) 1(1983), 145-165.
- [5] O. Reinov: *Approximation properties of order p and the existence of non- p -nuclear operators with p -nuclear second adjoints*, Doklady AN SSSR, 256(1981), 43-47.
- [6] A. Pietsch: *Operator ideals*, North-Holland, Deutscher Verlag der Wiss., Berlin, 1978.
- [7] P. Saphar: *Produits tensoriels d'espaces de Banach et classes d'applications lineaires*, Studia Math. 38(1970), 71-100.

OLEG REINOV:

DEPARTMENT OF MATHEMATICS AND MECHANICS, ST. PETERSBURG STATE UNIVERSITY, SAINT PETERSBURG, RUSSIA.

ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, 68-B, NEW MUSLIM TOWN, LAHORE 54600, PAKISTAN.

E-mail address: orein51@mail.ru

QAISAR LATIF:

ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, 68-B, NEW MUSLIM TOWN, LAHORE 54600, PAKISTAN.

² By [1], a Banach space X is said to have the *approximation property of type p* if for every Banach space Y , the finite rank operators $\mathcal{F}(Y, X)$ is dense in $\Pi_p(Y, X)$ in the λ_p -topology.