

# Congruences for an arithmetic function from 3-colored Frobenius partitions

Laizhong Song and Xinhua Xiong

Department of Mathematics  
China Three Gorges University, YiChang, Hubei 443002, P.R. China

Slz@ctgu.edu.cn and XinhuaXiong@fudan.edu.cn

## Abstract

Let  $a(n)$  defined by

$$\sum_{n=1}^{\infty} a(n)q^n := \prod_{n=1}^{\infty} \frac{1}{(1-q^{3n})(1-q^n)^3}.$$

In this note, we prove that for every nonnegative integer  $n$ ,

$$\begin{aligned} a(15n+6) &\equiv 0 \pmod{5}, \\ a(15n+12) &\equiv 0 \pmod{5}. \end{aligned}$$

As a corollary, we obtained some results of Ono [9].

**Keywords:** Congruence, modular form.

**AMS Classification:** 11F33, 11P83

## 1 Introduction

Arithmetic functions defined by infinite products have many beautiful congruence properties. A typical example is Euler's partition function  $p(n)$ , which is given by

$$\sum_{n=0}^{\infty} p(n)q^n = \prod_{n=1}^{\infty} \frac{1}{(1-q^n)}$$

In this note, Let  $a(n)$  defined by

$$\sum_{n=1}^{\infty} a(n)q^n := \prod_{n=1}^{\infty} \frac{1}{(1-q^{3n})(1-q^n)^3}.$$

we prove that

**Theorem 1.1.** *for every nonnegative integer  $n$ ,*

$$\begin{aligned} a(15n+6) &\equiv 0 \pmod{5}, \\ a(15n+12) &\equiv 0 \pmod{5}. \end{aligned}$$

As a corollary of our results, we obtain the following result which is due to Ono [9],

**Theorem 1.2.**

$$\begin{aligned} c\phi_3(45n + 7) &\equiv 0 \pmod{5}, \\ c\phi_3(45n + 22) &\equiv 0 \pmod{5}, \\ c\phi_3(45n + 37) &\equiv 0 \pmod{5}. \end{aligned}$$

Where  $c\phi_3(n)$  is the number of three-colored Frobenius partitions of  $n$ .

## 2 Preliminary Facts

This section contains the basic definitions and some properties of modular forms that we will use later. For more details see Koblitz [6].

Define  $\Gamma = Sl_2(\mathbb{Z})$ ,  $\Gamma_0(N) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid ad - bc = 1, c \equiv 0 \pmod{N} \right\}$ . Let  $M_k(\Gamma_0(N))$  denote the vector space of holomorphic modular forms on  $\Gamma_0(N)$  of weight  $k$ .

If  $f(z)$  in  $M_k(\Gamma_0(N))$  has the Fourier expansion  $f(z) = \sum_{n=0}^{\infty} a(n)q^n$ , for an integer  $t \mid N$ , define the U-operator by

$$f(z) | U(t) := \sum_{n=0}^{\infty} a(tn)q^n$$

It is a standard fact that  $f(z) | U(t) \in M_k(\Gamma_0(N))$ , if  $f(z) \in M_k(\Gamma_0(N))$ . Before starting the proof, we need some lemmas. The Dedekind Eta-function is defined by the infinite product

$$\eta(z) := q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n),$$

where  $q = e^{2\pi iz}$  and  $\text{Im}(z) > 0$ . A function  $f(z)$  is called an Eta-product if it can be written in the form of

$$f(z) = \prod_{\delta \mid N} \eta^{r_\delta}(\delta z)$$

where  $N$  and  $\delta$  are positive integers and  $r_\delta \in \mathbb{Z}$ .

**Lemma 2.1** ([4]). *If  $f(z) = \prod_{\delta \mid N} \eta^{r_\delta}(\delta z)$  is an Eta-product satisfying the following conditions:*

$$k = \frac{1}{2} \sum_{\delta \mid N} r_\delta \in 2\mathbb{Z}, \quad \sum_{\delta \mid N} \delta r_\delta \equiv 0 \pmod{24}, \quad \sum_{\delta \mid N} \frac{N}{\delta} r_\delta \equiv 0 \pmod{24},$$

and  $\prod_{\delta \mid N} \delta^{r_\delta} \in \mathbb{Q}^2$ . then  $f(z)$  satisfies

$$f\left(\frac{az + b}{cz + d}\right) = (cz + d)^k f(z)$$

for each  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$ .

**Lemma 2.2** ([8]). Let  $c, d$  and  $N$  be positive integers with  $d|N$  and  $(c, d) = 1$ . If  $f(z)$  is an Eta-product satisfying the conditions in Lemma 2.1 for  $N$ , then the order of vanishing of  $f(z)$  at the cusp  $\frac{c}{d}$  is

$$\frac{N}{24} \sum_{\delta|N} \frac{(d, \delta)^2 r_\delta}{(d, \frac{N}{d}) d \delta}.$$

**Lemma 2.3** ([10]). Let  $f(z) = \sum_{n=0}^{\infty} a(n)q^n$  be a modular form in  $M_k(\Gamma_0(N))$  and let  $d := \gcd(r, t)$ . If  $0 \leq r < t$ , then

$$f_{r,t}(z) = \sum_{n \equiv r \pmod{t}} a(n)q^n$$

is the Fourier expansion of a modular form in  $M_k(\Gamma_1(\frac{Nt^2}{d}))$

In order to state the last lemma, we need

**Definition** Let  $l$  be a positive integer and  $f = \sum_{n \geq N} a(n)q^n$  a formal power series in the variable  $q$  with rational integer coefficients. The  $l$ -adic order of  $f$  is defined by

$$\text{Ord}_l(f) = \inf\{n \mid a(n) \not\equiv 0 \pmod{l}\}$$

Sturm proved the following criterion for determining whether two modular forms are congruent for primes, One [10] noted the criterion holds for general integers as modulus.

**Lemma 2.4** ([14]). Let  $l$  be a positive integer and  $f(z), g(z) \in M_k(\Gamma_0(N))$  with rational integers satisfying

$$\text{ord}_l(f(z) - g(z)) \geq 1 + \frac{kN}{12} \prod_p \left(1 + \frac{1}{p}\right),$$

where the product is over the prime divisors  $p$  of  $N$ . Then  $f(z) \equiv g(z) \pmod{l}$ , i.e.,  $\text{ord}_l(f(z) - g(z)) = \infty$ .

### 3 Proof of Theorem 1.1

We construct modular forms to prove our theorem.

*Proof of Theorem 1.1.* Define the Eta-product

$$\begin{aligned} f(z) &= \frac{\eta^7(15z)\eta^9(45z)}{\eta^3(z)\eta(3z)} \\ &= q^{21} \left( \prod_{n=1}^{\infty} \frac{1}{(1-q^n)^3(1-q^{3n})} \right) \left( \prod_{n=1}^{\infty} (1-q^{15n})^7(1-q^{45n})^9 \right) \\ &=: \sum_{n \geq 21} r(n)q^n \end{aligned}$$

It easily can be seen that our Theorem 1.1 is equivalent to the two congruences: for every nonnegative integer  $n$ ,

$$r(15n+3) \equiv 0 \pmod{5} \text{ and } r(15n+12) \equiv 0 \pmod{5}.$$

Setting  $N = 45$ , by Lemma 2.1 and 2.2 it is easy to check that  $f(z) \in M_6(\Gamma_0(45))$ . In order to prove the equivalent congruences above, We use Lemma 2.3 and 2.4 To be more precise, define a modular form

$$f_{3,15}(z) := \sum_{n \equiv 3 \pmod{15}, n \geq 21} r(n)q^n$$

By Lemma 2.3 we find that  $f_{3,15}(z) \in M_6(\Gamma_1(3375))$ . By Lemma 2.4,  $r(15n+3) \equiv 0 \pmod{5}$  for every nonnegative integer  $n$  can be proved if it holds for the first 12150001 terms which is verified by machinery calculation. The another congruence is proved similarly.  $\blacksquare$

## 4 Proof of theorem 1.2

Recall that the three-colored partition function  $c\phi_3(n)$  has generating function

$$\sum_{n=0}^{\infty} c\phi_3(n)q^n = \sum_{n=0}^{\infty} p(n)q^n + 9q \prod_{n=1}^{\infty} \frac{(1-q^{9n})^3}{(1-q^{3n})(1-q^n)^3} := \sum_{n=0}^{\infty} (p(n) + \bar{c}\phi_3(n))q^n$$

Where  $p(n)$  is the ordinary partition function with the convention that  $p(n) = 0$ , if  $n \notin \mathbb{Z}$ . We will use modified Jacobi formula

$$q \prod_{n=1}^{\infty} (1-q^{9n})^3 = \sum_{n=0}^{\infty} (-1)^n (2n+1) q^{(9n^2+9n)/2+1}.$$

*Proof of Theorem 1.2.* By Jacobi formula above,

$$\begin{aligned} c\phi_3(45n+7) &= \bar{c}\phi_3(45n+7) \\ &= 9 \sum_{k \geq 0} (-1)^k (2k+1) (a(45n+7 - (1 + \frac{9k^2+9k}{2}))), \\ c\phi_3(45n+22) &= \bar{c}\phi_3(45n+22) \\ &= 9 \sum_{k \geq 0} (-1)^k (2k+1) (a(45n+22 - (1 + \frac{9k^2+9k}{2}))), \\ c\phi_3(45n+37) &= \bar{c}\phi_3(45n+22) \\ &= 9 \sum_{k \geq 0} (-1)^k (2k+1) (a(45n+37 - (1 + \frac{9k^2+9k}{2}))), \end{aligned}$$

Modulo 45,  $1 + (9k^2 + 9k)/2$  is 1, 10, or 28 and therefore  $45n + 7 - (1 + \frac{9k^2+9k}{2})$ ,  $45n + 22 - (1 + \frac{9k^2+9k}{2})$  and  $45n + 37 - (1 + \frac{9k^2+9k}{2})$  are congruent to  $45n + 6, 9, 12, 21, 24, 27, 36, 39, 42$ . The cases of  $45n + 9, 24, 39$  correspond to  $1 + (9k^2 + 9k)/2 \equiv 0 \pmod{45}$ , which is equivalent to  $k \equiv 2 \pmod{5}$ , we find that in this case the factors  $2k + 1 \equiv 0 \pmod{5}$ , Moreover  $45n + 6, 12, 21, 27, 36, 42$  are always the form of  $15n + 6$  or  $15n + 12$ . So by Theorem 1.1, Theorem 1.2 is proved.  $\blacksquare$

## References

- [1] G.E. Andrews, Generalized Frobenius partitions, *Mem. Amer. Math. Soc.* 49(1984), 301

- [2] William Y.C. Chen, Bernard L.S. Lin, Congruences for the number of cubic partitions derived from modular forms, arXiv:math.NT/0910.1263
- [3] D. Eichhorn and K. Ono, Congruences for partition functions, Proceedings for a Conference in Honor of Heini Halbertstam, 1996, 309-321.
- [4] B.Gordon and K. Hughies, Multiplicative properties of  $\eta$ -products, *Invent. Math.* 101(1990). 1-17
- [5] M.D. Hirschhorn and D.C. Hunt, A simple proof of the Ramanujan conjecture for powers of 5, *J. Reine Angew. Math.*, 326 (1981), 1–17.
- [6] N. Koblitz, "Introduction to Elliptic Curves and Modular Forms," Springer-Verlag, Berlin/New York, 1984
- [7] L. Kolitsch, A relationship between certain colored generalized Frobenius Partitions and ordinary partitions, *J. Number Th.* 33, (1989), 220-223.
- [8] G. Ligozat, "Courbes modulaires de genre 1," Vol 403, Bulletin de la Soc. Math. de France, 1975.
- [9] K. Ono, Congruences for Frobenius partitions, *J. Number Theory*, 57 (1996), 170–180.
- [10] K. Ono, Parity of the partition function in arithmetic progressions, *Journal fur die reine und angewandte Mathematik*, 472 (1996), 1–15.
- [11] K. Ono, Distribution of the partition function modulo  $m$ , *Ann. Math.*, 151 (2000), 293–307.
- [12] S. Ramanujan, Congruence properties of partitions, *Proc. Lond. Math. Soc. (2)*, 19 (1919), 207–210.
- [13] J. Sellers, New congruences for generalized Frobenius partitions with 2 or 3 colors, *Discrete Mathematics*, 131 (1994), 367–374.
- [14] J. Sturm, On the congruence of modular forms, *Lecture Notes in Math.* 1240 (1984), 275–280.