

**PARAQUATERNIONIC CR-SUBMANIFOLDS OF  
PARAQUATERNIONIC KÄHLER MANIFOLDS AND  
SEMI-RIEMANNIAN SUBMERSIONS**

STERE IANUȘ, STEFANO MARCHIAFAVA, GABRIEL EDUARD VÎLCU

ABSTRACT. In this paper we introduce paraquaternionic CR-submanifolds of almost paraquaternionic hermitian manifolds and state some basic results on their differential geometry. We also study a class of semi-Riemannian submersions from paraquaternionic CR-submanifolds of paraquaternionic Kähler manifolds.

*AMS Mathematics Subject Classification:* 53C15.

*Key Words and Phrases:* paraquaternionic Kähler manifold, foliation, CR-submanifold, semi-Riemannian submersion.

1. INTRODUCTION

The notion of CR-submanifold of a Kähler manifold was introduced by Bejancu [6] as a generalization both of totally real and of holomorphic submanifolds of Kähler manifolds. Then many papers appeared studying geometry of CR-submanifolds of Kähler manifolds and this notion was further extended to other ambient spaces; in the monographs [7, 12, 35] we can find the most significant results concerning CR-submanifolds. A class of examples of CR-submanifolds of manifolds endowed with different geometric structures is given in [30].

On the other hand, the paraquaternionic structures, firstly named quaternionic structures of second kind, have been introduced by P. Libermann in [26]. The differential geometry of manifolds endowed with this kind of structures is a very interesting subject and these manifolds have been intensively studied by many authors (see, e.g., [2, 3, 10, 14, 17, 23, 24, 36]). The study of submanifolds of a paraquaternionic Kähler manifold is also of interest and several types of such submanifolds we can find in the recent literature: paraquaternionic submanifolds [31], Kähler and para-Kähler submanifolds [2, 27], normal semi-invariant submanifolds [1, 8], lightlike submanifolds [19, 20],  $F$ -invariant submanifolds [32]. In this note we define a new class of submanifolds of paraquaternionic Kähler manifolds, which we call paraquaternionic CR-submanifolds, as a natural extension of CR-submanifolds in paraquaternionic setting.

The paper is organized as follows: in Section 2 we collect basic definitions, some formulas and results for later use. In Section 3 we introduce the concept of paraquaternionic CR-submanifold and show that on the normal bundle of a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold exist two complementary orthogonal distributions. In Section 4 we investigate the integrability of the distributions involved in the definition of a paraquaternionic CR-submanifold. In Section 5, following the same techniques as in [18], we study the canonical foliation induced on a paraquaternionic CR-submanifold; conditions

on total geodesicity are derived. We also obtain necessary and sufficient conditions for a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold to be a ruled submanifold with respect to the canonical foliation. In Section 6 we define the paraquaternionic CR-submersions (in the sense of Kobayashi [25]) as semi-Riemannian submersions from paraquaternionic CR-submanifolds onto an almost paraquaternionic hermitian manifold and obtain some properties concerning their geometry. In Section 7 we discuss curvature properties of fibers and base manifold for paraquaternionic CR-submersions.

## 2. PRELIMINARIES

Let  $M$  be a smooth manifold endowed with an *almost paraquaternionic structure*  $\sigma$ , that is, a rank-3 subbundle  $\sigma$  of  $End(TM)$  which admits a local basis  $\{J_1, J_2, J_3\}$  on a coordinate neighborhood  $U \subset M$  such that we have:

$$(1) \quad J_\alpha^2 = -\epsilon_\alpha Id, \quad J_\alpha J_{\alpha+1} = -J_{\alpha+1} J_\alpha = \epsilon_{\alpha+2} J_{\alpha+2}$$

where  $\epsilon_1 = 1$ ,  $\epsilon_2 = \epsilon_3 = -1$  and the indices are taken from  $\{1, 2, 3\}$  modulo 3. Moreover, the pair  $(M, \sigma)$  is said to be an *almost paraquaternionic manifold* and  $\{J_1, J_2, J_3\}$  is called a *canonical local basis* of  $M$ .

In an almost paraquaternionic manifold  $(M, \sigma)$  we take intersecting coordinate neighborhoods  $U$  and  $U'$ . Let  $\{J_1, J_2, J_3\}$  and  $\{J'_1, J'_2, J'_3\}$  be canonical local bases of  $\sigma$  in  $U$  and  $U'$  respectively. Then  $\{J'_1, J'_2, J'_3\}$  are linear combinations of  $\{J_1, J_2, J_3\}$  in  $U \cap U'$ :

$$(2) \quad J'_\alpha = \sum_{\beta=1}^3 a_{\alpha\beta} J_\beta, \quad \alpha = 1, 2, 3,$$

where  $a_{\alpha\beta}$  are functions in  $U \cap U'$ ,  $\alpha, \beta = 1, 2, 3$  and  $A = (a_{\alpha\beta})_{\alpha, \beta=1, 2, 3} \in SO(2, 1)$ .

Let  $(M, g)$  be a semi-Riemannian manifold and let  $\sigma$  be an almost paraquaternionic structure on  $M$ . The metric  $g$  is said to be *adapted* to the almost paraquaternionic structure  $\sigma$  if it satisfies:

$$(3) \quad g(J_\alpha X, J_\alpha Y) = \epsilon_\alpha g(X, Y), \quad \alpha \in \{1, 2, 3\},$$

for all vector fields  $X, Y$  on  $M$  and any local basis  $\{J_1, J_2, J_3\}$  of  $\sigma$ ; or, equivalently, if all endomorphisms of  $\sigma$  are skew-symmetric with respect to  $g$ . In this case,  $(M, \sigma, g)$  is said to be an *almost paraquaternionic hermitian manifold*. The existence of paraquaternionic hermitian structures on manifolds and tangent bundles has been recently investigated in [21].

It is easy to see that any almost paraquaternionic hermitian manifold is of dimension  $4m$ ,  $m \geq 1$ , and any adapted metric is necessarily of neutral signature  $(2m, 2m)$ .

Let  $\{J_1, J_2, J_3\}$  be a canonical local base of  $\sigma$  in a coordinate neighborhood  $U$  of an almost paraquaternionic hermitian manifold  $(M, \sigma, g)$ . If we denote by:

$$\Omega_{J_\alpha}(X, Y) = g(X, J_\alpha Y), \quad \alpha = 1, 2, 3$$

for any vector fields  $X$  and  $Y$ , then, by means of (2), we see that:

$$\Omega = \Omega_{J_1} \wedge \Omega_{J_1} - \Omega_{J_2} \wedge \Omega_{J_2} - \Omega_{J_3} \wedge \Omega_{J_3}$$

is a globally well-defined 4-form on  $M$ , called *the fundamental 4-form* of the manifold.

If  $(M, \sigma, g)$  is an almost paraquaternionic hermitian manifold such that the bundle  $\sigma$  is parallel with respect to the Levi-Civita connection  $\bar{\nabla}$  of  $g$ , then  $(M, \sigma, g)$  is said to be a paraquaternionic Kähler manifold. Equivalently, locally defined 1-forms  $\omega_1, \omega_2, \omega_3$  exist such that we have for all  $\alpha \in \{1, 2, 3\}$ :

$$(4) \quad \bar{\nabla}_X J_\alpha = -\epsilon_\alpha [\omega_{\alpha+2}(X)J_{\alpha+1} - \omega_{\alpha+1}(X)J_{\alpha+2}]$$

for any vector field  $X$  on  $M$ , where the indices are taken from  $\{1, 2, 3\}$  modulo 3 (see [17]). Moreover, it can be proved that an almost paraquaternionic hermitian manifold of dimension strictly greater than 4 is paraquaternionic Kähler if and only if  $\nabla\Omega = 0$  (see [22]).

We remark that any paraquaternionic Kähler manifold is an Einstein manifold, provided that  $\dim M > 4$  (see [10, 17, 23]).

Let  $(M, g)$  be a semi-Riemannian manifold and let  $N$  be an immersed submanifold of  $M$ . Then  $N$  is said to be a non-degenerate submanifold of  $M$  if the restriction of the semi-Riemannian metric  $g$  to  $TN$  is non-degenerate at each point of  $N$ . We denote by the same symbol  $g$  the semi-Riemannian metric induced by  $g$  on  $N$  and by  $TN^\perp$  the normal bundle to  $N$ .

For the rest of this section we will assume that the induced metric on  $N$  is non-degenerate.

Then we have the following orthogonal decomposition:

$$TM = TN \oplus TN^\perp.$$

Also, we denote by  $\bar{\nabla}$  and  $\nabla$  the Levi-Civita connection on  $M$  and  $N$ , respectively. Then the Gauss formula is given by:

$$(5) \quad \bar{\nabla}_X Y = \nabla_X Y + B(X, Y)$$

for any  $X, Y \in \Gamma(TN)$ , where  $B : \Gamma(TN) \times \Gamma(TN) \rightarrow \Gamma(TN^\perp)$  is the second fundamental form of  $N$  in  $M$ .

On the other hand, the Weingarten formula is given by:

$$(6) \quad \bar{\nabla}_X \xi = -A_\xi X + \nabla_X^\perp \xi$$

for any  $X \in \Gamma(TN)$  and  $\xi \in \Gamma(TN^\perp)$ , where  $-A_\xi X$  is the tangent part of  $\bar{\nabla}_X \xi$  and  $\nabla_X^\perp \xi$  is the normal part of  $\bar{\nabla}_X \xi$ ;  $A_\xi$  and  $\nabla^\perp$  are called the shape operator of  $N$  with respect to  $\xi$  and the normal connection, respectively. Moreover,  $B$  and  $A_\xi$  are related by:

$$(7) \quad g(B(X, Y), \xi) = g(A_\xi X, Y)$$

for any  $X, Y \in \Gamma(TN)$  and  $\xi \in \Gamma(TN^\perp)$  (see [29]).

If we denote by  $\bar{R}$  and  $R$  the curvature tensor fields of  $\bar{\nabla}$  and  $\nabla$  we have the Gauss equation:

$$(8) \quad \bar{R}(X, Y, Z, U) = R(X, Y, Z, U) - g(B(X, Z), B(Y, U)) + g(B(Y, Z), B(X, U)),$$

for all  $X, Y, Z, U \in \Gamma(TN)$ .

### 3. PARAQUATERNIONIC CR-SUBMANIFOLDS

**Definition 3.1.** Let  $N$  be an  $n$ -dimensional non-degenerate submanifold of an almost paraquaternionic hermitian manifold  $(M, \sigma, g)$ . We say that  $(N, g)$  is a *paraquaternionic CR-submanifold* of  $M$  if there exists a non-degenerate distribution

$\mathcal{D} : x \rightarrow \mathcal{D}_x \subseteq T_x N$  such that on any  $U \cap N$  we have:

i.  $\mathcal{D}$  is a paraquaternionic distribution, i.e.

$$(9) \quad J_\alpha \mathcal{D}_x = \mathcal{D}_x, \quad \alpha \in \{1, 2, 3\}$$

and

ii.  $\mathcal{D}^\perp$  is a totally real distribution, i.e.

$$(10) \quad J_\alpha \mathcal{D}_x^\perp \subset T_x^\perp N, \quad \alpha \in \{1, 2, 3\}$$

for any local basis  $\{J_1, J_2, J_3\}$  of  $\sigma$  on  $U$  and  $x \in U \cap N$ , where  $\mathcal{D}^\perp$  is the orthogonal complementary distribution to  $\mathcal{D}$  in  $TN$ .

**Definition 3.2.** A non-degenerate submanifold  $N$  of an almost paraquaternionic hermitian manifold  $(M, \sigma, g)$  is called a *paraquaternionic* (respectively, *totally real*) submanifold if  $\mathcal{D}^\perp = 0$  (respectively,  $\mathcal{D} = 0$ ). A paraquaternionic CR-submanifold is said to be *proper* if it is neither paraquaternionic nor totally real.

*Example 3.3.* i. The canonical immersion of  $P^n \mathbb{B}(c)$  into  $P^m \mathbb{B}(c)$ , where  $n \leq m$ , provides us a very natural example of paraquaternionic submanifold (see [27]).

ii. The real projective space  $P_s^n \mathbb{R}(\frac{c}{4})$  is a totally-real submanifold of the paraquaternionic projective space  $P^n \mathbb{B}(c)$ , where  $s \in \{0, \dots, n\}$  denotes the index of the manifold, defined as the dimension of the largest negative definite vector subspace of the tangent space.

iii. Let  $(M_1, g_1, \sigma_1)$  and  $(M_2, g_2, \sigma_2)$  be two paraquaternionic Kähler manifolds. If  $U_1$  and  $U_2$  are open subsets of  $M_1$  and  $M_2$  respectively, on which local basis  $\{J_1^{(1)}, J_2^{(1)}, J_3^{(1)}\}$  and  $\{J_1^{(2)}, J_2^{(2)}, J_3^{(2)}\}$  for  $\sigma_1$  and  $\sigma_2$  respectively, are defined, then the product manifold  $U = U_1 \times U_2$  can be endowed with an almost paraquaternionic hermitian non-Kähler structure  $(g, \sigma)$  (see [32]). Now, if  $N_1$  is a paraquaternionic submanifold of  $U_1$  and  $N_2$  is a totally-real submanifold of  $U_2$ , then  $N = N_1 \times N_2$  is a proper paraquaternionic CR-submanifold of the almost paraquaternionic hermitian manifold  $(U, g, \sigma)$ .

iv. A large class of examples of proper paraquaternionic CR-submanifolds can be constructed using the paraquaternionic momentum map [33] and the technique from [30]. Suppose that a Lie group  $G$  acts freely and isometrically on the paraquaternionic Kähler manifold  $(M, \sigma, g)$ , preserving the fundamental 4-form  $\Omega$  of the manifold. We denote by  $\mathfrak{g}$  the Lie algebra of  $G$ , by  $\mathfrak{g}^*$  its dual and by  $V$  the unique Killing vector field corresponding to a vector  $V^* \in \mathfrak{g}$ . Then there exists a unique section  $f$  of bundle  $\mathfrak{g}^* \otimes \sigma$  such that (see [33])

$$(11) \quad \nabla f_{V^*} = \theta_{V^*},$$

for all  $V^* \in \mathfrak{g}$ , where the section  $\theta_{V^*}$  of the bundle  $\Omega^1(\sigma)$  with values in  $\sigma$  is well defined globally by

$$\theta_{V^*}(X) = \sum_{\alpha=1}^3 \omega_\alpha(V, X) J_\alpha, \quad \forall X \in TM.$$

Moreover, the group  $G$  acts by isometries on the pre-image  $f^{-1}(0)$  of the zero-section  $0 \in \mathfrak{g}^* \otimes \sigma$ . Similarly as in [30], we have the decomposition

$$T_x(f^{-1}(0)) = T_x(G \cdot x) \oplus H_x, \quad \forall x \in M,$$

where  $G \cdot x$  represents the orbit of  $G$  through  $x$ , supposed to be non-degenerate, and  $H_x$  is the orthogonal complementary subspace of  $T_x(G \cdot x)$  in  $T_x(f^{-1}(0))$ . Because

$H_x$  is invariant under the action of  $\sigma$  and  $T_x(G \cdot x)$  is totally real, we can state now the following result.

*Proposition 3.4.* *If  $f^{-1}(0)$  is a smooth submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ , then  $f^{-1}(0)$  is a proper paraquaternionic CR-submanifold of  $M$ .*

We remark that, in general,  $f^{-1}(0)$  is not a differentiable submanifold of  $M$ , but always we can take a subset  $N \subset f^{-1}(0)$  which is invariant under the action of  $G$  and which is a submanifold of  $M$ . A particular example is given in [11], as a paraquaternionic version of the example constructed by Galicki and Lawson in [16]: if  $p$  and  $q$  are distinct and relatively prime natural numbers, then we have the action of the Lie group  $G = \{e^{jt} | t \in \mathbb{R}\}$  on  $P^2\mathbb{B}$  defined by

$$\phi_{p,q}(t) \cdot [u_0, u_1, u_2] := [e^{jqt}u_0, e^{jpt}u_1, e^{jpt}u_2],$$

where  $e^{jt} = \cos ht + j \sin ht$  and  $[u_0, u_1, u_2]$  are homogenous coordinates on  $P^2\mathbb{B}$ . We can see that this action is free, isometric and preserves the para-quaternionic structure on  $P^2\mathbb{B}$  and, moreover, we have that the pre-image by the momentum map  $f_{p,q} : P^2\mathbb{B} \rightarrow \text{Im}\mathbb{B}$  of the zero-section  $0 \in \text{Im}\mathbb{B}$  is (see also [33]):

$$f_{p,q}^{-1}(0) = \{[u_0, u_1, u_2] \in P^2\mathbb{B} | q\bar{u}_0ju_0 + p\bar{u}_1ju_1 + p\bar{u}_2ju_2 = 0\}.$$

Finally, we conclude that the subset  $N$  of the regular points of  $f_{p,q}^{-1}(0)$ , given by

$$N = \{[u_0, u_1, u_2] \in f_{p,q}^{-1}(0) | q^2|u_0|^2 + p^2|u_1|^2 + p^2|u_2|^2 \neq 0\}$$

is a proper paraquaternionic CR-submanifold of  $P^2\mathbb{B}$ .

**Definition 3.5.** Let  $N$  be a paraquaternionic CR-submanifold of an almost paraquaternionic hermitian manifold  $(M, \sigma, g)$ . Then we say that:

- i.  $N$  is  $\mathcal{D}$ -geodesic if  $B(X, Y) = 0, \forall X, Y \in \Gamma(\mathcal{D})$ ;
- ii.  $N$  is  $\mathcal{D}^\perp$ -geodesic if  $B(X, Y) = 0, \forall X, Y \in \Gamma(\mathcal{D}^\perp)$ ;
- iii.  $N$  is mixed geodesic if  $B(X, Y) = 0, \forall X \in \Gamma(\mathcal{D}), Y \in \Gamma(\mathcal{D}^\perp)$ ;
- iv.  $N$  is mixed foliated if  $N$  is mixed geodesic and  $\mathcal{D}$  is integrable.

We may easily prove the next result (see [4, 31]):

**Proposition 3.6.** *Any paraquaternionic submanifold of a paraquaternionic Kähler manifold is a totally geodesic paraquaternionic Kähler submanifold.*

By using this proposition, we deduce the next consequences.

**Corollary 3.7.** *Let  $(N, g)$  be a paraquaternionic submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . Then:*

- i.  $\dim N = 4n, n \geq 1$  and the signature of  $g|_{TN}$  is  $(2n, 2n)$ ;
- ii.  $N$  is an Einstein manifold, provided that  $\dim N > 4$ .

**Corollary 3.8.** *The paraquaternionic submanifolds of  $\mathbb{R}_{2m}^{4m}$  and of paraquaternionic projective space  $P^m\mathbb{B}$  are locally isometric with  $\mathbb{R}_{2n}^{4n}$  and  $P^n\mathbb{B}$ , respectively, where  $n \leq m$ .*

Next, let  $(N, g)$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . We put  $\nu_{\alpha x} = J_\alpha(D_x^\perp)$ ,  $\alpha \in \{1, 2, 3\}$ , and  $\nu_x^\perp = \nu_{1x} \oplus \nu_{2x} \oplus \nu_{3x}$ , and remark that  $\nu_{1x}, \nu_{2x}, \nu_{3x}$  are mutually orthogonal non-degenerate vector subspaces of  $T_x N^\perp$ , for any  $x \in U \cap N$ . We also note that the subspaces  $\nu_{\alpha x}$  depends on the choice of the local base  $(J_\alpha)_\alpha$ , while  $\nu_x^\perp$  does't depend from it.

**Proposition 3.9.** *Let  $(N, g)$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . Then we have:*

- i.  $J_\alpha(\nu_{\alpha x}) = \mathcal{D}_x^\perp$ ,  $\forall x \in U \cap N$ ,  $\alpha \in \{1, 2, 3\}$ ;
- ii.  $J_\alpha(\nu_{\beta x}) = \nu_{\gamma x}$ , for any even permutation  $(\alpha, \beta, \gamma)$  of  $(1, 2, 3)$  and  $x \in U \cap N$ ;
- iii. The mapping  $\nu^\perp : x \in N \rightarrow \nu_x^\perp$  defines a non-degenerate distribution of dimension  $3s$ , where  $s = \dim \mathcal{D}_x^\perp$ ;
- iv.  $J_\alpha(\nu_x) = \nu_x$ ,  $\forall x \in U \cap N$ ,  $\alpha \in \{1, 2, 3\}$ , where  $\nu$  is the complementary orthogonal distribution to  $\nu^\perp$  in  $TN^\perp$ .

*Proof.* The assertions i. and ii. are obvious from (1). The assertion iii. follows from the Definition 3.1, since  $J_\alpha$ ,  $\alpha \in \{1, 2, 3\}$ , are automorphisms of  $T_x N$ , for any  $x \in U \cap N$ . Concerning the proof of (iv.), firstly one can remark that  $\nu_x$  doesn't depend from the choice of the local base  $(J_\alpha)_\alpha$  and secondly, one could observe that the subspace  $TN_x^Q = \mathcal{D}_x \oplus \mathcal{D}_x^\perp \oplus \nu_x^\perp$  is the paraquaternionic subspace generated by  $TN_x$ , hence its orthogonal  $\nu$  is also a paraquaternionic subspace, i.e. closed under  $J_\alpha$ ,  $\alpha \in \{1, 2, 3\}$ . This completes the proof of the proposition.  $\square$

#### 4. INTEGRABILITY OF DISTRIBUTIONS

**Theorem 4.1.** *The distribution  $\mathcal{D}^\perp$  is integrable.*

*Proof.* For any  $U, V \in \Gamma(\mathcal{D}^\perp)$  and  $X \in \Gamma(\mathcal{D})$  we have:

$$\begin{aligned} g(\nabla_U V, X) &= g(\overline{\nabla}_U V, X) \\ &= \epsilon_\alpha g(J_\alpha \overline{\nabla}_U V, J_\alpha X) \\ &= \epsilon_\alpha g(\epsilon_\alpha [\omega_{\alpha+2}(U) J_{\alpha+1} V - \omega_{\alpha+1}(U) J_{\alpha+2} V] + \overline{\nabla}_U J_\alpha V, J_\alpha X) \\ &= \epsilon_\alpha g(\overline{\nabla}_U J_\alpha V, J_\alpha X) \end{aligned}$$

and by using (6) we obtain:

$$(12) \quad g(\nabla_U V, X) = -\epsilon_\alpha g(A_{J_\alpha V} U, J_\alpha X).$$

On the other hand, if we take  $U, V \in \Gamma(\mathcal{D}^\perp)$  and  $C \in \Gamma(TN)$  we have:

$$\begin{aligned} g(A_{J_\alpha V} U, C) &= -g(\overline{\nabla}_C J_\alpha V, U) \\ &= g(\epsilon_\alpha [\omega_{\alpha+2}(C) J_{\alpha+1} V - \omega_{\alpha+1}(C) J_{\alpha+2} V], U) - g(J_\alpha \overline{\nabla}_C V, U) \end{aligned}$$

hence

$$g(A_{J_\alpha V} U, C) = -g(J_\alpha \overline{\nabla}_C V, U).$$

Moreover, from  $C \cdot g(J_\alpha U, V) = 0$ , one has

$$g(J_\alpha \overline{\nabla}_C U, V) = g(J_\alpha \overline{\nabla}_C V, U).$$

We conclude:

$$(13) \quad g(A_{J_\alpha V} U, C) = g(A_{J_\alpha U} V, C).$$

From (12) and (13) we deduce that for any  $U, V \in \Gamma(\mathcal{D}^\perp)$  and  $X \in \Gamma(\mathcal{D})$  we have:

$$g(\nabla_U V - \nabla_V U, X) = 0,$$

which implies  $[U, V] \in \Gamma(\mathcal{D}^\perp)$ ,  $\forall U, V \in \Gamma(\mathcal{D}^\perp)$ . Thus  $\mathcal{D}^\perp$  is integrable.  $\square$

**Theorem 4.2.** *The paraquaternionic distribution  $\mathcal{D}$  is integrable if and only if  $N$  is  $\mathcal{D}$ -geodesic.*

*Proof.* Similarly as in above theorem, we obtain:

$$g(\nabla_X Y, U) = \epsilon_\alpha g(\overline{\nabla}_X J_\alpha Y, J_\alpha U)$$

for any  $X, Y \in \Gamma(\mathcal{D})$  and  $U \in \Gamma(\mathcal{D}^\perp)$ , and taking into account (5) we obtain:

$$(14) \quad g(\nabla_X Y, U) = \epsilon_\alpha g(B(X, J_\alpha Y), J_\alpha U).$$

If we suppose that  $N$  is  $\mathcal{D}$ -geodesic, from (14) we derive  $\nabla_X Y \in \Gamma(\mathcal{D})$ , which implies  $[X, Y] \in \Gamma(\mathcal{D})$ . Thus  $\mathcal{D}$  is integrable.

Conversely, if we suppose that  $\mathcal{D}$  is integrable, then the leaves are invariant of the paraquaternionic structure and so they are totally geodesic in  $N$ . In particular,  $N$  is  $\mathcal{D}$ -geodesic.  $\square$

*Remark 4.3.* If  $\mathcal{Q}$  is a non-degenerate distribution on a semi-Riemannian manifold  $(M, g)$ , then we can consider a well-defined  $\mathcal{Q}^\perp$ -valued vector field on  $N$ , called the mean curvature vector of  $\mathcal{Q}$  (see [9]), given by:

$$H^\mathcal{Q} = \frac{1}{q} \sum_{i=1}^q \theta_i h^\mathcal{Q}(E_i, E_i),$$

where  $h^\mathcal{Q}$  is the second fundamental forms of  $\mathcal{Q}$ ,  $q = \dim \mathcal{Q}$ ,  $\{E_1, \dots, E_q\}$  is a pseudo-orthonormal basis of  $\mathcal{Q}$  and  $\theta_i = g(E_i, E_i) \in \{-1, 1\}$ ,  $\forall i \in \{1, \dots, q\}$ .

The distribution  $\mathcal{Q}$  is said to be *minimal* if the mean curvature vector  $H^\mathcal{Q}$  of  $\mathcal{Q}$  vanishes identically.

**Theorem 4.4.** *The paraquaternionic distribution  $\mathcal{D}$  is minimal.*

*Proof.* For any  $X \in \Gamma(D)$  and  $U \in \Gamma(D^\perp)$  we obtain similarly as in above theorems:

$$(15) \quad g(\nabla_X X, U) = \epsilon_\alpha g(A_{J_\alpha U} J_\alpha X, X)$$

and

$$(16) \quad g(\nabla_{J_\alpha X} J_\alpha X, U) = -g(A_{J_\alpha U} X, J_\alpha X) = -g(A_{J_\alpha U} J_\alpha X, X),$$

for all  $\alpha \in \{1, 2, 3\}$ .

From (15) and (16) we deduce:

$$g(\nabla_X X + \epsilon_\alpha \nabla_{J_\alpha X} J_\alpha X, U) = 0$$

and so

$$(17) \quad h^\mathcal{D}(X, X) + \epsilon_\alpha h^\mathcal{D}(J_\alpha X, J_\alpha X) = 0, \quad \forall \alpha \in \{1, 2, 3\}.$$

From (17) it follows that

$$h^\mathcal{D}(X, X) = -h^\mathcal{D}(J_1 X, J_1 X) = -h^\mathcal{D}(J_2 J_3 X, J_2 J_3 X) = -h^\mathcal{D}(J_3 X, J_3 X) = -h^\mathcal{D}(X, X)$$

and so

$$(18) \quad h^\mathcal{D}(X, X) = 0, \quad \forall X \in \Gamma(D).$$

We obtain  $H^\mathcal{D} = 0$  and the assertion follows. We can remark that the relation (18) not imply  $h^\mathcal{D} = 0$ , since  $h^\mathcal{D}$  is not symmetric in general, unless  $\mathcal{D}$  is integrable.  $\square$

*Remark 4.5.* For any paraquaternionic CR-submanifold  $(N, g)$  of an almost paraquaternionic hermitian manifold  $(M^{4m}, \sigma, g)$ , having  $\dim \mathcal{D} = 4r$  and  $\dim \mathcal{D}^\perp = p$ , we can choose a local pseudo-orthonormal frame in  $M$ :

$\{e_1, \dots, e_r, e_{r+1}, \dots, e_{r+p}, e_{r+p+1}, \dots, e_m, J_1 e_1, \dots, J_1 e_m, J_2 e_1, \dots, J_2 e_m, J_3 e_1, \dots, J_3 e_m\}$   
such that restricted to  $N$ ,  $\{e_i, J_1 e_i, J_2 e_i, J_3 e_i\}_{i \in \{1, \dots, r\}}$  are in  $\mathcal{D}$  and  $\{e_{r+1}, \dots, e_{r+p}\}$  are in  $\mathcal{D}^\perp$ .

Let  $\{\omega^1, \dots, \omega^{4r}\}$  be the  $4r$  1-forms on  $N$  satisfying:

$$(19) \quad \omega^i(Z) = 0, \quad \omega_i(E_j) = \delta_{ij}, \quad i, j \in \{1, \dots, 4r\},$$

for any  $Z \in \Gamma(\mathcal{D}^\perp)$ , where  $E_j = e_j$ ,  $E_{r+j} = J_1 e_j$ ,  $E_{2r+j} = J_2 e_j$  and  $E_{3r+j} = J_3 e_j$ ,  $j \in \{1, \dots, r\}$ . Then  $\omega = \omega^1 \wedge \dots \wedge \omega^{4r}$  does not depend on the particular base  $\{J_1, J_2, J_3\}$  and defines a  $4r$ -form on  $N$ . Therefore, we have

$$(20) \quad d\omega = \sum_{i=1}^{4r} (-1)^i \omega^1 \wedge \dots \wedge d\omega^i \wedge \dots \wedge \omega^{4r}$$

and taking account of (19), we deduce that  $d\omega = 0$  if and only if

$$(21) \quad d\omega(Z_1, Z_2, X_1, \dots, X_{4r-1}) = 0$$

and

$$(22) \quad d\omega(Z_1, X_1, \dots, X_{4r}) = 0$$

for any  $Z_1, Z_2 \in \Gamma(\mathcal{D}^\perp)$  and  $X_1, \dots, X_{4r} \in \Gamma(\mathcal{D})$ . We remark now easily that (21) holds if and only if  $\mathcal{D}^\perp$  is integrable and (22) holds if and only if  $\mathcal{D}$  is minimal (see also [13]). On another hand, if  $\{\omega^{4r+1}, \dots, \omega^{4r+p}\}$  is the dual frame to the pseudo-orthonormal frame  $\{e_{r+1}, \dots, e_{r+p}\}$  of  $\mathcal{D}^\perp$ , we can define a  $p$ -form  $\omega^\perp$  on  $M$  by  $\omega^\perp = \omega^{4r+1} \wedge \dots \wedge \omega^{4r+p}$ . Similarly we find that  $\omega^\perp$  is closed if  $\mathcal{D}$  is integrable and  $\mathcal{D}^\perp$  is minimal.

Consequently, from Theorems 4.1 and 4.4, we obtain the following result.

*Theorem 4.6.* *Let  $N$  be a closed paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . Then the  $4r$ -form  $\omega$  is closed and defines a canonical de Rham cohomology class  $[\omega]$  in  $H^{4r}(M, \mathbb{R})$ . Moreover, this cohomology class is non-trivial if  $\mathcal{D}$  is integrable and  $\mathcal{D}^\perp$  is minimal.*

## 5. CANONICAL FOLIATIONS ON PARAQUATERNIONIC CR-SUBMANIFOLDS

Since the totally real distribution  $\mathcal{D}^\perp$  of a paraquaternionic CR-submanifold  $N$  of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  is always integrable we conclude that we have a foliation  $\mathfrak{F}^\perp$  on  $N$  with structural distribution  $\mathcal{D}^\perp$  and transversal distribution  $\mathcal{D}$  (see [9]). We say that  $\mathfrak{F}^\perp$  is the canonical totally real foliation on  $N$ .

**Theorem 5.1.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . The next assertions are equivalent:*

- i.*  $\mathfrak{F}^\perp$  is totally geodesic;
- ii.*  $B(X, Y) \in \Gamma(\nu)$ ,  $\forall X \in \Gamma(\mathcal{D}), Y \in \Gamma(\mathcal{D}^\perp)$ ;
- iii.*  $A_N X \in \Gamma(\mathcal{D}^\perp)$ ,  $\forall X \in \Gamma(\mathcal{D}^\perp), N \in \Gamma(\nu^\perp)$ ;
- iv.*  $A_N Y \in \Gamma(\mathcal{D}), \forall Y \in \Gamma(\mathcal{D}), N \in \Gamma(\nu^\perp)$ .



*Proof.* By using (3), (4), (5) and (6) we obtain for any  $X, Z \in \Gamma(\mathcal{D}^\perp)$  and  $Y \in \Gamma(\mathcal{D})$ :

$$\begin{aligned} g(J_\alpha(\nabla_X Z), Y) &= -g(\overline{\nabla}_X Z, J_\alpha Y) \\ &= g(\epsilon_\alpha[\omega_{\alpha+2}(X)J_{\alpha+1}Z - \omega_{\alpha+1}(X)J_{\alpha+2}Z] + \overline{\nabla}_X J_\alpha Z, Y) \\ &= g(-A_{J_\alpha Z}X + \nabla_X^\perp J_\alpha Z, Y) \\ &= -g(A_{J_\alpha Z}X, Y) \end{aligned}$$

and taking into account (7) we derive:

$$(23) \quad g(J_\alpha(\nabla_X Z), Y) = -g(B(X, Y), J_\alpha Z).$$

i.  $\Rightarrow$  ii. If  $\mathfrak{F}^\perp$  is totally geodesic, then  $\nabla_X Z \in \Gamma(\mathcal{D}^\perp)$ , for  $X, Z \in \Gamma(\mathcal{D}^\perp)$  and from (23) we derive:

$$g(B(X, Y), J_\alpha Z) = 0$$

and the implication is clear.

ii.  $\Rightarrow$  i. If we suppose  $B(X, Y) \in \Gamma(\mu)$ ,  $\forall X \in \Gamma(\mathcal{D})$ ,  $Y \in \Gamma(\mathcal{D}^\perp)$ , then from (23) we derive:

$$g(J_\alpha(\nabla_X Z), Y) = 0$$

and we conclude  $\nabla_X Z \in \Gamma(\mathcal{D}^\perp)$ . Thus  $\mathfrak{F}^\perp$  is totally geodesic.

ii.  $\Leftrightarrow$  iii. This equivalence is clear from (7).

iii.  $\Leftrightarrow$  iv. This equivalence is true because  $A_N$  is a self-adjoint operator.  $\square$

**Corollary 5.2.** *If  $N$  is a mixed-geodesic paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ , then the canonical totally real foliation  $\mathfrak{F}^\perp$  on  $N$  is totally geodesic.*

**Corollary 5.3.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  with  $\nu = 0$ . Then the canonical totally real foliation  $\mathfrak{F}^\perp$  on  $N$  is totally geodesic if and only if  $N$  is mixed geodesic.*

We note that the condition  $\nu = 0$  in Corollary 5.3 characterizes the following interesting class of paraquaternionic CR-submanifolds (compare with the definition in the complex case given in [35], pag. 78).

**Definition 5.4.** A paraquaternionic CR-submanifold  $N$  of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  is called *generic* if the restriction  $TM|_N$  to  $N$  of the tangent bundle of ambient manifold  $M$  is generated over the paraquaternions by the tangent bundle of  $N$ , i.e. if  $TN + \sigma TN = TM|_N$  or, equivalently, if  $\nu = 0$ .

*Remark 5.5.* The proper paraquaternionic CR-submanifolds given by Proposition 3.4 are generic because we have

$$T_x^\perp(f^{-1}(0)) = \sigma T_x(G \cdot x).$$

**Proposition 5.6.** *Let  $N$  be a generic paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . Then the paraquaternionic distribution  $\mathcal{D}$  is geodesic.*

*Proof.*  $N$  generic means that

$$TM|_N = \mathcal{D} \oplus \mathcal{D}^\perp \oplus \sigma \mathcal{D}^\perp$$

where

$$\nu^\perp = \sigma \mathcal{D}^\perp, \quad TN = \mathcal{D} \oplus \mathcal{D}^\perp.$$

From (3) and (14) we have

$$g(J_\alpha \nabla_X Y, J_\alpha U) = \epsilon_\alpha g(\nabla_X Y, U) = g(B(X, J_\alpha Y), J_\alpha U),$$

for all  $X, Y \in \Gamma(\mathcal{D})$ ,  $U \in \Gamma(\mathcal{D}^\perp)$  and since  $TN^\perp = \nu^\perp$ ,

$$(J_\alpha \nabla_X Y)^\perp = B(X, J_\alpha Y)$$

where  $W^\perp$  means the component of the vector  $W \in TN$  in the orthogonal to  $TN$ , that is in  $\nu^\perp$ .

More explicitly, we have

$$(J_\alpha \nabla_X^{\mathcal{D}} Y + J_\alpha \nabla_X^{\mathcal{D}^\perp} Y)^\perp = B(X, J_\alpha Y)$$

where  $\nabla_X^{\mathcal{D}} Y, \nabla_X^{\mathcal{D}^\perp} Y$  are the  $\mathcal{D}, \mathcal{D}^\perp$  components of  $\nabla_X Y$  respectively.

Equivalently, since  $J_\alpha \nabla_X^{\mathcal{D}} Y \in \mathcal{D}$ , we have

$$J_\alpha \nabla_X^{\mathcal{D}^\perp} Y = B(X, J_\alpha Y)$$

that is

$$(24) \quad \nabla_X^{\mathcal{D}^\perp} Y = -\epsilon_\alpha J_\alpha B(X, J_\alpha Y)$$

As a consequence, we have

$$\epsilon_\alpha J_\alpha B(X, J_\alpha Y) = \epsilon_\beta J_\beta B(X, J_\beta Y), \quad \forall \alpha, \beta$$

and, for  $J_\alpha J_\beta = \epsilon_\gamma J_\gamma$ ,

$$B(X, J_\alpha Y) = -\epsilon_\beta \epsilon_\gamma J_\gamma B(X, J_\beta Y).$$

Hence

$$(25) \quad B(X, Y) = -\epsilon_\alpha \epsilon_\beta J_\beta B(X, J_\alpha Y)$$

and also

$$(26) \quad B(X, J_\gamma Y) = J_\gamma B(X, Y)$$

since  $\epsilon_\gamma \epsilon_\alpha \epsilon_\beta = 1$ .

Note that from (26) and symmetry of  $B$  it follows

$$B(X, J_\gamma Y) = B(Y, J_\gamma X)$$

from which we deduce:

$$(27) \quad B(X, Y) = -\epsilon_\gamma B(J_\gamma X, J_\gamma Y).$$

By applying repeatedly the (27) we find

$$B(X, Y) = -B(J_1 X, J_1 Y) = -B(J_2 J_3 X, J_2 J_3 Y) = -B(J_3 X, J_3 Y) = -B(X, Y)$$

hence,

$$B = 0.$$

□

**Definition 5.7.** [9] A submanifold  $N$  of a semi-Riemannian manifold  $(M, g)$  is said to be a *ruled submanifold* if it admits a foliation whose leaves are totally geodesic submanifolds immersed in  $(M, g)$ .

**Definition 5.8.** A paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold which is a ruled submanifold with respect to the foliation  $\mathfrak{F}^\perp$  is called *totally real ruled paraquaternionic CR-submanifold*.

**Theorem 5.9.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . Then the following assertions are mutually equivalent:*

- i.  $N$  is a totally real ruled paraquaternionic CR-submanifold.*
- ii.  $N$  is  $\mathcal{D}^\perp$ -geodesic and:*

$$B(X, Y) \in \Gamma(\nu), \quad \forall X \in \Gamma(\mathcal{D}), \quad Y \in \Gamma(\mathcal{D}^\perp).$$

- iii. The subbundle  $\nu^\perp$  is  $\mathcal{D}^\perp$ -parallel, i.e:*

$$\nabla_X^\perp J_\alpha Z \in \Gamma(\nu^\perp), \quad \forall X, Z \in \Gamma(\mathcal{D}^\perp), \quad \alpha \in \{1, 2, 3\}$$

*and the second fundamental form satisfies:*

$$B(X, Y) \in \Gamma(\nu), \quad \forall X \in \Gamma(\mathcal{D}^\perp), \quad Y \in \Gamma(TN).$$

- iv. The shape operator satisfies:*

$$A_{J_\alpha Z} X = 0, \quad \forall X, Z \in \Gamma(\mathcal{D}^\perp), \quad \alpha \in \{1, 2, 3\}$$

*and*

$$A_N X \in \Gamma(\mathcal{D}), \quad \forall X \in \Gamma(\mathcal{D}^\perp), \quad N \in \Gamma(\nu).$$

*Proof.* i.  $\Leftrightarrow$  ii. This equivalence follows from Theorem 5.1 since for any  $X, Z \in \Gamma(\mathcal{D}^\perp)$  we have:

$$\begin{aligned} \bar{\nabla}_X Z &= \nabla_X Z + B(X, Z) \\ &= \nabla_X^{\mathcal{D}^\perp} Z + h^\perp(X, Z) + B(X, Z) \end{aligned}$$

and thus the leaves of  $\mathcal{D}^\perp$  are totally geodesic immersed in  $M$  if and only if  $h^\perp = 0$  and  $N$  is  $\mathcal{D}^\perp$ -geodesic.

- i.  $\Leftrightarrow$  iii.* If  $X, Z \in \Gamma(\mathcal{D}^\perp)$  and  $N \in \Gamma(\nu)$ , then we have:

$$\begin{aligned} g(\bar{\nabla}_X Z, N) &= \epsilon_\alpha g(J_\alpha \bar{\nabla}_X Z, J_\alpha N) \\ &= \epsilon_\alpha g(\epsilon_\alpha [\omega_{\alpha+2}(X) J_{\alpha+1} Z - \omega_{\alpha+1}(X) J_{\alpha+2} Z] + \bar{\nabla}_X J_\alpha Z, J_\alpha N) \\ &= \epsilon_\alpha g(-A_{J_\alpha Z} X + \nabla_X^\perp J_\alpha Z, J_\alpha N) \end{aligned}$$

and thus we obtain:

$$(28) \quad g(\bar{\nabla}_X Z, N) = \epsilon_\alpha g(\nabla_X^\perp J_\alpha Z, J_\alpha N).$$

Similarly we find:

$$(29) \quad g(\bar{\nabla}_X Z, U) = -\epsilon_\alpha g(B(X, J_\alpha U), J_\alpha Z), \quad \forall X, Z \in \Gamma(\mathcal{D}^\perp), \quad U \in \Gamma(\mathcal{D}).$$

On the other hand, from (5) we deduce for any  $X, Z, W \in \Gamma(\mathcal{D}^\perp)$ :

$$(30) \quad g(\bar{\nabla}_X Z, J_\alpha W) = g(B(X, Z), J_\alpha W).$$

But  $M$  is a totally real ruled paraquaternionic CR-submanifold iff  $\bar{\nabla}_X Z \in \Gamma(\mathcal{D}^\perp)$ ,  $\forall X, Z \in \Gamma(\mathcal{D}^\perp)$  and by using (28), (29) and (30) we deduce the equivalence.

- ii.  $\Leftrightarrow$  iv.* This equivalence follows from(7). □

**Corollary 5.10.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$ . If  $N$  is totally geodesic, then  $N$  is a totally real ruled paraquaternionic CR-submanifold.*

6. SEMI-RIEMANNIAN SUBMERSIONS FROM PARAQUATERNIONIC  
CR-SUBMANIFOLDS

Semi-Riemannian submersions were introduced by O'Neill [29]. Let  $(M, g)$  and  $(M', g')$  be two connected semi-Riemannian manifolds of index  $s$  ( $0 \leq s \leq \dim M$ ) and  $s'$  ( $0 \leq s' \leq \dim M'$ ) respectively, with  $s' \leq s$ . Roughly speaking, a semi-Riemannian submersion is a smooth map  $\pi : M \rightarrow M'$  which is onto and satisfies the following conditions:

- (i)  $\pi_{*|p} : T_p M \rightarrow T_{\pi(p)} M'$  is onto for all  $p \in M$ ;
- (ii) The fibres  $\pi^{-1}(p')$ ,  $p' \in M'$ , are semi-Riemannian submanifolds of  $M$ ;
- (iii)  $\pi_*$  preserves scalar products of vectors normal to fibres.

The vectors tangent to fibres are called vertical and those normal to fibres are called horizontal. We denote by  $\mathcal{V}$  the vertical distribution, by  $\mathcal{H}$  the horizontal distribution and by  $v$  and  $h$  the vertical and horizontal projection. An horizontal vector field  $X$  on  $M$  is said to be basic if  $X$  is  $\pi$ -related to a vector field  $X'$  on  $M'$ . It is clear that every vector field  $X'$  on  $M'$  has a unique horizontal lift  $X$  to  $M$  and  $X$  is basic.

*Remark 6.1.* If  $\pi : (M, g) \rightarrow (M', g')$  is a semi-Riemannian submersion and  $X, Y$  are basic vector fields on  $M$ ,  $\pi$ -related to  $X'$  and  $Y'$  on  $M'$ , then we have the next properties (see [28]):

- (i)  $g(X, Y) = g'(X', Y') \circ \pi$ ;
- (ii)  $h[X, Y]$  is a basic vector field and  $\pi_* h[X, Y] = [X', Y'] \circ \pi$ ;
- (iii)  $h(\nabla_X Y)$  is a basic vector field  $\pi$ -related to  $\nabla'_{X'} Y'$ , where  $\nabla$  and  $\nabla'$  are the Levi-Civita connections on  $M$  and  $M'$ ;
- (iv)  $[E, U] \in \Gamma(\mathcal{V})$ ,  $\forall U \in \Gamma(\mathcal{V})$  and  $\forall E \in \Gamma(TM)$ .

*Remark 6.2.* A semi-Riemannian submersion  $\pi : M \rightarrow M'$  determines, as well as in the Riemannian case (see [15]), two (1,2) tensor field  $T$  and  $A$  on  $M$ , by the formulas:

$$(31) \quad T(E, F) = h\nabla_{vE} vF + v\nabla_{vE} hF$$

and respectively:

$$(32) \quad A(E, F) = v\nabla_{hE} hF + h\nabla_{hE} vF$$

for any  $E, F \in \Gamma(TM)$ . We remark that for  $U, V \in \Gamma(\mathcal{V})$ ,  $T(U, V)$  coincides with the second fundamental form of the immersion of the fibre submanifolds and for  $X, Y \in \Gamma(\mathcal{H})$ ,  $A(X, Y) = \frac{1}{2}v[X, Y]$  characterizes the complete integrability of the horizontal distribution  $\mathcal{H}$ .

It is easy to see that  $T$  and  $A$  satisfy:

$$(33) \quad T(U, V) = T(V, U),$$

$$(34) \quad A(X, Y) = -A(Y, X)$$

$$(35) \quad \nabla_X Y = h\nabla_X Y + A(X, Y),$$

$$(36) \quad \nabla_U V = v\nabla_U V + T(U, V),$$

for any  $X, Y \in \Gamma(\mathcal{H})$  and  $U, V \in \Gamma(\mathcal{V})$ .

*Remark 6.3.* In [25], S. Kobayashi observed the next similarity between a Riemannian submersion and a CR-submanifold of a Kähler manifold: both involve two distributions (the vertical and horizontal distribution), one of them being integrable. Then he introduced the concept of CR-submersion, as a Riemannian submersion from a CR-submanifold to an almost hermitian manifold. Next, we'll consider CR-submersions from paraquaternionic CR-submanifolds of a paraquaternionic Kähler manifold.

**Definition 6.4.** Let  $N$  be a paraquaternionic CR-submanifold of an almost paraquaternionic hermitian manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost hermitian manifold. A semi-Riemannian submersion  $\pi : N \rightarrow M'$  is said to be a *paraquaternionic CR-submersion* if the following conditions are satisfied:

- i.  $\mathcal{V} = \mathcal{D}^\perp$ ;
- ii. For each  $p \in N$ ,  $\pi_* : \mathcal{D}_p \rightarrow T_{\pi(p)}M'$  is an isometry with respect to each complex and product structure of  $\mathcal{D}_p$  and  $T_{\pi(p)}M'$ , where  $T_{\pi(p)}M'$  denotes the tangent space to  $M'$  at  $\pi(p)$ .

**Proposition 6.5.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost paraquaternionic hermitian manifold. If  $\pi : N \rightarrow M'$  is a paraquaternionic CR-submersion, then*

$$(37) \quad h\nabla_X J_\alpha Y = J_\alpha h\nabla_X Y - \epsilon_\alpha [\omega_{\alpha+2}(X)J_{\alpha+1}Y - \omega_{\alpha+1}(X)J_{\alpha+2}Y]$$

$$(38) \quad A(X, J_\alpha Y) = J_\alpha \bar{\nu}B(X, Y)$$

$$(39) \quad \bar{\nu}B(X, J_\alpha Y) = J_\alpha A(X, Y)$$

$$(40) \quad \bar{h}B(X, J_\alpha Y) = J_\alpha \bar{h}B(X, Y)$$

for any  $X, Y \in \Gamma(\mathcal{D})$ , where  $\nabla$  is the Levi-Civita connection on  $N$  and  $\bar{h}, \bar{\nu}$  denote the canonical projections on  $\nu$  and  $\nu^\perp$ , respectively.

*Proof.* From Gauss equations and (35), we have

$$(41) \quad \bar{\nabla}_X Y = h\nabla_X Y + A(X, Y) + \bar{h}B(X, Y) + \bar{\nu}B(X, Y),$$

for any  $X, Y \in \Gamma(\mathcal{D})$ , where  $\bar{\nabla}$  is the Levi-Civita connection on  $M$ .

From (41) we obtain:

$$(42) \quad \begin{aligned} (\bar{\nabla}_X J_\alpha)Y &= h\nabla_X J_\alpha Y + A(X, J_\alpha Y) + \bar{h}B(X, J_\alpha Y) + \bar{\nu}B(X, J_\alpha Y) \\ &\quad - J_\alpha h\nabla_X Y - J_\alpha A(X, Y) - J_\alpha \bar{h}B(X, Y) - J_\alpha \bar{\nu}B(X, Y). \end{aligned}$$

On another hand, from (4) we have:

$$(43) \quad (\bar{\nabla}_X J_\alpha)Y = -\epsilon_\alpha [\omega_{\alpha+2}(X)J_{\alpha+1}Y - \omega_{\alpha+1}(X)J_{\alpha+2}Y] \in \Gamma(\mathcal{D}),$$

for any  $X, Y \in \Gamma(\mathcal{D})$ .

Comparing now (42) and (43) and identifying the components from  $\Gamma(\mathcal{D})$ ,  $\Gamma(\mathcal{D}^\perp)$ ,  $\Gamma(\nu)$  and  $\Gamma(\nu^\perp)$ , we obtain the wanted identities.  $\square$

**Proposition 6.6.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost paraquaternionic hermitian manifold. If  $\pi : N \rightarrow M'$  is a paraquaternionic CR-submersion, then*

$$(44) \quad \bar{\nu}\nabla_U^\perp J_\alpha V = J_\alpha \bar{\nu}\nabla_U V - \epsilon_\alpha [\omega_{\alpha+2}(U)J_{\alpha+1}V - \omega_{\alpha+1}(U)J_{\alpha+2}V]$$

$$(45) \quad J_\alpha T(U, V) = -hA_{J_\alpha V}U$$

$$(46) \quad J_\alpha \bar{v}B(U, V) = -vA_{J_\alpha V}U$$

$$(47) \quad \bar{h}\nabla_U^\perp J_\alpha V = J_\alpha \bar{h}B(U, V)$$

for any  $U, V \in \Gamma(\mathcal{D}^\perp)$ , where  $\nabla$  is the Levi-Civita connection on  $N$  and  $\nabla^\perp$  is the normal connection.

*Proof.* From Gauss equations and (36), we have:

$$(48) \quad \bar{\nabla}_U V = T(U, V) + v\nabla_U V + \bar{h}B(U, V) + \bar{v}B(U, V),$$

for any  $U, V \in \Gamma(\mathcal{D}^\perp)$ , where  $\bar{\nabla}$  is the Levi-Civita connection on  $M$ .

On the other hand, from Weingarten formula, we have:

$$(49) \quad \bar{\nabla}_U J_\alpha V = -hA_{J_\alpha V}U - vA_{J_\alpha V}U + \bar{h}\nabla_U^\perp J_\alpha V + \bar{v}\nabla_U^\perp J_\alpha V,$$

for any  $U, V \in \Gamma(\mathcal{D}^\perp)$ .

From (48) and (49) we deduce:

$$(50) \quad \begin{aligned} (\bar{\nabla}_U J_\alpha)V &= -hA_{J_\alpha V}U - vA_{J_\alpha V}U + \bar{h}\nabla_U^\perp J_\alpha V + \bar{v}\nabla_U^\perp J_\alpha V \\ &\quad - J_\alpha T(U, V) - J_\alpha v\nabla_U V - J_\alpha \bar{h}B(U, V) - J_\alpha \bar{v}B(U, V), \end{aligned}$$

On another hand, from (4) we have:

$$(51) \quad (\bar{\nabla}_U J_\alpha)V = -\epsilon_\alpha[\omega_{\alpha+2}(U)J_{\alpha+1}V - \omega_{\alpha+1}(U)J_{\alpha+2}V] \in \Gamma(\mu^\perp),$$

for any  $U, V \in \Gamma(\mathcal{D}^\perp)$ .

Comparing now (50) and (51) and identifying the components from  $\Gamma(\mathcal{D})$ ,  $\Gamma(\mathcal{D}^\perp)$ ,  $\Gamma(\nu)$  and  $\Gamma(\nu^\perp)$ , we obtain the wanted identities.  $\square$

**Theorem 6.7.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost paraquaternionic hermitian manifold. If  $\pi : N \rightarrow M'$  is a paraquaternionic CR-submersion, then  $(M', \sigma', g')$  is a paraquaternionic Kähler manifold.*

*Proof.* We have from (37):

$$(52) \quad h\nabla_X J_\alpha Y - J_\alpha h\nabla_X Y = -\epsilon_\alpha[\omega_{\alpha+2}(X)J_{\alpha+1}Y - \omega_{\alpha+1}(X)J_{\alpha+2}Y]$$

for any local basic vector fields  $X, Y$  on  $N$ .

We define the local 1-forms  $\omega'_\alpha$  on  $M'$  by:

$$(53) \quad \omega'_\alpha(X') \circ \pi = \omega_\alpha(X)$$

for any local vector field  $X'$  on  $M'$ ,  $\pi$ -related with an horizontal vector field  $X$  on  $N$ .

On the other hand, from the definition of a paraquaternionic CR-submersion, we deduce that for any local bases  $\{J'_1, J'_2, J'_3\}$  of  $\sigma'$  we have a corresponding local basis  $\{J_1, J_2, J_3\}$  of  $\sigma$  such that:

$$(54) \quad \pi_* \circ J_\alpha = J'_\alpha \circ \pi_*, \quad \alpha \in \{1, 2, 3\}.$$

Using (52), (53), (54) and Remark 6.1 we obtain:

$$(55) \quad (\nabla'_X J'_\alpha)Y = -\epsilon_\alpha[\omega'_{\alpha+2}(X')J'_{\alpha+1}Y' - \omega'_{\alpha+1}(X')J'_{\alpha+2}Y']$$

for any local vector fields  $X', Y'$  on  $M'$ ,  $\pi$ -related with two local basic vector fields  $X, Y$  on  $N$ , where  $\nabla'$  is the Levi-Civita connection on  $M'$ . Hence  $(M', \sigma', g')$  is a paraquaternionic Kähler manifold.  $\square$

**Theorem 6.8.** *Let  $N$  be a mixed foliated paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost paraquaternionic hermitian manifold. If  $\pi : N \rightarrow M'$  is a paraquaternionic CR-submersion, then  $N$  is locally a semi-Riemannian product of a paraquaternionic submanifold and a totally real submanifold of  $M$ . In particular, if  $N$  is complete and simply connected then it is a global semi-Riemannian product.*

*Proof.* Because  $N$  is a mixed foliated paraquaternionic CR-submanifold, in particular  $N$  is a mixed geodesic submanifold, and from (23) we obtain that the distribution  $\mathcal{D}^\perp$  is parallel. On another hand,  $\mathcal{D}$  being integrable, we have  $A(X, Y) = v\nabla_X Y = 0$ , for any  $X, Y \in \Gamma(\mathcal{D})$  and therefore the distribution  $\mathcal{D}^\perp$  is also parallel. Hence  $N$  is locally a semi-Riemannian product  $(N_1, g_1) \times (N_2, g_2)$ , where  $N_1$  and  $N_2$  are leaves of  $\mathcal{D}$  and  $\mathcal{D}^\perp$ .

Finally, if  $N$  is complete and simply connected, applying the decomposition theorem for semi-Riemannian manifolds (see [34]) we obtain the last part of the theorem.  $\square$

*Example 6.9.* We remarked in Section 3 that

$$N = \{[u_0, u_1, u_2] \in f_{p,q}^{-1}(0) | q^2|u_0|^2 + p^2|u_1|^2 + p^2|u_2|^2 \neq 0\}$$

is a proper paraquaternionic CR-submanifold of  $P^2\mathbb{B}$ . Moreover, the Lie group  $G = \{e^{jt} | t \in \mathbb{R}\}$  acts freely and isometrically on  $N$ . Using now the paraquaternionic Kähler reduction (see Theorem 5.2 from [33]) we obtain that the manifold  $M' = N/G$  equipped with the submersed metric (i.e the one  $g'$  which makes the projection  $\pi : (N, g) \rightarrow (M', g')$  a semi-Riemannian submersion) is again a paraquaternionic Kähler manifold with respect to the structure  $\sigma'$  induced on  $M'$  from the structure  $\sigma$  by the projection  $\pi$ . Moreover,  $\pi : N \rightarrow N/G$  is a paraquaternionic CR-submersion.

## 7. SOME CURVATURE PROPERTIES OF PARAQUATERNIONIC CR-SUBMERSIONS

Let  $(M, g)$  be a semi-Riemannian manifold. The sectional curvature  $K$  of a 2-plane in  $T_p M$ ,  $p \in M$ , spanned by  $\{X, Y\}$ , is defined by:

$$(56) \quad K(X \wedge Y) = \frac{R(X, Y, X, Y)}{g(X, X)g(Y, Y) - g(X, Y)^2}.$$

It is clear that the above definition makes sense only for non-degenerate planes, i.e. those satisfying  $Q(X \wedge Y) = g(X, X)g(Y, Y) - g(X, Y)^2 \neq 0$ .

If  $\pi : (N, g) \rightarrow (M', g')$  is a semi-Riemannian submersion, then the tensor fields  $A$  and  $T$  defined in the above Section play a fundamental role in expressing the curvatures of the  $(N, g)$ ,  $(M', g')$  and of any fibre  $(\pi^{-1}(p'), \hat{g}_{p'})$ ,  $p' \in M'$ , because we have the following formulas stated in [28]:

$$(57) \quad \begin{aligned} R(U, V, W, W') &= \hat{R}(U, V, W, W') - g(T(U, W), T(V, W')) \\ &\quad + g(T(V, W), T(U, W')), \end{aligned}$$

$$(58) \quad \begin{aligned} R(X, Y, Z, Z') &= R^*(X, Y, Z, Z') - 2g(A(X, Y), A(Z, Z')) \\ &\quad + g(A(Y, Z), A(X, Z')) - g(A(X, Z), A(Y, Z')), \end{aligned}$$

for any  $U, V, W, W' \in \Gamma(\mathcal{V})$  and  $X, Y, Z, Z' \in \Gamma(\mathcal{H})$ , where  $R$  is the Riemannian curvature of  $(N, g)$ ,  $\hat{R}$  is the Riemannian curvature of fibre and  $R^*(X, Y, Z, Z') = g(R^*(Z, Z')Y, X)$ ,  $R^*(\cdot, \cdot)$  being the (1,3)-tensor field on  $\Gamma(\mathcal{H})$  with values in  $\Gamma(\mathcal{H})$

which associates to any  $X, Y, Z \in \Gamma(\mathcal{H})$  and  $p \in N$ , the horizontal lift  $(R^*(X, Y)Z)_p$  of  $R'_{\pi(p)}(\pi_{*p}(X_p), \pi_{*p}(Y_p))\pi_{*p}(Z_p)$ ,  $R'$  denoting the Riemannian curvature of  $(M', g')$ .

**Theorem 7.1.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost paraquaternionic hermitian manifold. If  $\pi : N \rightarrow M'$  is a paraquaternionic CR-submersion, then the sectional curvatures of  $M$  and the fibres are related by:*

$$\begin{aligned} \overline{K}(U \wedge V) = \hat{K}(U \wedge V) & - \epsilon_\alpha \theta_U \theta_V [g(A_{J_\alpha U} U, A_{J_\alpha V} V) - g(A_{J_\alpha V} U, A_{J_\alpha U} V)] \\ (59) \quad & - \epsilon_\alpha \theta_U \theta_V [g(\overline{h}\nabla_U^\perp J_\alpha U, \overline{h}\nabla_V^\perp J_\alpha V) - g(\overline{h}\nabla_V^\perp J_\alpha U, \overline{h}\nabla_U^\perp J_\alpha V)] \end{aligned}$$

for any unit space-like or time-like orthogonally vector fields  $U, V \in \Gamma(\mathcal{D}^\perp)$  and  $\alpha \in \{1, 2, 3\}$ , where  $\theta_U = g(U, U) \in \{-1, 1\}$  and  $\theta_V = g(V, V) \in \{-1, 1\}$ .

*Proof.* From (57) and Gauss equation we have:

$$\begin{aligned} \overline{R}(U, V, W, W') & = \hat{R}(U, V, W, W') - g(T(U, W), T(V, W')) + g(T(V, W), T(U, W')) \\ (60) \quad & - g(B(U, W), B(V, W')) + g(B(V, W), B(U, W')), \end{aligned}$$

for any  $U, V, W, W' \in \Gamma(\mathcal{D}^\perp)$ .

From (56) we see that the sectional curvatures of  $M$  and the fibres for a 2-plane spanned by two unit space-like or time-like orthogonally vector fields  $U, V$  are:

$$(61) \quad \overline{K}(U \wedge V) = \theta_U \theta_V \overline{R}(U, V, U, V)$$

and, respectively,

$$(62) \quad \hat{K}(U \wedge V) = \theta_U \theta_V \hat{R}(U, V, U, V),$$

where  $\theta_U = g(U, U) \in \{-1, 1\}$  and  $\theta_V = g(V, V) \in \{-1, 1\}$ .

From (60), (61) and (62), using (33), we obtain:

$$\begin{aligned} \overline{K}(U \wedge V) & = \hat{K}(U \wedge V) - \theta_U \theta_V [g(T(U, U), T(V, V)) - g(T(U, V), T(U, V)) \\ (63) \quad & + g(B(U, U), B(V, V)) - g(B(U, V), B(U, V))] \end{aligned}$$

for any unit space-like or time-like orthogonally vector fields  $U, V \in \Gamma(\mathcal{D}^\perp)$ .

Finally, using (45), (46) and (47) in (63), we obtain (59).  $\square$

**Theorem 7.2.** *Let  $N$  be a paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost paraquaternionic hermitian manifold. If  $\pi : N \rightarrow M'$  is a paraquaternionic CR-submersion, then for any unit space-like or time-like horizontal vector field  $X$  one has:*

$$(64) \quad \overline{H}_\alpha(X) = H'_\alpha(\pi_* X) - 4g(\overline{v}B(X, X), \overline{v}B(X, X)) + 2g(\overline{h}B(X, X), \overline{h}B(X, X)),$$

for  $\alpha \in \{1, 2, 3\}$ , where  $\overline{H}_\alpha$  and  $H'_\alpha$  are the holomorphic sectional curvatures of  $M$  and  $M'$ , defined by  $\overline{H}_\alpha(X) = \overline{K}(X \wedge J_\alpha X)$  and  $H'_\alpha(X) = K'(X \wedge J_\alpha X)$ , respectively.

*Proof.* From (58) and Gauss equation we have:

$$\begin{aligned} \overline{R}(X, Y, Z, Z') & = R^*(X, Y, Z, Z') - 2g(A(X, Y), A(Z, Z')) \\ & + g(A(Y, Z), A(X, Z')) - g(A(X, Z), A(Y, Z')) \\ (65) \quad & + g(B(Y, Z), B(X, Z')) - g(B(Y, Z'), B(X, Z)), \end{aligned}$$

for any  $X, Y, Z, Z' \in \Gamma(\mathcal{D})$ .

From (3) and (56) we see that the holomorphic sectional curvature of  $M$  is:

$$(66) \quad \overline{H}_\alpha(X) = \epsilon_\alpha \overline{R}(X, J_\alpha X, X, J_\alpha X)$$



for any unit space-like or time-like vector field  $X \in \Gamma(\mathcal{D})$ .

On the other hand we can easily see that:

$$(67) \quad H'_\alpha(\pi_*X) = \epsilon_\alpha R^*(X, J_\alpha X, X, J_\alpha X)$$

for any unit space-like or time-like vector field  $X \in \Gamma(\mathcal{D})$ .

From (65), (66) and (67), using (34), we obtain:

$$(68) \quad \begin{aligned} \overline{H}_\alpha(X) &= H'_\alpha(\pi_*X) - 3\epsilon_\alpha g(A(X, J_\alpha X), A(X, J_\alpha X)) \\ &+ \epsilon_\alpha [g(B(X, J_\alpha X), B(X, J_\alpha X)) - g(B(X, X), B(J_\alpha X, J_\alpha X))]. \end{aligned}$$

From (34) and (39) one has:

$$0 = A(X, X) = -\epsilon_\alpha J_\alpha \overline{v}B(X, J_\alpha X)$$

and so we derive:

$$(69) \quad \overline{v}B(X, J_\alpha X) = 0.$$

Similarly, from (34), (38) and (39) we derive:

$$(70) \quad \overline{v}B(J_\alpha X, J_\alpha X) = \epsilon_\alpha \overline{v}B(X, X).$$

and from (40), using the symmetry of  $B$ , we obtain:

$$(71) \quad \overline{h}B(J_\alpha X, J_\alpha X) = -\epsilon_\alpha \overline{h}B(X, X).$$

Finally, using (69), (70) and (71) in (68) we obtain (64).  $\square$

**Corollary 7.3.** *Let  $N$  be a totally geodesic paraquaternionic CR-submanifold of a paraquaternionic Kähler manifold  $(M, \sigma, g)$  and  $(M', \sigma', g')$  be an almost paraquaternionic hermitian manifold. If  $\pi : N \rightarrow M'$  is a paraquaternionic CR-submersion one has:*

$$(72) \quad \overline{H}_\alpha(X) = H'_\alpha(\pi_*X),$$

for any unit space-like or time-like horizontal vector field  $X$ .

#### ACKNOWLEDGMENTS

We would like to express our sincere thanks to Professor Liviu Ornea for valuable discussions concerning this subject. S. Ianuș and G.E. Vilcu are supported by a CNCSIS PN2-IDEI grant, no. 525/2009. S. Marchiafava activity was done under the programs of G.N.S.A.G.A. of C.N.R.-Indam and "Riemannian metrics and differentiable structures" of MIUR (Italy).

#### REFERENCES

- [1] A. Al-Aqeel, A. Bejancu, *On normal semi-invariant submanifolds of paraquaternionic Kähler manifolds*, Toyama Math. J. 30 (2007), 63-75.
- [2] D. Alekseevsky, V. Cortés, *The twistor spaces of a para-quaternionic Kähler manifold*, Osaka J. Math., 45(1) (2008), 215-251.
- [3] D. Alekseevsky, Y. Kamishima, *Quaternionic and para-quaternionic CR structure on  $(4n+3)$ -dimensional manifolds*, Cent. Eur. J. Math., 2(5) (2004), 732-753.
- [4] D.V. Alekseevsky, S. Marchiafava, *Almost complex submanifolds of a quaternionic Kähler manifold*, Steps in Differential Geometry, Inst. Math. Inf. Debrecen 2001, 23-38.
- [5] M. Barros, B.-Y. Chen and F. Urbano, *Quaternion CR-submanifolds of quaternion manifolds*, Kodai Math. J., 4 (1981), 399-417.
- [6] A. Bejancu, *CR submanifolds of a Kaehler manifold I*, Proc. Am. Math. Soc., 69 (1978), 135-142.
- [7] A. Bejancu, *Geometry of CR-submanifolds*, Mathematics and its Applications (East European Series) 23, D. Reidel Publishing Co., Dordrecht, 1986.

- [8] A. Bejancu, *Normal semi-invariant submanifolds of paraquaternionic Kähler manifolds*, Kuwait J. Sci. Eng., 33(2) 2006, 33-46.
- [9] A. Bejancu and H. R. Farran, *Foliations and geometric structures*, Mathematics and Its Applications, Springer, 2006.
- [10] N. Blažič, *Para-quaternionic projective spaces and pseudo Riemannian geometry*, Publ. Inst. Math, 60 (1996), 101-107.
- [11] N. Blažič, S. Vukmirović, *Solutions of Yang-Mills equations on generalized Hopf bundles*, J. Geom. Phys. 41(1-2) (2002), 57-64.
- [12] B.-Y. Chen, *Geometry of submanifolds and its applications*, Science University of Tokyo, Tokyo, 1981.
- [13] B.-Y. Chen, *Cohomology of CR-submanifolds*, Ann. Fac. Sci. Toulouse 3, 167-172 (1981).
- [14] A.S. Dancer, H.R. Jørgensen, A.F. Swann, *Metric geometries over the split quaternions*, Rend. Sem. Mat. Univ. Pol. Torino, 63(2) (2005), 119-139.
- [15] M. Falcitelli, S. Ianuș, A.M. Pastore, *Riemannian submersions and related topics*, River Edge, NJ: World Scientific, 2004.
- [16] K. Galicki, B. Lawson, *Quaternionic reduction and quaternionic orbifolds*, Math. Ann. 282(1) (1988), 1-21.
- [17] E. Garcia-Rio, Y. Matsushita, R. Vasquez-Lorenzo, *Paraquaternionic Kähler manifolds*, Rocky Mt. J. Math., 31 (2001), 237-260.
- [18] S. Ianuș, A.M. Ionescu, G.E. Vilcu, *Foliations on quaternion CR-submanifolds*, Houston J. Math., 34(3) (2008), 739-751.
- [19] S. Ianuș, R. Mazzocco, G.E. Vilcu, *Real lightlike hypersurfaces of paraquaternionic Kähler manifolds*, Mediterr. J. Math., 3 (2006), 581-592.
- [20] S. Ianuș, G.E. Vilcu, *Hypersurfaces of paraquaternionic space forms*, J. Gen. Lie Theory Appl. 2(3) (2008), 175-179.
- [21] S. Ianuș, G.E. Vilcu, *Some constructions of almost para-hyperhermitian structures on manifolds and tangent bundles*, Int. J. Geom. Methods Mod. Phys., 5(6) (2008), 893-903.
- [22] A.M. Ionescu, G.E. Vilcu, *A note on paraquaternionic manifolds*, Missouri J. Math. Sci. 19(3) (2007), 213-218.
- [23] S. Ivanov, S. Zamkovoy, *Para-hermitian and para-quaternionic manifolds*, Differ. Geom. Appl., 23 (2005), 205-234.
- [24] S. Ivanov, I. Minchev, S. Zamkovoy, *Twistor and Reflector Spaces of Almost Para-Quaternionic Manifolds*, to appear in *Handbook of pseudo-Riemannian Geometry and Supersymmetry*, IRMA Lectures in Mathematics and Theoretical Physics.
- [25] S. Kobayashi, *Submersions of CR submanifolds*, Tôhoku Math. J. II. Ser. 39 (1987), 95-100.
- [26] P. Libermann, *Sur le problème d'équivalence de certaines structures infinitésimales*, Ann. Mat. Pura Appl., 36 (1954), 27-120.
- [27] S. Marchiafava, *Submanifolds of (para)-quaternionic Kähler manifolds*, Note Mat., 28 (2008), suppl. no. 1, 295-316.
- [28] B. O'Neill, *Fundamental equations of submersions*, Michigan Math. J. 39 (1966), 459-464.
- [29] B. O'Neill, *Semi-Riemannian geometry with applications to relativity*, Academic Press, New York-London, 1983.
- [30] L. Ornea, *CR-submanifolds. A class of examples*, Rev. Roum. Math. Pures Appl. 51(1) (2006), 77-85.
- [31] M. Vaccaro, *Kähler and para-Kähler submanifolds of a para-quaternionic Kähler manifold*, ph. d. thesis, Università degli Studi di Roma II "Tor Vergata", 2007.
- [32] G.E. Vilcu, *Submanifolds of an Almost Paraquaternionic Kähler Product Manifold*, Int. Math. Forum 2(15) (2007), 735-746.
- [33] S. Vukmirović, *Paraquaternionic reduction*, math.DG/0304424.
- [34] H. Wu, *On the de Rham decomposition theorem*, Illinois J. Math. 8 (1964), 291-311.
- [35] K. Yano, M. Kon, *CR submanifolds of Kählerian and Sasakian manifolds*, Progress in Mathematics 30, Birkhäuser, 1983.
- [36] S. Zamkovoy, *Geometry of paraquaternionic Kähler manifolds with torsion*, J. Geom. Phys., 57(1) (2006), 69-87.

Stere Ianuș

*University of Bucharest, Department of Mathematics,  
Str. Academiei, Nr. 14, Bucharest 70109, Romania*

e-mail: ianus@gta.math.unibuc.ro

Stefano Marchiafava

*Università di Roma "La Sapienza", Dipartimento di Matematica,  
P.le Aldo Moro N. 2, 00185 Rome, Italy*

e-mail: marchiafava@mat.uniroma1.it

Gabriel Eduard Țilcu<sup>1,2</sup>

<sup>1</sup>*University of Bucharest, Faculty of Mathematics and Computer Science,  
Research Center in Geometry, Topology and Algebra,  
Str. Academiei, Nr. 14, Sector 1, Bucharest 70109, Romania*

<sup>2</sup>*Petroleum-Gas University of Ploiești,  
Department of Mathematics and Computer Science,  
Bulevardul București, Nr. 39, Ploiești 100680, Romania*  
e-mail: gvilcu@mail.upg-ploiesti.ro