

Dwarf Novae in the Shortest Orbital Period Regime: II. WZ Sge Stars as the Missing Population near the Period Minimum

Makoto UEMURA¹, Taichi KATO², Daisaku NOGAMI³, and Takashi OHSUGI¹

¹*Astrophysical Science Center, Hiroshima University, Kagamiyama 1-3-1,
Higashi-Hiroshima 739-8526
uemuram@hiroshima-u.ac.jp*

²*Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502*

³*Kwasan Observatory, Kyoto University, Yamashina-ku, Kyoto 607-8471*

(Received 2010 January 4; accepted 2010 March 3)

Abstract

WZ Sge-type dwarf novae are characterized by long recurrence times of outbursts (~ 10 yr) and short orbital periods ($\lesssim 85$ min). A significant part of WZ Sge stars may remain undiscovered because of low outburst activity. Recently, the observed orbital period distribution of cataclysmic variables (CVs) has changed partly because outbursts of new WZ Sge stars have been discovered routinely. Hence, the estimation of the intrinsic population of WZ Sge stars is important for the study of the population and evolution of CVs. In this paper, we present a Bayesian approach to estimate the intrinsic period distribution of dwarf novae from observed samples. In this Bayesian model, we assumed a simple relationship between the recurrence time and the orbital period which is consistent with observations of WZ Sge stars and other dwarf novae. As a result, the minimum orbital period was estimated to be ~ 70 min. The population of WZ Sge stars exhibited a spike-like feature at the shortest period regime in the orbital period distribution. These features are consistent with the orbital period distribution previously predicted by population synthesis studies. We propose that WZ Sge stars and CVs with a low mass-transfer rate are excellent candidates for the missing population predicted by the evolution theory of CVs.

Key words: stars: novae, cataclysmic variables—stars: evolution

1. Introduction

Cataclysmic variables (CVs) are close binary systems containing a white dwarf and a Roche-lobe filling normal star. They have several subclasses, such as, dwarf novae (DNe), nova-like variables, novae, and magnetic CVs (Warner 1995). CVs are considered to be one of the final stages of the evolution of low mass binaries. The formation and evolution of low mass binaries can be explored through the orbital period (P_{orb}) distribution of CVs (for a review, see King 1988).

Stable mass-transfer processes in CVs are driven by angular momentum removal from the binaries. In short period CVs having $P_{\text{orb}} \lesssim 3$ hr, the driving mechanism is considered to be angular momentum removal associated with gravitational radiation (Paczynski 1981; Rappaport et al. 1982). A CV evolves toward a short- P_{orb} region by losing angular momentum. This phase is terminated when the core of the secondary star becomes degenerate. Then, the degeneracy pressure changes the mass–radius relationship of the secondary star. As a result, the CV evolves toward a long- P_{orb} regime after passing through the minimum P_{orb} (P_{min}).

The P_{orb} distribution near P_{min} has received attention, since the observed P_{orb} distribution has different characteristics from that expected from theoretical studies. It is well known that P_{min} is ~ 80 min in the observed CV population (Paczynski 1981; Rappaport et al. 1982).

However, the theoretically predicted P_{min} is 60–70 min, significantly shorter than the observed value (e.g. Kolb 1993). This is the so-called “period minimum problem”. According to population synthesis studies, most CVs have already passed through P_{min} . It has been proposed that the evolution time-scale is long for an evolutionary-advanced system with a low mass secondary star, in other words, the mass-transfer rate (\dot{M}) from the secondary star is low (e.g. Kolb 1993; Howell et al. 1997). As a result, a spike-like feature is expected to appear near P_{min} in the P_{orb} distribution due to the accumulation of systems. However, the observed distribution is rather flat. This is the so-called “period spike problem” (Kolb, Baraffe 1999; Renvoizé et al. 2002).

Thus, it is known that there is a “missing” population in the observed CVs with a P_{orb} range of 70–80 min when compared with the predicted population. Littlefair et al. (2008) have recently reported three CVs having a brown dwarf secondary, which are candidates for the missing population (Littlefair et al. 2006; Littlefair et al. 2007). This discovery indicates that CVs can actually reach the post- P_{min} regime as predicted by the theories, while the number of such systems is still not properly known.

It has been suggested that the problems can be reconciled by another mechanism of angular momentum removal in addition to gravitational radiation (Rezzolla et al. 2001), such as the effects of magnetic stellar wind braking (King et al. 2002), the circumbinary disk (Willems

et al. 2005), and a combination of magnetic propeller and accretion disk resonances (Matthews et al. 2006).

On the other hand, it is possible that the missing population exists, but remains undiscovered. According to the disk instability model for DNe, the recurrence time of superoutbursts (supercycle; T_s) is longer in a system with a lower \dot{M} (Osaki 1995). WZ Sge-type DNe are known to have long T_s ($\gtrsim 10$ yr), which indicate a low \dot{M} (Howell et al. 1995; Kato et al. 2001). It has been proposed that several WZ Sge stars, or CVs with a low \dot{M} have remained undiscovered because of their low activity (Meyer-Hofmeister et al. 1998). Since WZ Sge stars are found concentrating in a short P_{orb} regime, these objects are a candidate for the missing population (Patterson 1998). However, their intrinsic contribution to the whole CV population has been poorly understood because their T_s were uncertain and their quiescent luminosity is low. Gänsicke et al. (2009) have recently reported that the CV sample from the Sloan Digital Sky Survey (SDSS) database shows a period spike feature in the P_{orb} range of 80–86 min. The spike feature is a consequence of deep images and a homogeneously selected CV sample obtained by SDSS, which could detect intrinsically faint CVs with a low \dot{M} . Recently, in addition, T_s of several WZ Sge stars have been well determined owing to long monitoring of DNe by amateur observers (e.g. Waagen et al. 2007)

In this paper, we investigate the contribution of long- T_s DNe to the whole population of CVs by estimating their intrinsic population. In § 2, we demonstrate that the whole population of CVs has changed significantly due to a recent increase of the number of short P_{orb} CVs. In § 3, we estimate the population of DNe by considering the detection probability of superoutbursts using Bayesian analysis. We discuss the implications of our results to the study of CV evolution in § 5. In the final section, we summarize our findings.

2. Recent change in the observed P_{orb} distribution in CVs

Both the period minimum and spike problems arise from the discrepancy between the P_{orb} distribution predicted by the population synthesis studies and the observed one. In previous studies, the predicted distribution was occasionally compared with the distribution of all known CVs. The left panel of figure 1 shows the P_{orb} distribution of all known CVs with $P_{\text{orb}} < 130$ min discovered before 2003 (black) and 2008 (white). The samples were selected from Ritter & Kolb catalog¹: 184 sources in RKcat 7.0 for the 2003 sample and 312 sources in RKat 7.9 for the 2008 sample. Both magnetic and non-magnetic CVs are included in the samples. As can be seen in the figure, the fraction of short- P_{orb} systems in the 2008 sample is apparently larger than that in the 2003 sample. We applied the Kolmogorov–Smirnov (KS) test to evaluate whether the two samples differ significantly. We calculated KS probabilities, P_{KS} , from the 2003 and 2008 sam-




Fig. 1. P_{orb} distribution of CVs. Left panel: Sources in the 2003 sample (RKcat7.0, black histogram) and 2008 sample (RKcat7.9, white histogram). Right panel: Normalized cumulative distribution functions (CDF) of the 2003 (solid line) and the new samples discovered between 2003 and 2008 (dotted line).

Table 1. Results of Kolmogorov–Smirnov tests between the 2003 (RKcat ver. 7.0) and 2008 (RKcat ver. 7.9) samples.

sample	P_{KS}^*	N_1^\dagger	N_2^\dagger
only new objects	0.07	184	128
all objects	0.73	184	312

* P_{KS} : Kolmogorov–Smirnov probability.

† N_1, N_2 : the number of objects in the 2003 and 2008 samples.

ples, and also from the 2003 sample and 128 new sources discovered between 2003 and 2008. The results are listed in table 1. The test for the 2003 samples and the sample of new sources gives a low probability of 0.07, and based on this we conclude that the distributions of the samples differ significantly. For the test with the full 2008 sample, the probability is also low at 0.73 and hence it must be concluded that the 2003 and 2008 samples differ overall. The difference between the samples is also clear in the normalized cumulative distribution functions (CDF) of the 2003 sample and the new sources in the 2008 sample, as shown in the right panel of figure 1. These results demonstrate that the P_{orb} distribution of all known CVs has significantly changed recently.

This change is mainly caused by the outburst detection of new DNe and large spectroscopic surveys. First, outbursts of new and known WZ Sge stars have routinely been discovered recently and hence the number of known WZ Sge stars has rapidly increased (Shafter et al. 2007; Waagen et al. 2007; Templeton et al. 2007; Uemura et al. 2008; Soejima et al. 2009; Shugarov et al. 2008; Zharikov et al. 2008). The dramatic increase of newly recognized SU UMA-type and WZ Sge-type DNe is also apparent in tables in Kato et al. (2010), in which 14 WZ Sge-type DNe (out of a total of 32 systems; Ritter, Kolb 2003; Kato et al. 2010) was recognized after 2005. This increase of the outburst detection is attributed to the development of automatic sky monitor systems (e.g. Pojmanski 2002; Burd et al. 2005; and most recently, Drake et al. 2009), and improvements in the techniques and instruments of amateur observers (notably by K. Itagaki). The devel-

¹ (<http://www.mpa-garching.mpg.de/RKcat/>)

opment of online networks for transient phenomena has also played an important role in circulating information on WZ Sge stars, and enabled timely observation essential for identifying their nature (Kato et al. 2004).

Second, large spectroscopic surveys have also played an important role in increasing the number of CVs recently, for example, SDSS and the Hamburg Quasar Survey (HQS) (Szkody et al. 2002; Bade et al. 1998). Those large surveys allow us to investigate the P_{orb} distribution without complicated observation bias which should be considered in a sample containing all CVs discovered by various detection methods. The P_{orb} distribution of all known CVs would be same as before if the distribution of those new CVs is same as that of the previously known CVs. However, the distribution of the SDSS CVs has a period spike feature, and is different from the previously known distributions (Gänsicke et al. 2009).

Thus, it is evident that the period minimum and spike problems should be discussed not with the sample containing all known CVs, but with the sample in which observation bias for the P_{orb} distribution can be simply evaluated. In the next section, we collect outburst-selected DN samples, and estimate the intrinsic population of DNe without the complicated bias for the P_{orb} distribution.

3. Bayesian estimation of the intrinsic population of DNe

As well as new WZ Sge stars, outbursts of known WZ Sge stars have been observed routinely. Thereby, we can now estimate T_s more accurately for them. In Uemura et al. (2010) (hereafter, Paper I), we revised T_s of several DNe and found that WZ Sge stars form a major group in $P_{\text{orb}} \lesssim 86$ min. This implies that a DN enters the WZ Sge regime on an evolutionary course at $P_{\text{orb}} \sim 86$ min as \dot{M} decreases. Using T_s and P_{orb} from Paper I, we consider a statistical experiment to estimate the intrinsic P_{orb} distribution of DNe from observed systems based on a simple model for T_s . Using Bayesian analysis, we calculate the posterior distributions of parameters that characterize the P_{orb} distribution (e.g. Trotta 2008).

3.1. Overview of analysis

We first present an outline of the analysis. The models and samples are detailed in subsequent sections.

We consider a DN sample, $\mathbf{x} = \{P_{\text{orb},i}\}$, whose super-outbursts were detected at least once in a certain period of time. The intrinsic P_{orb} population of DNe can be reproduced from this outburst-selected sample if T_s can be described with a function of P_{orb} . The probability density function (PDF) of observed DNe, $Q(P_{\text{orb}})$, is given by the intrinsic period distribution, $I(P_{\text{orb}})$, and the selection effects arising from the observation process, $D(P_{\text{orb}})$, in the form $Q = D \cdot I/A_Q$, where A_Q is a normalization factor.

With a parameter set, \mathbf{y} , that characterizes I , the likelihood of \mathbf{x} , can be defined by

$$L(\mathbf{y} | \mathbf{x}) = Q(\mathbf{x} | \mathbf{y}) = \prod_i Q(P_{\text{orb},i} | \mathbf{y}). \quad (1)$$

Posterior distributions of parameters, $P(\mathbf{y} | \mathbf{x})$, are estimated with Bayes' theorem, as follows:

$$P(\mathbf{y} | \mathbf{x}) = \frac{L(\mathbf{y} | \mathbf{x})P(\mathbf{y})}{\sum L(\mathbf{y} | \mathbf{x})P(\mathbf{y})} = \frac{Q(\mathbf{x} | \mathbf{y})P(\mathbf{y})}{\sum Q(\mathbf{x} | \mathbf{y})P(\mathbf{y})}. \quad (2)$$

In our analysis, we used flat distributions for the prior probabilities of parameters: $P(\mathbf{y}) = \text{const.}$

3.2. Sample

Samples were selected from SU UMa-type DNe in Rkcat7.9. The ‘‘ASAS’’ sample consists of systems in which superoutbursts were recorded at least once in the ASAS-3 database between 1 Jan 2003 and 31 Dec 2007 (Pojmanski 2002). A superoutburst is defined as an outburst lasting for ≥ 5 d with a small fading rate ($\sim 0.1 \text{ mag d}^{-1}$). P_{orb} of our sample was limited to ≤ 130 min because our simple model of D cannot reproduce the features of a period gap of $2 \text{ hr} \lesssim P_{\text{orb}} \lesssim 3 \text{ hr}$ (for details, see the next section). We made another sample, called the ‘‘mixed’’ sample, using the procedure for data selection as for the ASAS sample, but using both the VSNET and ASAS-3 databases for searching superoutbursts (Kato et al. 2004).

We excluded two minor groups of objects in the shortest P_{orb} regime. The first group is ER UMa stars (ER UMa, IX Dra, RZ LMi, V1159 Ori, and DI UMa), which are characterized by atypically short T_s (20–50 d) despite a short P_{orb} (Kato, Kunjaya 1995). The other group was identified in Paper I for the first time and called ‘‘Group X’’. The members of this group have rather average T_s (~ 400 d) despite a short P_{orb} . These two groups were excluded from our sample because they may lead to an overestimate of the intrinsic population of short- P_{orb} systems having long T_s .

We obtained 42 and 146 sources for the ASAS and mixed samples. The ASAS sample is expected to be an ideal sample without complicated selection effects since the superoutbursts were detected through a single observation process defined by the ASAS system. The mixed sample may be affected by complicated selection effects. We calculated the KS probability of both samples. The obtained probability is unexpectedly high, 0.98, which means that there is no significant difference between those two samples. This implies that our method of data selection has the advantage of reducing selection bias. The P_{orb} distributions of these two samples are shown in figure 2.

3.3. Model of the P_{orb} distribution

The intrinsic P_{orb} distribution, I , must be characterized by at least two parameters, namely, one for the degree of accumulation near P_{min} and the other for the real P_{min} . We express I by the following formulae:

$$I(p) = \begin{cases} p^{-\alpha} e^{-\alpha/p} / A_I & (p \geq 1) \\ 0 & (p < 1) \end{cases} \quad (3)$$

$$p = P_{\text{orb}} - P_{\text{min}} (\text{min}), \quad (4)$$

where α is a parameter for the degree of the accumulation near P_{min} . These are the parameters of our Bayes model, that is, $\mathbf{y} = \{\alpha, P_{\text{min}}\}$. A large α yields a distribution with




Fig. 2. Distribution of P_{orb} of the DN samples. The black and white histograms indicate the ASAS and mixed samples, respectively. The dashed line indicates the best model PDF derived from our analysis (for details, see the text).

a prominent spike near P_{min} . The distribution becomes flat in the case of $\alpha = 0$. For a negative α , the distribution has a peak in the long- P_{orb} region. The function I has an extremum at $P_{\text{orb}} = P_{\text{min}} + 1$. A_I is a normalization factor. While this formula of I itself has no physical meaning, it can approximate the profile of the P_{orb} distribution below the period gap predicted by population synthesis studies, with high α ($\alpha \sim 1$) (e.g. Barker, Kolb 2003). This model should be used only below the period gap ($2 \text{ hr} \lesssim P_{\text{orb}} \lesssim 3 \text{ hr}$; Spruit, Ritter 1983), since it cannot reproduce the gap structure.

We considered a simple model for the selection effect, D . First, we considered that all the systems are pre- P_{min} CVs and neglected contributions from post- P_{min} CVs. The contribution of post- P_{min} CVs to our analysis is discussed in § 4.2. Second, we used a proposed model for the relationship between T_s and P_{orb} based on observations. In Paper I, we showed that most systems are ordinary SU UMa stars in a P_{orb} range of $\gtrsim 86$ min. The average T_s is 470 d in $86 \text{ min} \leq P_{\text{orb}} \leq 95$ min. On the other hand, in the shortest P_{orb} regime of $\lesssim 86$ min, the major group is WZ Sge stars. Based on these results in Paper I, we assumed the following for our model, D : i) T_s is constant at 470 d in $P_{\text{orb}} \geq P_{\text{crit}}$. ii) T_s becomes longer with shorter P_{orb} in $P_{\text{orb}} < P_{\text{crit}}$. iii) We set $P_{\text{crit}} = 86$ min.

The outburst detection probability, D_{outb} , is inversely proportional to T_s . Taking D_{outb} at $T_s = 470$ d to be unity, the above assumptions can be expressed as follows:

$$D_{\text{outb}}(P_{\text{orb}}) = \begin{cases} 1 & (P_{\text{orb}} \geq P_{\text{crit}}) \\ \left(\frac{P_{\text{orb}} - P_{\text{min}}}{P_{\text{crit}} - P_{\text{min}}} \right)^n & (P_{\text{orb}} < P_{\text{crit}}) \end{cases} \quad (5)$$

The parameter n can be estimated from the observed T_s listed in table 5 of Paper I. Using GW Lib ($P_{\text{orb}} = 76.7808$ min and $T_s = 8800$ d) and SW UMa ($P_{\text{orb}} = 81.8136$ min and $T_s = 1200$ d), for example, we obtained $n = 2.5$ and $P_{\text{min}} = 72.6$ min. We show D_{outb} calculated from the observed T_s of our sample with $P_{\text{min}} = 72.6$ min in figure 3. The dashed lines in the figure denote the models,




Fig. 3. Outburst detection probability, D_{outb} , as a function of $P_{\text{orb}} - P_{\text{min}}$. P_{min} is assumed to be 72.6 min in the figure (for more detail, see the text). The filled circles indicate SU UMa stars whose T_s is known. The dashed lines indicate the models with various n . Two objects, GW Lib and SW UMa are labeled. They are used to estimate a lower limit of n .

D_{outb} , with various n . This figure shows that the estimation of the parameters with GW Lib and SW UMa yields a lower-limit of n for our sample. As discussed in § 4.1, a larger n leads to a larger number of short- P_{orb} DNe. In the following analysis, we used $n = 2.0$, which gives a firm lower-limit for n , as can be seen from figure 3. n can be smaller than 2.0 only when $P_{\text{min}} > 72.6$ min. We check whether our estimation of n is consistent with the results of the analysis in § 4.1.

Finally, we considered the relationship between the absolute magnitude, M_V , at supermaximum and P_{orb} . Since the ASAS sample is a magnitude-limited sample, the selection bias caused by the P_{orb} dependence of M_V should be corrected. Under an assumption of a uniform space distribution of DNe, the number of detectable systems, D_{mag} , can be written as a function of M_V as

$$D_{\text{mag}}(P_{\text{orb}}) \propto 10^{-0.6M_V(P_{\text{orb}})}. \quad (6)$$

Harrison et al. (2004) reported the relationship between M_V at normal outburst maxima and P_{orb} . In addition to their relationship, Pretorius et al. (2007) proposed that WZ Sge stars have a constant M_V , independent of P_{orb} . We follow their models for M_V at the supermaxima of ordinary SU UMa stars ($M_{V,\text{SU}}$) and WZ Sge stars ($M_{V,\text{WZ}}$), that is:

$$\begin{aligned} M_{V,\text{SU}}(P_{\text{orb}}) &= C_1 - 0.383P_{\text{orb}} & (P_{\text{orb}} \geq P_{\text{crit}}) \\ M_{V,\text{WZ}}(P_{\text{orb}}) &= C_2 & (P_{\text{orb}} < P_{\text{crit}}) \end{aligned}, \quad (7)$$

where C_2 is a constant determined by $M_{V,\text{SU}}(P_{\text{crit}}) = M_{V,\text{WZ}}(P_{\text{crit}})$. The unit of P_{orb} is hour. The constant C_1 is not important in our analysis.

Finally, the model of the P_{orb} distribution of DNe is expressed by the PDF, Q , using the two parameters as follows:

$$Q(a, P_{\text{min}}) = D_{\text{mag}} \cdot D_{\text{outb}} \cdot I(a, P_{\text{min}}) / A_Q(a, P_{\text{min}}). \quad (8)$$




Fig. 4. Posterior probability distributions of α and P_{\min} . The solid and dotted lines were calculated from the ASAS and mixed samples, respectively. The medians and 68.3 % confidence levels of the parameters are also indicated.

3.4. Bayesian analysis and results

We calculated the likelihood of the sample using equation (1). We produced posterior probability distributions of α and P_{\min} using the Markov Chain Monte Carlo method (MCMC) (Metropolis 1953). The calculation procedure of MCMC is as follows: At the n -th step of MCMC, we obtain the likelihood, $L_n = L(a_n, P_{\min, n} | \mathbf{x})$. We then randomly move to another point in the parameter space of α and P_{\min} by adding random values drawn from a Gaussian distribution which has zero mean and dispersion chosen to efficiently sample the likelihood surface. As a result, we obtain L_{n+1} . We count the parameters of the $(n+1)$ -th step if $L_{n+1} > L_n$ or L_{n+1}/L_n is larger than a uniform random number between 0 and 1. Otherwise, the n -th parameters are substituted for the $(n+1)$ -th ones. After discarding the first 100 steps, we sample every 100 steps until the size of the sample reaches 10^5 . This cycle forms a set of samples for α and P_{\min} . We obtained ten sets of samples with different initial values of α and P_{\min} . We confirmed that the obtained ten samples had the same P_{orb} distribution. We merged them and finally obtained the posterior distributions.

Figure 4 shows the posterior distributions of α (left panel) and P_{\min} (right panel). Each panel contains both the results estimated from the ASAS and mixed samples. The median and 68.3% confidence level of α and P_{\min} are summarized in table 2 and are also indicated in the figure. As expected from the KS test in § 3.2, the parameters estimated from the ASAS sample agree with those from the mixed sample within the errors. The best model of Q for the ASAS sample is indicated by the dashed line in figure 2. The intrinsic P_{orb} distribution, I , is shown in figure 5.

As shown in table 2, α was estimated to be high, namely, the intrinsic P_{orb} distribution, I , has a spike near P_{\min} , as can be seen in figure 5. Short- P_{orb} systems in $P_{\text{orb}} < P_{\text{crit}} (= 86 \text{ min})$ account for 59 % of the total number of samples between $P_{\min} < P_{\text{orb}} < 130 \text{ min}$. The estimated P_{\min} is about 70 min, significantly shorter than the observed P_{\min} of about 78 min.

The estimated P_{orb} distribution, I , is analogous to those expected from population synthesis studies (e.g. Kolb 1993), in terms of the accumulation of systems near P_{\min} and the short P_{\min} of $\sim 70 \text{ min}$. We suggest that i) DNe

Table 2. Best parameters derived from Bayesian analysis.

Sample	n	P_{crit} (min)	M_V	α	P_{\min} (min)
ASAS	2.0	86.0	eq.(7)	$0.88^{+0.30}_{-0.33}$	$70.5^{+2.6}_{-4.6}$
all RKcat	2.0	86.0	eq.(7)	$0.86^{+0.15}_{-0.16}$	$71.1^{+1.2}_{-1.7}$
ASAS	1.0	86.0	eq.(7)	$0.61^{+0.29}_{-0.32}$	$71.3^{+1.9}_{-4.0}$
ASAS	3.0	86.0	eq.(7)	$1.07^{+0.34}_{-0.37}$	$66.2^{+3.8}_{-6.3}$
ASAS	2.0	82.0	eq.(7)	$0.60^{+0.27}_{-0.32}$	$71.3^{+2.5}_{-4.5}$
ASAS	2.0	90.0	eq.(7)	$1.15^{+0.32}_{-0.36}$	$69.5^{+2.6}_{-4.7}$
ASAS	2.0	86.0	const.	$0.64^{+0.33}_{-0.38}$	$68.6^{+3.0}_{-5.3}$
ASAS	0.2	86.0	const.	$0.10^{+0.25}_{-0.28}$	$73.0^{+1.2}_{-2.7}$




Fig. 5. Intrinsic P_{orb} distribution of DNe, I , estimated from the observed sample.

having long T_s are an excellent candidate for the missing population near P_{\min} , and ii) the observed P_{\min} is significantly longer than the real P_{\min} because of the low outburst activity of DNe near P_{\min} .

4. Discussion

4.1. Dependence of α and P_{\min} on the assumed model parameters

In the previous section, we assumed several parameters in our model to estimate α and P_{\min} , such as n , P_{crit} and M_V at supermaximum. Here, we discuss the dependence of the result on these parameters. We present α and P_{\min} estimated with several different sets of these parameters in table 2.

As can be seen in table 2, α is sensitive to both n and P_{crit} . Smaller n or P_{crit} leads to a smaller α . α also depends on the model of M_V . It is possible that M_V takes a form different from equation (7). According to Harrison et al. (2004), $M_{V, \text{SU}}$ for superoutbursts has a large dispersion and does not strictly follow the M_V - P_{orb} relationship for normal outbursts. This implies that the P_{orb} dependence of $M_{V, \text{SU}}$ is weak. Assuming $M_{V, \text{SU}} = M_{V, \text{WZ}} = \text{const.}$, we obtained a smaller α as shown in table 2. In all the above cases, α takes high values of $\gtrsim 0.6$

and a spike near P_{\min} appears in the P_{orb} distribution. The spike feature disappears only when n is extremely small. For example, we show the case of $n = 0.2$ in table 2. Such a small n is, however, unfavorable for the observed $D_{\text{outb}}-P_{\text{orb}}$ relationship, as shown in figure 3.

Table 2 indicates that P_{\min} is sensitive to n ; a small n yields a large P_{\min} . On the other hand, the estimated P_{\min} in table 2 are significantly smaller than the observed value, even in the extreme case of $n = 0.2$. Thus, the result is robust within the reasonable ranges of the model parameters.

As mentioned in § 3.3, we used $n = 2.0$, which is a lower limit of n in the case of $P_{\min} = 72.6$ min. n can be smaller than 2.0 only when $P_{\min} > 72.6$ min. As shown in table 2, all P_{\min} satisfies $P_{\min} < 72.6$ min, except for the extreme case of $n = 0.2$. Therefore, it is less likely that n is smaller than 2.0 in our model.

Our result highly depends on the assumption for the $D_{\text{outb}}-P_{\text{orb}}$ relationship. The assumption might be incorrect if a number of short- T_s systems remain undiscovered in the shortest P_{orb} regime, such as *Group X* identified in Paper I. The assumption for the $D_{\text{outb}}-P_{\text{orb}}$ relationship should thus be reconsidered if the intrinsic population of those objects is found to be much larger than the currently observed population. However, it is unlikely that there are significant undetected systems because the outburst detection probability must be high in such short- T_s systems and hence their outbursts should have already been discovered.

4.2. Implications for CV evolution

In this section, we discuss the implications of our result to CV evolution, considering the presence of post- P_{\min} CVs, magnetic CVs, and past theoretical and observational works in this field.

In the previous section, we regarded all systems as pre- P_{\min} CVs. However, it has been proposed that about 70% of CVs have already passed through P_{\min} (Kolb 1993; Howell et al. 1997). It is possible that all known DNe are pre- P_{\min} CVs, while most CVs have already passed through P_{\min} and remained undiscovered. In this case, the intrinsic P_{orb} distribution would have a taller spike near P_{\min} than that in figure 5 because of the contribution of the undiscovered post- P_{\min} CVs.

On the other hand, a fraction of the long- T_s DNe in our sample might actually be post- P_{\min} systems because post- P_{\min} systems presumably have quite low \dot{M} and long T_s . Even if most known WZ Sge stars are post- P_{\min} CVs, the spike structure in figure 5 is still expected to appear because the spike feature depends, in principle, on the number of long- T_s DNe near P_{\min} and is independent of the nature of the systems, namely, whether they are pre- or post- P_{\min} systems.

In contrast to the spike feature, the real P_{\min} is possibly sensitive to the number of pre- P_{\min} systems in the shortest P_{orb} regime. If the true fraction of post- P_{\min} systems is large in known WZ Sge stars, then it means that super-outbursts of the post- P_{\min} systems are detectable for us. If this is the case, we should be able to detect outbursts of

DNe below the observed P_{\min} . The estimated P_{\min} could then be close to the observed value.

The true intrinsic fraction of post- P_{\min} systems in the shortest P_{orb} regime is poorly known, while recent observational studies suggest the presence of post- P_{\min} systems. Townsley & Gänsicke (2009) investigated the temperature of the white dwarf in CVs as a probe of \dot{M} . Their result shows that there are anomalies having quite low temperatures of the white dwarf compared with most CVs in the shortest P_{orb} regime. The \dot{M} of those anomalies are possibly lower than the ordinary ones, and hence, suggesting that they are post- P_{\min} systems. Littlefair et al. (2008) reported three CVs in which the post- P_{\min} nature was dynamically confirmed (also see, Littlefair et al. 2006; Littlefair et al. 2007). Among known DNe, WZ Sge has the longest T_s and hence is a good candidate for post- P_{\min} systems. Steeghs et al. (2007), however, reported that its secondary mass is so high that WZ Sge is in fact a pre- P_{\min} system. The nature of WZ Sge implies that most DNe whose T_s are known are still in the pre- P_{\min} regime.

As mentioned in § 3.3, our model of D_{outb} assumed a decrease of \dot{M} toward P_{\min} in the region $P_{\text{orb}} < 86$ min. This was required from the observed T_s-P_{orb} relationship, as can be seen in figure 3. However, this is apparently inconsistent with theoretical calculations for the CV evolution, which predict a rather constant \dot{M} for pre- P_{\min} systems even near P_{\min} (e.g. Shafter 1992; Kolb 1993; Barker, Kolb 2003). If the theoretical prediction is true, we would expect many short- T_s systems in the shortest P_{orb} regime. The prediction is, however, unfavorable for the observed population, because there are only a few short- T_s systems in $P_{\text{orb}} \lesssim 86$ min, such as *Group X* in Paper I. This inconsistency implies that the current understanding of the secondary star is imperfect, since the theoretical $\dot{M}-P_{\text{orb}}$ relationship depends on the structure of the secondary star. Meyer, Meyer-Hofmeister (1999a) proposed that the magnetic activity of the secondary star reduces rapidly as it becomes degenerate, and thereby, the viscosity in the disk reduces and T_s increases. The observed T_s-P_{orb} relationship may be reproduced if T_s increases by this mechanism in $P_{\text{orb}} < 86$ min.

It has been reported that the P_{orb} distribution of magnetic CVs is similar to that of non-magnetic CVs (e.g. Willems et al. 2005). However, this discussion was based on samples obtained from various observation processes, such as samples in RKcat. As mentioned in section 2, such a sample is not suitable to be compared with the theoretical P_{orb} distribution obtained from population synthesis studies. Our result suggests that, in the case of DNe, the period minimum and spike problems are reconciled with a number of undiscovered WZ Sge stars. Since AM Her stars have no accretion disk, and thereby, experience no DN-type outbursts, our result also implies that another mechanism to reduce short- P_{orb} systems should work for magnetic CVs (e.g. Kolb 1995; Meyer, Meyer-Hofmeister 1999b). The reduce of magnetic braking in magnetic CVs may lead to a different P_{orb} distribution from non-magnetic CVs (Webbink, Wickramasinghe 2002; Araujo-Betancor et al. 2005).

Gänsicke et al. (2009) have recently reported the discovery of the period spike feature in the CV sample selected from the SDSS database. The appearance of the spike is in agreement with our result; we reproduced a spike based on observed outburst activity, while Gänsicke et al. (2009) obtained it directly from the deep survey. They discovered that CVs having a white-dwarf dominated spectrum form a major population in the CVs making the period spike feature. Their spectrum indicates that M of those CVs are quite low. Hence, they may be WZ Sge stars, and their outbursts could be observed with a long T_s . Actually, SDSS J080434.20+510349.2, a system having a white-dwarf dominated spectrum, experienced a WZ Sge-type superoutburst in 2005 (Pavlenko et al. 2007). The SDSS CVs having a white-dwarf dominated spectrum probably have the same nature as the objects that our analysis predicts to be present in the shortest P_{orb} regime.

On the other hand, the implications for the period minimum problem are different between our result and Gänsicke et al. (2009). In our result, the observed P_{min} is significantly longer than the real P_{min} because T_s is so long that the detection frequency is low near the real P_{min} . The SDSS images are so deep that we can expect to detect CVs with a low M near the real P_{min} . However, a sharp cut-off exists at $P_{\text{orb}} \sim 80$ min in the P_{orb} distribution of the SDSS sample and only a few sources were found in the P_{orb} range of 60–78 min. The period minimum problem, hence, remains unresolved in Gänsicke et al. (2009). The lack of the sources in the P_{orb} range 60–78 min may indicate that a fraction of known WZ Sge stars are post- P_{min} CVs, as discussed above. Alternatively, some CVs with a very low M may remain undiscovered even in the SDSS survey. Gänsicke et al. (2009) identified CVs based on spectral features of objects that were selected by several color criteria. A part of the color criteria plays a role in excluding single white dwarfs from the CV sample. The criteria, however, possibly excludes evolved CVs which have a quite low M , and thereby, have optical spectra dominated by the emission from a white dwarf.

5. Summary

We estimated the intrinsic P_{orb} distribution of DNe assuming a simple relationship between T_s and P_{orb} . The estimated distribution has a spike near P_{min} and a shorter P_{min} than the observed value, which was expected by the theory of binary evolution. We suggest that the nature of the “missing” population is DNe with long T_s near P_{min} .

We thank B. Gänsicke for his valuable comments and suggestions on this paper. This work was partly supported by a Grand-in-Aid from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (19740104).

Appendix 1. Outburst-selected DNe sample

Table 3 and 4 lists the objects of the ASAS and mixed sample used in our analysis described in section 3, respectively. The object name and P_{orb} were quoted from

Ritter & Kolb catalog 7.9 (Ritter, Kolb 2003).

References

- Araujo-Betancor, S., Gänsicke, B. T., Long, K. S., Beuermann, K., de Martino, D., Sion, E. M., & Szkody, P. 2005, *ApJ*, 622, 589
- Bade, N., Engels, D., Voges, W., Beckmann, V., Boller, T., Cordis, L., Dahlem, M., Englhauser, J., et al. 1998, *A&AS*, 127, 145
- Barker, J. & Kolb, U. 2003, *MNRAS*, 340, 623
- Burd, A., Cwiok, M., Czyrkowski, H., Dabrowski, R., Dominik, W., Grajda, M., Husejko, M., Jegier, M., et al. 2005, *New Astronomy*, 10, 409
- Drake, A. J., Djorgovski, S. G., Mahabal, A., Beshore, E., Larson, S., Graham, M. J., Williams, R., Christensen, E., et al. 2009, *ApJ*, 696, 870
- Gänsicke, B. T., Dillon, M., Southworth, J., Thorstensen, J. R., Rodríguez-Gil, P., Aungwerojwit, A., Marsh, T. R., Szkody, P., et al. 2009, *MNRAS*, 397, 2170
- Harrison, T. E., Johnson, J. J., McArthur, B. E., Benedict, G. F., Szkody, P., Howell, S. B., & Gelino, D. M. 2004, *AJ*, 127, 460
- Howell, S. B., Rappaport, S., & Politano, M. 1997, *MNRAS*, 287, 929
- Howell, S. B., Szkody, P., & Cannizzo, J. K. 1995, *ApJ*, 439, 337
- Kato, T., Imada, A., Uemura, M., Nogami, D., Maehara, H., Ishioka, R., Baba, H., Matsumoto, K., et al. 2010, *PASJ*, 61, S395
- Kato, T. & Kunjaya, C. 1995, *PASJ*, 47, 163
- Kato, T., Sekine, Y., & Hirata, R. 2001, *PASJ*, 53, 1191
- Kato, T., Uemura, M., Ishioka, R., Nogami, D., Kunjaya, C., Baba, H., & Yamaoka, H. 2004, *PASJ*, 56, 1
- King, A. R. 1988, *QJRAS*, 29, 1
- King, A. R., Schenker, K., & Hameury, J. M. 2002, *MNRAS*, 335, 513
- Kolb, U. 1993, *A&A*, 271, 149
- Kolb, U. 1995, in *Magnetic Cataclysmic Variables*, ed. D. A. H. Buckley, & B. Warner Vol. 85 of *Astronomical Society of the Pacific Conference Series*(Dordrecht: Kluwer Academic Publishers), 440
- Kolb, U. & Baraffe, I. 1999, *MNRAS*, 309, 1034
- Littlefair, S. P., Dhillon, V. S., Marsh, T. R., Gänsicke, B. T., Baraffe, I., & Watson, C. A. 2007, *MNRAS*, 381, 827
- Littlefair, S. P., Dhillon, V. S., Marsh, T. R., Gänsicke, B. T., Southworth, J., Baraffe, I., Watson, C. A., & Copperwheat, C. 2008, *MNRAS*, 388, 1582
- Littlefair, S. P., Dhillon, V. S., Marsh, T. R., Gänsicke, B. T., Southworth, J., & Watson, C. A. 2006, *Science*, 314, 1578
- Matthews, O. M., Wheatley, P. J., Wynn, G. A., & Truss, M. R. 2006, *MNRAS*, 372, 1593
- Metropolis, N. 1953, *The journal of chemical physics*, 21, 1087
- Meyer, F. & Meyer-Hofmeister, E. 1999a, *A&A*, 341, L23
- Meyer, F. & Meyer-Hofmeister, E. 1999b, *A&A*, 346, L13
- Meyer-Hofmeister, E., Meyer, F., & Liu, B. F. 1998, *A&A*, 339, 507
- Osaki, Y. 1995, *PASJ*, 47, 47
- Paczynski, B. 1981, *Acta Astronomica*, 31, 1
- Patterson, J. 1998, *PASP*, 110, 1132
- Pavlenko, E., Shugarov, S. Y., Katysheva, N. A., Nogami, D., Nakajima, K., Maehara, H., Andreev, M., Shimansky, V., et al. 2007, in *ASP Conf. Ser. 372*, *European Workshop on*

Table 3. Members of the ASAS sample.

Object	P_{orb} (min)						
GW Lib	76.8	J0233–1047	78.9	J0137–0912	79.7	V1108 Her	81.9
WX Cet	83.9	J1112–3538	84.1	CC Scl	84.5	2219+1824	86.3
V1040 Cen	86.8	AQ Eri	87.8	J0918–2942	88.8	J1025–1542	89.3
V436 Cen	90.0	EK TrA	90.5	OY Car	90.9	J1600–4846	91.3
J1536–0839	91.6	J0232–3717	93.2	AK Cnc	93.7	GO Com	94.8
RZ Sge	98.3	DT Oct	104.5	V368 Pe	105.1	VZ Pyx	105.6
CC Cnc	105.9	J1556–0009	106.7	VW Hyi	107.0	Z Cha	107.3
QW Ser	107.4	WX Hyi	107.7	RZ Leo	109.5	DH Aql	111.4
J0219–3045	112.9	CU Vel	113.0	J0549–4921	115.5	V877 Ara	116.7
TU Crt	118.2	TY PsA	121.1	KK Tel	121.7	V364 Peg	122.4
YZ Cnc	125.0	GZ Cnc	126.9				

Table 4. Members of the mixed sample.

Object	P_{orb} (min)						
J0329+1250	76.0	GW Lib	76.8	J0222+4122	78.9	J0233–1047	78.9
J0137–0912	79.7	SS LMi	80.2	J1021+2349	80.5	J0025+1217	80.7
V455 And	81.1	AL Com	81.6	SW UMa	81.8	J1839+2604	81.9
J1959+2242	83.5	WX Cet	83.9	J1112–3538	84.1	CC Scl	84.5
FL TrA	84.8	J0804+5103	85.0	V585 Lyr	85.4	2219+1824	86.3
J0807+1138	86.4	CI UMa	86.4	DV Dra	86.4	RX Vol	86.5
MM Sco	86.7	V1040 Cen	86.8	V1028 Cyg	86.9	UZ Boo	87.4
AQ Eri	87.8	V1454 Cyg	87.8	XZ Eri	88.1	V4140 Sgr	88.5
J0918–2942	88.8	J1025–1542	89.3	V1141 Aql	89.3	V402 And	89.4
V2051 Oph	89.9	V436 Cen	90.0	BC UMa	90.2	HO Del	90.3
EK TrA	90.5	TV Crv	90.6	J1227+5139	90.7	OY Car	90.9
J1600–4846	91.3	J1536–0839	91.6	MR UMa	91.6	Var 79 Peg	91.7
DO Vul	92.2	J1653+2010	92.7	J0232–3717	93.2	UV Per	93.5
CT Hya	93.5	AK Cnc	93.7	AO Oct	94.1	DM Lyr	94.2
GO Com	94.8	V551 Sgr	94.9	BC Dor	95.2	FQ Mon	95.8
SX LMi	96.8	V701 Tau	96.9	SS UMi	97.6	NSV 4838	97.6
BZ UMa	97.9	J0824+4931	97.9	KS UMa	97.9	V337 Cyg	97.9
RZ Sge	98.3	TY Psc	98.4	IR Gem	98.5	V699 Oph	98.6
V1208 Tau	99.1	V1504 Cyg	100.1	CY UMa	100.2	RU Hor	100.4
BB Ari	101.2	VW CrB	101.8	PU Per	102.7	FO And	103.1
AW Sge	104.3	TU Tri	104.3	DT Oct	104.5	OU Vir	104.7
V1212 Tau	105.1	V368 Peg	105.1	KV And	105.6	VZ Pyx	105.9
CC Cnc	105.9	AY Lyr	106.1	IY UMa	106.4	J1556–0009	106.7
VW Hyi	107.0	Z Cha	107.3	QW Ser	107.4	J0242–2802	107.4
BE Oct	107.6	WX Hyi	107.7	RZ Leo	109.5	QY Per	109.6
0417+7445	109.9	SU UMa	109.9	V630 Cyg	110.0	J1730+6247	110.2
V1113 Cyg	110.3	AW Gem	110.6	HS Vir	110.7	EG Aqr	110.9
TT Boo	110.9	V344 Pav	110.9	DH Aql	111.4	TY Vul	111.9
V503 Cyg	111.9	PV Per	112.0	V359 Cen	112.2	V660 Her	112.7
BZ Cir	112.9	J0219–3045	112.9	CU Vel	113.0	J2100+0044	113.8
NSV 14652	113.8	BR Lup	114.5	NSV 9923	114.6	J2258–0949	115.2
J0549–4921	115.5	RX Cha	116.6	V877 Ara	116.7	AB Nor	117.1
CP Dra	117.5	V369 Peg	117.6	TU Crt	118.2	HV Aur	118.5
EF Peg	120.5	TY PsA	121.1	BF Ara	121.7	KK Tel	121.7
V452 Cas	121.8	V364 Peg	122.4	DV UMa	123.6	V419 Lyr	124.4
YZ Cnc	125.0	DM Dra	125.3	V344 Lyr	126.1	GZ Cnc	126.9
GX Cas	128.2	UV Gem	128.2				

- White Dwarfs, ed. R. Napiwotzki, & M. R. Burleigh (San Francisco: ASP), 511
- Pojmanski, G. 2002, *Acta Astron.*, 52, 397
- Pretorius, M. L., Knigge, C., & Kolb, U. 2007, *MNRAS*, 374, 1495
- Rappaport, S., Joss, P. C., & Webbink, R. F. 1982, *ApJ*, 254, 616
- Renvoizé, V., Baraffe, I., Kolb, U., & Ritter, H. 2002, *A&A*, 389, 485
- Rezzolla, L., Uryū, K. ō., & Yoshida, S. 2001, *MNRAS*, 327, 888
- Ritter, H. & Kolb, U. 2003, *A&A*, 404, 301
- Shafter, A. W. 1992, *ApJ*, 394, 268
- Shafter, A. W., Coelho, E. A., & Reed, J. K. 2007, *PASP*, 119, 388
- Shugarov, S. Y., Volkov, I. M., & Chochol, D. 2008, *Information Bulletin on Variable Stars*, 5862, 1
- Soejima, Y., Imada, A., Nogami, D., Kato, T., & Monard, B. 2009, *PASJ*, 61, 395
- Spruit, H. C. & Ritter, H. 1983, *A&A*, 124, 267
- Steeeghs, D., Howell, S. B., Knigge, C., Gänsicke, B. T., Sion, E. M., & Welsh, W. F. 2007, *ApJ*, 667, 442
- Szkody, P., Anderson, S. F., Agüeros, M., Covarrubias, R., Bentz, M., Hawley, S., Margon, B., Voges, W., et al. 2002, *AJ*, 123, 430
- Templeton, M., Waagen, E. O., Poyner, G., Gobet, F., Rodriguez, D., Gualdoni, C., Koff, R., Waddill, G., et al. 2007, *Central Bureau Electronic Telegrams*, 1053, 1
- Townsley, D. M. & Gänsicke, B. T. 2009, *ApJ*, 693, 1007
- Trotta, R. 2008, *Contemporary Physics*, 49, 71
- Uemura, M., Arai, A., Kato, T., Maehara, H., Nogami, D., Kubota, K., Moritani, Y., Imada, A., et al. 2010, *PASJ*, 62, 187
- Uemura, M., Arai, A., Krajci, T., Pavlenko, E., Shugarov, S. Y., Katysheva, N. A., Goranskij, V. P., Maehara, H., et al. 2008, *PASJ*, 60, 227
- Waagen, E. O., Schmeer, P., Stubbings, R., & Pearce, A. 2007, *IAU Circ.*, 8829, 1
- Warner, B. 1995, *Cataclysmic Variable Stars* (Cambridge University Press)
- Webbink, R. F. & Wickramasinghe, D. T. 2002, *MNRAS*, 335, 1
- Willems, B., Kolb, U., Sandquist, E. L., Taam, R. E., & Dubus, G. 2005, *ApJ*, 635, 1263
- Zharikov, S. V., Tovmassian, G. H., Neustroev, V. V., Michel, R., Zurita, C., Echevarría, J., Bikmaev, I. F., Pavlenko, E. P., et al. 2008, *A&A*, 486, 505