## Learning as a phenomenon occurring in a critical state

Lucilla de Arcangelis \* <sup>† ‡</sup> and Hans J. Herrmann <sup>†</sup>

<sup>†</sup> Institute Computational Physics for Engineering Materials, ETH, Schafmattstr.6, 8093 Zürich, CH, and <sup>‡</sup>Department of Information Engineering and CNISM, Second University of Naples, 81031 Aversa (CE), Italy

Submitted to Proceedings of the National Academy of Sciences of the United States of America

Recent physiological measurements have provided clear evidence about scale-free avalanche brain activity and EEG spectra, feeding the classical enigma of how such a chaotic system can ever learn or respond in a controlled and reproducible way. Models for learning, like neural networks or perceptrons, have traditionally avoided strong fluctuations. Conversely, we propose that brain activity having features typical of systems at a critical point, represents a crucial ingredient for learning. We present here a study which provides novel insights toward the understanding of the problem. Our model is able to reproduce quantitatively the experimentally observed critical state of the brain and, at the same time, learns and remembers logical rules including the exclusive OR (XOR), which has posed difficulties to several previous attempts. We implement the model on a network with topological properties close to the functionality network in real brains. Learning occurs via plastic adaptation of synaptic strengths and exhibits universal features. We find that the learning performance and the average time required to learn are controlled by the strength of plastic adaptation, in a way independent of the specific task assigned to the system. Even complex rules can be learned provided that the plastic adaptation is sufficiently slow.

learning | scale-free network | self-organized criticality | scaling

Spontaneous activity is an important property of the cerebral cortex that can have a crucial role in information processing and storage. Recently it has been shown that a



Fig. 1. One neuronal avalanche in the scale-free network. 40 neurons are connected by directed bonds (direction indicated by the arrow at one edge), representing the synapses. The size of each neuron is proportional to the number of in-connections, namely the number of dendrites. The two red neurons are the two input sites, whereas the black neuron is the output. Connections involved in the avalanche propagation are shown in red, whereas inactive connections are black.

novel spatio-temporal form of spontaneous activity is neuronal avalanches, which can involve from a few to a very large number of neurons. These bursts of firing neurons have been first observed [1, 2] in organotypic cultures from coronal slices of rat cortex, where the size and duration of neuronal avalanches follow power law distributions with very stable exponents. The presence of a power law behaviour is the typical feature of a system acting in a critical state [3], where large fluctuations are present and the response does not have a characteristic size. The same critical behaviour, namely the same power law exponents, has been recently measured also in vivo from superficial layers of cortex in anesthetized rats during early postnatal development [4], and awake adult rhesus monkeys [5], using micro-electrode array recordings. Results confirm that indeed spontaneous cortical activity adjusts in a critical state where the spatio-temporal organization of avalanches is scale invariant. Moreover, the investigation on the spontaneous activity of dissociated neurons from different networks as rat hippocampal neurons [6], rat embryos [7] or leech ganglia [6], has also confirmed the robustness of this scaling behaviour. In all these cases, the emergence of power law distributions has been interpreted in terms of self-organized criticality (SOC) [8]. The term SOC usually refers to a mechanism of slow energy accumulation and fast energy redistribution driving the system toward a critical state, where the avalanche extensions and durations do not have a characteristic size.

The understanding of the fundamental relations between electro-physiological activity and brain organization with respect to performing even simple tasks, is a long-standing fascinating question. A number of theoretical models [9] have been proposed for learning, from the simple perceptron [10] to Attractor Neural Networks (ANNs) [11] of artificial twostate neurons [12]. In these models the state of the "brain" is the snapshot of the ensemble of the individual states of all neurons, which explores phase space following an appropriate dynamics and eventually recovers memories. The ability of the brain to self-organize connections in an efficient way is a crucial ingredient in biologically plausible models. The breakthrough of Hebbian plasticity, postulating synapse strengthening for correlated activity at the pre- and post-synaptic neuron and synapse weakening for decorrelated activity, triggered the development of algorithms for neuronal learning and memory, as, for instance, "reinforcement learning" [13] or error backpropagation [14], leading for the first time to the XOR rule learning. Recent results have shown that extremal dynamics, where only the neuron with the largest input fires, and

The authors declare no conflict of interest

<sup>\*</sup> To whom correspondence should be addressed. Email:  $idearcangelis@na.infn.it_{i}$  @2007 by The National Academy of Sciences of the USA

uniform negative feedback are sufficient ingredients to learn the following task: to identify the right connection between an input and an output node [15, 16]. Similarly, low activity probabilistic firing rules, where again a single neuron fires at each step of the iteration, together with a uniform negative feedback plastic adaptation acting on time scales slower than the neuron firing time scale, enables learning the XOR rule without error back-propagation [17]. Both results suggest that the system learns by mistakes, namely depression rather than enhancement of synaptic strength is the crucial mechanism for learning. However, in both studies a single neuron fires at each step of the evolution, not in complete agreement with recent experimental discoveries. Cooperative effects leading to self-organization and learning are completely neglected in the aforementioned approaches.

Operating at a critical level, far from an uncorrelated subcritical or a too correlated supercritical regime, may optimize information management and transmission in real brains [18, 1, 19, 20], as recently confirmed by experiments [21]. Moreover, a recent study of visual perceptual learning has evidenced that training to a specific task induces dynamic changes in the functional connectivity able to "sculpt" the spontaneous activity of the resting human brain and to act as a form of "system memory" [22]. It is therefore tempting to investigate the role that critical behaviour plays in the most important task of neuronal networks, namely learning and memory. The emergence of a critical state with the same critical behaviour found experimentally has been recently reproduced by a neuronal network model based on SOC ideas [23, 24]. The model implements several physiological properties of real neurons: a continuous membrane potential, firing at threshold, synaptic plasticity and pruning. Extensive numerical studies on regular, small world and scale free networks have shown that indeed the system exhibits a robust critical behaviour. The distributions of avalanche size and duration scale with exponents independent of model parameters and in excellent agreement with experimental data (Fig.2). More precisely, the distribution of avalanche sizes, measured experimentally either in terms of number of active electrodes or summed local field potentials in a micro-electrode array [1, 2], decreases with an exponent -1.5, whereas the distribution of avalanche temporal durations decreases with an exponent close to -2.0. A critical avalanche activity has been also found on fully connected [25] and random networks [26]. Moreover, the temporal signal for electrical activity and the power spectrum of the resulting time series have been compared with EEG data [23, 24]. The spectrum exhibits a power law behaviour,  $P(f) \sim f^{-0.8}$ , with an exponent in good agreement with EEG medical data [27] and physiological signal spectra for other brain controlled activities [28]. This model therefore seems to capture many of the essential ingredients of spontaneous activity, as measured in cortical networks.

Here we study the learning performance of a neuronal network acting in a critical state. The response of the system to external stimuli is therefore scale-free, i.e. no characteristic size in the number of firing neurons exists. The approach reproduces closely the physiological mechanisms of neuronal behaviour and is implemented on a plausible network having topological properties similar to the brain functionality network. Neuronal activity is a collective process where all neurons at threshold can fire and self-organize an efficient path for information transmission. Plastic adaptation is introduced via a non uniform negative feedback procedure with no error back propagation.

## The model

We consider N neurons positioned at random in a two dimensional space. Each neuron is characterized by the potential  $v_i$ . Connections among neurons are established by assigning to each neuron i a random out-going connectivity degree,  $k_{out_i}$ . The distribution of the number of out-connections is then chosen in agreement with the experimentally measured properties of the functionality network [29] in human adults. Functional magnetic resonance imaging has indeed shown that this network has universal scale free properties, namely it exhibits a scaling behaviour  $n(k_{out}) \propto k_{out}^{-2}$ , independent of the different tasks performed by the patient. We adopt this distribution for the number of pre-synaptic terminals of each neuron, over the range of possible values between  $k_{out}^{min}$  and  $k_{out}^{max} = 100$ , as in experimental data. Two neurons are then connected according to a distance dependent probability,  $p(r) \propto e^{-r/r_0}$ , where r is their spatial distance [30] and  $r_0$  a typical edge length. To each synaptic connection we then assign an initial random strength  $g_{ij}$ , where  $g_{ij} \neq g_{ji}$ , and an excitatory or inhibitory character, with a fraction  $p_{in}$  of inhibitory synapses. An example of such a network is shown in Fig.1.

The firing dynamics implies that, whenever at time t the value of the potential at a site i is above a certain threshold  $v_i \ge v_{\text{max}} = 6.0$ , approximately equal to -55mV for real neurons, the neuron sends action potentials leading to the production of an amount of neurotransmitter proportional to  $v_i$ . As a consequence, the total charge released by a neuron is proportional to the number of synaptic connections,  $q_i \propto v_i k_{out_i}$ . Each connected neuron receives charge in proportion to the strength of the synapse  $g_{ij}$ 

$$v_j(t+1) = v_j(t) \pm \frac{q_i(t)}{k_{in_j}} \frac{g_{ij}(t)}{\sum_k g_{ik}(t)}$$
[1]

where  $k_{in_j}$  is the in-degree of neuron j and the sum is extended to all out-going connections of i. In Eq.(1) it is as-



**Fig. 2.** Demonstration of the critical behaviour of the neuronal network avalanche activity. The distribution of the sizes of the avalanches of firing neurons, n(S) (circles), follows a power law behaviour with an exponent  $1.5 \pm 0.1$  (dashed line). The size is measured as the number of firing neurons. The distribution of avalanche durations, n(T) (squares), exhibits a power law behaviour with an exponent  $2.2 \pm 0.2$  (dot-dashed line), followed by an exponential cutoff. Data are obtained for 40 realizations of a network of 4000 neurons with  $p_{in} = 0.05$ .

sumed that the received charge is distributed over the surface of the soma of the post-synaptic neuron, proportional to the number of in-going terminals  $k_{in_j}$ . The plus or minus sign in Eq.(1) is for excitatory or inhibitory synapses, respectively. After firing a neuron is set to a zero resting potential and in a refractory state lasting  $t_{ref} = 1$  time step, during which it is unable to receive or transmit any charge. We wish to stress that the unit time step in Eq.(1) does not correspond to a real time scale, it is simply the time unit for charge propagation from one neuron to the connected ones. In real system this time could vary and be as large as 100 ms for longer firing periods. The synaptic strengths have initially a random value  $g_{ij} \in [0.5, 1.0]$ , whereas the neuron potentials are uniformly distributed random numbers between  $v_{\rm max} - 1$  and  $v_{\rm max}$ . Moreover, a small random fraction (10%) of neurons is chosen to be boundary sites, with a potential fixed to zero, playing the role of sinks for the charge.

In order to start activity we identify input neurons at which the imposed signal is applied and the output neuron at which the response is monitored. These nodes are randomly placed inside the network under the condition that they are not boundary sites and they are mutually separated on the network by  $k_d$  nodes.  $k_d$  represents the chemical distance on the network and plays the role of the number of hidden layers in a perceptron. We test the ability of the network to learn different rules: AND, OR, XOR and a random rule RAN which associates to all possible combinations of binary states at three inputs a random binary output. More precisely, the AND, OR and XOR rules are made of three input-output relations (we disregards the double zero input which is a trivial test leading to zero output), whereas the RAN rule with three input sites implies a sequence of seven input-output relations. A single learning step requires the application of the entire sequence of states at the input neurons, monitoring the state of the output neuron. For each rule the binary value 1 is identified with the output neuron firing, namely the neuron membrane potential at a value greater or equal to  $v_{\max}$  at some time during the activity. Conversely, the binary state 0 at the output neuron corresponds to the physiological state of a real neuron which has been depolarized by incoming ions but fails to reach the firing threshold membrane potential during the entire avalanche propagation. Once the input sites are stimulated, their activity may bring to threshold other neurons and therefore lead to avalanches of firings. We impose no restriction on the number of firing neurons in the propagation and let the avalanche evolve to its end according to Eq.(1). If at the end of the avalanche the propagation of charge did not reach the output neuron, we consider that the state of the system was unable to respond to the given stimulus, and as a consequence to learn. We therefore increase uniformly the potential of all neurons by units of a small quantity,  $\beta = 0.01$ , until the configuration reaches a state where the output neuron is first perturbed. We then compare the state of the output neuron with the desired output. Namely we follow the evolution in phase space of the initial state of the system and verify if the non-ergodic dynamics has led to an attractor associated with the right answer.

Plastic adaptation. Plastic adaptation is applied to the system according to a non uniform negative feedback algorithm. Namely, if the output neuron is in the correct state according to the rule, we keep the value of synaptic strengths. Conversely, if the response is wrong we modify the strengths of those synapses involved in the information propagation by  $\pm \alpha/d_k$ , where  $d_k$  is the chemical distance of the presynaptic neuron from the output neuron. Here  $\alpha$  represents the ensemble of all possible physiological factors influencing synaptic plasticity. The sign of the adjustment depends on the mistake made by the system: If the output neuron fails to be in a firing state we increase the used synapses by a small additive quantity proportional to  $\alpha$ . Synaptic strengths are instead decreased by if the expected output 0 is not fulfilled. Once the strength of a synapse is below an assigned small value  $g_t = 10^{-4}$ , we remove it, i.e. set its strength equal to zero, which corresponds to the so-called *pruning*. This ingredient is very important as since decades the crucial role of selective weakening and elimination of unneeded connections in adult learning has been recognized [31, 32]. The synapses involved in the signal propagation and responsible for the



**Fig. 3.** Demonstration of the learning ability of the neuronal network. (*Left*) Percentage of configurations learning the XOR rule as function of the number of learning steps for different plastic adaptation strengths  $\alpha$  (decreasing from bottom to top). Data are for 400 realizations of a network with N = 1000 neurons,  $p_{in} = 0.1$ ,  $r_0 = 15$ ,  $k_{min} = 3$  and  $k_d = 5$ . (*Right*) Collapse of the curves by rescaling the number of learning steps by the characteristic learning time  $\tau = 1/\alpha$  and the percentage of success by  $\alpha^{-0.05}$ .



**Fig. 4.** Demonstration of the scaling behaviour of learning time and success. *(Top)* Scaling of the average learning time for different rules as function of  $\alpha$ . *(Bottom)* Scaling of the asymptotic percentage of configurations learning different rules as function of  $\alpha$ . Data are obtained for  $k_{min} = 3$ ,  $k_d = 5$  and  $p_{in} = 0.1$ . Both quantities follow a power law with an exponent independent of the rule and the percentage of inhibitory synapses.

wrong answer, are therefore not adapted uniformly but inversely proportional to the chemical distance from the output site. Namely, synapses directly connected to the output neuron receive the strongest adaptation  $\pm \alpha$ . This adaptation rule intends to mimic the feedback to the wrong answer triggered locally at the output site and propagating backward towards the input sites. This could be the case, for instance, of some hormones strongly interfering with learning and memory, as dopamine suppressing LTD [34] or adrenal hormones enhancing LTD [35]. Moreover a new class of messenger molecules as nitric oxide has been found to have an important role in plastic adaptation [36]. For all these agents, released at the output neuron, the concentration is reduced with the distance from the origin. In our neuronal network simulation this non uniform adaptation has a crucial role since it prevents, in case of successive wrong positive answers, synapses directly connected to the input sites to decrease excessively, hindering any further signal transmission. This plastic adaptation is a non-Hebbian form of plasticity and can be interpreted as a subtractive form of synaptic scaling [33], where synapses are changed by an amount independent of their strength. The procedure mimics the performance of a *good critic* who does not tell the system which neurons should have fired or not. However it tells more than just "right" or "wrong", it expresses an evaluation on the type of error. Finally, we wish to stress that this model naturally sets the system in a critical state and therefore the study of the response of the system in a subcritical or supercritical state requires the introduction of additional parameters. We can however suppose that in both cases learning becomes a more difficult task. For instance, in a subcritical state, being the size of neuronal avalanches smaller, it would be more complex to generate a firing state in the output site. Conversely, in a supercritical state it would be more difficult to generate a non-firing state in the output site.

## **Results and Discussion**

We analyse the ability of the system to learn the different rules. Fig. 3 shows the fraction of configurations learning the XOR rule versus the number of learning steps for different values of the plastic adaptation strength  $\alpha$ . We notice that the larger the value of  $\alpha$  the sooner the system starts to learn the rule, however the final percentage of learning configurations is lower. The final rate of success increases as the strength of plastic adaptation decreases. This result is due to the highly non linear dynamics of the model, where firing activity is an all or none event controlled by the threshold. Very slow plastic adaptation allows then to tune finely the role of the neurons involved in the propagation and eventually recover the right answer. Moreover, very slow plastic adaptation also makes the system more stable with respect to noise, since too strong synaptic changes may perturb excessively the evolution hampering the recovery of the right answer. The dependence of the learning success on the plasticity strength is found consistently for different values of the parameters  $k_d$ ,  $k_{min}$  and  $p_{in}$ , where a higher percentage of success is observed in systems with no inhibitory synapses. Moreover, the dependence on the plastic adaptation  $\alpha$  is a common feature of all tested rules. Data indicate that the easiest rule to learn is OR, where a 100% percentage of success can be obtained. AND and XOR present similar difficulties and lead both to a percentage of final success around 80%, whereas the most difficult rule to learn is the RAN rule with three inputs where only 50% of final success is obtained. This different performance is mainly due to the higher number of inputs, since the system has to organized a more complex path of connections leading to the output site.

The most striking result is that all rules give a higher percentage of success for weaker plastic adaptation. Indeed this result is in agreement with recent experimental findings on visual perceptual learning, where better performances are measured when minimal changes in the functional network occur as a result of learning [22]. We characterize the learning ability of a system for different rules by the average learning time, i.e. the average number of times a rule must be applied to obtain the right answer, and the asymptotic percentage of learning configurations. This is determined as the percentage of learning configurations at the end of the teaching routine, namely after  $10^6$  applications of the rule. Fig. 4 shows that the average learning time scales as  $\tau \propto 1/\alpha$  for all rules and independently of parameter values. Since some configurations never learn and do not contribute to the average learning time, we also evaluate the median learning time which exhibits the same scaling behaviour as the average learning time. The asymptotic percentage of success increases by decreasing  $\alpha$  as a very slow power law,  $\propto \alpha^{-0.05}$ . Since this quantity has a finite upper bound equal to unity, this scaling suggests that in a finite, even if very long, time any configuration could learn the rule by applying an extremely slow plastic adaptation. It is interesting to notice that a larger fraction of systems with no inhibitory synapses finds the right answer and the average learning time for these systems is slightly shorter. We understand this result by considering that for only excitatory synapses the system more easily selects a path of strong enough synapses connecting inputs and output sites and giving the right answer. Conversely, the presence of inhibitory synapses may lead to frustration in the system as not all local interactions contribute in the optimal way to provide the right answer and the system has to find alternative paths. We check this scaling behaviour by appropriately rescaling the axes in Fig. 3. The curves corresponding to different  $\alpha$ values indeed all collapse onto a unique scaling function. Similar collapse is observed for the OR, AND and RAN rules and for different parameters  $k_d$ ,  $k_{min}$  and  $p_{in}$ . In fact, two different cases of distributions of inhibitory synapses, one in which they are chosen randomly among all synapses, the other where certain randomly chosen neurons have all outgoing synapses inhibitory, provide equivalent results. The learning dynamics shows therefore universal properties, independent of the details of the system or the specific task assigned.

The learning behaviour is sensitive to the number of neurons involved in the propagation of the signal, and therefore depends on the distance between input and output neurons and the level of connectivity in the system. We then investigate the effect of the parameters  $k_d$  and  $k_{min}$  on the performance of the system. Fig. 5 shows the percentage of configurations learning the XOR rule for different minimum values of the neuron out degree. Systems with larger  $k_{min}$  have a larger average number of synapses per neuron, producing a more branched network. The presence of several alternative paths facilitates information transmission from the in-

puts to the output site. However, the participation of more branched synaptic paths in the learning process may delay the time the system first gives the right answer. As expected the performance of the system improves as the minimum outconnectivity degree increases, with the asymptotic percentage of success scaling as ~  $k_{min}^{0.4}$ . The dependence of the learning performance on the level of connectivity is confirmed by the analysis of systems with different number of neurons N, the same out-degree distribution and the same set of parameters. We verify that larger systems exhibit better performances. In larger systems, in fact, the number of hubs, i.e. highly connected neurons, increases improving the overall level of connectivity. Indeed, the existence of complex patterns of activation has been recently recognized as very important in linking together large scale networks in visual perceptual learning[22].

On the other hand, also the chemical distance between the input and output sites has a very important role, as the number of hidden layers in a perceptron. Indeed, as  $k_d$  becomes larger (Fig. 5), the length of each branch in a path involved in the learning process increases. As a consequence, the system needs a higher number of tests to first give the right answer and a lower fraction of configurations learns the rule after the same number of steps. The percentage of learning configurations after  $10^6$  applications is found, as expected, to decrease as  $\sim k_d^{-0.3}$  and similar behaviour is detected for the OR, AND and RAN rules.

Learning stability and Memory. The existence of systems that are unable to learn, even after many learning steps, raises intriguing questions about the learning dynamics. We question what happens when a second chance is given to the configurations failing the right answer. We then restart the learning routine after imposing a small change in the initial configuration of voltages. This small perturbation leads to about 25%more configurations learning the rule. The initial state of the system can therefore influence the ability to learn, especially for complex rules as XOR or RAN. On the other hand, the analysis of the out-degree distribution in configurations which did and did not give the right answer indicates that "dumb" configurations tend to have less highly connected nodes than the "smart" ones. Namely, giving repeatedly wrong positive answers leads to pruning of several synapses. This affects in particular the highly connected neurons, which have a crucial role in identifying the right synaptic learning path. Finally we test the ability of the configurations that do learn to remember the right answer once the initial configuration is changed. The memory performance of the system is expected to depend on the intensity of the variation imposed, namely on the extension of the basin of the attraction of states leading to the right answer. The system is able to recover the right answer in more than 50% of the configurations if a very small perturbation (of the order of  $10^{-3}$ ) is applied to all neurons or else a larger one (of the order of  $10^{-2}$ ) to 10% of neurons. The system has a different memory ability depending on the rule: almost all configurations remember OR, whereas typically 80% remember AND and at most 70% the XOR rule.

## Conclusions

In conclusion, we investigate the learning ability of a model able to reproduce the critical avalanche activity as observed for spontaneous activity in in vitro and in vivo cortical networks. The ingredients of the model are close to most functional and topological properties of real neuronal networks. The implemented learning dynamics is a cooperative mechanism where all neurons contribute to select the right answer and negative feedback is provided in a non-uniform way. Despite the complexity of the model and the high number of degrees of freedom involved at each step of the iteration, the system can learn successfully even complex rules as XOR or a random rule with three inputs. In fact, since the system acts in a critical state, the response to a given input can be highly flexible, adapting more easily to different rules. The analysis of the dependence of the performance of the system on the average connectivity confirms that learning is a truly collective process, where a high number of neurons may be involved and the system learns more efficiently if more branched paths are possible. The role of the plastic adaptation strength, considered as a constant parameter in most studies, provides a striking new result: The neuronal network has a "universal" learning dynamics, even complex rules can be learned provided that the plastic adaptation is sufficiently slow. This important requirement for plastic adaptation is confirmed by recent experimental results [22] showing that the learning performance, in humans trained to a specific visual task, improves when minimal changes occur in the functionality network. Stronger modifications of the network do not necessarily lead to better results.



**Fig. 5.** Demonstration of the dependence of the learning performance on the minimum connectivity degree and the chemical distance between input and output sites. (*Left*) Percentage of configurations learning the XOR rule as function of the number of learning steps, for  $\alpha = 0.005$  in systems with  $k_d = 5$  and different values of  $k_{min}$  (increasing from bottom to top). (*Right*) Percentage of configurations learning the XOR rule as function of the number of learning steps, for  $\alpha = 0.005$  in systems with  $k_{min} = 3$  and different values of  $k_d$  (decreasing from bottom to top). Data are for 400 realizations of a network with N = 1000 neurons,  $p_{in} = 0.1$ .

- Beggs JM, Plenz D (2003) Neuronal avalanches in neocortical circuits. J Neurosci 23: 11167-11177.
- Beggs JM, Plenz D (2004) Neuronal avalanches are diverse and precise activity patterns that are stable for many hours in cortical slice cultures. J Neurosci 24: 5216-5229.
- Stanley HE (1971) Introduction to Phase Transitions and Critical Phenomena (Oxford Univ Press, New York).
- Gireesh ED, Plenz D (2008) Neuronal avalanches organize as nested theta- and beta/gamma - oscillations during development of cortical layer 2/3. Proc Natl Acad Sci USA 105: 7576-7581.
- Petermann T, Thiagarajan TC, Lebedev MA, Nicolelis MAL, Chialvo DR, Plenz D (2009) Spontaneous cortical activity in awake monkeys composed of neuronal avalanches. *Proc Natl Acad Sci USA* 106: 15921-15926.
- Mazzoni A, Broccard FD, Garcia-Perez E, Bonifazi P, Ruaro ME, Torre V (2007) On the dynamics of spontaneous activity in neuronal networks. *PLoS ONE* 2(5): e439, 1-12, doi:10.1371/journal.pone.0000439.
- Pasquale V, Massobrio P, Bologna LL, Chiappalone M, Martinoia S (2008) Self-organization and neuronal avalanches in networks of dissociated cortical neurons. *Neuroscience* 153: 1354-1369.
- Bak P (1996) How nature works. The science of self-organized criticality (Springer, New York).
- 9. Amit D (1989) Modeling brain function ( Cambridge University Press, Cambridge).
- Rosenblatt F (1958) The perceptron: A probabilistic model for information storage and organization in the brain. *Psycological review* 65: 386-408.
- Hopfield J (1982) Neural networks and physical systems with emergent collective computational abilities. Proc Natl Acad Sci USA 79: 2554-2558.
- McCulloch WS, Pitts W (1943) A logical calculus of the ideas immanent in nervous activity. Bull Math Biophys 5: 115-133.
- Barto AG, Sutton RS, Anderson CW (1983) Neuron-like adaptive elements that can solve difficult learning control problems. *IEEE Trans Syst Man Cybern* 15: 835-846.
- Rumelhart DE, Hinton GE, Williams RJ (1986) Learning representations by back-propagating errors. *Nature* 323: 533-536.
- 15. Bak P, Chialvo DR (2001) Adaptive learning by extremal dynamics and negative feedback. Physical Review E 63: 0319121-12
- Chialvo DR, Bak P (1999) Learning from mistakes. Neuroscience 90: 1137-1148.
- Klemm K, Bornholdt S, Schuster HG (2000) Beyond Hebb: Exclusive-OR and biological learning. *Physical Review Letters* 84: 3013-3016.
- Turing AM (1957) Computing machines and intelligence. Mind 59: 433-460.
- Kinouchi O, Copelli M (2006) Optimal dynamical range of excitable networks at criticality. Nat Phys 2: 348-351.

- Eurich CW, Herrmann JM, Ernst UA (2002) Finite- size effects of avalanche dynamics. *Physical Review E* 66:0661371-15.
- Shew WL, Yang H, Petermann T, Roy R, Plenz D (2009) Neuronal avalanches imply maximum dynamic range in cortical networks at criticality. J Neurosci 29: 155595-15600.
- Lewis CM, Baldassarre A, Committeri G, Romani GL, Corbetta M (2009) Learning sculpts the spontaneous activity of the resting human brain. Proc Natl Acad Sci USA 106: 17558-17563.
- de Arcangelis L, Perrone-Capano C, Herrmann HJ (2006) Selforganized criticality model for brain plasticity. *Physical Review Letters* 96: 0281071-4.
- Pellegrini GL, de Arcangelis L, Herrmann HJ, Perrone-Capano C (2007) Activity-dependent neural network model on scale-free networks. *Physical Review E* 76: 0161071-9.
- Levina A, Herrmann JM, Geisel T(2007) Dynamical synapses causing self-organized criticality in neural networks. Nat Phys 3: 857-860
- Teramae JN, Fukai T (2007) Local cortical circuit model inferred from power-law distributed neuronal avalanches. J Comput Neurosci 22: 301-321.
- Novikov E, Novikov A, Shannahoff-Khalsa D, Schwartz B, Wright J (1997) Scale-similar activity in the brain. *Physical Review E* 56: R2387-R2389.
- Hausdorff JM, Ashkenazy Y, Peng CK, Ivanov PC, Stanley EH Goldberger AL (2001) When human walking becomes random walking: fractal analysis and modeling of gait rhythm fluctuations. *Physica A* 302: 138-147.
- Eguiluz VM, Chialvo DR, Cecchi GA, Baliki M, Apkarian AV (2005) Scale-free brain functional networks. *Physical Review Let*ters 94: 0181021-4.
- Roerig B, Chen B (2002) Relationships of local inhibitory and excitatory circuits to orientation preference maps in ferret visual cortex. *Cerebral Cortex* 12: 187-198.
- 31. Young JZ (1964) A model of the Brain (Clarendon, Oxford)
- 32. Changeux JP (1985) Neuronal Man: The Biology of Mind (Oxford University Press, New York)
- Abbott LF, Nelson SB (2000) Synaptic plasticity: taming the beast. Nature Neuroscience 3: 1178-1183.
- Otmakhova NA, Lisman JE (1998) D1/D5 Dopamine receptors inhibit depotentiation at CA1 synapses via cAMP-dependent mechanism. J Neurosci 18: 1270-1279.
- 35. Coussens CM, Kerr DS, Abraham WC (1997) Glucocorticoid receptor activation lowers the threshold for NMDA-receptordependent homosynaptic long-term depression in the hippocampus through activation of voltage-dependent Calcium channels. J Neurophysiol 78: 1-9.
- Reyes-Harde M, Empson R, Potter BVL, Galione A, Stanton PK (1999) Evidence of a role for cyclic ADP-ribose in long-term synaptic depression in hippocampus. *Proc Natl Acad Sci* 96: 4061-4066.