Smooth and Peaked Solitons of the CH equation

Darryl D. Holm¹ and Rossen I. Ivanov^{1,2}

PACS numbers: 47.10.+g, 52.65.Kj, 45.20.Jj, 47.65.+a, 02.40.Yy, 11.10.Ef, 05.45.-a

Keywords: Solitons, Peakons, Hamilton's principle, Integrable Hamiltonian systems, Inverse Spectral Transform (IST), Singular solutions, Shallow water waves, Euler's equations, Geodesic motion, Diffeomorphisms, Sobolev norms, Momentum maps, Compactons

Abstract

The relations between smooth and peaked soliton solutions are reviewed for the Camassa-Holm (CH) shallow water wave equation in one spatial dimension. The canonical Hamiltonian formulation of the CH equation in action-angle variables is expressed for solitons by using the scattering data for its associated isospectral eigenvalue problem, rephrased as a Riemann-Hilbert problem. The momentum map from the action-angle scattering variables $T^*(\mathbb{T}^N)$ to the flow momentum (\mathfrak{X}^*) provides the Eulerian representation of the N-soliton solution of CH in terms of the scattering data and squared eigenfunctions of its isospectral eigenvalue problem. The dispersionless limit of the CH equation and its resulting peakon solutions are examined by using an asymptotic expansion in the dispersion parameter. The peakon solutions of the dispersionless CH equation in one dimension are shown to generalize in higher dimensions to peakon wave-front solutions of the EPDiff equation whose associated momentum is supported on smoothly embedded subspaces. The Eulerian representations of the singular solutions of both CH and EPDiff are given by the (cotangent-lift) momentum maps arising from the left action of the diffeomorphisms on smoothly embedded subspaces.

¹Department of Mathematics, Imperial College London. London SW7 2AZ, UK. d.holm@imperial.ac.uk, r.ivanov@imperial.ac.uk

²School of Mathematical Sciences, Dublin Institute of Technology, Kevin Street, Dublin 8, Ireland, rivanov@dit.ie

Contents

1	Shallow water background for the CH equation	2
2	Soliton Solutions of CH equation from Inverse Scattering2.1Inverse scattering for the KdV equation	5 5 9 10
3	Momentum map formulation with action-angle variables	11
4	Peakons 4.1 Peakons: the singular solution ansatz 4.2 Integrable peakon dynamics of CH	12 12 13
5	Peakon limit of the CH soliton solutions	15
6	Superintegrability of the peakon system	17
7	Other singular solutions: the dispersionless b -equation	18
8	Euler-Poincaré theory in higher dimensions	19
9	Three open problems	24

1 Shallow water background for the CH equation

Euler's equations for irrotational incompressible ideal fluid motion under gravity with a free surface have an asymptotic expansion for shallow water waves that contains two small parameters, ϵ and δ^2 , with ordering $\epsilon \geq \delta^2$. These small parameters are $\epsilon = a/h_0$ (the ratio of wave amplitude to mean depth) and $\delta^2 = (h_0/l_x)^2$ (the squared ratio of mean depth to horizontal length, or wavelength). Euler's equations are made non-dimensional by introducing $x = l_x x'$ for horizontal position, $z = h_0 z'$ for vertical position, $t = (l_x/c_0)t'$ for time, $\eta = a\eta'$ for surface elevation and $\varphi = (gl_x a/c_0)\varphi'$ for velocity potential, where $c_0 = \sqrt{gh_0}$ is the mean wave speed and g is the constant gravity. The quantity $\sigma = \sigma'/(h_0\rho c_0^2)$ is the dimensionless Bond number, in which ρ is the mass density of the fluid and σ' is its surface tension, both of which are taken to be constants. After dropping primes, this asymptotic expansion yields the nondimensional Korteweg-de Vries (KdV) equation for the horizontal velocity variable $u = \varphi_x(x, t)$ at *linear* order in the small dimensionless ratios ϵ and δ^2 , as the left hand side of

$$u_t + u_x + \frac{3\epsilon}{2}uu_x + \frac{\delta^2}{6}(1 - 3\sigma)u_{xxx} = O(\epsilon\delta^2).$$
(1)

Here, partial derivatives are denoted using subscripts, and boundary conditions are u = 0 and $u_x = 0$ at spatial infinity on the real line. The famous $sech^2(x - ct)$ traveling wave solutions (the solitons) for KdV (1) arise in a balance between its (weakly) nonlinear steepening and its third-order linear dispersion, when the quadratic terms in ϵ and δ^2 on its right hand side are neglected.

On the right hand side of equation (1), a normal form transformation due to Kodama [40] has been used to remove the other possible quadratic terms of order $O(\epsilon^2)$ and $O(\delta^4)$. The remaining quadratic correction terms in the KdV equation (1) may be collected at order $O(\epsilon\delta^2)$. These terms may be expressed, after introducing a "momentum variable,"

$$m = u - \nu \delta^2 u_{xx} \,, \tag{2}$$

and neglecting terms of *cubic* order in ϵ and δ^2 , as

$$m_t + m_x + \frac{\epsilon}{2}(um_x + b\,mu_x) + \frac{\delta^2}{6}(1 - 3\sigma)u_{xxx} = 0\,.$$
(3)

In the momentum variable $m = u - \nu \delta^2 u_{xx}$, the parameter ν is given by [21]

$$\nu = \frac{19 - 30\sigma - 45\sigma^2}{60(1 - 3\sigma)}.$$
(4)

Thus, the effects of δ^2 -dispersion also enter the nonlinear terms. After restoring dimensions in equation (3) and rescaling velocity u by (b+1), the following "b-equation" emerges,

$$m_t + c_0 m_x + u m_x + b \, m u_x + \Gamma u_{xxx} = 0 \,, \tag{5}$$

where $m = u - \alpha^2 u_{xx}$ is the dimensional momentum variable, and the constants α^2 and Γ/c_0 are squares of length scales. When $\alpha^2 \to 0$, one recovers KdV from the *b*-equation (5), up to a rescaling of velocity. Any value of the parameter $b \neq -1$ may be achieved in equation (5) by an appropriate Kodama transformation [21].

As we have emphasized, the values of the coefficients in the asymptotic analysis of shallow water waves at quadratic order in their two small parameters only hold, modulo the Kodama normalform transformations. Hence, these transformations may be used to advance the analysis and thereby gain insight, by optimizing the choices of these coefficients. The freedom introduced by the Kodama transformations among asymptotically equivalent equations at quadratic order in ϵ and δ^2 also helps to answer the perennial question, "Why are integrable equations so ubiquitous when one uses asymptotics in modeling?" Namely, there may be special values of the free parameters in the normal-form transformations of the asymptotics that allow integrability.

Integrable cases of the *b*-equation (5). The cases b = 2 and b = 3 are special values. For these values, the *b*-equation becomes completely integrable as a Hamiltonian system. For b = 2, equation (5) specializes to the integrable CH equation of Camassa and Holm [9]. The case b = 3 in (5) recovers the integrable DP equation of Degasperis and Procesi [19]. These two cases exhaust the integrable candidates for (5), as was shown using Painlevé analysis. The *b*-family of equations (5) was also shown in [46] to admit the symmetry conditions necessary for integrability, only in the cases b = 2 for CH and b = 3 for DP.

The *b*-equation (5) with b = 2 was first derived in Camassa and Holm [9] by using asymptotic expansions directly in the Hamiltonian for Euler's equations governing inviscid incompressible flow in the shallow water regime. In this analysis, the CH equation was shown to be bi-Hamiltonian and thereby was found to be completely integrable by the inverse scattering transform (IST) on the real line. This development of IST for CH equation (5) with b = 2 is discussed in Section 2.

Camassa and Holm [9] also discovered the remarkable peaked soliton (peakon) solutions of (58,59) for the CH equation on the real line, given by (5) in the case b = 2. The peakons arise as

solutions of (5), when $c_0 = 0$ and $\Gamma = 0$ in the absence of linear dispersion. Peakons move at a speed equal to their maximum height, at which they have a sharp peak (jump in derivative). The single peakon solution is

$$u(x,t) = ce^{-|x-ct|/\alpha},\tag{6}$$

Unlike the KdV soliton, the peakon speed is independent of its width (α). Periodic peakon solutions of CH were treated in Alber *et al.* [3]. There, the sharp peaks of periodic peakons were associated with billiards reflecting at the boundary of an elliptical domain. These billiard solutions for the periodic peakons arise from geodesic motion on a tri-axial ellipsoid, in the limit that one of its axes shrinks to zero length.

The CH equation was found after its derivation as a shallow water equation in [9] to fit into a class of integrable equations derived previously by using hereditary symmetries in Fokas and Fuchssteiner [23]. See Fuchssteiner [25] for an insightful history of how the shallow water equation (5) in the integrable case with b = 2 relates to the mathematical theory of hereditary symmetries.

Equation (5) with b = 2 was recently re-derived as a shallow water equation by using asymptotic methods in three different approaches in Dullin *et al.* [21], in Fokas and Liu [24] and also in Johnson [37]. These three derivations all used different variants of the method of asymptotic expansions for shallow water waves in the absence of surface tension. Only the derivation in Dullin *et al.* [21] took advantage of the parametric freedom in the Kodama normal-form transformations of the asymptotic expansion results at quadratic order.

The effects of the parameter b on the solutions of equation (5) were investigated in Holm and Staley [35], where b was treated as a bifurcation parameter, in the limiting case when the linear dispersion coefficients are set to $c_0 = 0$ and $\Gamma = 0$. This limiting case allows a variety of special solutions for different ranges of the values of b, in which the two nonlinear terms in equation (5) balance each other in the *absence* of linear dispersion.

Since its first appearance in [9], the CH equation has been the centre of a confluence of scientific endeavors that includes water waves, integrable systems, PDE analysis, asymptotics, geometry and Lie groups. This confluence has led to a continued interest and opportunities for contributions from many different fields in mathematics. The interest in CH solutions may be measured by noticing that the entry 'peakon' has acquired many thousands of *Google hits*.

Plan of the paper

This paper aims to review some of the geometric highlights of recent work on the CH equation. It is certainly not exhaustive. It mainly focuses on comparing the soliton theory for smooth CH solutions with the peakon theory for its singular solutions in the dispersionless limit. Section 2 briefly explains the application of the method of the inverse scattering transform (IST) for obtaining the soliton solutions of CH equation. The set of scattering data is introduced and the formulation of the inverse scattering as a Riemann-Hilbert problem is outlined. The solution is expressed via the scattering data in a form that admits the peakon limit in the sections that follow. In Section 3 the map between the action-angle variables (expressed via the scattering data) and the momentum of the CH solution is formulated as a momentum map from the symplectic action-angle variables to the dual of the Lie algebra of smooth vector fields on the real line. This is a Poisson map, but the noncanonical bi-Hamiltonian structure of the CH equation which led to the discovery of its isospectral problem in [9] and its geometrical significance as geodesic flow on the diffeomorphism group are not discussed here. In Section 4 we introduce the peakons as singular solutions that appear in dispersionless limit. The N-peakon solution is governed by a finite-dimensional integrable

dynamical system. Section 5 presents the multi-peakon solution as a limiting case of the CH multi-soliton solution and points out the similarity between the dynamics of the peakon system and the well known Toda lattice. In Section 6 we comment briefly on the existence of additional integrals of motion of the peakon system, a property known as *superintegrability*. The two-peakon system is analysed explicitly. Section 7 presents the compacton and pulson solutions, which are singular solutions, similar to peakons that can also be represented by the singular momentum map for peakons discussed in [33]. In Section 8 we deal with generalizations of the CH equation in higher dimensions. These also allow for singular solutions. Although such multidimensional generalizations are unlikely to be integrable, numerical studies show that their solutions are stable and interact elastically in collisions. In Section 9 we comment briefly on some limitations of our present discussion and point out three open problems for further research.

2 Soliton Solutions of CH equation from Inverse Scattering

2.1 Inverse scattering for the KdV equation

One of the most significant results in the theory of nonlinear partial differential equations was the development of Gardner, Greene, Kruskal and Miura [28, 29] of a method for the exact solution of the initial-value problem for the KdV equation. Prior to their work the only known exact solutions of KdV were the travelling wave solutions. The idea is based on a representation of the equation $(u_t + 6uu_x + u_{xxx} = 0)$ as a compatibility condition of two eigenvalue problems with a time-independent spectral parameter λ :

$$\Psi_{xx} + u\Psi = \lambda\Psi \tag{7}$$

$$\Psi_t = (u_x + \gamma)\Psi - (4\lambda + 2u)\Psi_x \tag{8}$$

where γ is an arbitrary constant. The method is conceptually analogous in many ways to the Fourier transform method for solving linear equations. This method is now known as Inverse Scattering Transform (IST). It was recast by Lax [41] in a general framework that allows the IST to be used for solving other nonlinear PDEs. An important consequence is the relationship between the discrete eigenvalues and the characteristics of the solitons that emerge asymptotically. Another feature of soliton collisions is the preservation of soliton identities after the interaction (asymptotically in time). For example, collisions of KdV solitons only result in a phase shift from the positions where they would have been without the interaction.

The KdV was formulated as a completely integrable Hamiltonian system in a work by Faddeev and Zakharov [55], the Hamiltonian form was also noted by Gardner [27].

Reviews of IST may be found, for example, in Ablowitz et al. [1], Dubrovin *et al.* [20], Novikov *et al.* [47]. For discussions of other related bi-Hamiltonian equations, see [19].

2.2 Inverse scattering for CH solitons with dispersion

In this section we outline the application of the IST for the CH equation. We use the form with a linear dispersion term

$$u_t - u_{xxt} + 2\omega u_x + 3uu_x - 2u_x u_{xx} - uu_{xxx} = 0, (9)$$

where ω is a real constant. The equation in this form also appears as a model of unidirectional propagation of shallow water waves over a flat bottom [9, 37, 38] as well as that of axially symmetric waves in a hyperelastic rod [18]. It can be obtained from the *b*-equation (5) with b = 2 via Galilean transformation that removes the u_{xxx} term.

Equation (9) admits a Lax pair formulation [9]

$$\Psi_{xx} = \frac{1}{4} \left(1 - (m+\omega) \right) \Psi \tag{10}$$

$$\Psi_t = -\left(\frac{2}{\lambda} + u\right)\Psi_x + \frac{u_x}{2}\Psi + \gamma\Psi$$
(11)

where γ is an arbitrary constant. We will use this freedom for a proper normalization of the eigenfunctions.



Figure 1: A Gaussian initial condition for the CH equation breaks up into an ordered train of solitons as time evolves (the time direction being vertical). The soliton train eventually wraps around the periodic domain, thereby allowing the leading solitons to overtake the slower emergent solitons from behind in collisions that cause phase shifts, as discussed in [9].

In our further considerations m will be a Schwartz class function, $\omega > 0$ and $m(x, 0) + \omega > 0$. Then $m(x, t) + \omega > 0$ for all t [11]. Let us introduce a new spectral parameter k, such that

$$\lambda(k) = \frac{1}{\omega} \left(1 + 4k^2 \right). \tag{12}$$

The spectrum of the problem (10) under these conditions is described in [10]. The continuous spectrum in terms of k corresponds to k – real. The discrete spectrum (in the upper half plane) consists of finitely many points $k_n = i\kappa_n$, n = 1, ..., N where κ_n is real and $0 < \kappa_n < 1/2$.

For all real $k \neq 0$ a basis in the space of solutions of (10) can be introduced, $\psi(x, k)$ and $\bar{\psi}(x, \bar{k})$, fixed by its asymptotic when $x \to \infty$ [10]:

$$\psi(x,k) = e^{-ikx} + o(1), \qquad x \to \infty$$
(13)

Another basis can be introduced, $\varphi(x, k)$ and $\overline{\varphi}(x, \overline{k})$ fixed by its asymptotic when $x \to -\infty$:

$$\varphi(x,k) = e^{-ikx} + o(1), \qquad x \to -\infty \tag{14}$$

and the relation between the two bases is [13]

$$\varphi(x,k) = a(k)\psi(x,k) + b(k)\bar{\psi}(x,k).$$
(15)

where

$$|a(k)|^{2} - |b(k)|^{2} = 1.$$
(16)

The quantity $\mathcal{R}(k) = b(k)/a(k)$ is called the reflection coefficient, without any danger of confusing b(k) with the bifurcation parameter b introduced in (5). All of the required information about the scattering, i.e. a(k) and b(k), is provided by $\mathcal{R}(k)$ for

All of the required information about the scattering, i.e. a(k) and b(k), is provided by $\mathcal{R}(k)$ for k > 0 only [15]. It is sufficient to know $\mathcal{R}(k)$ only on the half line k > 0, since $\bar{a}(k) = a(-k)$, $\bar{b}(k) = b(-k)$ and thus $\mathcal{R}(-k) = \bar{\mathcal{R}}(k)$.

The constant γ in (11) can be chosen for each eigenfunction in such a way that a(k) does not depend on t and is a generating function of the integrals of motion [15]. At the points of the discrete spectrum, a(k) has simple zeroes [10], therefore φ and $\bar{\psi}$ are linearly dependent (15):

$$\varphi(x, i\kappa_n) = b_n \bar{\psi}(x, -i\kappa_n). \tag{17}$$

In other words, the discrete spectrum is simple, there is only one (real) eigenfunction $\varphi^{(n)}(x)$, corresponding to each eigenvalue $i\kappa_n$, and we can take this eigenfunction to be

$$\varphi^{(n)}(x) \equiv \varphi(x, i\kappa_n) \tag{18}$$

The asymptotic behaviour of $\varphi^{(n)}$, according to (14), (17) is

$$\varphi^{(n)}(x) = e^{\kappa_n x} + o(e^{\kappa_n x}), \qquad x \to -\infty; \tag{19}$$

$$\varphi^{(n)}(x) = b_n e^{-\kappa_n x} + o(e^{-\kappa_n x}), \qquad x \to \infty.$$
(20)

The sign of b_n obviously depends on the number of the zeroes of $\varphi^{(n)}$. Suppose that

$$0 < \kappa_1 < \kappa_2 < \ldots < \kappa_N < 1/2.$$

Then from the oscillation theorem for the Sturm-Liouville problem [7], $\varphi^{(n)}$ has exactly n-1 zeroes. Therefore

$$b_n = (-1)^{n-1} |b_n|. (22)$$

The set

$$\mathcal{S} \equiv \{ \mathcal{R}(k) \quad (k > 0), \quad \kappa_n, \quad R_n = b_n / ia'(i\kappa_n), \quad n = 1, \dots N \}$$
(23)

is called the *scattering data*. The Hamiltonians for the CH equation in terms of the scattering data are presented in [15]. The time evolution of the scattering data is [13]:

$$\mathcal{R}(k,t) = \mathcal{R}(k,0)e^{-\frac{4ik}{\lambda}t}.$$
(24)

$$R_n(t) = R_n(0) \exp\left(\frac{4\kappa_n}{\lambda_n}t\right), \qquad (25)$$

where $R_n(t)$ is always a positive quantity [13] and $\lambda_n = \lambda(\kappa_n)$.

The scattering coefficient a(k) is analytic for Im k > 0 with asymptotic [15]

$$e^{i\beta k}a(k) \to 1, \qquad |k| \to \infty.$$
 (26)

where β is a constant (integral of motion):

$$\beta = \int_{-\infty}^{\infty} \left(\sqrt{1 + \frac{m(x)}{\omega}} - 1 \right) dx.$$
(27)

The asymptotics of the Jost solutions (or, rather the following quantities, depending on the Jost solutions) for $|k| \to \infty$ have the form [13]

$$\underline{\psi}(x,k) \equiv \psi(x,k)e^{\frac{iky}{\sqrt{\omega}}} = X_0(x) + \frac{X_1(x)}{k} + \frac{X_2(x)}{k^2} + \dots,$$
(28)

$$\underline{\varphi}(x,k) \equiv \varphi(x,k)e^{ik(\frac{y}{\sqrt{\omega}}+\beta)} = X_0(x) + \frac{\dot{X}_1(x)}{k} + \frac{\dot{X}_2(x)}{k^2} + \dots$$
(29)

where $X_0(x) = [\omega/(m(x) + \omega)]^{1/4} > 0$,

$$y(x) = \kappa \left[x + \int_{\infty}^{x} (X_0^{-2}(\tilde{x}) - 1) d\tilde{x} \right].$$
(30)

Moreover, (28) is analytic for Im k < 0, (29) is analytic for Im k > 0 [13].

From (15) with (28) and (29) we obtain

$$\frac{\underline{\varphi}(x,k)}{e^{ik\beta}a(k)} = \underline{\psi}(x,k) + \mathcal{R}(k)\underline{\overline{\psi}}(x,k)e^{2iky(x)/\sqrt{\omega}}.$$
(31)

The function $\underline{\varphi}(x,k)/(e^{ik\beta}a(k))$ is analytic for Im k > 0, $\underline{\psi}(x,k)$ is analytic for Im k < 0. Thus, (31) represents an additive **Riemann-Hilbert Problem** (**RHP**) with a jump on the real line, given by $\mathcal{R}(k)\overline{\psi}(x,k)e^{2iky(x)/\sqrt{\omega}}$.

The solution of this RHP follows a standard pattern, see [13] for more details. The solution simplifies considerably in the case $\mathcal{R}(k) \equiv 0$. This is the N-soliton solution:

$$\underline{\psi}(x,k) = X_0(x) - i \sum_{n=1}^{N} \frac{R_n(t) e^{-2\kappa_n y/\sqrt{\omega}} \underline{\psi}(x, -i\kappa_n)}{i\kappa_n - k}, \quad \text{Im } k < 0,$$
(32)

$$\frac{\underline{\varphi}(x,k)}{e^{ik\beta}a(k)} = X_0(x) - i\sum_{n=1}^N \frac{R_n(t)e^{-2\kappa_n y/\sqrt{\omega}}\underline{\psi}(x,-i\kappa_n)}{i\kappa_n - k}, \quad \text{Im } k > 0.$$
(33)

From (32) one has a linear system for the quantities $\psi(x, -i\kappa_n, t)$ with a solution

$$\underline{\psi}(x, -i\kappa_n, t) = X_0(x) \sum_{p=1}^N A_{np}^{-1}[y, t], \quad n = 1, \dots, N.$$
(34)

where

$$A_{pn}[y,t] \equiv \delta_{pn} + \frac{R_n(t)e^{-2\kappa_n y/\sqrt{\omega}}}{\kappa_p + \kappa_n}.$$
(35)

Taking k = -i/2 in (32) produces

$$e^{-\frac{1}{2}(x-\frac{y}{\sqrt{\omega}})} \equiv \underline{\psi}(x,-i/2) = X_0(x) \left(1 - \sum_{n,p=1}^N \frac{R_n(t)e^{-2\kappa_n y/\sqrt{\omega}}}{\kappa_n + \frac{1}{2}} A_{np}^{-1}[y,t] \right).$$
(36)

The substitution $\kappa = i/2$ in (33) with a(i/2) = 1 gives

$$e^{\frac{1}{2}(x-\frac{y}{\sqrt{\omega}})} \equiv \frac{\underline{\varphi}(x,i/2)}{e^{-\beta/2}a(i/2)} = X_0(x) \left(1 - \sum_{n,p=1}^N \frac{R_n(t)e^{-2\kappa_n y/\sqrt{\omega}}}{\kappa_n - \frac{1}{2}} A_{np}^{-1}[y,t]\right).$$
(37)

From (36) and (37) there follows a parametric representation

$$x = X(y,t) \equiv \frac{y}{\sqrt{\omega}} + \ln \frac{f_+}{f_-}, \tag{38}$$

$$f_{\pm} \equiv 1 - \sum_{n,p=1}^{N} \frac{R_n(t)e^{-2\kappa_n y/\sqrt{\omega}}}{\kappa_n \mp \frac{1}{2}} A_{np}^{-1}[y,t].$$
(39)

2.3 Parametric form of the dispersive CH soliton solution

From (30) and (38) one can compute the solution in parametric form

$$u(X(y,t),t) = X_t(y,t), \qquad x = X(y,t),$$
(40)

where X(y,t) is given in terms of the scattering data in (38), (39).

Upon introducing new notations

$$\xi_j = 2\kappa_j \left(-\frac{y}{\sqrt{\omega}} + \frac{2t}{\lambda_j} + x_{j0} \right)$$
(41)

$$x_{j0} = \frac{1}{2\kappa_j} \ln \frac{R_j(0)}{2\kappa_j} \tag{42}$$

$$\phi_j = \ln \frac{1 - 2\kappa_j}{1 + 2\kappa_j} \tag{43}$$

$$\gamma_{ij} = \ln\left(\frac{\kappa_i - \kappa_j}{\kappa_i + \kappa_j}\right)^2 \tag{44}$$

one can rewrite the expression for f_{\pm} (39) in the form [44, 48, 49, 50]

$$f_{\pm} \equiv \sum_{\mu=0,1} \exp\left[\sum_{i=1}^{N} \mu_i(\xi_i \mp \phi_i) + \sum_{1 \le i < j \le j} \mu_i \mu_j \gamma_{ij}\right].$$

$$(45)$$

The solution for m can be obtained from (40). First we notice that

$$\frac{\partial u(X(y,t),t)}{\partial y} = u_x(X(y,t),t)X_y.$$

and also

$$\frac{\partial u(X(y,t),t)}{\partial y} = \frac{\partial X_t}{\partial y} = X_{ty}.$$

Thus $u_x(X(y,t),t) = X_{ty}/X_y$. Similarly,

$$u_{xx}(X(y,t),t) = \frac{1}{X_y} \left(\frac{X_{ty}}{X_y}\right)_y$$

and

$$m(X(t,y),t) = u(X(t,y),t) - u_{xx}(X(t,y),t) = X_t - \frac{1}{X_y} \left(\frac{X_{ty}}{X_y}\right)_y;$$

$$m(x,t) = \int_{-\infty}^{\infty} P(y,t)\delta(x - X(y,t))dy$$
(46)

with
$$P(y,t) = X_t X_y - \left(\frac{X_{ty}}{X_y}\right)_y,$$
 (47)

$$u(x,t) = \frac{1}{2} \int_{-\infty}^{\infty} P(y,t) \exp(-|x - X(y,t)|) \mathrm{d}y.$$
 (48)

As we shall see in Section 8, this representation is also useful in the study of multidimensional solutions.

2.4 Relation to KdV hierarchy

The spectral problem (10) is gauge equivalent to a standard Sturm-Liouville problem, well known from the KdV hierarchy, cf. (7), with short notation $q = m + \omega$:

$$-\Phi_{yy} + U(y)\Phi = \mu\Phi, \qquad \mu = \frac{\lambda}{4} - \frac{1}{4\omega},$$

$$\Phi(y) = q^{1/4}\Psi, \qquad \frac{dy}{dx} = \sqrt{q},$$
(49)

$$U(y) = \frac{1}{4q(y)} + \frac{q_{yy}(y)}{4q(y)} - \frac{3q_y^2(y)}{16q^2(y)} - \frac{1}{4\omega}.$$
 (50)

Note that (49) leads to two possible expressions for the change of the variables in the Liouville transformation:

$$y = \sqrt{\omega}x + \int_{-\infty}^{x} (\sqrt{q(x')} - \sqrt{\omega}) dx' + \text{const},$$
 (51)

$$y = \sqrt{\omega}x + \int_{\infty}^{x} (\sqrt{q(x')} - \sqrt{\omega}) dx' + \text{const.}$$
 (52)

These two possibilities, (51), (52) are only consistent iff

$$\int_{-\infty}^{\infty} (\sqrt{q(x)} - \sqrt{\omega}) \mathrm{d}x = \text{constant},$$

which is always the case, since the integral under question is (up to a multiplier) the Casimir function β (27).

The matching of the CH hierarchy to KdV hierarchy requires solving the Ermakov-Pinney equation (50) [10], which is not straightforward and leads to the same solution (40) in parametric form.

3 Momentum map formulation with action-angle variables

The canonical Poisson brackets for the scattering data of the CH equation are computed in [15] where also the action-angle variables are expressed in terms of the scattering data. Let us consider the action variable for the N-soliton solution. (These considerations can be extended easily to the variables of the continuous spectrum.) The angle variable is $\Phi_n = \ln R_n(t)$; it is linear in time t and $\dot{\Phi}_n = 4\kappa_n/\lambda_n$. Let us introduce $\Lambda_n := 4\kappa_n/\lambda_n$ into Hamilton's principle $\delta S = 0$, with

$$S[u, \Phi_n, \Pi_n] = \int \left(\ell[u] + \sum_{n=1}^N \Pi_n \left(\dot{\Phi}_n - \Lambda_n[u] \right) \right) dt.$$

Here the Lagrange multiplier Π_n enforces the action-angle relation for the CH scattering data as a \mathbb{T}^N shift of the angles Φ_n at constant angular frequencies Λ_n , with $n = 1, \ldots, N$. The stationary variation is

$$0 = \delta S = \int \left(\left(\frac{\delta \ell}{\delta u} - \sum_{n=1}^{N} \prod_{n} \frac{\delta \Lambda_{n}}{\delta u} \right) \delta u + \sum_{n=1}^{N} \left(\dot{\Phi}_{n} - \Lambda_{n}[u] \right) \delta \prod_{n} - \sum_{n=1}^{N} \dot{\Pi}_{n} \delta \Phi_{n} \right) dt.$$

Since by definition $m = \delta \ell / \delta u$ is the momentum, $\delta S = 0$ implies the Eulerian representation

$$m = \frac{\delta \ell}{\delta u} = \sum_{n=1}^{N} \prod_{n} \frac{\delta \Lambda_{n}}{\delta u}, \quad \text{with}$$
(53)

$$\dot{\Phi}_n = \Lambda_n[u]$$
 and $\dot{\Pi}_n = 0.$ (54)

Relation (53) is the momentum map

$$(\Phi_n, \Pi_n) \in T^* \mathbb{T}^N \to m \in \mathfrak{X}^*$$

for the toral \mathbb{T}^N action (54) on the angles Φ_n at constant angular frequencies Λ_n . This momentum map from the action-angle scattering variables $T^*(\mathbb{T}^N)$ to the flow momentum $\mathfrak{X}^*(\mathbb{R})$ (dual to the smooth vector fields $\mathfrak{X}(\mathbb{R})$ on the real line) provides the Eulerian representation of the *N*soliton solution of CH in terms of the scattering data and squared eigenfunctions of its isospectral eigenvalue problem. Momentum maps for Hamiltonian dynamics are reviewed in [43], for example.

By using the spectral quantities of the N-soliton solution (recall: $\lambda_n = (1 - 4\kappa_n^2)/\omega$) one may express the variation of the spectrum with respect to the CH solution in terms of the squaredeigenfunctions of the isospectral problem as [15]

$$\frac{\delta\Lambda_n}{\delta m(x,t)} = \frac{(1+4\kappa_n^2)}{2\omega\kappa_n\lambda_n} R_n(t) [\bar{\psi}(x,-i\kappa_n,t)]^2,$$

in which $\psi(x, -i\kappa_n, t)$ is the eigenfunction that belongs to eigenvalue λ_n , see (17).

On the other hand, the expansion of u(x,t) over squares of eigenfunctions is given by [14]

$$u(x,t) = \sum_{n=1}^{N} \frac{4\kappa_n}{\omega \lambda_n^2} R_n(t) [\bar{\psi}(x,-i\kappa_n,t)]^2.$$

Consequently,

$$m(x,t) = \sum_{n=1}^{N} \frac{4\kappa_n}{\omega \lambda_n^2} R_n(t) (1-\partial^2) [\bar{\psi}(x,-i\kappa_n,t)]^2,$$

or

$$m(x,t) = \sum_{n=1}^{N} \prod_{n} J_n(x,t) \,$$

where Π_n and $J_n(x,t)$ denote explicitly

$$\Pi_n = \frac{8\kappa_n^2}{\lambda_n(1+4\kappa_n^2)} = \frac{2\Lambda_n\kappa_n}{1+4\kappa_n^2},$$
(55)

$$J_n(x,t) \equiv \frac{\delta \Lambda_n}{\delta u(x,t)}$$
(56)

$$= \frac{(1+4\kappa_n^2)}{2\omega\kappa_n\lambda_n}R_n(t)(1-\partial^2)[\bar{\psi}(x,-i\kappa_n,t)]^2.$$
(57)

Thus, the momentum map (53) from the action-angle variables under going dynamics (54) to the Eulerian representation of the momentum for the CH solution is expressed in terms of the scattering data and squared eigenfunctions of its N-soliton isospectral eigenvalue problem. Perhaps not unexpectedly, this momentum map may be applied to the action-angle representation of the solution of any integrable Hamiltonian PDE.

4 Peakons

4.1 Peakons: the singular solution ansatz

Camassa and Holm [9] discovered the "peakon" solitary traveling wave solution for a shallow water wave,

$$u(x,t) = ce^{-|x-ct|/\alpha},\tag{58}$$

whose fluid velocity u is a function of position x on the real line and time t. The peakon traveling wave moves at a speed equal to its maximum height, at which it has a sharp peak (jump in derivative). Peakons are an emergent phenomenon, solving the initial value problem for a partial differential equation derived by an asymptotic expansion of Euler's equations using the small parameters of shallow water dynamics. Peakons are *nonanalytic* solitons, which superpose as

$$u(x,t) = \frac{1}{2} \sum_{a=1}^{N} p_a(t) e^{-|x-q_a(t)|/\alpha} =: \frac{1}{2} \sum_{a=1}^{N} p_a(t) g(x-q_a(t))/\alpha),$$
(59)

for sets $\{p\}$ and $\{q\}$ satisfying canonical Hamiltonian dynamics. Peakons arise for shallow water waves in the limit of zero linear dispersion in one dimension. Peakons satisfy a partial differential equation (PDE) arising from Hamilton's principle for geodesic motion on the smooth invertible maps (diffeomorphisms) with respect to the H^1 Sobolev norm of the fluid velocity. Peakons generalize to higher dimensions, as well. We explain how peakons were derived in the context of shallow water asymptotics and describe some of their remarkable mathematical properties.

Peakons were first found as singular soliton solutions of the completely integrable CH equation. This is equation (5) with b = 2, now rewritten in terms of the velocity, as

$$u_t + c_0 u_x + 3u u_x + \Gamma u_{xxx} = \alpha^2 (u_{xxt} + 2u_x u_{xx} + u u_{xxx}).$$
(60)

Peakons were found in [9] to arise in the absence of linear dispersion. That is, they arise when $c_0 = 0$ and $\Gamma = 0$ in CH (60). Specifically, peakons are the individual terms in the peaked *N*-soliton solution of CH (60) for its velocity, in the absence of linear dispersion. Each term in the sum (59) is a solition with a sharp peak at its maximum. Hence, the name "peakon." Expressed using its momentum, $m = (1 - \alpha^2 \partial_x^2)u$, the peakon velocity solution (59) of dispersionless CH becomes a sum over a delta functions, supported on a set of points moving on the real line. Namely, the peakon velocity solution (59) implies

$$m(x,t) = \alpha \sum_{a=1}^{N} p_a(t)\delta(x - q_a(t)), \qquad (61)$$

because of the relation $(1 - \alpha^2 \partial_x^2) e^{-|x|/\alpha} = 2\alpha \delta(x)$. These solutions satisfy the *b*-equation (5) for any value of *b*, provided $c_0 = 0$ and $\Gamma = 0$. As we shall discuss later, the peakon momentum relation (61) is again a momentum map.

Thus, peakons are singular momentum solutions of the dispersionless b-equation, although they are not stable for every value of b. From numerical simulations [35], peakons are conjectured to be stable for b > 1. In the integrable cases b = 2 for CH and b = 3 for DP, peakons are stable singular soliton solutions. The spatial velocity profile $e^{-|x|/\alpha}/(2\alpha)$ of each separate peakon in (59) is the Green's function for the Helmholtz operator on the real line, with vanishing boundary conditions at spatial infinity. Unlike the KdV soliton, whose speed and width are related, the width of the peakon profile is set by its Green's function, independently of its speed.

4.2 Integrable peakon dynamics of CH

Substituting the peakon solution ansatz (59) and (61) into the dispersionless CH equation,

$$m_t + um_x + 2mu_x = 0$$
, with $m = u - \alpha^2 u_{xx}$, (62)

yields Hamilton's canonical equations for the dynamics of the discrete set of peakon parameters $p_a(t)$ and $q_a(t)$,

$$\dot{q}_a(t) = \frac{\partial h_N}{\partial p_a} \quad \text{and} \quad \dot{p}_a(t) = -\frac{\partial h_N}{\partial q_a},$$
(63)

for a = 1, 2, ..., N, with Hamiltonian given by [9],

$$h_N = \frac{1}{4} \sum_{a,b=1}^N p_a \, p_b \, e^{-|q_a - q_b|/\alpha} \,. \tag{64}$$

Or explicitly,

$$\dot{q}_a = \frac{1}{2} \sum_{b=1}^{N} p_b e^{-|q_a - q_b|/\alpha}$$
(65)

$$\dot{p}_{a} = \frac{p_{a}}{2\alpha} \sum_{b=1}^{N} p_{b} e^{-|q_{a}-q_{b}|/\alpha} \operatorname{sgn}(q_{a}-q_{b})$$
(66)

Thus, one finds that the points $x = q_a(t)$ in the peakon solution (59) move with the flow of the fluid velocity u at those points, since $u(q_a(t), t) = \dot{q}_a(t)$. This means the $q_a(t)$ are Lagrangian coordinates. Moreover, the singular momentum solution (61) is the Lagrange-to-Euler map for an invariant manifold of the dispersionless CH equation (62). On this finite-dimensional invariant manifold for the partial differential equation (62), the dynamics is canonically Hamiltonian.

With Hamiltonian (64), the canonical equations (63) for the 2N canonically conjugate peakon parameters $p_a(t)$ and $q_a(t)$ were interpreted in [9] as describing geodesic motion on the N-dimensional Riemannian manifold whose co-metric is $g^{ab}(\{q\}) = e^{-|q_a-q_b|/\alpha}$. Moreover, the canonical geodesic equations arising from Hamiltonian (64) comprise an integrable system for any number of peakons N. This integrable system was studied in [9] for solutions on the real line, and in [3, 16] and references therein, for spatially periodic solutions.

The integrals generated by the action variables in terms of the coordinates can be recovered as $tr(L^k)$, where L is the Lax operator for the peakon system [9].

Being a completely integrable Hamiltonian soliton equation, the continuum CH equation (60) has an associated isospectral eigenvalue problem, discovered in [9] for any values of its dispersion parameters c_0 and Γ . Remarkably, when $c_0 = 0$ and $\Gamma = 0$, this isospectral eigenvalue problem has a purely *discrete* spectrum. Moreover, in this case, each discrete eigenvalue corresponds precisely to the time-asymptotic velocity of a peakon. This discreteness of the CH isospectrum in the absence of linear dispersion implies that *only* the singular peakon solutions (61) emerge asymptotically in time, in the solution of the initial value problem for the dispersionless CH equation (62). This is borne out in numerical simulations of the dispersionless CH equation (62), starting from a smooth initial distribution of velocity [26, 35].

Figure 2 shows the emergence of peakons from an initially Gaussian velocity distribution and their subsequent elastic collisions in a periodic one-dimensional domain. This figure demonstrates that singular solutions dominate the initial value problem and, thus, that it is imperative to go beyond smooth solutions for the CH equation; the situation is similar for the EPDiff equation.

Peakons as mechanical systems. Governed by canonical Hamiltonian equations, each N-peakon solution can be associated with a mechanical system of moving particles. Calogero et al. [8] further extended the class of mechanical systems of this type. The r-matrix approach was applied to the Lax pair formulation of the N-peakon system for CH by Ragnisco and Br-uschi [52], who also pointed out the connection of this system with the classical Toda lattice. A discrete version of the Adler-Kostant-Symes factorization method was used by Suris [53] to study a discretization of the peakon lattice, realized as a discrete integrable system on a certain Poisson submanifold of gl(N) equipped with an r-matrix Poisson bracket. Beals *et al.* [6] used the Stieltjes theorem on continued fractions and the classical moment problem for studying multi-peakon solutions of the CH equation. Generalized peakon systems are described for any simple Lie algebra by Alber *et al.* [3].



Figure 2: A Gaussian initial condition for the CH equation breaks up into an ordered train of peakons as time evolves (the time direction being vertical). The peakon train eventually wraps around the periodic domain, thereby allowing the leading peakons to overtake the slower emergent peakons from behind in collisions that cause phase shifts, as discussed in [9].

5 Peakon limit of the CH soliton solutions

The limit $\omega \to 0$ in the N-soliton solution u(x, t) produces the N-peakon solution (59). The limiting procedure is described in detail in [45]. Due to (12) one can write for the discrete eigenvalues $k_n = i\kappa_n$

$$2\kappa_j = (1 - \omega\lambda_j)^{1/2} = 1 - \frac{1}{2}\omega\lambda_j + \dots$$
 (67)

The solution (38) depends explicitly on κ_j (67) and the limit can be computed with (67) by taking $\omega \to 0$ and keeping the eigenvalue λ_j constant. The result is the expression (59) with

$$p_i = \frac{4D_{N-i+1}^{(0)}D_{N-i}^{(2)}}{D_{N-i+1}^{(1)}D_{N-i}^{(1)}}, \qquad (i = 1, 2, ..., N),$$
(68)

$$q_i = \alpha \ln \left[\frac{2D_{N-i+1}^{(0)}}{D_{N-i}^{(2)}} \right], \qquad (i = 1, 2, ..., N),$$
(69)

where

$$D_n^{(m)} = \sum_{1 \le i_1 < i_2 < \dots < i_n \le N} \Delta_n(i_1, i_2, \dots, i_n) (\lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_n})^m R_{i_1} R_{i_2} \dots R_{i_n},$$

$$n = 1, 2, \dots, N.$$

$$\Delta_n(i_1, i_2, ..., i_n) = \prod_{1 \le l < m \le n} (\lambda_{i_l} - \lambda_{i_m})^2, \qquad (n \ge 2),$$
(70)

$$R_i(t) = R_i(0)e^{\frac{2}{\lambda_i}t}, \qquad (x_{i0} = \ln R_i(0)).$$
 (71)

The quantities $D_n^{(m)}$ are called **Hankel determinants**. By definition $D_0^{(m)} = 1$. In general, Hankel determinant is a determinant of $n \times n$ matrix of the form $D_n^{(m)} \equiv \det(a_{ij}^{(m)})$ where a_{ij} are elements of a sequence, i.e.

$$a_{ij}^{(m)} = A_{i+j+m-2}$$

In this particular case

$$A_l = \sum_{i=1}^N \lambda_i^l R_i(t).$$
(72)

Similarity of peakon lattice and Toda lattice Hankel determinants appear in the solutions of other integrable systems, e.g. Toda lattice, e.g. see [31]. The Toda equation [54]

 $\frac{\mathrm{d}p_n}{\mathrm{d}t} = e^{q_{n+1}-q_n} - e^{q_n-q_{n-1}},$ $\frac{\mathrm{d}q_n}{\mathrm{d}t} = p_n, \quad n \in \mathbb{Z}$

is one of the most important integrable systems. We have a finite chain with N nodes under the fixed ends boundary conditions $q_0 = -q_{N+1} = \infty$. The solution is in the form

$$q_n(t) = q_1(0) + \ln \frac{D_n^{(0)}}{D_{n-1}^{(0)}}, \tag{73}$$

$$q_1(0) = -\frac{1}{N} \ln D_N^{(0)}(0)$$
 is a constant. (74)

The Hankel determinants are obtained by a similar sequence,

$$A_l = \sum_{i=1}^N \lambda_i^l R_i(t), \tag{75}$$

$$R_i(t) = R_i(0)e^{-\lambda_i t}, (76)$$

where λ_i are N different constants (eigenvalues of the Lax matrix) and the quantities $R_i(0)$, $i = 1, \ldots, N$ represent another set N of constants.

Due to its simple form, the N-peakon solution can be used as an approximation of the N-soliton CH solution when the dispersion term is small (and the term $2\omega u_x$ can be neglected). Similarly, the Toda chain with complex dynamical variables (the so-called Complex Toda Chain – CTC) provides an approximation for the N-soliton solution of the Nonlinear Schrödinger Equation (NLS)

$$iu_t + \frac{1}{2}u_{xx} + |u|^2 u = 0,$$

see [32, 30, 31] for more details. Such an approximation is called adiabatic approximation and means that the N-soliton solution consists of N well separated solitons

$$u(x,t) \approx \sum_{k=1}^{N} \frac{2\nu_k e^{i[2\mu_k(x-\xi_k(t)+\delta_k(t)]]}}{\cosh(2\nu_k(x-\xi_k(t)))},$$

i.e. the overlap between the solitons is small. The variables $q_n(t)$ of the CTC are related to the NLS solitons parameters by

$$q_k(t) = -2\nu_0\xi_k(t) + i(2\mu_0\xi_k(t) - \delta_k(t)) + \text{const}$$

where ξ_k , δ_k , $\mu_k = \frac{1}{2}\dot{\xi}_k$ and $\nu_k = (\frac{1}{2}\dot{\delta}_k - \mu_k^2)^{1/2}$ characterize the center-of-mass position, the phase, velocity and amplitude respectively of the k-th soliton in the chain; ν_0 and μ_0 are the average amplitude and velocity of the soliton train. ξ_k and δ_k can be obtained as the real and imaginary parts of $q_k(t)$. Such soliton trains and their asymptotic behavior appear to be important for the needs of soliton based fiber optics communications.

6 Superintegrability of the peakon system

Suppose we have an integrable system with 2N- dimensional phase space which in terms of the Action-Angle (canonical) variables can be represented as

$$\dot{\Lambda}_n = 0, \qquad \dot{\Phi}_n = \Lambda_n, \qquad n = 1, 2, \dots, N, \tag{77}$$

or, if there exists a bracket such that 1

$$\{\Phi_n, \Lambda_l\} = \frac{1}{2}\delta_{nl}, \qquad \{\Phi_n, \Phi_l\} = \{\Lambda_n, \Lambda_l\} = 0, \tag{78}$$

the system is Hamiltonian with a Hamiltonian

$$h_N = \Lambda_1^2 + \dots \Lambda_N^2. \tag{79}$$

The integrals Λ_n , n = 1, 2, ..., N are clearly in involution, which guarantees the integrability of the system. There is however another set of integrals:

$$I_{j} = (\Phi_{j} - \Phi_{j+1})(\Lambda_{1} + \ldots + \Lambda_{N}) - (\Lambda_{j} - \Lambda_{j+1})(\Phi_{1} + \ldots + \Phi_{N})$$

$$j = 1, 2, \dots, N - 1.$$
 (80)

If the 'action' variables Λ_n are all different, the set (80) is functionally independent from the set Λ_n , n = 1, 2, ..., N. In addition, the integrals (80) form another set of N-integrals in involution together with H. Due to the existence of two sets of functionally independent integrals in involution such systems are termed superintegrable.

An example of such system is Toda lattice, see the discussion in [2]. The peakon system is also superintegrable. The canonical variables in terms of the scattering data for CH equation can be used in the peakon limit: $\Lambda_n = 2/\lambda_n$, $\Phi_n = \ln R_n(t)$, the Hamiltonian (79) is also a peakon limit $(\omega \to 0)$ of the N-soliton Hamiltonian [15]

$$H_N(\omega) = \omega^2 \sum_{n=1}^{N} \left(\ln \frac{1 - 2\kappa_n}{1 + 2\kappa_n} + \frac{4\kappa_n (1 + 4\kappa_n^2)}{(1 - 4\kappa_n^2)^2} \right).$$

 $^{^{1}}$ The 1/2 coefficient in the definition of the bracket appears in order to match it to the Poisson bracket used in the CH Hamiltonian formulation.

The Poisson bracket for the CH peakon solution is

$$\{A, B\} \equiv -\int_{-\infty}^{\infty} \frac{\delta A}{\delta m} (m\partial + \partial m) \frac{\delta B}{\delta m} \mathrm{d}x.$$
(81)

and the scattering data satisfy (78) with respect to (81), see [15] for the details.

More interesting are the integrals (80). From (69) and (68) one can recover these integrals in coordinate form. For example, when N = 2 we have

$$I_1 = \ln \frac{\sqrt{J} + p_1 - p_2}{\sqrt{J} - p_1 + p_2} + \frac{\sqrt{J}}{p_1 + p_2} \left(\frac{q_1 + q_2}{\alpha} + \ln \frac{p_1}{p_2}\right),$$
(82)

where
$$J = (p_1 - p_2)^2 + 4p_1 p_2 e^{-|q_1 - q_2|/\alpha}$$
. (83)

Note that I_1 depends on both combinations $q_1 + q_2$ and $q_1 - q_2$ as well as the momentum variables. The Hamiltonian h_2 (which depends only on $q_1 - q_2$ and the momentum variables) and I_1 form a complete system of integrals in involution. The integration of the 2-peakon system with these integrals can be performed as follows. First, one can express q_1 and q_2 in terms of I_1 , h_2 and the momentum variables: $q_i = q_i(p_1, p_2, I_1, h_2)$. Next,

$$\dot{q}_i = \frac{\partial q_i}{\partial p_1} \dot{p}_1 + \frac{\partial q_i}{\partial p_2} \dot{p}_2.$$
(84)

The substitution of \dot{q}_i , \dot{p}_i from (65) (66) to (84) produces an algebraic equation that gives, say p_2 as a function of p_1 . Then (66) is an ODE for p_1 of the form $\dot{p}_1 = f(p_1, I_1, h_2)$. Clearly, from practical viewpoint it is much more convenient to work with the other system of integrals in involution: h_2 and the conserved momentum $P = p_1 + p_2$. Note that $J = 4h_2 - 3P^2$ is itself an integral.

7 Other singular solutions: the dispersionless b-equation

Pulsons: Generalizing the peakon solutions of the dispersionless *b*-equation for other Green's functions. The Hamiltonian h_N in equation (64) depends on the Green's function for the relation between velocity *u* and momentum *m*. However, the singular momentum solution ansatz (61) is *independent* of this Green's function. Thus, as discovered in Fringer and Holm [26], The singular momentum solution ansatz (61) for the dispersionless equation,

$$m_t + um_x + 2mu_x = 0, \quad with \quad u = g * m, \tag{85}$$

provides an invariant manifold on which canonical Hamiltonian dynamics occurs, for any choice of the Green's function g relating velocity u and momentum m by the convolution u = g * m.

The fluid velocity solutions corresponding to the singular momentum ansatz (61) for equation (85) are the *pulsons*. Pulsons are given by the sum over N velocity profiles determined by the Green's function g, as

$$u(x,t) = \sum_{a=1}^{N} p_a(t)g(x,q_a(t)).$$
(86)

Again for (85), the singular momentum ansatz (61) results in a finite-dimensional invariant manifold of solutions, whose dynamics is canonically Hamiltonian. The Hamiltonian for the canonical dynamics of the 2N parameters $p_a(t)$ and $q_a(t)$ in the "pulson" solutions (86) of equation (85) is

$$h_N = \frac{1}{2} \sum_{a,b=1}^N p_a \, p_b \, g(q_a, q_b) \,. \tag{87}$$

Again for the pulsons, the canonical equations for the invariant manifold of singular momentum solutions provide a phase-space description of geodesic motion, this time with respect to the cometric given by the Green's function g. Mathematical analysis and numerical results for the dynamics of these pulson solutions are given in [26]. These results describe how the collisions of pulsons (86) depend upon their shape.

Compactons in the $1/\alpha^2 \to 0$ **limit of CH.** As mentioned earlier, in the limit that $\alpha^2 \to 0$, the CH equation (60) becomes the KdV equation. In the opposite limit that $1/\alpha^2 \to 0$ CH becomes the Hunter-Zheng equation [36]

$$(u_t + uu_x)_{xx} = \frac{1}{2}(u_x^2)_x$$
 (Hunter-Zheng)

This equation has "compacton" solutions, whose collision dynamics was studied numerically and put into the present context in [26]. The corresponding Green's function satisfies $-\partial_x^2 g(x) = 2\delta(x)$, so it has the triangular shape, g(x) = 1 - |x| for |x| < 1, and vanishes otherwise, for $|x| \ge 1$. That is, the Green's function in this case has compact support; hence, the name "compactons" for these pulson solutions, which as a limit of the integrable CH equations are true solitons, solvable by IST.

Pulson solutions of the dispersionless b-equation. Holm and Staley [35] give the pulson solutions of the traveling wave problem and their elastic collision properties for the dispersionless b-equation,

$$m_t + um_x + b \, mu_x = 0, \quad \text{with} \quad u = g * m, \tag{88}$$

with any (symmetric) Green's function g and for any value of the parameter b. Numerically, pulsons and peakons are both found to be stable for b > 1, [35]. The reduction to noncanonical Hamiltonian dynamics for the invariant manifold of singular momentum solutions (61) of the other integrable case b = 3 with peakon Green's function $g(x, y) = e^{-|x-y|/\alpha}$ is found in [19].

8 Euler-Poincaré theory in higher dimensions

Generalizing the peakon solutions of the CH equation to higher dimensions. In [35], weakly nonlinear analysis and the assumption of columnar motion in the variational principle for Euler's equations were found to produce the two-dimensional generalization of the dispersionless CH equation (62). This generalization is the Euler-Poincaré (EP) equation [34] for the Lagrangian consisting of the kinetic energy,

$$\ell = \frac{1}{2} \int \left[|\mathbf{u}|^2 + \alpha^2 (\operatorname{div} \mathbf{u})^2 \right] dx dy, \qquad (89)$$

in which the fluid velocity \mathbf{u} is a two-dimensional vector. Evolution generated by kinetic energy in Hamilton's principle results in geodesic motion, with respect to the velocity norm $\|\mathbf{u}\|$, which is provided by the kinetic energy Lagrangian. For ideal incompressible fluids governed by Euler's equations, the importance of geodesic flow was recognized by Arnold [4] for the L^2 norm of the fluid velocity. The EP equation generated by any choice of kinetic energy norm without imposing incompressibility is called "EPDiff," for "Euler-Poincaré equation for geodesic motion on the diffeomorphisms." EPDiff is given by [34]

$$\left(\frac{\partial}{\partial t} + \mathbf{u} \cdot \nabla\right) \mathbf{m} + \nabla \mathbf{u}^T \cdot \mathbf{m} + \mathbf{m}(\operatorname{div} \mathbf{u}) = 0, \qquad (90)$$

with momentum density $\mathbf{m} = \delta \ell / \delta \mathbf{u}$, where $\ell = \frac{1}{2} ||\mathbf{u}||^2$ is given by the kinetic energy, which defines a norm in the fluid velocity $||\mathbf{u}||$, yet to be determined. By design, this equation has no contribution from either potential energy, or pressure. It conserves the velocity norm $||\mathbf{u}||$ given by the kinetic energy. Its evolution describes geodesic motion on the diffeomorphisms with respect to this norm [34]. An alternative way of writing the EPDiff equation (90) in either two, or three dimensions is,

$$\frac{\partial}{\partial t}\mathbf{m} - \mathbf{u} \times \operatorname{curl} \mathbf{m} + \nabla(\mathbf{u} \cdot \mathbf{m}) + \mathbf{m}(\operatorname{div} \mathbf{u}) = 0.$$
(91)

This form of EPDiff involves all three differential operators, curl, gradient and divergence. For the kinetic energy Lagrangian ℓ given in (89), which is a norm for *irrotational* flow (with curl $\mathbf{u} = 0$), we have the EPDiff equation (90) with momentum $\mathbf{m} = \delta \ell / \delta \mathbf{u} = \mathbf{u} - \alpha^2 \nabla (\text{div } \mathbf{u})$.

EPDiff (90) may also be written intrinsically as

$$\frac{\partial}{\partial t} \frac{\delta \ell}{\delta \mathbf{u}} = -\operatorname{ad}_{\mathbf{u}}^* \frac{\delta \ell}{\delta \mathbf{u}}, \qquad (92)$$

where ad^* is the L^2 dual of the ad-operation (commutator) for vector fields. See [5, 43] for additional discussions of the beautiful geometry underlying this equation.

Reduction to the dispersionless CH equation in 1D. In one dimension, the EPDiff equation (90-92) with Lagrangian ℓ given in (89) simplifies to the dispersionless CH equation (62). The dispersionless limit of the CH equation appears, because we have ignored potential energy and pressure.

Strengthening the kinetic energy norm to allow for circulation. The kinetic energy Lagrangian (89) is a norm for irrotational flow, with curl $\mathbf{u} = 0$. However, inclusion of rotational flow requires the kinetic energy norm to be strengthened to the H^1_{α} norm of the velocity, defined as

$$\ell = \frac{1}{2} \int \left[|\mathbf{u}|^2 + \alpha^2 (\operatorname{div} \mathbf{u})^2 + \alpha^2 (\operatorname{curl} \mathbf{u})^2 \right] dx dy$$
$$= \frac{1}{2} \int \left[|\mathbf{u}|^2 + \alpha^2 |\nabla \mathbf{u}|^2 \right] dx dy = \frac{1}{2} ||\mathbf{u}||^2_{H^1_{\alpha}}.$$
(93)

Here we assume boundary conditions that give no contributions upon integrating by parts. The corresponding EPDiff equation is (90) with $\mathbf{m} \equiv \delta \ell / \delta \mathbf{u} = \mathbf{u} - \alpha^2 \Delta \mathbf{u}$. This expression involves inversion of the familiar *Helmholtz* operator in the (nonlocal) relation between fluid velocity and momentum density. The H^1_{α} norm $\|\mathbf{u}\|^2_{H^1_{\alpha}}$ for the kinetic energy (93) also arises in three dimensions for turbulence modeling based on Lagrangian averaging and using Taylor's hypothesis that the turbulent fluctuations are "frozen" into the Lagrangian mean flow [22].

Generalizing the CH peakon solutions to n dimensions. Building on the peakon solutions (59) for the CH equation and the pulsons (86) for its generalization to other traveling-wave shapes in [26], Holm and Staley [35] introduced the following measure-valued singular momentum solution ansatz for the n-dimensional solutions of the EPDiff equation (90):²

$$\mathbf{m}(\mathbf{x},t) = \sum_{a=1}^{N} \int \mathbf{P}^{a}(s,t) \,\delta\big(\mathbf{x} - \mathbf{Q}^{a}(s,t)\big) ds.$$
(94)

These singular momentum solutions, called "diffeons," are vector density functions supported in \mathbb{R}^n on a set of N surfaces (or curves) of codimension (n-k) for $s \in \mathbb{R}^k$ with k < n. They may, for example, be supported on sets of points (vector peakons, k = 0), one-dimensional filaments (strings, k = 1), or two-dimensional surfaces (sheets, k = 2) in three dimensions.

Figure 3 shows the results for the EPDiff equation when a straight peakon segment of finite length is created initially moving rightward (East). Because of propagation along the segment in adjusting to the condition of zero speed at its ends and finite speed in its interior, the initially straight segment expands outward as it propagates and curves into a peakon "bubble."



Figure 3: A peakon segment of finite length is initially moving rightward (East). Because its speed vanishes at its ends and it has fully two-dimensional spatial dependence, it expands into a peakon "bubble" as it propagates. (The colors indicate speed: red is highest, yellow is less, blue low, grey zero.)

Figure 4 shows an initially straight segment whose velocity distribution is exponential in the transverse direction, but is wider than α for the peakon solution. This initial velocity distribution evolves under EPDiff to separate into a train of curved peakon "bubbles," each of width α . This example illustrates the emergent property of the peakon solutions in two dimensions. This phenomenon is observed in nature, for example, as trains of internal wave fronts in the south China Sea [42].

²These solutions represent smooth embeddings $\text{Emb}(\mathbb{R}^k, \mathbb{R}^n)$ with k < n. In contrast, the similar expression (46) for the solutions represent smooth functions $\mathbb{R} \to \mathbb{R}$.



Figure 4: An initially straight segment of velocity distribution whose exponential profile is wider than the width α for the peakon solution will break up into a train of curved peakon "bubbles," each of width α . This example illustrates the emergent property of the peakon solutions in two dimensions.

Substitution of the singular momentum solution ansatz (94) into the EPDiff equation (90) implies the following integro-partial-differential equations (IPDEs) for the evolution of the parameters $\{\mathbf{P}\}$ and $\{\mathbf{Q}\}$,

$$\frac{\partial}{\partial t} \mathbf{Q}^{a}(s,t) = \sum_{b=1}^{N} \int \mathbf{P}^{b}(s',t) G(\mathbf{Q}^{a}(s,t) - \mathbf{Q}^{b}(s',t)) ds',$$

$$\frac{\partial}{\partial t} \mathbf{P}^{a}(s,t) = -\sum_{b=1}^{N} \int \left(\mathbf{P}^{a}(s,t) \cdot \mathbf{P}^{b}(s',t) \right)$$

$$\frac{\partial}{\partial \mathbf{Q}^{a}(s,t)} G\left(\mathbf{Q}^{a}(s,t) - \mathbf{Q}^{b}(s',t) \right) ds'.$$
(95)

Importantly for the interpretation of these solutions, the coordinates $s \in \mathbb{R}^k$ turn out to be *Lagrangian* coordinates. The velocity field corresponding to the momentum solution ansatz (94) is given by

$$\mathbf{u}(\mathbf{x},t) = G * \mathbf{m}$$

= $\sum_{b=1}^{N} \int \mathbf{P}^{b}(s',t) G(\mathbf{x} - \mathbf{Q}^{b}(s',t)) ds',$ (96)



Figure 5: A single collision is shown involving reconnection as the faster peakon segment initially moving Southeast along the diagonal expands, curves and obliquely overtakes the slower peakon segment initially moving rightward (East). This reconnection illustrates one of the collision rules for the strongly two-dimensional EPDiff flow.

for $\mathbf{u} \in \mathbb{R}^n$. When evaluated along the curve $\mathbf{x} = \mathbf{Q}^a(s, t)$, this velocity satisfies,

$$\mathbf{u}(\mathbf{Q}^{a}(s,t),t) = \sum_{b=1}^{N} \int \mathbf{P}^{b}(s',t) G(\mathbf{Q}^{a}(s,t) - \mathbf{Q}^{b}(s',t)) ds'$$
$$= \frac{\partial \mathbf{Q}^{a}(s,t)}{\partial t}.$$
(97)

Consequently, the lower-dimensional support sets defined on $\mathbf{x} = \mathbf{Q}^{a}(s,t)$ and parameterized by coordinates $s \in \mathbb{R}^{k}$ move with the fluid velocity. This means the $s \in \mathbb{R}^{k}$ are Lagrangian coordinates. Moreover, equations (95) for the evolution of these support sets are canonical Hamiltonian equations,

$$\frac{\partial}{\partial t}\mathbf{Q}^{a}(s,t) = \frac{\delta H_{N}}{\delta \mathbf{P}^{a}}, \qquad \frac{\partial}{\partial t}\mathbf{P}^{a}(s,t) = -\frac{\delta H_{N}}{\delta \mathbf{Q}^{a}}.$$
(98)

The corresponding Hamiltonian function $H_N : (\mathbb{R}^n \times \mathbb{R}^n)^N \to \mathbb{R}$ is,

$$H_N = \frac{1}{2} \iint \sum_{a,b=1}^{N} \left(\mathbf{P}^a(s,t) \cdot \mathbf{P}^b(s',t) \right)$$
$$G\left(\mathbf{Q}^a(s,t), \mathbf{Q}^b(s',t) \right) \, ds \, ds' \,. \tag{99}$$

This is the Hamiltonian for geodesic motion on the cotangent bundle of a set of curves $\mathbf{Q}^{a}(s,t)$ with respect to the metric given by G. This dynamics was investigated numerically in [35] to which we refer for more details of the solution properties. One important result found *numerically*

in [35] is that only codimension-one singular momentum solutions appear to be stable under the evolution of the EPDiff equation. Thus, we have

Stability for codimension-one: the singular momentum solutions of EPDiff are stable, as points on the line (peakons), as curves in the plane (filaments, or wave fronts), or as surfaces in space (sheets).

Proving this stability result analytically remains an outstanding problem. The stability of peakons on the real line is proven in [17].

Reconnections in oblique overtaking collisions of peakon wave fronts. Figures 5 shows results of oblique wave front collisions producing reconnections for the EPDiff equation in two dimensions. Figure 5 shows a single oblique overtaking collision, as a faster expanding peakon wave front overtakes a slower one and reconnects with it at the collision point.

The peakon solution ansatz is a momentum map. As shown in [33], the solution expressions (61) in one dimension and (94) in higher dimensions may be interpreted as equivariant momentum maps, from the cotangent bundle of the smooth embeddings of lower dimensional sets $\mathbb{R}^s \subset \mathbb{R}^n$, to the dual of the Lie algebra of vector fields defined on these sets. (Momentum maps for Hamiltonian dynamics are reviewed in [43], for example.) The result that the singular solution ansatz (94) is a momentum map helps to organize the theory, to explain previous results and to suggest new avenues of exploration. This geometric feature underlies the remarkable reduction properties of the CH and EPDiff equations, and *explains* why they must be Lie-Poisson Hamiltonian equations. This is because of the general fact that equivariant momentum maps are Poisson maps. This geometric feature also underlies the singular momentum solution (94) and its associated velocity (96) which generalize the peakon solutions, both to higher dimensions and to arbitrary kinetic energy metrics. As we saw in Section 3, the soliton solution (53) is also a momentum map. This soliton momentum map may be expected to apply in the action-angle representation of the solution of any integrable Hamiltonian PDE. Its further properties will be studied in detail elsewhere.

9 Three open problems

(1.) Throughout this discussion the solutions u(x,t) were confined to be functions in the Schwartz class, $\omega > 0$. The situation when the condition $m(x,0) + \omega > 0$ on the initial data does not hold is more complicated and requires separate analysis [39, 10, 12]. In general, it leads to wave-breaking [12]. An attempt at developing the inverse scattering theory for this case has been made by Kaup [39], who suggested applying the inverse scattering approach separately in each interval where $m(x,t) + \omega$ is of the same sign. The problem however is how to join solutions that are valid in different intervals.

(2.) The peakon solution (68) was obtained from the soliton solution under the assumption $m(x, 0) + \omega > 0$. Thus, all p_k are of the same sign, since all the eigenvalues λ_n in this case are positive. However, one can formally use the same solution with eigenvalues of various signs to model p_k of various signs (mixture of peakons and 'anti-peakons') and thus to study peakon – anti-peakon interactions, see e.g. [51]. The result is that the multi-peakon interaction (including anti-peakons) in general decomposes into a sequence of pairwise collisions [26]. The collision of a single peakon-anti-peakon pair was studied already in [9]. When the eigenvalues are of mixed signs, the Hankel determinants in the denominator of (68) may develop singularities for finite values of

t. This 'peakon-breaking' phenomenon is apparently the analog of the wave-breaking mentioned earlier when $\omega \to 0$. This needs to be investigated further.

(3.) Stability for EPDiff singular momentum solutions, that is, proving their stability analytically, remains an outstanding problem.

Acknowledgments. DDH is grateful to R. Camassa, J. E. Marsden, T. S. Ratiu and A. Weinstein for their collaboration, help and inspiring discussions over the years. DDH gratefully acknowledges partial support by the Royal Society of London's Wolfson Award scheme. RII acknowledges funding from a Marie Curie Intra-European Fellowship. Both authors thank J. Percival and M. F. Staley for providing figures.

References

 Ablowitz M J and Clarkson P A 1991 Solitons, Nonlinear Evolution Equations and Inverse Scattering (Cambridge: Cambridge University Press)

Ablowitz M J and Segur H 1981 Solitons and the Inverse Scattering Transform (Philadelphia: SIAM)

- [2] Agrotis M, Damianou P and Christodulos S 2006 Physica A 365 235
- [3] Alber M S, Camassa R, Fedorov Y, Holm D D and Marsden J E 1999 Phys. Lett. A 264 171;
 Ibid 2001 Comm. Math. Phys. 221 197;
 Alber M S, Camassa R, Holm D D and Marsden J E 1994 Lett. Math. Phys. 32 137
 Alber M S, Camassa R and Gekhtman M 2000 CRM Proc. and Lecture Notes 25 1
- [4] Arnold V I 1966 Ann. Inst. Fourier, Grenoble 16 319
- [5] Arnold V I and Khesin B A 1998 Topological Methods in Hydrodynamics (New York: Springer)
- [6] Beals R, Sattinger D H and Szmigielski J 1999 Inverse Problems 15 L1 Ibid 2000 Adv. in Math. 154 229
 Ibid 2001 Commun. Pure and Appl. Math. 54 91
- Birkhoff G and Rota G-C 1969 Ordinary differential equations (Waltham: Blaisdell Publishing Company)
- [8] Calogero F 1995 Phys. Lett. A 201 306
 Calogero F and Francoise J-P 1996 J. Math. Phys. 37 2863
- [9] Camassa R and Holm D D 1993 Phys. Rev. Lett. **71** 1661; ArXiv: patt-sol/9305002;
 Camassa R, Holm D D and Hyman J M 1994 Adv. Appl. Mech. **31** 1
- [10] Constantin A 2001 Proc. R. Soc. Lond. A457 953
- [11] Constantin A 2005 J. Math. Phys. 46 023506
- [12] Constantin A and Escher J 1998 Acta Mathematica 181 229
- [13] Constantin A, Gerdjikov V S and Ivanov R I 2006 Inv. Problems 22 2197; arXiv:nlin/0603019v2 [nlin.SI]

- [14] Constantin A, Gerdjikov V S and Ivanov R I 2007 Inv. Problems 23 1565; arXiv:nlin.SI/0707.2048
- [15] Constantin A and Ivanov, R I 2006 Lett. Math. Phys. 76 93; arXiv: nlin.SI/0602049
- [16] Constantin A and McKean H P 1999 Commun. Pure Appl. Math. 52 949
- [17] Constantin A and Strauss W 2000 Commun. Pure Appl. Math. 53 603
- [18] Dai H-H 1998 Acta Mech. **127** 193
- [19] Degasperis A and Procesi M 1999 Asymptotic integrability, In: Symmetry and Perturbation Theory, ed A Degasperis and G Gaeta (Singapore: World Scientific) p 23;
 Degasperis A, Holm D D and Hone A N W 2002 Theoret. and Math. Phys. 133 1463
- [20] Dubrovin B A 1981 Russ. Math. Surv. 36 11;
 Dubrovin B A, Novikov S P and Krichever I M 1985 Integrable Systems. I In: Itogi Nauki i Tekhniki. Sovr. Probl. Mat. Fund. Naprav. 4 (Moscow: VINITI) (Russian); 1989 Encyclopaedia of Mathematical Sciences 4 (Berlin: Springer) (Engl. Transl.)
- [21] Dullin H R, Gottwald G A and Holm D D 2001 Phys. Rev. Lett. 87 194501-04; ArXiv: nlin.CD/0104004;
 Ibid 2003 Fluid Dyn. Res. 33 73;
 Ibid 2004 Physica D 190 1
- [22] Foias C, Holm D D and Titi E S 2001 Physica D 152 505
- [23] Fokas A S and Fuchssteiner B 1981 Nonlinear Anal. TMA 5 423
- [24] Fokas A S and Liu Q M 1996 Phys. Rev. Lett. 77 2347
- [25] Fuchssteiner B 1996 *Physica D* **95** 229
- [26] Fringer O and Holm D D 2001 Physica D 150 237; ArXiv:solv-int/9903007
- [27] Gardner C S 1971 J. Mathematical Phys. 12 1548
- [28] Gardner C S, Greene J M, Kruskal M D and Miura R M 1967 Phys. Rev. Lett. 19 1095
- [29] Gardner C S, Greene J M, Kruskal M D and Miura R M 1974 Comm. Pure Appl. Math. 27 97
- [30] Gerdjikov V S, Evstatiev E G and Ivanov R I 1998 J. Phys. A **31** 8221; ArXiv:solv-int/9712004
- [31] Gerdjikov V S, Evstatiev E G and Ivanov R I 2000 J. Phys. A 33 975
- [32] Gerdjikov V S, Kaup D J, Uzunov I M and Evstatiev E G 1996 Phys. Rev. Lett. 77 3943
- [33] Holm D D and Marsden J E 2004 Momentum Maps and Measure-valued Solutions (Peakons, Filaments and Sheets) for the EPDiff Equation. In: The Breadth of Symplectic and Poisson Geometry, Progr. Math. 232 ed J E Marsden and T S Ratiu (Boston: Birkhäuser) p 203
- [34] Holm D D, Marsden J E and Ratiu T S 1998 Adv. in Math. 137 1; ArXiv: chao-dyn/9801015;
 Ibid 1998 Phys. Rev. Lett. 349 4173

- [35] Holm D D and Staley M F 2003 Phys. Lett. A 308 437
 Ibid 2003 SIAM J. Appl. Dyn. Syst. 2 323
- [36] Hunter J K and Zheng Y 1994 Physica D 79 361
- [37] Johnson R S 2002 J. Fluid Mech. 455 63
- [38] Johnson R S 2003 Fluid Dynamics Research 33 97
- [39] Kaup D J 2006 Stud. Appl. Math. 117 149
- [40] Kodama Y 1985 Phys. Lett. A 107 245
 Ibid 1985 Phys. Lett. A 112 193
 Ibid 1987 Phys. Lett. A 123 276
- [41] Lax P D 1968 Comm. Pure Appl. Math. 21 467
- [42] Liu A K, Chang Y S, Hsu M-K and Liang N K 1998 J. Geophys. Res. 103 7995
- [43] Marsden J E and Ratiu T S 1999 Introduction to Mechanics and Symmetry, in Texts in Applied Mathematics 17 1994, Second Edition, 1999 (Springer-Verlag)
- [44] Matsuno Y 2005 J. Phys. Soc. Japan 74 1983; ArXiv:nlin.SI/0504055
- [45] Matsuno Y 2007 J. Phys. Soc. Japan **76** 034003; ArXiv:nlin/0701051
- [46] Mikhailov A V and Novikov V S 2002 J. Phys. A 35 4775
- [47] Novikov S P, Manakov S V, Pitaevskii L P and Zakharov V E 1984 Theory of solitons: the inverse scattering method (New York: Plenum) (translated from Russian)
- [48] Parker A 2004 Proc. R. Soc. Lond. A 460 2929
- [49] Parker A 2005 Proc. R. Soc. Lond. A 461 3611
- [50] Parker A 2005 Proc. R. Soc. Lond. A 461 3893
- [51] Parker A 2008 Chaos, solitons and Fractals **35** 220
- [52] Ragnisco O and Bruschi M 1996 Physica A 228 150
- [53] Suris Y B 1996 Phys. Lett. A **217** 321
- [54] Toda M 1967 J. Phys Soc. Japan 22 431
- [55] Zakharov V E and Faddeev L D 1971 Funkz. Anal. Prilož. 5 18 (Russian) 1971 Func. Anal. Appl. 5 280 (English translation)