Neutrino Masses and a TeV Scale Seesaw Mechanism

Wei Chao^{*}

Center for High Energy Physics, Peking university, Beijing 100871, China

Abstract

A simple extension of the Standard Model providing TeV scale seesaw mechanism is presented. Beside the Standard Model particles and right-handed Majorana neutrinos, the model contains a singly charged scalar, an extra Higgs doublet and three vector like singly charged fermions. In our model, Dirac neutrino mass matrix raises only at the loop level. Small but non-zero Majorana neutrino masses come from integrating out heavy Majorana neutrinos, which can be at the TeV scale. The phenomenologies of the model are investigated, including scalar mass spectrum, neutrino masses and mixings, lepton flavor violations, heavy neutrino magnetic moments as well as possible collider signatures of the model at the LHC.

^{*}Electronic address: chaow@pku.edu.cn

I. INTRODUCTION

The observation of neutrino oscillations [1-4] has revealed that neutrinos have small but non-zero masses and lepton flavors are mixed, which can not be accommodated in the Standard Model (SM) without introducing extra ingredients. As such, neutrino physics offer an exciting window into new physics beyond the SM. Perhaps the most attractive approach towards understanding the origin of small neutrino masses is using the dimensionfive weinberg operator [5]:

$$\frac{1}{4} \kappa_{gf} \overline{\ell_{Lc}^C}^g \varepsilon_{cd} \phi_d \ell_{Lb}^f \varepsilon_{ba} \phi_a + \text{h.c.} , \qquad (1)$$

which comes from integrating out new superheavy particles.

A simple way to obtain the operator in Eq. (1) is through the Type-I seesaw mechanism [6], in which three right-handed neutrinos with large Majorana masses are introduced to the SM. Then three active neutrinos may acquire tiny Majorana masses through the Type-I seesaw formula, i.e., the mass matrix of light neutrinos is given by $M_{\nu} = -M_D M_R^{-1} M_D^T$, where M_D is the Dirac mass matrix linking left-handed light neutrinos to right-handed heavy neutrinos and M_R is the mass matrix of heavy Majorana neutrinos. Actually, there are three tree-level seesaw scenarios (namely type-I, Type-II [7] and Type-III [8] seesaw mechanisms) and one loop-level seesaw scenario (namely Ma [9] model), which may lead to the effective operator in Eq. (1).

Although seesaw mechanisms can work naturally to generate Majorana neutrino masses, they lose direct testability on the experimental side. A direct test of seesaw mechanism would involve the detection of these heavy seesaw particles at a collider and the measurement of their Yukawa couplings with the electroweak doublets. In the canonical seesaw mechanism, heavy seesaw particles turn out to be too heavy, i.e., $10^{14\sim16}$ GeV, to be experimentally accessible. One straightforward way out is to lower the seesaw scale "by hand" down to the TeV scale, an energy frontier to be explored by the Large Hadron Collider (LHC). However this requires the structural cancellation between the Yukawa coupling texture and the heavy Majorana mass matrix, i.e. $M_D M_R^{-1} M_D^T \approx 0$ [10–15] at the tree level, and is thus unnatural!

To solve this unnaturalness problem, we propose a novel TeV-scale seesaw mechanism in this paper. The model includes, in addition to the SM fields and right-handed Majorana neutrinos, a charged scalar singlet, an extra Higgs doublet and three vector like singly charged fermions. Due to Z_2 discrete flavor symmetry, right-handed Majorana neutrinos don't couple to left-handed lepton doublets, such that Dirac mass matrix only raises at the loop level and is comparable with the charged lepton mass matrix. This drives down the seesaw scale to the TeV, and thus the model is detectable at the LHC.

The paper is organized as follows: In section II, we describe our model. Section III is devoted to investigate the phenomenologies of the model, including neutrino masses and mixings, lepton flavor violations, transition magnetic moments of heavy Majorana neutrinos as well as possible collider signatures. We conclude in Section IV. An alternative settings to the model is presented in appendix A.

II. THE MODEL

In our model, we extend the SM by introducing three right-handed Majorana neutrinos N_R , three singly charged vector-like fermion S_L, S_R , an extra Higgs doublet H_n , a singly charged scalar Φ as well as discrete Z_2 flavor symmetry. The Z_2 charges for these fields is given in table I. Due to Z_2 symmetry, right-handed neutrinos don't couple to SM Higgs.

TABLE I: Z_2 charges of particles.fields ℓ_L e_R N_R S_L S_R H H_n Φ

ielas	ℓ_L	e_R	N_R	\mathcal{S}_L	\mathcal{S}_R	Н	H_n	Ψ
Z_2	+1	+1	-1	-1	+1	+1	+1	+1

As a result the new lagrangian can be written as

$$\mathcal{L}_{N} = V(H, H_{n}, \Phi) - \overline{\ell_{L}} Y_{S} H S_{R} - \overline{S_{L}} M_{S} S_{R} - Y_{N} \overline{S_{L}} \Phi N_{R} - \frac{1}{2} \overline{N_{R}^{C}} M_{R} N_{R} + \text{h.c.} , \qquad (2)$$

where Y_S and Y_N are new Yukawa couplings, M_S and M_R are mass matrices of S and N_R , respectively. Z_2 symmetry is explicitly broken by $\overline{S}_L M_S S_R$ term. It can be recovered by adding an extra scalar singlet η , with Z_2 charge -1 and Yukawa coupling $\overline{S}_L \eta S_R$. We will not consider Yukawa couplings $\overline{\ell}_L H_n e_R$ and $\overline{\ell}_L H_n S_R$, which can be forbidden by another Z'_2 symmetry. The following is the full Higgs potential:

$$V = -m_1^2 H^{\dagger} H - m_2^2 H_n^{\dagger} H_n - m_3^2 \Phi^{\dagger} \Phi + \lambda_1 (H^{\dagger} H)^2 + \lambda_2 (H_n^{\dagger} H_n)^2 + \lambda_3 (H^{\dagger} H) (H_n^{\dagger} H_n) + \frac{\lambda_4}{4} (H^{\dagger} H_n + H_n^{\dagger} H)^2 - \frac{\lambda_5}{4} (H^{\dagger} H_n - H_n^{\dagger} H)^2 + \lambda_6 (\Phi^{\dagger} \Phi) (H^{\dagger} H) + \lambda_7 (\Phi^{\dagger} \Phi) (H_n^{\dagger} H_n) + [\lambda_n \Phi (H^T i \sigma_2 H_n) + \text{h.c.}] .$$
(3)

We define $\langle H \rangle = v_1/\sqrt{2}$ and $\langle H_n \rangle = v_2/\sqrt{2}$. After imposing the conditions of global minimum, one finds that

$$v_1^2 = \frac{\lambda_2 m_1^2 - \lambda_o m_2^2}{\lambda_1 \lambda_2 - \lambda_o^2} ; \qquad v_2^2 = \frac{\lambda_1 m_2^2 - \lambda_o m_1^2}{\lambda_1 \lambda_2 - \lambda_o^2} , \qquad (4)$$

where $\lambda_o = 1/2(\lambda_3 + \lambda_4)$.

In the basis (h^-, h_n^-, S^-) , we can derive the mass matrix for charged scalars:

$$M_{\rm C} = \begin{pmatrix} -\lambda_4 v_2^2 & \lambda_4 v_1 v_2 & \sqrt{2}\lambda_n v_2 \\ \lambda_4 v_1 v_2 & -\lambda_4 v_1^2 & -\sqrt{2}\lambda_n v_1 \\ \sqrt{2}\lambda_n v_2 & -\sqrt{2}\lambda_n v_1 & \rho \end{pmatrix} , \qquad (5)$$

where $\rho \equiv -2m_3^2 + \lambda_6 v_1^2 + \lambda_7 v_2^2$. $M_{\rm C}$ can be diagonalized by the 3×3 unitary transformation V: $V^{\dagger}M_{\rm C}V^* = {\rm diag}(M_{G^+}, M_{H^+}, M_{S^+})$. The mass eigenvalues for these charged scalars are then

$$M_{G^{+}} = 0;$$

$$M_{H^{+}} = \frac{1}{2} \left| B - \sqrt{B^{2} + 4(v_{1}^{2} + v_{2}^{2})(\rho\lambda_{4} + 2\lambda_{n}^{2})} \right|;$$

$$M_{S^{+}} = \frac{1}{2} \left| B + \sqrt{B^{2} + 4(v_{1}^{2} + v_{2}^{2})(\rho\lambda_{4} + 2\lambda_{n}^{2})} \right|,$$
(6)

where $B \equiv \rho - \lambda_4 (v_1^2 + v_2^2)$. Here G^+ is the SM goldstone boson. We also derive the mass matrix for CP-even scalars in the basis $(h, h_n)^T$ and CP-odd scalars in the basis $(G, G_n)^T$:

$$M_{\rm N} = \begin{pmatrix} 2\lambda_1 v_1^2 & \lambda_o v_1 v_2 \\ \lambda_o v_1 v_2 & 2\lambda_2 v_2^2 \end{pmatrix} ; \qquad M_{\rm G} = \begin{pmatrix} -\lambda_s v_2^2 & \lambda_s v_1 v_2 \\ \lambda_s v_1 v_2 & -\lambda_s v_1^2 \end{pmatrix} , \tag{7}$$

$$/2(\lambda_4 - \lambda_5).$$

where $\lambda_s = 1/2(\lambda_4 - \lambda_5)$

We also derive the masses for gauge bosons, which are $M_W^2 = g^2(v_1^2 + v_2^2)/4$ and $M_Z^2 = g^2(v_1^2 + v_2^2)/4 \cos^2 \theta_w$, separately. Such that electroweak precision observable $\rho \equiv M_W^2/M_Z^2 \cos^2 \theta_w = 1$ in our theory. Our scalar field sector is similar to that in Zee model [16]. We present in appendix A a different setting to the particle contents, by replacing scalar singlet with triplet.

III. PHENOMENOLOGICAL ANALYSIS

In this section, we devote to investigate some phenomenological implications of our model. We focus on (A) neutrino masses and mixings; (B)lepton flavor violations; (C) electromagnetic properties and (D) collider signatures of heavy majorana neutrinos, which will be deployed in the following:

A. Neutrino masses and lepton mixing martrix

In our model, there is no Dirac neutrino mass term at the tree level. However we can derive a small Dirac neutrino mass matrix at the loop level. The relevant feynman diagram is shown in Fig. 1.



FIG. 1: One-loop correction to the Dirac neutrino mass matrix.

A direct calculation results in

$$(M_D)_{ab}^{\text{loop}} = \frac{\lambda_n \langle H_n \rangle}{32\pi^2} (Y_S)_{ac} (Y_N^{\dagger})_{cb} M_{S_c} \mathcal{F} \left(M_H^2, M_{\Phi}^2, M_{S_c}^2 \right) , \qquad (8)$$

where the one loop function appearing in upper equation is given by

$$\mathcal{F}(m_1^2, m_2^2, m^2) = \frac{1}{m_1^2 - m_2^2} \left(\frac{\ln \beta_1}{\beta_1 - 1} - \frac{\ln \beta_2}{\beta_2 - 1} \right) , \qquad (9)$$

with $\beta_i = m^2/m_i^2$. When $m_1 = m_2 = m$, \mathcal{F} reduces to $1/2m^2$.

Here M_D^{loop} is the Dirac neutrino mass matrix linking the left and right hand neutrinos, which only raises at the loop level in our model. If neutrinos are Dirac particles, then Eq. (8) is just neutrino mass formula, whose predication must be consistent with present neutrino oscillation data. In this paper, we assume that neutrinos are Majorana particles, i.e., left-handed and right-handed neutrinos have different mass eigenvalues. Then three active neutrino masses can be generated from seesaw mechanism. In this case, we can write down the 6 × 6 neutrino mass matrix:

$$\mathcal{M} = \begin{pmatrix} 0 & M_D^{\text{loop}} \\ M_D^{\text{loop}T} & M_R \end{pmatrix} , \qquad (10)$$

which can be diagonalized by the unitary transformation $\mathcal{U}^{\dagger}\mathcal{M}\mathcal{U}^{*} = \widehat{\mathcal{M}}$; or explicitly,

$$\begin{pmatrix} V & R \\ S & T \end{pmatrix}^{\dagger} \begin{pmatrix} 0 & M_D^{\text{loop}} \\ M_D^{\text{loop}T} & M_R \end{pmatrix} \begin{pmatrix} V & R \\ S & T \end{pmatrix}^* = \begin{pmatrix} \hat{M}_{\nu} & 0 \\ 0 & \hat{M}_N \end{pmatrix} .$$
(11)

Given $M_D^{\text{loop}} \ll M_R$, the light Majorana neutrino mass formula is then $M_{\nu} = -M_D^{\text{loop}} M_R^{-1} M_D^{\text{loop}^T}$. Notice that Dirac neutrino mass matrix is suppressed by loop factor, we assume $M_D^{\text{loop}} \sim \mathcal{O}(\text{MeV})$, which will not cause any fine-tune problem. Then, to generate electron-volt scale active neutrino masses, heavy Majorana neutrinos would be of the order of several hundred GeV.

We also obtain the charged lepton mass matrix in the basis $(E_L, S_L)^T$,

$$M_{\ell} = \begin{pmatrix} M_E & M_C \\ 0 & M_S \end{pmatrix} = \begin{pmatrix} \mathbf{I} & M_C M_S^{-1} \\ 0 & \mathbf{I} \end{pmatrix} \begin{pmatrix} M_E & 0 \\ 0 & M_S \end{pmatrix} , \qquad (12)$$

where $M_E = v/\sqrt{2}Y_E$ and $M_C = v/\sqrt{2}Y_S$.

According to Eqs. (11) and (12), we may derive the lepton mixing matrix (MNS), which comes from the mismatch between the diagonalizations of the neutrino mass matrix and charged lepton mass matrix, i.e., $U = V_e^{\dagger} V_{\nu}$:

$$U \approx (\mathbf{1} - \frac{1}{2} |M_C M_S^{-1}|^2) V .$$
(13)

As a result, the effective charged and neutral current interactions for charged leptons can be written as

$$-\mathcal{L}_{\rm CC} \approx \frac{g}{\sqrt{2}} \overline{e_{\alpha}} \gamma^{\mu} P_L U_{\alpha i} \nu_i W_{\mu} + \text{h.c.} ; \qquad (14)$$

$$-\mathcal{L}_{\rm NC} \approx \frac{g}{\cos\theta_w} \overline{e_\alpha} \gamma^\mu \left[(UU^\dagger)_{\alpha\beta} (-\frac{1}{2} + \sin^2\theta_w) P_L + \delta_{\alpha\beta} \sin^2\theta_w P_R \right] e_\beta Z_\mu .$$
(15)

The MNS matrix in Eq. (13) is non-unitary, which is mainly because the large mixing between charged leptons and heavy vector like fermions. To a better degree of accuracy, we have $UU^{\dagger} \approx \mathbf{1} - |M_C M_S^{-1}|^2$. A global analysis of current neutrino oscillation data and precision electroweak data (e.g., on the invisible width of the Z^0 boson, universality tests and rare decays) has yield quite strong constraints on the unitarity of U. Translating the numerical results of Refs. [17–20] into the restriction on $|M_C M_S^{-1}|^2$, we have

$$|M_C M_S^{-1}|^2 = \begin{pmatrix} <1.1 \cdot 10^{-2} & <7.0 \cdot 10^{-5} & <1.6 \cdot 10^{-2} \\ <7.0 \cdot 10^{-5} & <1.0 \cdot 10^{-2} & <1.0 \cdot 10^{-2} \\ <1.6 \cdot 10^{-2} & <1.0 \cdot 10^{-2} & <1.0 \cdot 10^{-2} \end{pmatrix} ,$$
(16)

at the 90% confidence level. In addition, interactions in Eqs. (14) and (15) will lead to tree level lepton flavor violations (as can be seen in Eq. (15)) and "zero distance" effects [17] in neutrino oscillations, which can be verified in the future long baseline neutrino oscillation experiments.

B. Lepton flavor violations (LFV)

Notice that the emergence of big unitary violation of MNS matrix can lead to observable LFV effects. In this subsection, we investigate constraints on parameter space from LFV processes.

In our model, $\ell_i \to 3\ell_j$ may occur at the tree level, just like the case in type III seesaw mechanism. The branching ratios for the $\mu \to 3e$ can be given by

$$BR(\mu \to e^- e^+ e^-) = |UU^{\dagger}|^2_{e\mu} \left[|UU^{\dagger}|_{ee} (-1 + 2\sin^2 \theta_w)^2 + 4\sin^4 \theta_w \right] \Omega .$$
(17)

Here Ω is the final states phase space integration

$$\Omega = \int_{z_d}^{z_u} \sqrt{1 - \frac{4z_0}{z}} \left[-2z^2 + (1 + 3z_0)z - 4z_0(1 + z_0) + \frac{2z_0(1 - z_0)^2}{z} \right] \lambda^{\frac{1}{2}}(1, z_0, z) dz , \quad (18)$$

where $z_d = 4z_0$, $z_u = (1 - \sqrt{z_0})^2$, $z_0 = m_e^2/m_\mu^2$ and $\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz$.

Radiative decays, i.e., $\ell_i \to \ell_j + \gamma$ occur at one-loop level. The branching ratios for these processes can be written as

$$BR(\ell_{\beta} \to \ell_{\alpha} \gamma) = \frac{3\pi\alpha}{16G_{F}^{2}m_{\beta}^{4}} \left| \sum_{i} Y_{S\alpha i} Y_{S\beta i}^{*} \mathcal{S}_{i} \right|^{2} BR(\ell_{\beta} \to \nu_{\beta} \ell_{\alpha} \bar{\nu}_{\alpha}) , \qquad (19)$$

with

$$S_{\rho} = \frac{\sqrt{z_{\alpha} z_{\beta}} (1 - z_h) - 2z_{\beta} (1 - z_h) + 2\sqrt{z_{\alpha} z_{\beta}}}{(1 - z_h)^2} - \frac{z_{\beta} (1 - z_h) - \sqrt{z_{\alpha} z_{\beta}}}{(1 - z_h)^3} \ln z_h , \qquad (20)$$

where $z_\alpha=m_\alpha^2/M_{Si}^2$; $z_\beta=m_\beta^2/M_{Si}^2$; $z_h=M_h^2/M_{Si}^2$.

In Fig. 2 (a), we plot $BR(\mu \to 3e)$ as function of $|UU^{\dagger}|_{e\mu}^2$. The horizon line stands for current experimental constraints. Our result shows that, to meet the experimental data, $|UU^{\dagger}|_{e\mu}^2$ must lie below 8.5×10^{-6} . Assuming that there is only one generation vector like fermion, we plot, in Fig. 2 (b), $BR(\mu \to e\gamma)/|Y_SY_S^{\dagger}|_{e\mu}^2$ as function of M_S by setting $M_h = 120$ GeV. We find that, to get big Yukawa coupling Y_S , M_S must lie around 270 GeV or be heavier than several TeV.

C. Electromagnetic properties of heavy Majorana neutrinos

The electromagnetic properties of Majorana neutrinos show up, in a quantum field theory, as its interaction with the photon, and is described by the following effective interaction



FIG. 2: Branching ratios for (a) $\mu \to 3e$ and (b) $\mu \to e\gamma$.

vertex: $\mathcal{L}_{\text{eff}} = \bar{\psi}\Gamma_{\mu}\psi A^{\mu}$. The most general matrix element of \mathcal{L}_{eff} between two one-particle states, i.e., $\langle p', s' | J_{\mu}(0) | p, s \rangle = \bar{u}_{s'}(p')\Gamma_{\mu}u_s(p)$, which is consistent with the Lorentz invariance, can be written as

$$\bar{u}(p_2, s_2)\Gamma^{\mu}u(p_1, s_1) = \bar{u}(p_2, s_2) \left[\mathcal{E}(q^2)\gamma^{\mu} - \mathcal{M}(q^2)i\sigma^{\mu\nu}q^{\nu} + \mathcal{H}(q^2)q^{\mu} \right] u(p_1, s_1) + \bar{u}(p_2, s_2) \left[\mathcal{G}(q^2)\gamma^{\mu}\gamma^5 - \mathcal{T}(q^2)i\sigma^{\mu\nu}q^{\nu}\gamma^5 + \mathcal{S}(q^2)q^{\mu}\gamma^5 \right] u(p_1, s_1) \quad , \quad (21)$$

where $q = p_2 - p_1$. $2M\mathcal{M}(0)$ and $2M\mathcal{T}(0)$ correspond to the magnetic moment and electro dipole moment of heavy neutrinos, respectively.

Due to the Majorana nature, the magnetic moment of heavy Majorana neutrinos is zero. There is only transition magnetic moment for them. In the model considered, we have four diagrams contributing to the transition magnetic moment, which are depicted in Fig. 3. The Yukawa interactions of heavy Majorana neutrinos with Φ and S can be rewritten in the following way

$$\frac{1}{2} \left[\overline{N_{\alpha}^{C}} \Phi^{-} (Y^{T})_{\alpha i} P_{R} S_{i}^{C} + \overline{N_{\alpha}} \Phi^{+} (Y^{\dagger})_{\alpha i} P_{L} S_{i} \right] + \frac{1}{2} \left[\overline{S_{i}} Y_{i\alpha} \Phi^{-} P_{R} N_{\alpha} + \overline{S_{i}^{C}} (Y^{*})_{i\alpha} \Phi^{+} P_{L} N_{\alpha}^{C} \right]$$
(22)

through which we can derive relevant feynman rules. Assuming that heavy Majorana neutrinos are nearly degenerate, i.e., $M_{\alpha} \approx M_{\beta} \approx M$, we derive the transition magnetic moment for heavy Majorana neutrinos

$$a_{\alpha\beta}^{N} = \frac{M^{2}}{64\pi^{2}} \left[(Y_{N}^{\dagger})_{\beta i} (Y_{N})_{i\alpha} - (Y_{N}^{T})_{\beta i} (Y_{N})_{i\alpha}^{*} \right] \left[\mathcal{I}(M_{\Phi}^{2}, M^{2}, M_{i}^{2}) - \mathcal{I}(M_{i}^{2}, M^{2}, M_{\Phi}^{2}) \right] , \quad (23)$$

with

$$\mathcal{I}(A, B, C) = \int dx \frac{x(1-x)^2}{(1-x)A + x(x-1)B + xC} \, ,$$



FIG. 3: Feynman diagrams contributing to heavy Majorana neutrino transition magnetic moment. where M_i and M_{Φ} are the mass eigenvalues of heavy vector-like fermion S and scalar Φ , respectively.

Now, we turn to some numerical analysis. As shown in Eq. (2), $Y_N \overline{S_L} \Phi N_R$ is totally the interaction of new fields beyond the SM, so that there is no experimental constraint on Y_N except $\mathcal{O}(Y_N) < \sqrt{4\pi}$ (to satisfy the perturbation theory). We plot, in Fig. 4, $|a_{\alpha\beta}^N/[(Y_N^{\dagger})_{\beta i}(Y_N)_{i\alpha} - (Y_N^T)_{\beta i}(Y_N)_{i\alpha}^*]|$ as function of M_N . Assuming $\mathcal{O}([(Y_N^{\dagger})_{\beta i}(Y_N)_{i\alpha} - (Y_N^T)_{\beta i}(Y_N)_{i\alpha}]) \sim 1$, We can find that the transition magnetic moment of heavy Majorana neutrinos can be of $\mathcal{O}(10^{-2})$ for special parameter settings. Our result in Eq. (23) is different from that in Ref. [22] for not considering the Yukawa coupling $\overline{N_R^C} \Phi S_R$, which is forbidden by the Z_2 symmetry in our model. Given the large electromagnetic form factors, heavy Majorana neutrinos can be produced at the LHC through the electromagnetic interaction.

D. Collider signatures

We switch to comment on the collider signatures of our model. All of the new particles introduced in the model lie around several hundred GeV. Singly charged scalar and vector like fermions can be produced through the electromagnetic interaction at the LHC. The most promising production channel may be $pp \rightarrow S^+S^- \rightarrow \ell^+\ell^-jjjj$ for heavy charged



FIG. 4: Numerical illustration for the transition magnetic moment of heavy Majorana neutrinos as function of M_N . We assume there is only one generation vector-like fermion S and choose $(M_{\Phi}, M_i) = (350, 150) \text{ (GeV)}$ for solid line, (300, 200) (GeV) for dashed line and (300, 250) (GeV) for dotted line.

fermions and $pp \to \Phi^+ \Phi^- \to jjjj$ for heavy charged scalar. The production cross sections for these charged particles at the LHC (with $\sqrt{s} = 14$ TeV) are about several fb when heavy particle masses lie around 300 GeV [21, 22]. The large transition magnetic moment can help to produce the heavy Majorana neutrinos at the LHC. Its signatures are similar to that in Type-III seesaw model [23–25]. The only distinguish is that, heavy neutrinos can not be produced through weak interactions and must be produced in pair in our model.

IV. CONCLUDING REMARKS

In this paper, we have proposed a novel TeV-scale seesaw mechanism. One salient feature of our model is that Dirac neutrino mass matrix raises only at the loop level. As a result, the heavy Majorana neutrinos can be several hundred GeV. Another salient feature is that heavy Majorana neutrinos can get large electromagnetic form factors, through which they can be produced and detected at the LHC. We have derived light Majorana neutrino mass formula and calculated constraints on parameter space from LFV processes. At last we have discussed the signatures of heavy fermions (vector-like fermions and heavy Majorana neutrinos) and scalar at the LHC.

Acknowledgments

The author thanks to Tong Li, Yi Liao and Shu Luo for useful discussion. This work was supported in part by the National Natural Science Foundation of China.

Appendix A: An alternative setting to the model

Beside the model presented in section II, we can extending the SM with different particle contents, which may lead to the same TeV seesaw mechanism. For example, we can substitute S_L, S_R, Φ , with vector like fermion triplets Σ_L, Σ_R and scalar triplet Δ . In this case the lagrangian can be written as

$$\mathcal{L} = \mathcal{L}_{\rm SM} - \overline{\ell_L} Y_{\Psi} \tilde{H} \Sigma_R - \overline{\Sigma_L} M_{\Sigma} \Sigma_R - Y_N \operatorname{Tr}[\Psi_L \Delta] N_R + V(H, \Delta) .$$
(A1)

Here the weak hypercharge of the Ψ and Σ are zero. Ψ_L , Φ and N_R are odd, while the other fields are even under Z_2 transformation.

The Higgs potential can be written as

$$V = \frac{1}{2}m_H^2 H^{\dagger}H + \frac{1}{2}m_{\Delta}\text{Tr}[\Delta^{\dagger}\Delta] + \frac{1}{4}\lambda(H^{\dagger}H)^2 + \lambda_X H^{\dagger}\Sigma i\sigma_2 H + \cdots , \qquad (A2)$$

where dots denote Higgs potential terms we don't concern.

- [1] SNO Collaboration, Q. P. Ahamed *et al.*, Phys. Rev. Lett. **89**, 011301 (2002).
- [2] For a review, see: C. K. Jung et al., Ann. Rev. Nucl. Part. Sci. 51, 451 (2001).
- [3] KamLAND Collaboration, K. Eguchi et al., Phys. Rev. Lett. 90, 021802 (2003).
- [4] K2K Collaboration, M. H. Ahn *et al.*, Phys. Rev. Lett. **90**, 041801 (2003).
- [5] S. Weinberg, Phys. Rev. Lett. **43**, 1566 (1979).
- [6] P. Minkowski, Phys. Lett. B 67, 421 (1977); T. Yanagida, in Workshop on Unified Theories, KEK report 79-18 p.95 (1979); M. Gell-Mann, P. Ramond, R. Slansky, in Supergravity (North Holland, Amsterdam, 1979) eds. P. van Nieuwenhuizen, D. Freedman, p.315; S. L. Glashow, in 1979 Cargese Summer Institute on Quarks and Leptons (Plenum Press, New York, 1980) eds. M. Levy, J.-L. Basdevant, D. Speiser, J. Weyers, R. Gastmans and M. Jacobs, p.687; R. Barbieri, D. V. Nanopoulos, G. Morchio and F. Strocchi, Phys. Lett. B 90, 91 (1980);

R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. 44, 912 (1980); G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B 181, 287 (1981).

- [7] W. Konetschny and W. Kummer, Phys. Lett. B 70, 433 (1977); T. P. Cheng and L. F. Li, Phys. Rev. D 22, 2860 (1980); G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B 181, 287 (1981); J. Schechter and J. W. F. Valle, Phys. Rev. D 22, 2227 (1980); R. N. Mohapatra and G. Senjanovic, Phys. Rev. D 23, 165 (1981).
- [8] R. Foot, H. Lew, X. G. He and G. C. Joshi, Z. Phys. C 44, 441 (1989).
- [9] E. Ma and U. Sarkar, Phys. Rev. Lett. 80, 5716 (1998).
- [10] M. Y. Keung and G. Senjanovic, Phys. Rev. Lett. 50, 1427 (1983); A. Pilaftsis, Z. Phys. C 55, 275 (1992); B. Bajc, M. Nemevsek and G. Senjanovic, Phys. Rev. D 76, 055011 (2007).
- [11] J. Bernabeu, A. Santamaria, J. Vidal, A. Mendez, and J.W.F. Valle, Phys. Lett. B 187, 303 (1987); W. Buchmuller and D. Wyler, Phys. Lett. B 249, 458 (1990); W. Buchmuller and C. Greub, Nucl. Phys. B 363, 345 (1991); A. Datta and A. Pilaftsis, Phys. Lett. B 278, 162 (1992); G. Ingelman and J. Rathsman, Z. Phys. C 60, 243 (1993); C.A. Heusch and P. Minkowski, Nucl. Phys. B 416, 3 (1994); D. Tommasini, G. Barenboim, J. Bernabeu, and C. Jarlskog, Nucl. Phys. B 444, 451 (1995).
- [12] J. Gluza, Acta Phys. Polon. B 33, 1735 (2002); J. Kersten and A.Yu Smirnov, Phys. Rev. D 76, 073005 (2007); X. G. He, S. Oh, J. Tandean and C. C. Wen, Phys. Rev. D 80, 073012 (2009).
- [13] T. Han and B. Zhang, Phys. Rev. Lett. 97, 171804 (2006).
- [14] F.del Aguila, J.A. Aguilar-Saavedra, A.M. de la Ossa, and M. Meloni, Phys. Lett. B 613, 170 (2005); F.del Aguila and J.A. Aguilar-Saavedra, JHEP 0505, 026 (2005); F.del Aguila, J. A. Aguilar-Saavedra, and R. Pittau, JHEP 0710, 047 (2007); N. Haba, S. Matsumoto and K. Yoshioka, Phys. Lett. B 677, 291 (2009).
- [15] W. Chao, S. Luo, Z.Z. Xing, and S. Zhou, Phys. Rev. D 77, 016001 (2008); W. Chao, Z. Si,
 Z.Z. Xing, and S. Zhou, Phys. Lett. B 666, 451 (2008); Z. Z. Xing, Phys. Lett. B 679, 255 (2009); W. Chao, Z. Si, Y. J. Zheng and S. Zhou, Phys. Lett. B 683, 26 (2010).
- [16] A. Zee, Phys. Lett. B **93**, 389 (1980), Erratum-ibid.B **95**, 461 (1980).
- [17] S. Antusch, C. Biggio, E. Fernandez-Martinez, M. B. Gavela and J. Lopez-Pavon, JHEP 0610, 084 (2006).
- [18] E. Fernandez-Martinez, M. B. Gavela, J. Lepez-Pavon and O. Yasuda, Phys. Lett. B 649, 427

(2007).

- [19] Z. Z. Xing, Phys. Lett. B 660, 515 (2008).
- [20] A. Abada, C. Biggio, F. Bonnet, B. Gavela and T. Hambye, JHEP, 0712, 061 (2007).
- [21] A. Aparici, K. Kim, A. Santamaria and J. Wudka, Phys. Rev. D 80, 013010 (2010).
- [22] A. Aparici, K. Kim, A. Santamaria and J. Wudka, arXiv: 0911.4103 [hep-ph].
- [23] R. Franceschini, T. Hambye and A. Strumia, Phys. Rev. D 78, 033002 (2008).
- [24] F. del Agulia and J. A. Aguilar-Saavedra, Nucl. Phys. B 813, 22 (2009).
- [25] Tong Li and X. G. He, Phys. Rev. D 80. 093003 (2009).