Spreading speeds and traveling waves for non-cooperative integro-difference systems

Haiyan Wang

Division of Mathematical and Natural Sciences Arizona State University Phoenix, AZ 85069-7100, USA E-mail: wangh@asu.edu

Carlos Castillo-Chavez

Mathematics, Computational and Modeling Sciences Center, Arizona State University; PO Box 871904, Tempe, AZ, 85287, USA School of Human Evolution and Social Change, Arizona State University, Tempe, AZ 85282, USA Santa Fe Institute, SFI, 1399 Hyde Park Road, Santa Fe, NM, 87501 E-Mail: ccchavez@asu.edu

Abstract

The development of macroscopic descriptions for the joint dynamics and behavior of large heterogeneous ensembles subject to ecological forces like dispersal remains a central challenge for mathematicians and biological scientists alike. Over the past century, specific attention has been directed to the role played by dispersal in shaping plant communities, or on the dynamics of marine open-ocean and intertidal systems, or on biological invasions, or on the spread of disease, to name a few. Mathematicians and theoreticians, starting with the efforts of researchers that include Aronson, Fisher, Kolmogorov, Levin, Okubo, Skellam, Slobodkin, Weinberger and many others, set the foundation of a fertile area of research at the interface of ecology, mathematics, population biology and evolutionary biology. Integrodifference systems, the subject of this manuscript, arise naturally in the study of the spatial dispersal of organisms whose local population dynamics are prescribed by models with discrete generations. The brunt of the *mathematical* research has focused on the the study of existence traveling wave solutions and characterizations of the spreading speed particularly, in the context of cooperative systems. In this paper, we characterize the spreading speed for a large class of non cooperative systems, all formulated in terms of integrodifference equations, by the convergence of initial data to wave solutions. In this setting, the spreading speed is characterized as the slowest speed of a family of non-constant traveling wave solutions. Our results are applied to a specific non-cooperative competition system in detail.

Key words: traveling waves, cooperative systems, spreading speed, discrete-time systems, Schauder fixed point theorem, dispersal, biological invasions, epidemics 1991 MSC: 92D40, 92D25

1 Introduction

Dispersal has long been identified as a major contributor to the shaping of biological communities. Efforts to build macroscopic descriptions for the joint dynamics of large heterogeneous ensembles subject to evolutionary forces are of central concern to ecologists and evolutionary biologists alike. This powerful theme was set in "motion" by fundamental ecological contributions of Skellam (1951) [\[24\]](#page-27-0), Kierstad and Slobodkin (1953) [\[9\]](#page-26-0), Levin and Paine (1974) [\[12\]](#page-26-1), Okubo (1980) [\[13\]](#page-26-2) and others. A glance at the literature will highlight the study of the role of dispersal in contexts that include, for example, plant communities, marine open-ocean and intertidal systems, biological invasions, and epidemics.

Dispersal models given by integrodifference equations arise in population biology from the study of the spatial dynamics of organisms with discrete primarily non-overlapping (but also possibly overlapping generations [\[25\]](#page-27-1)) local dynamics and dispersal processes modeled via re-distribution kernels. Early models for invasive species were primarily modeled by nonlinear reaction-diffusion equations such as Fisher's Equation [\[5\]](#page-26-3) in the context of populations with overlapping generations. The primary motivation (emphasis) was on the characterization of the speed of propagation of an invading species in an unoccupied habita. The pioneering work of Fisher [\[5\]](#page-26-3) and Kolmogorov, Petrowski, and Piscounov $[10]$, Aronson and Weinberger $[1,2]$ $[1,2]$ focused on the rigorous classification of the concept of speed of propagation for continuous time and discrete time models. Weinberger [\[28\]](#page-27-2) and Lui [\[19\]](#page-27-3) research helped expand the mathematical foundation for the theory of spreading speeds and traveling waves for cooperative operators, through their analysis of traveling waves via the convergence of initial data to wave solutions. Naturally, the mathematical challenges posed by the study of integrodifference equation models has attracted the attention of mathematicians and recently, Weinberger, Lewis and Li have made additional significant contributions [\[26,](#page-27-4)[14,](#page-26-7)[15,](#page-27-5)[27\]](#page-27-6). The mathematical analyses of these systems are also enhancing our understanding of biological invasions of introduced species like weeds or pests in terrestrial systems and/or the study of the impact of dominant alien species in freshwater.

Mathematical challenges remain because of the presence of biological forces tied in to overcompensation [\[11\]](#page-26-8). The pervasiveness of overcompensation in biological systems implies that integrodifference equations models are in general non-cooperative. Fortunately, relevant mathematical work has been carried out in such a setting. For example, Thieme [\[22\]](#page-27-7) showed, in the context of a general model with non-monotone growth functions, that the asymptotic spreading speed could still be obtained with the aid of carefully constructed monotone functions. Recently, Hsu and Zhao [\[8\]](#page-26-9), Li, Lewis and Weinberger [\[16\]](#page-27-8) extended the theory of spreading speeds in the context of non-monotone integrodifference equations and their extension relied on the construction of two monotone operators with appropriate properties and the application of fixed point theorems in Banach spaces. This approach was also used in Ma [\[21\]](#page-27-9) and Wang [\[23\]](#page-27-10) to establish the existence of traveling wave solutions of reaction-diffusion equations. Our results on the speed of propagation for noncooperative systems in the context of integrodifference equations rely on the approaches used in the study of monotone systems in Weinberger et al. [\[26\]](#page-27-4).

In order to motivate the mathematical results established in this manuscript, we proceed to formulate the key question in the context of a specific model, namely Hassell and Comins' model [\[6\]](#page-26-10) . Their two-dimensional nonlinear discrete system models the dynamics of two competing species that reproduce at discrete generations. Specifically, the model focuses on the growth and spread of two species whose population densities at time n and spatial location x are $X_n(x)$ and $Y_n(x)$. This model is a natural extension of the classical one dimensional Ricker's "scramble" competition model. The non-spatial interference competition model is given by the following system of coupled nonlinear difference equations:

$$
X_{n+1}(x) = X_n(x)e^{r_1 - X_n(x) - \sigma_1 Y_n(x)}
$$

\n
$$
Y_{n+1}(x) = Y_n(x)e^{r_2 - Y_n(x) - \sigma_2 X_n(x)}
$$
\n(1.1)

where $r_1, r_2, \sigma_1, \sigma_2$ are all positive constants.

Next, a discrete model where individuals interact locally according to Model [\(1.1\)](#page-2-0) but where the possibility of individuals dispersing to a different site is modeled with the aid of the kernel $k_i(y)$. The following system of coupled nonlinear integrodifference equations arise from the incorporation of redistribution kernels k_i , $i = 1, 2$:

$$
X_{n+1}(x) = \int_{\mathbb{R}} k_1(x - y) X_n(y) e^{r_1 - X_n(y) - \sigma_1 Y_n(y)} dy
$$

\n
$$
Y_{n+1}(x) = \int_{\mathbb{R}} k_2(x - y) Y_n(y) e^{r_2 - Y_n(y) - \sigma_2 X_n(y)} dy
$$
\n(1.2)

The re-distribution kernel $k_i(y)$ describe the dispersal of X, Y , which it is assumed to depend upon the signed distance $x - y$ connecting the location of "birth" y and the "settlement" location x; $k_i(y)$ is a homogenous probability kernel that satisfies $\int_{\infty}^{\infty} k_i(y) dy = 1$.

The above system is in general non-cooperative. It is in the context of noncooperative systems like those described by the model above that our results for non-cooperative systems will be formulated, stated and illustrated. The explicit mathematical formulation of the model requires the introduction of additional notation.

 $\beta, \beta^{\pm}, F, F^{\pm}, r, u, v$ are used to denote vectors in \mathbb{R}^{N} or N-vector valued functions while x, y, ξ are used to denote variables in R. The use of $u = (u^i)$ and $v = (v^i) \in \mathbb{R}^N$ allow us to define $u \geq v$ whenever $u^i \geq v^i$ for all i; and $u \gg v$ whenever $u^i > v^i$ for all i. We further define for any $r = (r^i) \gg 0, r \in \mathbb{R}^N$ the R^N -interval

$$
[0,r] = \{u : 0 \le u \le r, u \in \mathbb{R}^N\} \subseteq \mathbb{R}^N
$$

and

$$
\mathcal{C}_r = \{ u = (u^1, ..., u^N) : u^i \in C(\mathbb{R}, \mathbb{R}), 0 \le u^i(x) \le r^i, x \in \mathbb{R}, i = 1, ..., N \},\
$$

where $C(\mathbb{R}, \mathbb{R})$ is the set of all continuous functions from \mathbb{R} to \mathbb{R} .

Now, we focus on $\mathcal{C}_{\beta^+}, \beta^+$ >> 0. We now consider the system of integrodifference equations where

$$
u_{n+1} = \mathcal{Q}[F(u_n)];\tag{1.3}
$$

 $u_n = (u_n^i) \in C_{\beta^+}, F(u) = (f_i(u));$

$$
\mathcal{Q}[F(u)] = (\mathcal{Q}^i[F(u)]);
$$

$$
\mathcal{Q}^i[F(u)](x) = \int_{\mathbb{R}} k_i(x - y) f_i(u(y)) dy;
$$

 $u_n(x)$ is the density of individuals at point x and time n; $F(u)$ is the densitydependent fecundity; and $k_i(x - y)$ (dispersal kernel) models the dispersal of u, which it is assumed to depend on the signed distance $x - y$ between the location of "birth" y and the location of "settlement" x. Here, $k_i(x - y)$ is a probability kernel that satisfies $\int_{\infty}^{\infty} k_i(x)dx = 1$.

The integro-difference Equation [\(1.3\)](#page-3-0) models the reproduction and dispersal of a time-synchronized species in which all individuals first undergo reproduction and then redistribute their offspring before reproduction occurs once again.

Specifically, in this manuscript, the spreading speed of rather general non cooperative systems [\(1.3\)](#page-3-0) is characterized as the slowest speed of a family of non-constant traveling wave solutions of [\(1.3\)](#page-3-0). The main theorem, Theorem [2.2,](#page-6-0) generalizes relevant results, already established in the context of cooperative systems, to non-cooperative systems. The details are collected in the following sections.

2 Non-cooperative Systems' Results

The focus of our analysis is on the characterization of the speeds of propagation for [\(1.3\)](#page-3-0) when the system is non-cooperative. As it is typical in mathematics, we make use of the existing results for cooperative systems $([26])$ $([26])$ $([26])$ and for this purpose, we introduce additional notation and assumptions. First, the existence of two additional monotone operators F^{\pm} with the properties that the first lies above F and the second below F is assumed. The use of this approach is motivated by the work on non-monotone equations carried out in [\[22,](#page-27-7)[8](#page-26-9)[,16](#page-27-8)[,21,](#page-27-9)[23\]](#page-27-10). F^{\pm} can be "constructed" via piecewise functions made up of "pieces' of F and appropriate constants. If F is monotone, then $F^{\pm} = F$. The feasibility of our analysis depends on whether or not the components of our problem meet the following assumptions:

(H1) For $i = 1, ..., N$, $k_i(\tau) \geq 0$ is integrable on \mathbb{R} , $k_i(\tau) = k_i(-\tau)$, $\tau \in$ $(-\infty, +\infty)$, and

$$
\int_{\mathbb{R}} k_i(\tau) d\tau = 1, \int_{\mathbb{R}} k_i(\tau) e^{\lambda \tau} d\tau < +\infty,
$$

for all $\lambda > 0$.

(H2)(i) Given that $F : [0, \beta^+] \to [0, \beta^+]$ is a continuous, twice piecewise continuous differentiable function with $0 \ll \beta^- = (\beta_i^-) \leq \beta \leq \beta^+ = (\beta_i^+),$ it is assumed that there exist continuous, twice piecewise continuous differentiable functions $F^{\pm} = (f_i^{\pm}) : [0, \beta^+] \rightarrow [0, \beta^+]$ such that for $u \in [0, \beta^+]$

$$
F^{-}(u) \le F(u) \le F^{+}(u).
$$

- (ii) $F(0) = 0, F(\beta) = \beta$ and there is no other positive equilibrium of $\mathcal{Q}[F]$ between 0 and β (that is, there is no constant $v \neq \beta$ such that $F(v) =$ $v, 0 \ll v \leq \beta$). $F^{\pm}(0) = 0, F^{\pm}(\beta^{\pm}) = \beta^{\pm}$ and there is no other positive equilibrium of $\mathcal{Q}[F^{\pm}]$ between 0 and β^{\pm} .
- (iii) F^{\pm} are nondecreasing functions on their domains with $F^{\pm}(u) = F(u)$ if $u \in [0, \beta^+]$ are sufficiently small.

Assumptions (H1-H2) are not sufficient to achieve to characterize the speeds of propagation for (1.3) . We need the additional assumption $(H3)$ which include a requirement that the operator grows less than its linearization along the particular function $\nu_{\mu}e^{-\mu x}$ (see [\[26\]](#page-27-4)). Assumption H3, to be explicitly formulated below, is satisfied by many biological systems of interest (as it will be highlighted in our example) and hence it does not severely handicap the value of our results.

First we recall that a matrix is irreducible if it is not similar to a lower triangular matrix with two blocks via a permutation (See, $(7,26)$) as we will make us of a theorem of Frobenius that states that any nonzero irreducible matrix with nonnegative entries has a unique principal positive eigenvalue with a corresponding principal eigenvector "made up" of strictly positive coordinates. The formulation of Assumption (H3) requires, following the approach in [\[26\]](#page-27-4), that for each $\mu > 0$, the $N \times N$ matrix B_{μ} that results from the linearization of (1.3) at 0, namely

$$
B_{\mu} = (b_{\mu}^{i,j}) = \left(\partial_j f_i(0) \int_{\mathbb{R}} k_i(s) e^{\mu s} ds\right)
$$
\n(2.4)

where $b_{\mu}^{i,j}$ is the (i, j) entry of the matrix, is in Frobenius form. In our case, this is always possible albeit it may require the reordering of the coordinates, of the matrix B_{μ} which is made up of irreducible diagonal blocks (An irreducible matrix consists of the single diagonal block which is the matrix itself). We let $\lambda(\mu)$ denote therefore the largest principal eigenvalue of the diagonal blocks (See $|7,26|$). In conclusion:

- $(H3)(i)$ Assume that B_μ is in Frobenius form, that the first diagonal block includes the largest principal eigenvalue, $\lambda(\mu)$, and that B_{μ} has a positive eigenvector $\nu_{\mu} = (\nu_{\mu}^{i}) >> 0$ corresponding to $\lambda(\mu)$. Further assume that $\lambda(0) > 1$.
	- (ii) For each $\mu > 0$ and $\alpha > 0$, we let $v^{\pm} = (v_i^{\pm}) = (\min\{\beta_i^{\pm}, \nu_{\mu}^i \alpha\})$, and assume that

$$
F^{\pm}(v^{\pm}) \le B_0 v^{\pm}.
$$

(iii) For every sufficiently larger positive integer k , there is a small constant vector $\omega = (\omega^i) >> 0$ such that

$$
F(u) \ge (1 - \frac{1}{k})B_0u \ u \in [0, \omega],
$$

It follows from (H1) that $\lambda(\mu)$ is an even function. In fact, it was shown by Lui [\[19\]](#page-27-3) that $\ln \lambda(\mu)$ is a convex function and therefore, $\ln \lambda(\mu)$ achieves its minimum at $\mu = 0$ and $\lambda(0) > 1$ which implies that $\ln \lambda(\mu) > 0$.

The statement in Proposition [2.1](#page-5-0) below which is critical to our analysis, involves the following function of the largest principal eigenvalue $\lambda(\mu)$

$$
\Phi(\mu) = \frac{1}{\mu} \ln \lambda(\mu) > 0.
$$

Proposition 2.1 Assume that $(H1) - (H3)$ hold. Then

 $1 \Phi(\mu) \rightarrow \infty$ as $\mu \rightarrow 0$; 2 $\Phi(\mu)$ is decreasing as $\mu = 0^+$; $3 \Phi'(\mu)$ changes sign at most once on $(0, \infty)$ $4 \Phi(\mu)$ has a minimum $c^* > 0$. 5 For each $c > c^*$, there exist $\Lambda_c > 0$ and $\gamma \in (1, 2)$ such that

$$
\Phi(\Lambda_c) = c, \ \ \Phi(\gamma \Lambda_c) < c.
$$

Parts (1)-(4) of Proposition [2.1](#page-5-0) are essentially due to Lui [\[19\]](#page-27-3). The proof that $c^* > 0$ is carried out in the Appendix. Since $\lambda(\mu)$ is a simple root of the characteristic equation of an irreducible block, it can also be shown that $\lambda(\mu)$ is twice continuously differentiable on $\mathbb R$. Part(5) is a direct consequence of the results stated in Parts $(1)-(4)$.

A traveling wave solution u_n of (1.3) is defined as a solution of the form $u_n(\xi) = u(\xi - cn), u \in C(\mathbb{R}, \mathbb{R}^N)$. The theorems that guarantee the existence of traveling wave solutions for cooperative systems has been established (e.g. [\[15\]](#page-27-5)). It also has been established that the asymptotic spreading speed can be characterized as the speed of the slowest non-constant traveling wave solution for monotone operators [\[15,](#page-27-5)[29,](#page-27-11)[28\]](#page-27-2) and for non-monotone scalar equations [\[8](#page-26-9)[,16\]](#page-27-8).

In this paper, we shall guarantee the existence of the spreading speed and show that the spreading speed can be characterized as the speed of the slowest member of a family of non-constant traveling wave solutions of [\(1.3\)](#page-3-0). Our analysis of the asymptotic behavior of traveling wave solutions for monotone operator enables us to establish the existence of traveling wave solutions for non monotone operators. The following theorem ebcapsulates our main results.

Theorem 2.2 Assume $(H1) - (H3)$ hold. Then the following statements are valid:

(i) For any $u_0 \in C_\beta$ with compact support and $0 \le u_0 \ll \beta$, the solution u_n of [\(1.3\)](#page-3-0) satisfies

$$
\lim_{n \to \infty} \sup_{|x| \ge nc} u_n(x) = 0, \text{ for } c > c^*
$$

(ii) For any strictly positive vector $\omega \in \mathbb{R}^N$, there is a positive R_{ω} with the property that if $u_0 \in C_\beta$ and $u_0 \geq \omega$ on an interval of length $2R_\omega$, then the solution $u_n(x)$ of (1.3) satisfies

$$
\beta^{-} \le \liminf_{n \to \infty} \inf_{|x| \le nc} u_n(x) \le \beta^{+}, \text{ for } 0 < c < c^*
$$

(iii) For each $c > c^*$ [\(1.3\)](#page-3-0) admits a traveling wave solution $u(\xi - cn) =$ $(u^{i}(\xi - cn))$ such that $0 \lt \lt u(\xi) \leq \beta, \xi \in \mathbb{R}$,

$$
\beta_i^- \le \liminf_{\xi \to -\infty} u^i(\xi) \le \limsup_{\xi \to -\infty} u^i(\xi) \le \beta_i^+
$$

 $\lim_{\xi \to \infty} u^i(\xi) = 0, i = 1, ..., N$ and

$$
\lim_{\xi \to \infty} u^i(\xi) e^{\Lambda_c \xi} = \nu^i_{\Lambda_c}.
$$
\n(2.5)

If, in addition, F is nondecreasing on \mathcal{C}_{β} , then u is nonincreasing on \mathbb{R} .

(iv) For $c = c^*$ [\(1.3\)](#page-3-0) admits a nonconstant traveling wave solution $u(\xi - c n) =$ $(u^{i}(\xi - c n))$ such that $0 \leq u(\xi) \leq \beta, \xi \in \mathbb{R}$,

$$
\beta_i^- \le \liminf_{\xi \to -\infty} u^i(\xi) \le \limsup_{\xi \to -\infty} u^i(\xi) \le \beta_i^+.
$$

(v) For $0 < c < c^*$ [\(1.3\)](#page-3-0) does not admit a traveling wave solution $u_n(\xi) =$ $u(\xi - c n)$ such that $u \in \mathcal{C}_{\beta}$ with $\liminf_{\xi \to -\infty} u(\xi) > 0$ and $u(+\infty) = 0$.

Remark 2.3 The proof of these results is carried out in the sections below. In order to get a better estimate for the traveling wave solution u for non cooperative systems, the analysis requires the selection of two function F^{\pm} close enough in order. F^{\pm} can be built via piecewise functions that use pieces of F and appropriate constants as is is explicitly shown in the example at the end of this paper. The smallest monotone function above F and the largest monotone function below F are the most natural choices of F^{\pm} as long as they satisfy all the needed requirements. See [\[22](#page-27-7)[,8,](#page-26-9)[16\]](#page-27-8) for a discussion in the context of scalar cases and [\[30\]](#page-28-0) for a discussion of this approach in the context of partially cooperative reaction-diffusion system. In particular, when F is monotone, $F^{\pm} = F, \beta^{\pm} = \beta$.

We shall establish Theorem [2.2](#page-6-0) in Sections [3](#page-7-0) and [4.](#page-9-0)

3 Spreading Speeds

Our results on the speed of propagation for non-cooperative systems make use of Theorem [3.1](#page-7-1) below which collects the properties of the spreading speed c^* for monotone systems as established in Weinberger, Lewis and Li [\[26\]](#page-27-4). Theorem [3.1](#page-7-1) extends the related spreading results in Lui [\[19\]](#page-27-3) to systems of monotone recursion operators with more than two equilibria. The spreading results for monotone recursion operators with delays are established in Liang and Zhao [\[18\]](#page-27-12) where the operators support two equilibria. The operator at the center of this manuscript may support more than two equilibria with one lying at the boundary as in [\[26\]](#page-27-4) (see Section [5\)](#page-18-0).

Theorem 3.1 (Weinberger, Lewis and Li [\[26\]](#page-27-4) [Lemma 2.2, Theorem 3.1]) Assume $(H1) - (H3)$ hold. Further assume that $f^{i}(x), i = 1, ..., N$ is **nonde**creasing. Then the following statements are valid:

(i) For any $u_0 \in \mathcal{C}_{\beta}$ with compact support and $0 \leq u_0 \ll \beta$, the solution $u_n(x)$ of [\(1.3\)](#page-3-0) satisfies

$$
\lim_{n \to \infty} \sup_{|x| \ge nc} u_n(x) = 0, \text{ for } c > c^*
$$

(ii) For any strictly positive vector $\omega \in \mathbb{R}^N$, there is a positive R_{ω} with the property that if $u_0 \in C_\beta$ and $u_0 \geq \omega$ on an interval of length $2R_\omega$, then the solution $u_n(x)$ of (1.3) satisfies

$$
\liminf_{n \to \infty} \inf_{|x| \le nc} u_n(x) = \beta, \text{ for } 0 < c < c^*
$$

It is clear that $\mathcal{Q}[F^{\pm}]$ are monotone (order preserving) on \mathcal{C}_{β^+} . That is, if $u, v \in \mathcal{C}_{\beta^+}$ and $u(x) \leq v(x), x \in \mathbb{R}$, then

 $\mathcal{Q}[F^{\pm}(u)](x) \leq \mathcal{Q}[F^{\pm}(v)](x), x \in \mathbb{R}.$

Further, for $u = (u^i) \in \mathcal{C}_{\beta^+}$ and $x \in \mathbb{R}$, we have

$$
f_i^-(u(x)) \le f_i(u(x)) \le f_i^+(u(x)), i = 1, ..., N.
$$

and therefore

$$
\mathcal{Q}[F^-(u)](x) \le \mathcal{Q}[F(u)](x) \le \mathcal{Q}[F^+(u)](x), \ x \in \mathbb{R}.
$$

We are now able to establish Part (i) and (ii) of Theorem [2.2](#page-6-0) by following essentially the proof for the scalar cases found in [\[8,](#page-26-9)[16\]](#page-27-8).

Proof of Parts (i) and (ii) of Theorem [2.2.](#page-6-0)

Part (i). For a given $u_0 \in \mathcal{C}_{\beta}$ with compact support, let u_n be the *n*-th iteration of $\mathcal{Q}[F]$ starting from u_0 and let u_n^+ be the *n*-th iteration of $\mathcal{Q}[F^+]$ starting from u_0 . By (H2), we have

$$
0 \le u_n(x) \le u_n^+(x), x \in \mathbb{R}, n > 0.
$$

Thus for any $c > c^*$, it follows from Theorem [3.1](#page-7-1) (i) that

$$
\lim_{n \to \infty} \sup_{|x| \ge nc} u_n^+(x) = 0,
$$

and hence

$$
\lim_{n \to \infty} \sup_{|x| \ge nc} |u_n(x)| = 0,
$$

Part (ii). Let u_n be the *n*-th iteration of $\mathcal{Q}[F]$ starting from u_0 and u_n^+ the *n*-th iteration of $\mathcal{Q}[F^+]$ starting from u_0 . Let $v_0^i = \min\{u_0^i, \beta_i^-\}, i = 1, ..., N$. Then $v_0 = (v_0^i) \in \mathcal{C}_{\beta^{-}}$. Letting u_n^- denote the *n*-th iteration of $\mathcal{Q}[F^-]$ starting from v_0 and observing that $v_0 \leq u_0$ and $\beta^- \leq \beta \leq \beta^+$, from (H2), we have that

$$
u_n^{-}(x) \le u_n(x) \le u_n^{+}(x), x \in \mathbb{R}, n > 0.
$$

Theorem [3.1](#page-7-1) (ii) states that for any strictly positive constant ω , there is a positive R_{ω} (choose the larger one between the R_{ω} for F^{+} and the R_{ω} for F^{-})

with the property that if $u_0 \geq \omega$ on an interval of length $2R_{\omega}$. Hence, it follows that the solutions $u_n^{\pm}(x)$ satisfy

$$
\liminf_{t \to \infty} \inf_{|x| \le t c} u^{\pm}(x, t) = \beta^{\pm}, \text{ for } 0 < c < c^*.
$$

Thus for any $c < c^*$, it follows from Theorem [3.1](#page-7-1) (ii) that

$$
\liminf_{n \to \infty} \inf_{|x| \le nc} u_n^{\pm}(x) = \beta^{\pm},
$$

and consequently, that

$$
\beta^{-} \leq \liminf_{n \to \infty} \inf_{|x| \geq nc} u_n(x) \leq \beta^{+}.
$$

 \Box

4 Characterization of c^* as the slowest speeds of traveling waves

As it was noted above, a solution of (1.3) is a traveling wave of speed c if has the form $u_n(x) = u(x - cn)$, where $u \in C(\mathbb{R}, \mathbb{R}^N)$ and, of course, if it satisfies [\(1.3\)](#page-3-0). By substituting this form into (1.3), we see that $u(\xi)$ must satisfy the following system of equations.

$$
u(\xi) = Q_c[F(u)](\xi) = (Q_c^i[F(u)](\xi)) := Q[F(u)](\xi + c)
$$
\n(4.6)

In this section we complete the proof of Theorem [2.2](#page-6-0) (iii), (iv) and (v), that is, the portion of our main result that characterizes the spread speed c^* as the speed of the slowest member of a family of non-constant traveling wave solutions. This is an extension of prior results for monotone operators [\[29,](#page-27-11)[28](#page-27-2)[,15\]](#page-27-5) and for non-monotone scalar equations [\[8](#page-26-9)[,16\]](#page-27-8).

4.1 Upper and lower solutions

In this subsection, we shall verify ϕ^+ and ϕ^- defined below are upper and lower solutions of [\(4.6\)](#page-9-1) respectively. These solutions are only continuous on R. Upper and lower solutions of this type have been frequently used in the literatures (see Diekmann [\[4\]](#page-26-12), Weinberger [\[29\]](#page-27-11), Lui [\[19\]](#page-27-3), Weinberger, Lewis and Li [\[26\]](#page-27-4), Rass and Radcliffe [\[20\]](#page-27-13), Weng and Zhao [\[31\]](#page-28-1), more recently by Ma [\[21\]](#page-27-9)). In particular, the explicit use of upper vector-valued solutions can be traced to the work in [\[19](#page-27-3)[,26](#page-27-4)[,20,](#page-27-13)[31\]](#page-28-1); for lower vector-valued solutions, in the context of multi-type epidemic models, to the work in [\[20\]](#page-27-13); and in [\[31\]](#page-28-1) in the

context of multi-type SIS epidemic models. Our construction of ϕ^+ and ϕ^- , the upper and lower solutions of [\(4.6\)](#page-9-1), is motivated by the research in these references. The details follow below.

Let $c > c*, 1 < \gamma < 2, q > 1$ and recall the definitions of Λ_c and $\gamma \Lambda_c$ as utilized in Proposition [2.1.](#page-5-0) The corresponding positive eigenvectors ν_{Λ_c} and $\nu_{\gamma\Lambda_c}$ of B_μ for the eigenvalues λ_μ when $\mu = \Lambda_c$, $\gamma\Lambda_c$ can therefore be identified.

Define

$$
\phi^+(\xi) = (\phi_i^+),
$$

where

$$
\phi_i^+ = \min\{\beta_i, \nu^i_{\Lambda_c}e^{-\Lambda_c\xi}\},\ \xi\in\mathbb{R};
$$

and

$$
\phi^-(\xi) = (\phi_i^-),
$$

where

$$
\phi_i^- = \max\{0, \nu_{\Lambda_c}^i e^{-\Lambda_c \xi} - q \nu_{\gamma \Lambda_c}^i e^{-\gamma \Lambda_c \xi}\}, \ \xi \in \mathbb{R}.
$$

It is clear that if $\xi \leq$ $\frac{\ln \frac{\beta_i}{\nu_{\Lambda_c}^*}}{-\Lambda_c}$ then $\phi_i^+(\xi) = \beta_i$ and if $\xi >$ $rac{\ln \frac{k_i}{\nu_{\Lambda_c}^i}}{-\Lambda_c}$ then $\phi_i^+(\xi) =$ $\nu_{\Lambda_c}^i e^{-\Lambda_c \xi}$. Similarly, if $\xi \leq \ln(q \frac{\nu_{\gamma \Lambda_c}^i}{\nu_{\Lambda_c}^i}) \frac{1}{(\gamma - 1)}$ $\frac{1}{(\gamma-1)\Lambda_c}$ then $\phi_i^-(\xi) = 0$ and if $\xi > \ln(q\frac{\nu_{\gamma\Lambda_c}^i}{\nu_{\Lambda_c}^i})\frac{1}{(\gamma-1)}$ $(\gamma-1)\Lambda_c$ then $\phi_i^-(\xi) = \nu_{\Lambda_c}^i e^{-\Lambda_c \xi} - q \nu_{\gamma \Lambda_c}^i e^{-\gamma \Lambda_c \xi}$.

We choose $q > 1$ large enough so that

$$
\frac{\ln(q\frac{\nu_{\gamma\Lambda_c}^i}{\nu_{\Lambda_c}^i})}{(\gamma-1)\Lambda_c} > \frac{\ln \frac{\beta_i}{\nu_{\Lambda_c}^i}}{-\Lambda_c}
$$

and therefore

$$
\phi_i^+(\xi) > \phi_i^-(\xi), \xi \in \mathbb{R}.
$$

We now verify, two lemmas below, that ϕ^+ and ϕ^- are upper and lower solutions of [\(4.6\)](#page-9-1) respectively. In particular, it is assumed that Lemma [4.1](#page-10-0) is valid when F is monotone. In this case, $F^{\pm} = F, \beta^{\pm} = \beta$.

Lemma 4.1 Assume F is monotone and $(H1) - (H3)$ hold. For any $c > c^*$, then ϕ^+ is an upper solution of $\mathcal{Q}_c[F]$. That is

$$
\mathcal{Q}_c[F(\phi^+)](\xi) \le \phi^+(\xi), \xi \in \mathbb{R}.
$$

PROOF. Let $\xi_i^* =$ $\frac{\partial h}{\partial t} \frac{\partial i}{\partial t}}{\partial t}$. Then $\phi_i^+(\xi) = \beta_i$ if $\xi \leq \xi_i^*$, and $\phi_i^+(\xi) = \nu_{\Lambda_c}^i e^{-\Lambda_c \xi}$ if $\xi > \xi_i^*$. Note that $\phi_i^+(\xi) \leq \nu_{\Lambda_c}^i e^{-\Lambda_c \xi}, \xi \in \mathbb{R}$.

In view of (H3) we have, for $\xi \in \mathbb{R}$

$$
f_i(\phi^+(\xi)) \le \sum_{j=1}^N \partial_j f_i(0) \phi_i^+(\xi) \le \sum_{j=1}^N \partial_j f_i(0) \nu_{\Lambda_c}^j e^{-\Lambda_c \xi}
$$

Thus, for $\xi \in \mathbb{R}$, in view of [\(2.4\)](#page-5-1), (H3), Proposition [2.1,](#page-5-0) we obtain that

$$
\mathcal{Q}^{i}[F(\phi^{+})](\xi+c) \leq e^{-\Lambda_c(\xi+c)} \sum_{j=1}^{N} \nu_{\Lambda_c}^{j} b_{\Lambda_c}^{i,j}
$$

$$
= e^{-\Lambda_c(\xi+c)} \lambda(\Lambda_c) \nu_{\Lambda_c}^{i}
$$

$$
= e^{-\Lambda_c(\xi+c)} e^{\Lambda_c \Phi(\Lambda_c)} \nu_{\Lambda_c}^{i}
$$

$$
= \nu_{\Lambda_c}^{i} e^{-\Lambda_c(\xi+c)} e^{\Lambda_c c}
$$

$$
= \nu_{\Lambda_c}^{i} e^{-\Lambda_c \xi}.
$$

$$
(4.7)
$$

On the other hand, since $\phi_i^+(\xi) \leq \beta_i$, $i = 1, ..., N$, we have for $\xi \in \mathbb{R}$

$$
\mathcal{Q}^i[F(\phi^+)](\xi+c) \le \beta_i. \tag{4.8}
$$

Thus, we have for $\xi \in \mathbb{R}$

$$
\mathcal{Q}_c^i[F(\phi^+)](\xi) = \mathcal{Q}^i[F(\phi^+)](\xi + c) \le \phi_i^+(\xi). \tag{4.9}
$$

This completes the proof of Lemma [4.1.](#page-10-0)

 \Box

In order to verify the lower solutions, we need the following estimate for F which can be derived from a simply application of the Taylor's Theorem for multi-variable functions (see [\[23\]](#page-27-10)).

Lemma 4.2 Assume (H1−H2) hold. There exist positive constants D_{ij} , i, j = $1, ..., N$ such that

$$
f_i(u) \ge \sum_{j=1}^N \partial_j f_i(0)u^j - \sum_{j=1}^N D_{ij}(u^j)^2, \ \ u = (u^i), u^i \in [0, \beta_i^+], i = 1, ..., N.
$$

Lemma 4.3 Assume $(H1) - (H3)$ hold. For any $c > c^*$ if q (which is independent of ξ) is sufficiently large then ϕ^- is a lower solution of $\mathcal{Q}_c[F]$. That is

$$
\mathcal{Q}_c[F(\phi^-)](\xi) \ge \phi^-(\xi), \xi \in \mathbb{R}.
$$

PROOF. Again let $\xi_i^* = \ln(q \frac{\nu_{\gamma \Lambda_c}^i}{\nu_{\Lambda_c}^i}) \frac{1}{(\gamma - 1)}$ $\frac{1}{(\gamma-1)\Lambda_c}$. Hence if $\xi \leq \xi_i^*$ then $\phi_i^-(\xi) = 0$ while if $\xi > \xi_i^*$ then $\phi_i^-(\xi) = \nu_{\Lambda_c}^i e^{-\Lambda_c \xi} - q \nu_{\gamma \Lambda_c}^i e^{-\gamma \Lambda_c \xi}$. It is easy to see that

$$
\nu_{\Lambda_c}^i e^{-\Lambda_c \xi} \ge \phi^-(\xi) \ge \nu_{\Lambda_c}^i e^{-\Lambda_c \xi} - q \nu_{\gamma \Lambda_c}^i e^{-\gamma \Lambda_c \xi}, \ \xi \in \mathbb{R}, i = 1, ..., N. \tag{4.10}
$$

For $\xi \in \mathbb{R}$, in view of Lemma [4.2,](#page-11-0) we have, for $\xi \in \mathbb{R}, i = 1, ..., N$,

$$
f^{i}(\phi^{-}(\xi)) \geq \sum_{j=1}^{N} \partial_{j} f_{i}(0) \phi_{j}^{-}(\xi) - \sum_{j=1}^{N} D_{ij}(\phi_{j}^{-}(\xi))^{2}
$$

\n
$$
\geq \sum_{j=1}^{N} \partial_{j} f_{i}(0) \nu_{\Lambda_{c}}^{j} e^{-\Lambda_{c}\xi} - q \sum_{j=1}^{N} \partial_{j} f_{i}(0) \nu_{\gamma\Lambda_{c}}^{j} e^{-\gamma\Lambda_{c}\xi}
$$
(4.11)
\n
$$
- \widehat{M}_{i} e^{-2\Lambda_{c}\xi}
$$

where

$$
\widehat{M}_i = \sum_{j=1}^{N} D_{ij} (\nu_{\Lambda_c}^j)^2 > 0.
$$
\n(4.12)

Now we are able to estimate $\mathcal{Q}[\phi^-]$ for $\xi \ge \min_i \xi_i^*, i = 1, ..., N$ as in [\(4.7\)](#page-11-1)

$$
\mathcal{Q}^{i}[F(\phi^{-})](\xi+c) \geq e^{-\Lambda_{c}(\xi+c)} \sum_{j=1}^{N} \nu_{\Lambda_{c}}^{j} b_{\Lambda_{c}}^{i,j} - q e^{-\gamma \Lambda_{c}(\xi+c)} \sum_{j=1}^{N} \nu_{\gamma \Lambda_{c}}^{j} b_{\gamma \Lambda_{c}}^{i,j}
$$

\n
$$
- \widehat{M}_{i} e^{-2\Lambda_{c}(\xi+c)} \int_{\mathbb{R}} k_{i}(y) e^{2\Lambda_{c}y} dy
$$

\n
$$
= \nu_{\Lambda_{c}}^{i} e^{-\Lambda_{c}(\xi+c)} e^{\Lambda_{c}\Phi(\Lambda_{c})} - q \nu_{\gamma \Lambda_{c}}^{i} e^{-\gamma \Lambda_{c}(\xi+c)} e^{\gamma \Lambda_{c}\Phi(\gamma \Lambda_{c})}
$$

\n
$$
- \widehat{M}_{i} e^{-2\Lambda_{c}(\xi+c)} \int_{\mathbb{R}} k_{i}(y) e^{2\Lambda_{c}y} dy
$$

\n
$$
= \nu_{\Lambda_{c}}^{i} e^{-\Lambda_{c}\xi} - q \nu_{\gamma \Lambda_{c}}^{i} e^{-\gamma \Lambda_{c}\xi} e^{\gamma \Lambda_{c}(\Phi(\gamma \Lambda_{c})-c)}
$$

\n
$$
- \widehat{M}_{i} e^{-2\Lambda_{c}(\xi+c)} \int_{\mathbb{R}} k_{i}(y) e^{2\Lambda_{c}y} dy
$$

\n
$$
= \nu_{\Lambda_{c}}^{i} e^{-\Lambda_{c}\xi} - q \nu_{\gamma \Lambda_{c}}^{i} e^{-\gamma \Lambda_{c}\xi}
$$

\n
$$
+ q \nu_{\gamma \Lambda_{c}}^{i} e^{-\gamma \Lambda_{c}\xi} - q \nu_{\gamma \Lambda_{c}}^{i} e^{-\gamma \Lambda_{c}\xi} e^{\gamma \Lambda_{c}(\Phi(\gamma \Lambda_{c})-c)}
$$

\n
$$
- \widehat{M}_{i} e^{-2\Lambda_{c}(\xi+c)} \int_{\mathbb{R}} k_{i}(y) e^{2\Lambda_{c}y} dy
$$

\n
$$
= \phi_{i}^{-}(\xi) + e^{-\gamma \Lambda_{c}\xi} \left(q \nu_{\gamma \Lambda_{c}}^{i}(1 - e^{\gamma \Lambda_{c}(\Phi(\gamma \Lambda_{c})-c)} \right)
$$

\n
$$
- \widehat{
$$

For $\xi \ge \min_i \xi_i^*, e^{(\gamma - 2)\Lambda_c \xi}$ are bounded above. Finally, from [\(4.13\)](#page-12-0) and the fact that $\Phi(\gamma \Lambda_c) < c$, we conclude that there exists $q > 0$, which is independent of ξ , such that, for $\xi \geq \xi_i^*$

$$
\mathcal{Q}^i[F(\phi^-)](\xi+c) \ge \nu_{\Lambda_c}^i e^{-\Lambda_c \xi} - q \nu_{\gamma \Lambda_c}^i e^{-\gamma \Lambda_c \xi}.
$$
\n(4.14)

And since $\phi^-(\xi) = 0$ for $\xi < \xi_i^*, i = 1, ..., N$

$$
Q_c^i[F(\phi^-)](\xi) = Q^i[F(\phi^-)](\xi + c) \ge \phi_i^-(\xi), \xi \in \mathbb{R}.
$$

This completes the proof. \Box

4.2 Proof of Theorem [2.2](#page-6-0) (iii) with monotonicity of F

Theorems that guarantee the existence of traveling wave solutions for cooperative systems have been established (e.g. $[15]$). In this section, it is assumed that F is nondecreasing on $[0, \beta]$ and from this assumption, we proceed to established Theorem [2.2.](#page-6-0) As we shall see, our analysis of the asymptotic behavior of traveling wave solutions for monotone operator enable us also to prove the existence of traveling wave solutions for non monotone operators.

In order to achieve the last goal, we must introduce/define the Banach space

$$
\mathcal{B}_{\rho} = \{ u = (u^{i}) : u_{i} \in C(\mathbb{R}), \quad \sup_{\xi \in \mathbb{R}} |u_{i}(\xi)| e^{\rho \xi} < \infty, i = 1, ..., N \}
$$

which comes equipped with the weighted norm

$$
||u||_{\rho} = \sum_{i=1}^{N} \sup_{\xi \in \mathbb{R}} |u_i(\xi)| e^{\rho \xi},
$$

where $C(\mathbb{R})$ is the set of all continuous functions on \mathbb{R} and ρ is a positive constant such that $\rho < \Lambda_c$. It follows that $\phi^+ \in \mathcal{B}_{\rho}$ and $\phi^- \in \mathcal{B}_{\rho}$. We now consider the following set

$$
\mathcal{A}_{\rho} = \{ u : u \in \mathcal{B}_{\rho}, \phi^{-}(\xi) \le u(\xi) \le \phi^{+}(\xi), \xi \in \mathbb{R} \}
$$

It is clear that $\mathcal{A}_{\rho} \subseteq \mathcal{C}_{\beta}$. We proceed to show that $\mathcal{Q}_{c}[F]$ is a continuous map of the bounded set \mathcal{A}_{ρ} into a compact set (standard procedure as it can be seen in [\[21](#page-27-9)[,8](#page-26-9)[,23\]](#page-27-10)).

Lemma 4.4 Assume (H1)−(H3) hold. Then $Q_c[F] : A_\rho \to B_\rho$ is continuous with the weighted norm $\lVert . \rVert_{\rho}$.

PROOF. By the assumptions, let $M > 0$ be the Lipschitz constant of all f_i

on
$$
[0, \beta_i^+]
$$
, $i = 1, ..., N$. For any $u = (u^i), v = (v^i) \in A_\rho$, we have, for $\xi \in \mathbb{R}$

$$
|Q_c^i[F(u)](\xi) - Q_c^i[F(v)](\xi)|e^{\rho\xi}
$$

\n
$$
= \int_{\mathbb{R}} k_i(\xi + c - y)|f_i(u(y)) - f_i(v(y))|dy e^{\rho\xi}
$$

\n
$$
\leq M \int_{\mathbb{R}} k(\xi + c - y)|u(y) - v(y)|e^{\rho y}e^{-\rho y}dy e^{\rho\xi}
$$

\n
$$
\leq M \int_{\mathbb{R}} k(\xi + c - y)e^{-\rho y}e^{\rho\xi}dy||u(y) - v(y)||_{\rho}
$$

\n
$$
= M \int_{\mathbb{R}} k_i(y)e^{\rho(y-\xi-c)}e^{\rho\xi}dy||u(y) - v(y)||_{\rho}
$$

\n
$$
= e^{-\rho c}M \int_{\mathbb{R}} k_i(y)e^{\rho y}dy||u(y) - v(y)||_{\rho}.
$$
 (4.15)

and

$$
\|\mathcal{Q}_c[F(u)] - \mathcal{Q}_c[F(v)]\|_{\rho} \leq \left(e^{-\rho c} M \sum_{i=1}^N \int_{\mathbb{R}} k_i(y) e^{\rho y} dy\right) \|u - v\|_{\rho}.
$$

Thus, $\mathcal{Q}[u]$ is continuous. We now show that the set $\mathcal{Q}_c[F](\mathcal{A})$ is relatively compact in \mathcal{B}_{ρ} . If $u \in \mathcal{A}_{\rho}, \xi \in \mathbb{R}$ and δ , we have, $i = 1, ..., N$

$$
|Q_c^i[F(u)](\xi + \delta) - Q_c^i[F(u)](\xi)|
$$

\n
$$
\leq \int_{\mathbb{R}} |k_i(\xi + \delta + c - y) - k_i(\xi + c - y)|f_i(u(y))dy
$$

\n
$$
\leq R_i \int_{\mathbb{R}} |k_i(\xi + \delta + c - y) - k_i(\xi + c - y)|dy
$$

\n
$$
= R_i \int_{\mathbb{R}} |k_i(\delta + y) - k_i(y)|dy
$$
\n(4.16)

and the fact that $\int_{\mathbb{R}} |k_i(\delta + y) - k_i(y)| dy \to 0$ if $\delta \to 0$ implies

$$
\lim_{\delta \to 0} (\mathcal{Q}_c^i[F(u)](\xi + \delta) - \mathcal{Q}_c^i[F(u)](\xi)) = 0 \tag{4.17}
$$

uniformly for all $u \in A_{\rho}, \xi \in \mathbb{R}, i = 1, ..., N$. Take any sequence $(u_n) = ((u_n^i)) \in$ \mathcal{A}_{ρ} and let $v_n = (v_n^i) = \mathcal{Q}_c[F(u_n)]$. From [\(4.17\)](#page-14-0), (v_n) is uniformly bounded on R and uniformly equicontinuous. For $I_k = [-k, k]$, $k \in \mathbb{N}$, by Ascoli's theorem and the standard diagonal process, we can construct subsequences (u_{n_k}) of (u_n) such that there is a function $v = (v^i), v_i \in C(-\infty, \infty), i = 1, ..., N$ and $(v_{n_k}) = (Q_c[F(u_{n_k})])$ uniformly converges to v on each I_k for $k \in \mathbb{N}$. Now we need to show that $v \in A_\rho$ and $||v_{n_k} - v||_\rho \to 0$ as $n_k \to \infty$. It follows from Lemmas [4.1](#page-10-0) and [4.3,](#page-11-2) $\phi_i^-(\xi) \le v_i(\xi) \le \phi_i^+(\xi)$, $i = 1, ..., N$ for all $\xi \in \mathbb{R}$, and therefore $v \in \mathcal{A}_{\rho}$. Note that

$$
\lim_{\xi \to \pm \infty} (\phi_i^+(\xi) - \phi_i^-(\xi))e^{\rho \xi} = 0, i = 1, ..., N.
$$

For any $\epsilon > 0$, we can find $K_0 > 0$ such that if $|\xi| > K_0$, then, for all $k \in \mathbb{N}$

$$
|v_{n_k}^i(\xi) - v^i(\xi)|e^{\rho \xi} \le (\phi_i^+(\xi) - \phi_i^-(\xi))e^{\rho \xi} < \epsilon, i = 1, ..., N.
$$

On the other hand, on $[-I_k, I_k]$, (v_{n_k}) uniformly converges to v. Thus there exists a $N_1 > 0$ such that, for $n_k > N_1$

$$
|v_{n_k}^i(\xi) - v^i(\xi)|e^{\rho \xi} < \epsilon, \ \xi \in [-K_0, K_0], i = 1, ..., N
$$

Consequently, if $n_k > N_1$, the following inequality is true for all $\xi \in \mathbb{R}$

$$
|v_{n_k}^i(\xi) - v^i(\xi)|e^{\rho \xi} < \epsilon, i = 1, ..., N.
$$

Thus $||v_{n_k} - v||_\rho \to 0$ as $n_k \to \infty$. \Box

Now we are in a position to prove Theorem [2.2](#page-6-0) when F is monotone.

Define the following iteration

$$
u_1 = (u_1^i) = \mathcal{Q}_c[F(\phi^+)], \ u_{n+1} = (u_n^i) = \mathcal{Q}_c[F(u^n)], n > 1.
$$
 (4.18)

From Lemmas [4.1,](#page-10-0) [4.3,](#page-11-2) and the fact that F is nondecreasing, u_n is nondecreasing on R, it follows that

$$
\phi_i^-(\xi) \le u_{n+1}^i(\xi) \le u_n^i(\xi) \le \phi_i^+(\xi), \xi \in \mathbb{R}, n \ge 1, n = 1, ..., N.
$$

By Lemma [4.4](#page-13-0) and monotonicity of (u_n) , there is $u \in \mathcal{A}$ such that $\lim_{n\to\infty}||u_n-\hat{u}|$ $||u||_{\rho} = 0.$ Lemma [4.4](#page-13-0) implies that $\mathcal{Q}[u] = u$. Furthermore, u is nonincreasing. It is clear that $\lim_{\xi \to \infty} u^i(\xi) = 0, i = 1, ..., N$. Assume that $\lim_{\xi \to -\infty} u_i(\xi) =$ $\hat{k}^i, i = 1, ..., N$ $\hat{k}^i > 0, i = 1, ..., N$ because of $u \in A_\rho$. Applying the dominated convergence theorem, we get $\hat{k}_i = f_i(\hat{k})$. By (H2), $\hat{k} = \beta$. Finally, note that

$$
\nu_{\Lambda_c}^i(e^{-\Lambda_c \xi} - q e^{-\gamma \Lambda_c \xi}) \le u^i(\xi) \le \nu_{\Lambda_c}^i e^{-\Lambda_c \xi}, \xi \in \mathbb{R}.
$$

We immediately obtain that

$$
\lim_{\xi \to \infty} u^i(\xi) e^{\Lambda_c \xi} = \nu^i_{\Lambda_c}, i = 1, ..., N. \tag{4.19}
$$

This completes the proof of Theorem [2.2](#page-6-0) when F is monotone.

4.3 Proof of Theorem [2.2](#page-6-0) (iii)

PROOF. In this section, the assumption that F is monotone is *dropped*. As in Section [4,](#page-9-0) both $\mathcal{Q}_c[F^+]$ and $\mathcal{Q}_c[F^-]$ are monotone. Note that $F, F^+, F^$ have the same linearization at the origin. In view of Section [4,](#page-9-0) there exists a nonincreasing fixed point $u_{-} = (u_{-}^i) \in \mathcal{C}_{\beta^{-}}$ of $\mathcal{Q}_c[F^-]$ such that

$$
\mathcal{Q}_c[F^-(u^-)] = u^-
$$

and $\lim_{\xi \to -\infty} u_i^-(\xi) = \beta_i^-, i = 1, ..., N$, and $\lim_{\xi \to \infty} u_i^-(\xi) = 0, i = 1, ..., N$. Furthermore, $\lim_{\xi \to \infty} u_i^-(\xi) e^{\Lambda_c \xi} = \nu_{\Lambda_c}^i, i = 1, ..., N$. Let

$$
\widetilde{\phi^+}(\xi) = (\widetilde{\phi_i^+}),
$$

where

$$
\phi_i^+ = \min \{ \beta_i^+, \nu_{\Lambda_c}^i e^{-\Lambda_c \xi} \}, \ \xi \in \mathbb{R}, i = 1, ..., N.
$$

According to Lemma [4.1,](#page-10-0) ϕ^+ is an upper solution of $\mathcal{Q}_c[F^+]$. Also note that if β^+ is replaced with β^- , $\phi^+(\xi)$ is an upper solution of $\mathcal{Q}_c[F^-]$. By the construction of $u^i_-(\xi)$, it then follows that

$$
u_{-}(\xi) \leq \widetilde{\phi^{+}}(\xi), \xi \in \mathbb{R}
$$

Now let

$$
\mathcal{D}_{\rho} = \{ u : u = (u^i) \in \mathcal{B}_{\rho}, u^i_{-}(\xi) \le u^i(\xi) \le \widetilde{\phi_i^+}(\xi), \xi \in (-\infty, \infty), i = 1, ..., N \},\
$$

where \mathcal{B}_{ρ} is defined in Section [4.](#page-9-0) It is clear that \mathcal{D}_{ρ} is a bounded nonempty closed convex subset in \mathcal{B}_{ρ} . Furthermore, we have, for any $u = (u^{i}) \in \mathcal{D}$

$$
u_{-} = \mathcal{Q}_c[F^{-}(u_{-})] \leq \mathcal{Q}_c[F^{-}(u)] \leq \mathcal{Q}_c[F(u)] \leq \mathcal{Q}_c[F^{+}(u)] \leq \mathcal{Q}_c[F^{+}(\widetilde{\phi^+})] \leq \widetilde{\phi^+}.
$$

Therefore, $\mathcal{Q}_c[F] : \mathcal{D}_{\rho} \to \mathcal{D}_{\rho}$. Note that the proof of Lemmas [4.4](#page-13-0) does not need the monotonicity of F^- . In the same way as in Lemmas [4.4,](#page-13-0) we can show that $\mathcal{Q}_c[F^-]: \mathcal{D}_{\rho} \to \mathcal{B}_{\rho}$ is continuous and maps bounded sets into compact sets. Therefore, the Schauder Fixed Point Theorem shows that the operator $\mathcal{Q}_c[F]$ has a fixed point u in \mathcal{D}_ρ , which is a traveling wave solution of [\(1.3\)](#page-3-0) for $c > c^*$. Since $u_i^-(\xi) \leq u^i(\xi) \leq \phi_i^+(\xi), \xi \in (-\infty, \infty), i = 1, ..., N$, it is easy to see that for $i = 1, ..., N$, $\lim_{\xi \to \infty} u_i(\xi) = 0$, $\lim_{\xi \to \infty} u^i(\xi) e^{\Lambda_c \xi} = \nu_{\Lambda_c}^i$,

$$
\beta_i^- \le \liminf_{\xi \to -\infty} u^i(\xi) \le \limsup_{\xi \to -\infty} u^i(\xi) \le \beta_i^+
$$

and $0 < u^i_-(\xi) \leq u^i(\xi) \leq \beta_i^+, \xi \in (-\infty, \infty)$. \Box

4.4 Proof of Theorem [2.2](#page-6-0) (iv)

PROOF. The proof in this subsection follows the approach found in [\[3,](#page-26-13)[8\]](#page-26-9). We make use of the results in Theorem [2.2](#page-6-0) (iii). Hence, for each $m \in \mathbb{N}$, we choose $c_m > c^*$ such that $\lim_{m \to \infty} c_m = c^*$. According to Theorem [2.2](#page-6-0) (iii), for each c_m there is a traveling wave solution $u_m = (u_m^i)$ of (1.3) such that

$$
u_m = \mathcal{Q}[F(u_m)](\xi + c_m).
$$

and

$$
\beta_i^- \le \liminf_{\xi \to -\infty} u_m^i(\xi) \le \limsup_{\xi \to -\infty} u_m^i(\xi) \le \beta_i^+, i = 1, ..., N.
$$

As it was shown in (4.16) , (u_m) is equicontinuous and uniformly bounded on R. Hence, the Ascoli's theorem implies that there is vector valued continuous function $u = (u^i)$ on $\mathbb R$ and subsequence (u_{m_k}) of (u_m) such that

$$
\lim_{m \to \infty} u_m(\xi) = u(\xi)
$$

uniformly in ξ on any compact interval of R. Further, the use of the dominated convergence theorem guarantees that we have

$$
u = \mathcal{Q}[F(u)](\xi + c^*).
$$

Thus u is a traveling solution of (1.3) for $c = c^*$ and satisfies

$$
\beta_i^- \leq \liminf_{\xi \to -\infty} u^i(\xi) \leq \limsup_{\xi \to -\infty} u^i(\xi) \leq \beta_i^+, i = 1, ..., N
$$

Because of the translation invariance of u_m , we always can assume that $u_m(0) \leq$ 1 $\frac{1}{2}\beta^-$ for all m. Consequently u is not a constant traveling solution of [\(1.3\)](#page-3-0). \Box

4.5 Proof of Theorem [2.2](#page-6-0) (v)

PROOF. The proof of this subsection follows that in [\[8,](#page-26-9)[16\]](#page-27-8). Suppose, by contradiction, that for some $c \in (0, c^*)$, [\(1.3\)](#page-3-0) has a traveling wave $u_n(x) =$ $u(x - cn)$ such that $u \in \mathcal{C}_{\beta}$ with $\liminf_{x \to -\infty} u(x) > 0$ and $u(+\infty) = 0$. Thus $u(x)$ can be larger than a positive vector with arbitrary length. It follows from Theorem [2.2](#page-6-0) (ii)

$$
\liminf_{n \to \infty} \inf_{|x| \leq nc} u_n(x) \geq \beta^-, \text{ for } 0 < c < c^*
$$

Let $\hat{c} \in (c, c^*)$ and $x = \hat{c}n$. Then

$$
\lim_{n \to \infty} u((\hat{c} - c)n) = \lim_{n \to \infty} u_n(\hat{c}n) \ge \liminf_{n \to \infty} \inf_{|x| \le n\hat{c}} u_n(x) \ge \beta^-.
$$

However,

$$
\lim_{n \to \infty} u((\hat{c} - c)n) = u(\infty) = 0,
$$

which is a contradiction. \Box

5 An example

Hassell and Comins [\[6\]](#page-26-10) as noted in the first section of this manuscript analyze a discrete time model for the dynamics of two competing species. Their model describes the growth and spread of two population densities at time n and location x under an interference competition regime. If the two densities are denoted by $X_n(x)$ and $Y_n(x)$ then the model is given by the set the nonlinear coupled difference equations [\(1.1\)](#page-2-0). The addition of the possibility of dispersal via the re-distribution kernel $k_i(x - y)$ leads to Model [\(1.2\)](#page-2-1) also described in the introductory section of this manuscript.

Model [\(1.2\)](#page-2-1) has four equilibria $(0, 0), (0, r_2), (r_1, 0)$ and

$$
(\frac{r_1 - \sigma_1 r_2}{1 - \sigma_1 \sigma_2}, \frac{r_2 - \sigma_2 r_1}{1 - \sigma_1 \sigma_2}).
$$

The change of variables

$$
p = X, q = r_2 - Y
$$

allows to convert system [\(1.2\)](#page-2-1) into the following coupled system of integrodifference equations

$$
p_{n+1}(x) = \int_{\mathbb{R}} k_1(x - y) f(p_n(y), q_n(y)) dy
$$

\n
$$
q_{n+1}(x) = \int_{\mathbb{R}} k_2(x - y) g(p_n(y), q_n(y)) dy.
$$
\n(5.20)

where

$$
f(p,q) = h(p)e^{r_1 - \sigma_1 r_2 + \sigma_1 q}
$$

$$
g(p,q) = r_2 - (r_2 - q)e^{q - \sigma_2 p}
$$

$$
h(p) = pe^{-p}
$$

It is clear that [\(1.2\)](#page-2-1) and [\(5.20\)](#page-18-1) are not monotone systems. Theorem [2.2](#page-6-0) can be applied to Model [\(5.20\)](#page-18-1) guaranteeing the existence of traveling waves as well as results on the speed of propagation. In fact, we can derive the corresponding results for Model [\(1.2\)](#page-2-1) in the context of our example, Model [\(5.20\)](#page-18-1). A straightforward calculation shows that Model [\(5.20\)](#page-18-1) has four equilibria $(0, 0), (0, r_2), (r_1, r_2)$ and

$$
(\frac{r_1 - \sigma_1 r_2}{1 - \sigma_1 \sigma_2}, \sigma_2 \frac{r_1 - \sigma_1 r_2}{1 - \sigma_1 \sigma_2}).
$$
\n(5.21)

If it is further assumed that $r_1 > 1, r_2 < 1$, and $\sigma_1 < 1, \sigma_2 > 1, \sigma_1 \sigma_2 < 1$ then

$$
\left(\frac{r_1 - \sigma_1 r_2}{1 - \sigma_1 \sigma_2}, \sigma_2 \frac{r_1 - \sigma_1 r_2}{1 - \sigma_1 \sigma_2}\right) \gg (r_1, r_2). \tag{5.22}
$$

Thus [\(5.20\)](#page-18-1) has no other positive equilibrium (p, q) between $(0, 0)$ and (r_1, r_2) with $p > 0$ and $q > 0$. Observe that 1 is the maximum point of $h(p)$. That is, $h(p)$ is not monotone on [0, r_1]. Further simple calculations show that

$$
g_p(p,q) = \sigma_2(r_2 - q)e^{q - \sigma_2 p} \ge 0, \text{ for } q \in [0, r_2]
$$

$$
g_q(p,q) = (1 - r_2 + q)e^{q - \sigma_2 p} \ge 0
$$

Theorem [2.2](#page-6-0) can now be used to guarantee the existence of a spreading speed and traveling wave solutions of the nonmonotone system [\(5.20\)](#page-18-1). We summarize the results obtained in the context of this example in Theorem [5.1.](#page-19-0) In fact, assumption $r_2 < e^{r_1-1-e^{r_1-1}}$ can be relaxed as long as [\(5.38\)](#page-24-0) $(r_2 \le t_0)$ holds.

Theorem 5.1 Let $0 < r_2 < 1 < r_1, 0 < \sigma_1 < 1 < \sigma_2, \sigma_1 \sigma_2 < 1, r_2 <$ $e^{r_1-1-e^{r_1-1}}$ and $\sigma_2 e^{r_1-1-e^{r_1-1}} > r_2$. Assume that k_1, k_2 satisfy (H1) and $\int_{\mathbb{R}} k_2(s)e^{\mu s}ds \leq \int_{\mathbb{R}} k_2(s)e^{\mu s}ds$ for $\mu > 0$. Then the conclusions of Theorem [2.2](#page-6-0) hold for [\(5.20\)](#page-18-1).

In order to use Theorem [2.2,](#page-6-0) we define the upper monotone function

$$
h^{+}(p) = \begin{cases} h(p), & 0 \le p \le 1, \\ h(1) = e^{-1}, & 1 \le p. \end{cases}
$$

and corresponding monotone systems with h^+

$$
p_{n+1}(x) = \int_{\mathbb{R}} k_1(x - y) f^{+}(p_n(y), q_n(y)) dy
$$

\n
$$
q_{n+1}(x) = \int_{\mathbb{R}} k_2(x - y) g(p_n(y), q_n(y)) dy.
$$
\n(5.23)

where

$$
f^+(p,q) = h^+(p)e^{r_1 - \sigma_1 r_2 + \sigma_1 q}
$$

Note that $g(p,q) = q$ has only two possible solutions $q = r_2$ and $q = \sigma_2 p$. Based on this observation, it is clear that (5.23) has three equilibria $(0,0), (0, r_2), (e^{r_1-1}, r_2)$. If [\(5.23\)](#page-19-1) has another positive equilibrium (p^*, q^*) , it must satisfy $p^* > 1$ (otherwise, (p^*, q^*) is (5.21) , which contradicts (5.22)) and therefore

$$
e^{r_1 - \sigma_1 r_2 + \sigma_1 \sigma_2 p^* - 1} = p^*
$$

\n
$$
q^* = \sigma_2 p^*.
$$
\n(5.24)

We will use a simply inequality to estimate e^x :

$$
e^x \ge x + 1, x \in \mathbb{R}.
$$

Thus $p^* = e^{r_1 - \sigma_1 r_2 + \sigma_1 \sigma_2 p^* - 1} \ge r_1 - \sigma_1 r_2 + \sigma_1 \sigma_2 p^*$, we have

$$
p^* \ge \frac{r_1 - \sigma_1 r_2}{1 - \sigma_1 \sigma_2} > r_1,
$$

which further implies that $p^* > e^{r_1-1}$ as $\sigma_2 r_1 > r_2$. Again since $\sigma_2 r_1 > r_2$, we have

$$
q^* = \sigma_2 p^* > r_2.
$$

Thus [\(5.23\)](#page-19-1) has on other positive equilibrium between $(0,0)$ and (e^{r_1-1}, r_2) .

There is a $t_0 \in (0,1]$ such that $h(t_0) = h(e^{r_1-1})$ and define

$$
h^{-}(p) = \begin{cases} h(p), & 0 \le p \le t_0, \\ h(t_0), & t_0 \le p \le e^{r_1 - 1}. \end{cases}
$$

and corresponding lower monotone system

$$
p_{n+1}(x) = \int_{\mathbb{R}} k_1(x - y) f^-(p_n(y), q_n(y)) dy
$$

\n
$$
q_{n+1}(x) = \int_{\mathbb{R}} k_2(x - y) g(p_n(y), q_n(y)) dy.
$$
\n(5.25)

where

$$
f^-(p,q) = h^-(p)e^{r_1 - \sigma_1 r_2 + \sigma_1 q}
$$

Then

 $h^{\pm}(p)$

$$
0 < h^{-}(p) \le h(p) \le h^{+}(p) \le h'(0)p, p \in (0, e^{r_1 - 1}]
$$
\n
$$
\pm(p) = h(p), \text{ on a small right neighborhood of } 0, \quad h^{-}(0) = h^{+}(0) = 0.
$$

Again we observe that $g(p,q) = q$ has only two possible solutions $q = r_2$ and $q = \sigma_2 p$. In view of the fact that $h(t_0) = t_0 e^{-t_0} = h(e^{r_1-1}) = e^{r_1-1}e^{-e^{r_1-1}}$, we can calculate that [\(5.25\)](#page-20-0) has three equilibria $(0,0), (0, r_2)$ and (t_1, r_2) where $t_1 = e^{2r_1 - 1 - e^{r_1 - 1}}$. We shall prove

$$
t_1 < r_1,\tag{5.26}
$$

see Fig. [1.](#page-21-0)

Indeed, [\(5.26\)](#page-20-1) is equivalent to the following

$$
2r_1 - 1 - e^{r_1 - 1} < \ln r_1.
$$

Fig. 1. The curve below is $t_1 = e^{2r_1 - 1 - e^{r_1 - 1}}$ and the line above is r_1 . t_1 can be larger than 1, but is always less than r_1 .

Let $l(r_1) = 2r_1 - 1 - e^{r_1-1} - \ln r_1$. Then $l(1) = 0$ and

$$
l'(r_1) = 2 - e^{r_1 - 1} - \frac{1}{r_1} \le 2 - r_1 - \frac{1}{r_1} < 0, \text{ for } r_1 > 1.
$$

This verifies [\(5.26\)](#page-20-1). Since $0 < t_0 < 1 < r_1$, the following inequality holds

$$
0 < e^{r_1 - 1 - e^{r_1 - 1}} < t_0 = e^{t_0} e^{r_1 - 1} e^{-e^{r_1 - 1}} < e^{2r_1 - 1 - e^{r_1 - 1}} = t_1.
$$
 (5.27)

If (p^*, q^*) is a another positive equilibrium of [\(5.25\)](#page-20-0), it must satisfy $p^* > t_0$ (otherwise, (p^*, q^*) is (5.21) , which contradicts (5.22)) and therefore

$$
e^{r_1 - 1}e^{-e^{r_1 - 1}}e^{r_1 - \sigma_1 r_2 + \sigma_1 \sigma_2 p^*} = p^*
$$

\n
$$
q^* = \sigma_2 p^*.
$$
\n(5.28)

Since $r_1 > \sigma_1 r_2$ and $p^* > 0$, we have $p^* > e^{r_1-1-e^{r_1-1}}$. In view of the assumption that $\sigma_2 e^{r_1-1-e^{r_1-1}} > r_2$, we have

$$
q^* > r_2,
$$

and therefore from [\(5.28\)](#page-21-1)

$$
p^* > e^{2r_1 - 1 - e^{r_1 - 1}} = t_1,
$$

Thus [\(5.25\)](#page-20-0) has on other positive equilibrium between $(0, 0)$ and (t_1, r_2) and

$$
(0,0) << (t_1, r_2) \le (r_1, r_2) \le (e^{r_1 - 1}, r_2).
$$

This verifies (H2) for [\(5.20\)](#page-18-1).

Fig. 2. The construction of h^+ and h^- . The red curve is h.

The matrix in (2.4) for (5.20) is

$$
B_{\mu} = (a_{\mu}^{i,j}) = \begin{pmatrix} e^{r_1 - \sigma_1 r_2} \int_{\mathbb{R}} k_1(s) e^{\mu s} ds & 0\\ r_2 \sigma_2 \int_{\mathbb{R}} k_2(s) e^{\mu s} ds & (1 - r_2) \int_{\mathbb{R}} k_2(s) e^{\mu s} ds \end{pmatrix}
$$
(5.29)

Since $e^{r_1 - \sigma_1 r_2} > 1 > 1 - r_2$, the principle eigenvalue for the matrix is

$$
\lambda(\mu) = e^{r_1 - \sigma_1 r_2} \int_{\mathbb{R}} k_1(s) e^{\mu s} ds
$$

and the corresponding positive eigenvector

$$
\eta_{\mu} = \begin{pmatrix} \nu_{\mu}^{(1)} \\ \nu_{\mu}^{(2)} \end{pmatrix} = \begin{pmatrix} \frac{e^{r_1 - \sigma_1 r_2} \int_{\mathbb{R}} k_1(s)e^{\mu s}ds - (1 - r_2) \int_{\mathbb{R}} k_2(s)e^{\mu s}ds}{r_2 \sigma_2 \int_{\mathbb{R}} k_2(s)e^{\mu s}ds} \\ 1 \end{pmatrix} . \tag{5.30}
$$

Because $\int_{\mathbb{R}} k_2(s)e^{\mu s}ds \leq \int_{\mathbb{R}} k_1(s)e^{\mu s}ds$, canceling $\int_{\mathbb{R}} k_2(s)e^{\mu s}ds$ in $\nu_{\mu}^{(1)}$ leads to

$$
\nu_{\mu}^{(1)} \ge \frac{e^{r_1 - \sigma_1 r_2} - (1 - r_2)}{r_2 \sigma_2}
$$
\n
$$
\ge \frac{1}{\sigma_2} + \frac{e^{r_1 - \sigma_1 r_2} - 1}{r_2 \sigma_2}
$$
\n
$$
\ge \frac{1}{\sigma_2}
$$
\n(5.31)

It is clear now $(H3)(i)$ holds. Now we can verify $(H3)(ii)$ for (5.23) . Let

$$
(p,q) = (\min\{e^{r_1-1}, \nu_\mu^{(1)}\alpha\}, \min\{r_2, \alpha\}), \alpha > 0.
$$

Since $e^{q-\sigma_2 p} > 1 + q - \sigma_2 p$, we need to show that

$$
h^{+}(p)e^{r_{1}-\sigma_{1}r_{2}+\sigma_{1}q} \leq e^{r_{1}-\sigma_{1}r_{2}}p
$$

\n
$$
r_{2} - (r_{2} - q)e^{q-\sigma_{2}p} \leq r_{2}\sigma_{2}p + (1 - r_{2})q + q(q - \sigma_{2}p)
$$

\n
$$
\leq r_{2}\sigma_{2}p + (1 - r_{2})q
$$
\n(5.32)

Therefore, it is easy to see that we only need to verify that

$$
q \leq \sigma_2 p \tag{5.33}
$$

and

$$
\frac{h^+(p)}{p} \le e^{-\sigma_1 q} \tag{5.34}
$$

For [\(5.33\)](#page-23-0), we need to consider the two cases: $p = e^{r_1-1}$ and $p = \nu_{\mu}^{(1)}\alpha$. If $p=e^{r_1-1}$, then

$$
q \le r_2 \le \sigma_2 e^{r_1 - 1} \tag{5.35}
$$

which is true by the assumptions. If $p = \nu_{\mu}^{(1)} \alpha$, then

$$
q\leq \alpha\leq \sigma_2\nu_\mu^{(1)}\alpha
$$

which is true because of (5.31) . In order to verify (5.34) , first assume that $p \in [0,1],$ then $h^+(p) = pe^{-p}$ and $p = \nu_{\mu}^{(1)} \alpha$ since $e^{r_1-1} > 1$. Noting that $e^{-\sigma_1 \alpha} \leq e^{-\sigma_1 q}$, it suffices to verify

$$
e^{-\nu^{(1)}_\mu\alpha}\leq e^{-\sigma_1\alpha}
$$

which is true because of [\(5.31\)](#page-22-0) and $\sigma_2 \sigma_1 < 1$. For the case that $p \geq 1$, then $h^+(p) = e^{-1}$. Again noting that

$$
e^{-r_2} \le e^{-\sigma_1 r_2} \le e^{-\sigma_1 q},\tag{5.36}
$$

it suffices to verify

$$
\frac{e^{-1}}{p} \le e^{-1} \le e^{-r_2}
$$

which holds because $r_2 < 1$.

It remains to verify $(H3)(ii)$ for (5.25) . Let

$$
(p,q) = (\min\{t_1, \nu_\mu^{(1)}\alpha\}, \min\{r_2, \alpha\}), \alpha > 0.
$$

For [\(5.33\)](#page-23-0), we need to consider the two cases: $p = t_1$ and $p = \nu_{\mu}^{(1)} \alpha$. If $p = t_1$, then

$$
q \le r_2 \le \sigma_2 t_1 = \sigma_2 e^{2r_1 - 1 - e^{r_1 - 1}} \tag{5.37}
$$

which is true by the assumptions. If $p = \nu_{\mu}^{(1)} \alpha$, then

$$
q \le \alpha \le \sigma_2 \nu_{\mu}^{(1)} \alpha
$$

which is true because of (5.31) .

Now we need to verify (5.34) (with h^+ being replaced by h^-) for (5.25) . If $0 < p < t_0$, then $h^-(p) = pe^{-p}$ and $p = \nu_\mu^{(1)} \alpha$ because of [\(5.27\)](#page-21-2). Noting that $e^{-\sigma_1\alpha} \leq e^{-\sigma_1 q}$, it suffices to verify

$$
e^{-\nu^{(1)}_\mu\alpha}\leq e^{-\sigma_1\alpha}
$$

which is true because of [\(5.31\)](#page-22-0) and $\sigma_2 \sigma_1 < 1$. For the case that $p \geq t_0$, then $h^-(p) = h(t_0)$. And from the definition of h^- and [\(5.36\)](#page-23-2), it suffices to verify

$$
\frac{h^-(p)}{p} \le \frac{h(t_0)}{t_0} = e^{-t_0} \le e^{-r_2}.\tag{5.38}
$$

[\(5.38\)](#page-24-0) holds since, from [\(5.27\)](#page-21-2) and the assumption,

$$
e^{-t_0} \le e^{-e^{r_1 - 1 - e^{r_1 - 1}}} \le e^{-r_2}.\tag{5.39}
$$

To verify $(H3)(iii)$, we note that, from Lemma [4.2,](#page-11-0) for sufficiently larger k, there is a small $\omega >> 0$, if $0 \leq (p, q) \leq \omega$,

$$
f(p,q) \ge f(p,0) \ge (1 - \frac{1}{k})e^{r_1 - \sigma_1 r_2}p
$$

and

$$
g(p,q) \ge (1 - \frac{1}{k})r_2 \sigma_2 p + (1 - \frac{1}{k})(1 - r_2)q.
$$

Thus we verify all the conditions (H1-H3) and conclude the proof of Theorem [5.1.](#page-19-0)

6 Conclusions

Integrodifference systems arise naturally in the study of the dispersal of populations, including interacting populations, composed of organisms that reproduce locally via discrete generations before dispersing. The brunt of the mathematical research has focused on the the study of the existence of traveling wave solutions and characterizations of the spreading speed in the context of cooperative systems. In this paper, we characterize the spreading speed for a large class of non cooperative systems, formulated in terms of integrodifference equations, via the convergence of initial data to wave solutions. The spreading speed is characterized as the slowest speed of a family of non-constant traveling wave solutions. The results are applied to the non-cooperative competitive system proposed by Hassell and Comins (1976) [\[6\]](#page-26-10). We are in the process of applying the approach outlined in the last Section of this manuscript to ecological and epidemiological systems where the local dynamics are naturally non-cooperative with the hope of increasing our understanding of the role of dispersal in the shaping communities where the local dynamics are more realistic than those previously supported by the mathematical theory.

7 Appendix

Proof of Proposition [2.1](#page-5-0) (iv). If $\Phi(\mu) = \frac{1}{\mu} \ln \lambda(\mu)$ achieves its minimum at a finite μ , then $c^* = \min_{\mu>0} \Phi(\mu) > 0$. Now let $c^* = \lim_{\mu \to \infty} \frac{1}{\mu}$ $\frac{1}{\mu} \ln \lambda(\mu)$. We recall that $\lambda(\mu)$ is the principal eigenvalue of its largest diagonal irreducible block and it has a positive eigenvector, see [\[7\]](#page-26-11). Thus there exists a positive constant $\delta > 0$ and a positive integer $i \leq N$ such that

$$
\lambda(\mu) \ge \delta \int_{\mathbb{R}} k_i(x) e^{\mu x} dx
$$

Thus

$$
c^* \ge \lim_{\mu \to \infty} \frac{1}{\mu} \ln \left(\delta \int_{\mathbb{R}} k_i(x) e^{\mu x} dx \right)
$$

Let

$$
\Psi(\mu) = \frac{\int_{\mathbb{R}} x k_i(x) e^{\mu x} dx}{\int_{\mathbb{R}} k_i(x) e^{\mu x} dx} \ \mu \ge 0.
$$

Then by the L'Hopital's rule we have

$$
c^* \geq \lim_{\mu \to \infty} \Psi(\mu)
$$

Differentiation of Ψ and rearrangement of terms show

$$
\Psi'(\mu) = \frac{\int_{\mathbb{R}} \left(x - \Psi(\mu)\right)^2 k_i(x) e^{\mu x} dx}{\int_{\mathbb{R}} k_i(x) e^{\mu x} dx} > 0, \mu \ge 0,
$$

also see Weinberger [\[29\]](#page-27-11). Note that $\Psi(0) = 0$ and therefore, $c^* \geq \lim_{\mu \to \infty} \Psi(\mu) >$ $0. \Box$

8 Acknowledgments

We like to thank Professor Bintuan Li for bringing [\[17\]](#page-27-14) to our attention which also investigated the minimum speed of [\(1.2\)](#page-2-1). This project have been partially supported by grants from the National Science Foundation (NSF - Grant DMS - 0502349), the National Security Agency (NSA - Grant H98230- 06-1-0097), the Alfred T. Sloan Foundation and the Office of the Provost at Arizona State University.

References

- [1] D. G. Aronson and H. F. Weinberger, Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation, in Partial Differential Equations and Related Topics, J. A. Goldstein, ed., Lecture Notes in Mathematics Ser. 446, Springer-Verlag, Berlin, 1975, pp. 549.
- [2] D. G. Aronson and H. F. Weinberger, Multidimensional nonlinear diffusion arising in population dynamics, Adv. Math., 30 (1978), pp. 3376.
- [3] K. Brown and J. Carr, Deterministic epidemic waves of critical velocity, Math. Proc. Cambridge Philos. Soc. 81 (1977) 431-433.
- [4] O. Diekmann, Thresholds and traveling waves for the geographical spread of an infection. J. Math. Biol. 6(1978) 109-130.
- [5] R. Fisher, The wave of advance of advantageous genes. Ann. of Eugenics, 7(1937) 355 369.
- [6] M. Hassell and H. Comins, Discrete time models for two-species competition. Theoretical Population Biology, 9(1976),202-221.
- [7] R. Horn, C. Johnson, Matrix Analysis. Cambridge: University Press, Cambridge 1985
- [8] S. Hsu and X-Q. Zhao, Spreading speeds and traveling waves for nonmonotone integrodifference equations, SIAM J. Math. Anal. 40(2008) 776-789.
- [9] Kierstad, H and L.B' Slobodkin (1953). "The size of water masses containing plankton blooms," J. Mar. Res. 12: 141-147
- [10] A. Kolmogorov, Petrovsky, N.I. Piscounov, Etude de lequation de la diffusion avec croissance de la quantite de matiere et son application a un probleme biologique. Bull. Moscow Univ. Math. Mech., 1(6), 126 (1937)
- [11] M. Kot, Discrete-time traveling waves: Ecological Examples. J. of Math. Biol., 30(1992) 413-436.
- [12] Levin, S. A. and R.T.Paine. "Disturbance, Patch Formation, and Community Structure," Proc.Nat. Acad. Sci. USA Vol. 71, No. 7, pp. 2744-2747, July 1974
- [13] Okubo, A., 1980:Diffusion and Ecological Problems: Mathematical Models. Springer-Verlag,Berlin.
- [14] M. Lewis, B. Li and H. Weinberger, Spreading speed and linear determinacy for two-species competition models, Journal of Mathematical Biology, 45(2002) 219-233.
- [15] B. Li, H. Weinberger, M. Lewis, Spreading speeds as slowest wave speeds for cooperative systems. Math. Biosciences. 196 (2005), no. 1, 82-98.
- [16] B. Li, M. Lewis and H. Weinberger, Existence of traveling waves for integral recursions with nonmonotone growth functions, Journal of Mathematical Biology, Journal of Mathematical Biology, 58(2009) 323-338.
- [17] B. Li, Personal communication.
- [18] X. Liang and X.-Q. Zhao, Asymptotic speeds of spread and traveling waves for monotone semiflows with applications, Communications on Pure and Applied Math., 60(2007), 1-40. Erratum: 61(2008), 137-138.
- [19] R. Lui, Biological growth and spread modeled by systems of recursions. I. Mathematical theory. Math. Biosciences 93 (1989), no. 2, 269295.
- [20] L. Rass and J. Radcliffe, Spatial deterministic epidemics, Povidence, American Mathematical Society, 2003.
- [21] S. Ma, Traveling waves for non-local delayed diffusion equations via auxiliary equation, Journal of Differential Equations, 237 (2007) 259-277.
- [22] H. R. Thieme, Density-Dependent Regulation of Spatially Distributed Populations and their Asymptotic speed of Spread. J. of Math. Biol., 8 (1979) 173-187.
- [23] H. Wang, On the existence of traveling waves for delayed reaction-diffusion equations, Journal of Differential Equations, 247(2009) 887-905.
- [24] Skellam,J.G.(1951), "Random dispersal in theoretical populations," Biometrika, 38,196-218.
- [25] Rios-Soto, K.R., Castillo-Chavez, C., Neubert, M., Titi, E.S., and A-A Yakubu. Epidemic Spread in Populations at Demographic Equilibrium. In: Mathematical Studies on Human Disease Dynamics: Emerging Paradigms and Challenges. Gumel A., Castillo-Chavez, C., Clemence, D.P. and R.E. Mickens, American Mathematical Society, pp. 297- 310, Vol. 410, (2006).
- [26] H. F. Weinberger, M. A. Lewis and B. Li, Analysis of linear determinacy for spread in cooperative models, J. Math. Biol. 45(2002) 183-218.
- [27] H. F. Weinberger, M. A. Lewis and B. Li, Anomalous spreading speeds of cooperative recursion systems, J. Math. Biol. 55(2007) 207-222.
- [28] H. F. Weinberger, Long-time behavior of a class of biological models. SIAM J. Math. Anal., 13 (1982) 353-396.
- [29] H. F. Weinberger, Asymptotic behavior of a model in population genetics. In Nonlinear Partial Differential Equations and Applications, ed. J. M. Chadam Lecture Notes in Mathematics, Volume 648, pages 47-96. Springer-Verlag, Berlin, 1978.
- [30] H. F. Weinberger, K. Kawasaki and N. Shigesada, Spreading speeds for a partially cooperative 2-species reaction-diffusion model, discrete and continuous dynamical systems, 23(2009), 10871098.
- [31] P. Weng, X-Q, Zhao, Spreading speed and traveling waves for a multi-type SIS epidemic model, Journal of Differential Equations, 229(2006) 270-296.