

New definite integrals and a two-term dilogarithm identity

F. M. S. Lima

Institute of Physics, University of Brasilia, P.O. Box 04455, 70919-970, Brasilia, DF, Brazil

Abstract

Among the several proofs known for $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/6$, the one by Beukers, Calabi, and Kolk involves the evaluation of $\int_0^1 \int_0^1 1/(1-x^2y^2) dx dy$. It starts by showing that this double integral is equivalent to $\frac{3}{4} \sum_{n=1}^{\infty} 1/n^2$, and then a non-trivial *trigonometric* change of variables is applied which transforms that integral into $\iint_T 1 du dv$, where T is a triangular domain whose area is simply $\pi^2/8$. Here in this note, I introduce a hyperbolic version of this change of variables and, by applying it to the above integral, I find exact closed-form expressions for $\int_0^{\infty} [\sinh^{-1}(\cosh u) - u] du$, $\int_{\alpha}^{\infty} [u - \cosh^{-1}(\sinh u)] du$, and $\int_{\alpha/2}^{\infty} \ln(\tanh u) du$, where $\alpha = \sinh^{-1}(1)$. From the latter integral, I also derive a two-term dilogarithm identity.

Keywords: Euler sums, Riemann zeta function, Double integrals, Hyperbolic functions, Dilogarithm identities

2010 MSC: 40C10, 11M06, 33B30

1. Introduction

The Riemann zeta function $\zeta(s)$ is defined, for complex values of s , $\Re(s) > 1$, by $\zeta(s) := \sum_{n=1}^{\infty} 1/n^s$. For integer values of s , $s > 1$, the first zeta value is $\zeta(2) = \sum_{n=1}^{\infty} 1/n^2$. Euler was the first to estimate the numerical value of this series to more than 5 decimal places (1735), as well as to determine an exact closed-form expression for it (1740), as given by [1]:

Lemma 1 (Euler's result for $\zeta(2)$). $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/6$.

From Euler's time onward, this result has been proved in several forms, from elementary to complex ones [2, 3]. Among these proofs, at least two involve the evaluation of a double integral over the unit square and these are the ones in

Email address: fabio@fis.unb.br (F. M. S. Lima)

which we are interested here. Let us briefly describe these two proofs for a better exposition of the hyperbolic approach and its consequences.

PROOF (APOSTOL'S PROOF). The short proof by Apostol (1983) can be found in Ref. [4]. He starts by expanding the integrand of Beukers' integral $I := \int_0^1 \int_0^1 \frac{1}{1-xy} dx dy$ as a geometric series, in order to show that it is equivalent to

$$\int_0^1 \int_0^1 (1 + xy + x^2y^2 + \dots) dx dy = \int_0^1 \left(1 + \frac{y}{2} + \frac{y^2}{3} + \dots \right) dy = \sum_{n=1}^{\infty} \frac{1}{n^2},$$

the interchange of integrals and sums being justifiable by the fact that each integrand is nonnegative and the sums converge absolutely (see the only corollary of Theorem 7.16 in Ref. [5]).¹ By applying a simple linear change of variables corresponding to the rotation of the coordinate axes through the angle $\pi/4$ radians, as given by²

$$x = \frac{u-v}{\sqrt{2}} \quad \text{and} \quad y = \frac{u+v}{\sqrt{2}},$$

directly to the Beukers' integral, one finds [4, 6]:

$$I = 4 \int_0^{\sqrt{2}/2} \frac{\arctan\left(\frac{u}{\sqrt{2-u^2}}\right)}{\sqrt{2-u^2}} du + 4 \int_{\sqrt{2}/2}^{\sqrt{2}} \frac{\arctan\left(\frac{\sqrt{2}-u}{\sqrt{2-u^2}}\right)}{\sqrt{2-u^2}} du.$$

Now, the substitution $u = \sqrt{2} \sin \theta$, followed by the use of the trigonometric identity $\arctan(\sec \theta - \tan \theta) = \pi/4 - \theta/2$ on the second integral, yields $I = \pi^2/18 + \pi^2/9$. Therefore, $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/18 + \pi^2/9 = \pi^2/6$. \square

The other proof, in which we are more interested here, was given by Beukers, Calabi, and Kolk (1993) [7].

PROOF (BCK PROOF). Similarly to Apostol's proof, Beukers, Calabi, and Kolk (BCK) start by showing that

$$\begin{aligned} K &:= \int_0^1 \int_0^1 \frac{1}{1-x^2y^2} dx dy = \int_0^1 \int_0^1 \sum_{n=0}^{\infty} x^{2n} y^{2n} dx dy \\ &= \sum_{n=0}^{\infty} \int_0^1 \int_0^1 x^{2n} y^{2n} dx dy = \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2}. \end{aligned} \quad (1)$$

Again, as each integrand is nonnegative the absolute convergence allows the interchange of the sum and the integral. Since $\sum_{n=1}^{\infty} 1/(2n)^2 = \frac{1}{4} \sum_{n=1}^{\infty} 1/n^2$,

¹Note that $1/(1-xy)$ is a positive real number at every point in the unit square, the only exception being the point $(1,1)$, where it is undefined. Of course, usual methods for treating improper integrals apply to this case.

²For this change of variables, it is easy to see that $|J| = 1$, J being the Jacobi determinant $\frac{d(x,y)}{d(u,v)} = \begin{vmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{vmatrix}$.

then $\sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = \frac{3}{4} \sum_{n=1}^{\infty} \frac{1}{n^2}$. They then evaluate the unit square integral that defines K by applying the following non-trivial, trigonometric change of variables [7]:

$$x = \frac{\sin u}{\cos v}, \quad y = \frac{\sin v}{\cos u}.$$

For this change of variables, one easily finds that $|J| = 1 - \tan^2 u \tan^2 v = 1 - x^2 y^2$, which reduces the integral for K to $\iint_T 1 \, du \, dv$ where T is the domain in the uv -plane defined by $\{(u, v) : u \geq 0, v \geq 0, u + v \leq \pi/2\}$ [2, 6, 7]. This double integral exactly evaluates to $\pi^2/8$, i.e. the area of the triangle corresponding to T , which implies Lemma 1. \square

Here in this work, I introduce a hyperbolic version of BCK change of variables for double integrals over the unit square. By applying this new change of variables to $\int_0^1 \int_0^1 1/(1 - x^2 y^2) \, dx \, dy$, I derive exact closed-form expressions for some definite integrals. From one of these integrals, I deduce a closed-form expression for $\text{Li}_2(\sqrt{2} - 1) + \text{Li}_2(1 - \sqrt{2}/2)$.

2. A hyperbolic version of BCK change of variables

Let us introduce the following *hyperbolic version* of BCK change of variables for double integrals over the unit square:

$$x = \frac{\sinh u}{\cosh v} \quad \text{and} \quad y = \frac{\sinh v}{\cosh u}. \quad (2)$$

The corresponding Jacobi determinant is:

$$J = \begin{vmatrix} \frac{\cosh u}{\cosh v} & -\frac{\sinh u \sinh v}{\cosh^2 v} \\ -\frac{\sinh u \sinh v}{\cosh^2 u} & \frac{\cosh v}{\cosh u} \end{vmatrix} = 1 - \tanh^2 u \tanh^2 v.$$

As $0 \leq x y \leq 1$ in the unit square, then $1 - x^2 y^2 \geq 0$, then $|J|$ is also equal to $1 - \tanh^2 u \tanh^2 v$. Therefore

$$\begin{aligned} \int_0^1 \int_0^1 F(x, y) \, dx \, dy &= \iint_S F(x(u, v), y(u, v)) |J(u, v)| \, du \, dv \\ &= \iint_S \tilde{F}(u, v) (1 - \tanh^2 u \tanh^2 v) \, du \, dv, \end{aligned} \quad (3)$$

where S is the cusped hyperbolic ‘quadrilateral’ on the right-hand side of Fig. 1. We are assuming that this change of variables is a C^1 diffeomorphism, which is reasonable since the functions $x = x(u, v)$ and $y = y(u, v)$ given in Eq. (2), as well as

$$u = u(x, y) = \cosh^{-1} \sqrt{\frac{1+x^2}{1-x^2 y^2}}$$

and

$$v = v(x, y) = \cosh^{-1} \sqrt{\frac{1+y^2}{1-x^2y^2}},$$

which define the *inverse* change,³ are continuously differentiable in all points of their domains (see the hachured regions in Fig. 1), the only exception being point $(x, y) = (1, 1)$. As indicated there in Fig. 1, S is bounded by the axes u and v , the curve $v = f(u) = \sinh^{-1}(\cosh u)$, above the diagonal $v = u$, and the curve $v = g(u) = \cosh^{-1}(\sinh u)$, below the diagonal. The parameter α can be readily evaluated by noting that it is the abscissa of the point where $v = g(u)$ intersects the u -axis. There in that point, one has $\cosh^{-1}(\sinh \alpha) = 0$, thus $\sinh \alpha = 1$. Since $\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1})$, $\forall x \in \mathbb{R}$, then

$$\alpha = \sinh^{-1}(1) = \ln(1 + \sqrt{2}). \quad (4)$$

3. New definite integrals and a dilogarithm identity

Let us now apply our hyperbolic change of variables to the unit square integral K , as defined in Eq. (1).

Theorem 1 (First definite integral).

$$\int_0^\infty [\sinh^{-1}(\cosh u) - u] du = \frac{\pi^2}{16}.$$

PROOF. From the BCK proof, one knows that $\int_0^1 \int_0^1 1/(1-x^2y^2) dx dy = \pi^2/8$. By combining this result with (3), one has

$$\begin{aligned} \frac{\pi^2}{8} &= \iint_S \frac{1}{1 - \tanh^2 u \tanh^2 v} (1 - \tanh^2 u \tanh^2 v) du dv \\ &= \iint_S 1 du dv = \text{area of } S. \end{aligned}$$

The area of the domain S can be equalized to a single definite integral by noting that S is symmetric with respect to the diagonal $v = u$ (see Fig. 1). Then, by taking only the upper half of S into account, one has

$$\text{area of } S = 2 \int_0^\infty \int_u^{f(u)} 1 dv du = 2 \int_0^\infty [\sinh^{-1}(\cosh u) - u] du,$$

which leads to the desired result. \square

³Note that the real function $\sinh^{-1} x$ is the inverse of the *bijective* function $\sinh x$ for all $x \in \mathbb{R}$, as usual. However, since the function $\cosh x$ is not bijective over the entire real domain, we can reduce its domain to the nonnegative reals and then *define* its inverse as $\cosh^{-1} x := \ln(x + \sqrt{x^2 - 1})$, $\forall x \geq 1$.

Theorem 2 (Second definite integral).

$$\int_{\alpha}^{\infty} [u - \cosh^{-1}(\sinh u)] du = \frac{\pi^2}{16} - \frac{1}{2} \ln^2(1 + \sqrt{2}).$$

PROOF. Similarly to the previous proof, by taking the lower half of S into account and noting that it is bounded by the u -axis, the diagonal $v = u$, and the curve $v = g(u)$, as seen in Fig. 1, one has, from Eq. (3), that:

$$\text{area of } S = 2 \left[\frac{\alpha^2}{2} + \int_{\alpha}^{\infty} \int_{g(u)}^u 1 dv du \right] = \alpha^2 + 2 \int_{\alpha}^{\infty} [u - \cosh^{-1}(\sinh u)] du.$$

Again, since the area of S is $\pi^2/8$, one finds

$$\int_{\alpha}^{\infty} [u - \cosh^{-1}(\sinh u)] du = \frac{1}{2} \left(\frac{\pi^2}{8} - \alpha^2 \right) = \frac{\pi^2}{16} - \frac{\alpha^2}{2}.$$

□

Theorem 3 (Third definite integral).

$$\int_{\alpha/2}^{\infty} \ln(\tanh z) dz = \frac{1}{4} \ln^2(1 + \sqrt{2}) - \frac{\pi^2}{16}.$$

PROOF. From Theorem 1, we know that the area of S evaluates to

$$2 \int_0^{\infty} \int_u^{f(u)} 1 dv du = \frac{\pi^2}{8},$$

where $f(u) = \sinh^{-1}(\cosh u)$. Let us rotate the coordinate axes by the angle $\pi/4$ rad, as in Apostol's proof. Since "pure" rotations about the origin always yield an unitary Jacobi determinant, one has:

$$\begin{aligned} \frac{\pi^2}{8} &= 2 \int_0^{\alpha/\sqrt{2}} \int_0^X 1 dY dX + 2 \int_{\alpha/\sqrt{2}}^{\infty} \int_0^{h(X)} 1 dY dX \\ &= 2 \int_0^{\alpha/\sqrt{2}} X dX + 2 \int_{\alpha/\sqrt{2}}^{\infty} h(X) dX \\ &= \frac{\alpha^2}{2} + 2 \int_{\alpha/\sqrt{2}}^{\infty} h(X) dX, \end{aligned} \tag{5}$$

where

$$u = \frac{X - Y}{\sqrt{2}}, \quad v = \frac{X + Y}{\sqrt{2}},$$

and $h(X)$ is determined by changing the variables u and v on the equation $v = f(u)$ to the new variables X and Y , as follows. Since $v = f(u) = \sinh^{-1}(\cosh u)$, then $\sinh v = \cosh u$, so

$$\sinh \left(\frac{X + Y}{\sqrt{2}} \right) = \cosh \left(\frac{X - Y}{\sqrt{2}} \right).$$

By making use of the addition and subtraction formulas for hyperbolic functions, as well as of the substitutions $X' := X/\sqrt{2}$ and $Y' := Y/\sqrt{2}$, the above equality expands to

$$\sinh X' \cosh Y' + \sinh Y' \cosh X' = \cosh X' \cosh Y' - \sinh X' \sinh Y'.$$

By dividing both sides by $\cosh X' \cosh Y'$, one finds

$$\tanh X' + \tanh Y' = 1 - \tanh X' \cdot \tanh Y',$$

which can be written as

$$\tanh Y' = \frac{1 - \tanh X'}{1 + \tanh X'}. \quad (6)$$

Now, for simplicity, let us define $t := \tanh X'$. After some algebra, Eq. (6) becomes

$$Y' = \frac{1}{2} \ln \left(\frac{1 + \frac{1-t}{1+t}}{1 - \frac{1-t}{1+t}} \right) = \frac{1}{2} \ln \left(\frac{1}{t} \right) = -\frac{1}{2} \ln t,$$

which means that

$$Y = h(X) = -\frac{\sqrt{2}}{2} \ln \left[\tanh \left(\frac{X}{\sqrt{2}} \right) \right].$$

By putting this function on the last integral of Eq. (5), one has

$$\frac{\pi^2}{8} = \frac{\alpha^2}{2} - \sqrt{2} \int_{\alpha/\sqrt{2}}^{\infty} \ln \left[\tanh \left(\frac{X}{\sqrt{2}} \right) \right] dX.$$

The simple substitution $z = X/\sqrt{2}$ promptly reduces this equation to

$$\int_{\alpha/2}^{\infty} \ln(\tanh z) dz = \frac{\alpha^2}{4} - \frac{\pi^2}{16}. \quad (7)$$

□

After this proof, we make the substitution $t = \tanh z$ in the integral in Eq. (7), above. When the result is expressed in terms of the dilogarithm function, an interesting identity arises.

Theorem 4 (Dilogarithm identity). *Let us define $\text{Li}_2(z) := \sum_{k=1}^{\infty} z^k/k^2$, which is valid for all complex z with $|z| \leq 1$ [8]. Then*

$$\text{Li}_2(\sqrt{2} - 1) + \text{Li}_2\left(1 - \frac{1}{\sqrt{2}}\right) = \frac{\pi^2}{8} - \frac{\ln^2(1 + \sqrt{2})}{2} - \frac{1}{8} \ln^2 2.$$

PROOF. By applying the substitution $t = \tanh z$ ($dt = \operatorname{sech}^2 z dz$) to the definite integral of Theorem 3, in the previous section, one finds

$$\frac{\alpha^2}{4} - \frac{\pi^2}{16} = \int_{\tanh(\alpha/2)}^1 \frac{\ln t}{1-t^2} dt = \int_{\sqrt{2}-1}^1 \frac{\ln t}{1-t^2} dt, \quad (8)$$

since

$$\tanh\left(\frac{\alpha}{2}\right) = \tanh\left(\ln \sqrt{1+\sqrt{2}}\right) = \frac{\sqrt{1+\sqrt{2}} - \frac{1}{\sqrt{1+\sqrt{2}}}}{\sqrt{1+\sqrt{2}} + \frac{1}{\sqrt{1+\sqrt{2}}}} = \frac{\sqrt{2}}{2+\sqrt{2}} = \sqrt{2}-1.$$

By expanding the integrand in Eq. (8) in partial fractions, one has

$$\frac{\alpha^2}{4} - \frac{\pi^2}{16} = \frac{1}{2} \int_{\sqrt{2}-1}^1 \frac{\ln t}{1+t} dt - \frac{1}{2} \int_1^{\sqrt{2}-1} \frac{\ln t}{1-t} dt. \quad (9)$$

These integrals can be reduced to special values of the dilogarithm function if one adopts the more general integral definition for this function [8], namely

$$\operatorname{Li}_2(z) := - \int_0^z \frac{\ln(1-s)}{s} ds,$$

valid for $z \in \mathbb{C} \setminus [1, \infty)$.⁴ Note that, by expanding the logarithm in powers of z in this integral definition, one finds $\operatorname{Li}_2(z) = \sum_{k=1}^{\infty} z^k/k^2$, which agrees with the basic definition stated in the theorem for all complex z with $|z| \leq 1$ [8]. By taking into account this summation form, it is clear that $\operatorname{Li}_2(1) = \pi^2/6$, which implies that

$$\operatorname{Li}_2(z) = \frac{\pi^2}{6} - \int_1^z \frac{\ln(1-s)}{s} ds.$$

From this integral form, it is easy to deduce that

$$\int_1^z \frac{\ln t}{1-t} dt = \operatorname{Li}_2(1-z)$$

and

$$\int_z^1 \frac{\ln t}{1+t} dt = -\operatorname{Li}_2(-z) - \ln z \ln(z+1) - \frac{\pi^2}{12}.$$

By putting $z = \sqrt{2}-1$ in these integrals and then substituting the resulting dilogarithms in Eq. (9), one finds

$$\begin{aligned} \frac{\alpha^2}{4} - \frac{\pi^2}{16} &= \frac{1}{2} \left[-\operatorname{Li}_2(1-\sqrt{2}) - \ln(\sqrt{2}-1) \ln \sqrt{2} - \frac{\pi^2}{12} \right] - \frac{1}{2} \operatorname{Li}_2(2-\sqrt{2}) \\ &= -\frac{\operatorname{Li}_2(1-\sqrt{2})}{2} - \frac{\operatorname{Li}_2(2-\sqrt{2})}{2} - \frac{1}{4} \ln(\sqrt{2}-1) \ln 2 - \frac{\pi^2}{24}. \end{aligned} \quad (10)$$

⁴We are considering here the principal branch of the dilogarithm function $\operatorname{Li}_2(z)$, defined by taking the principal branch of $\ln z$, for which it has a cut along the negative real axis, with $|\arg(z)| < \pi$. This defines the principal branch of $\operatorname{Li}_2(z)$ as a single-valued function in the complex plane cut along the real axis, from 1 to $+\infty$.

Now, by applying the Euler reflection formula for $\text{Li}_2(z)$ about $z = \frac{1}{2}$, which reads (see Eq. (1.11) in Ref. [8])

$$\text{Li}_2(1-z) = -\text{Li}_2(z) - \ln z \ln(1-z) + \frac{\pi^2}{6},$$

to $\text{Li}_2(2-\sqrt{2})$ in Eq. (10), one finds

$$\text{Li}_2(\sqrt{2}-1) - \text{Li}_2(1-\sqrt{2}) = \frac{\alpha^2}{2} - \frac{\pi^2}{8} + \frac{\pi^2}{4} - \ln^2(\sqrt{2}-1). \quad (11)$$

By applying the Landen's formula, namely (see Eq. (1.12) in Ref. [8])

$$\text{Li}_2(z) = -\text{Li}_2\left(\frac{z}{z-1}\right) - \frac{1}{2} \ln^2(1-z),$$

which is valid for all complex z , $z \notin [1, \infty)$, to $\text{Li}_2(1-\sqrt{2})$, one has

$$\text{Li}_2(1-\sqrt{2}) = -\text{Li}_2\left(1-\frac{1}{\sqrt{2}}\right) - \frac{1}{8} \ln^2 2.$$

Finally, by substituting this result on Eq. (11), one finds

$$\text{Li}_2(\sqrt{2}-1) + \text{Li}_2\left(1-\frac{1}{\sqrt{2}}\right) = \frac{\pi^2}{8} - \frac{\alpha^2}{2} - \frac{1}{8} \ln^2 2.$$

□

It should be mentioned that neither Mathematica (release 7) nor Maple (release 13) are able to generate any closed-form expression for this sum. Indeed, this identity is not found in standard bibliographic sources for dilogarithms [8, 9], though it resembles the identity

$$\mathcal{L}(\sqrt{2}-1) + \mathcal{L}\left(1-\frac{1}{\sqrt{2}}\right) = \frac{3}{4},$$

where $\mathcal{L}(x) := \frac{6}{\pi^2} [\text{Li}_2(x) + \frac{1}{2} \ln x \ln(1-x)]$ is the *normalized Roger's dilogarithm*, as found by Bytsko (1999) [10].

It seems plausible to extend the results obtained here in this paper to other known unit square integrals, such as $\int_0^1 \int_0^1 \ln(xy)/(1-x^2y^2) dx dy = -\frac{7}{4} \zeta(3)$, $\int_0^1 \int_0^1 \ln x/(1-x^2y^2) dx dy = -\frac{7}{8} \zeta(3)$, $\int_0^1 \int_0^1 \ln(xy)/(1-xy) dx dy = -2 \zeta(3)$, $\int_0^1 \int_0^1 \ln(1-xy)/(1-xy) dx dy = -\zeta(3)$, and many others (see, e.g., Ref. [11]). Similarly, it should also be interesting to include a third dimension in the hyperbolic change of variables in order to investigate some triple integrals over the unit cube, such as

$$\int_0^1 \int_0^1 \int_0^1 \frac{1}{1-xyz} dx dy dz = \zeta(3) \quad \text{and} \quad \int_0^1 \int_0^1 \int_0^1 \frac{1}{1-x^2y^2z^2} dx dy dz = \frac{7}{8} \zeta(3).$$

Extensive work in both directions is already ongoing by the author.

Acknowledgments

The author acknowledges a postdoctoral fellowship from CNPq (Brazilian agency) during the course of this work.

References

- [1] L. Euler, *De summis serierum reciprocarum*, Opera Omnia **14** (1740) 73–86.
- [2] D. Kalman, Six ways to sum a series, Coll. Math. J. **24** (1993) 402–421.
- [3] R. Chapman, Evaluating $\zeta(2)$. Available at:
<http://secamlocal.ex.ac.uk/people/staff/rjchapma/etc/zeta2.pdf>
- [4] T. M. Apostol, A proof that Euler missed: evaluating $\zeta(2)$ the easy way, Math. Intelligencer **5:3** (1983) 59–60.
- [5] W. Rudin, *Principles of Mathematical Analysis*, 3rd ed., McGraw-Hill, New York, 1976.
- [6] M. Aigner and G. M. Ziegler, *Proofs from THE BOOK*, 4th ed., Springer, New York, 2010. Chap. 8.
- [7] F. Beukers, J. A. C. Kolk, and E. Calabi, Sums of generalized harmonic series and volumes, Nieuw Arch. Wiskd. **11** (1993) 217–224.
- [8] L. Lewin, *Polylogarithms and Associated Functions*, Elsevier, New York, 1981.
- [9] D. Zagier, *The Dilogarithm Function*. In: *Frontiers in Number Theory, Physics, and Geometry II*, Ed. by P. Cartier, P. Moussa, B. Julia, and P. Vanhove. Springer, New York, 2007. Pages 3–61.
- [10] A. G. Bytsko, Two-term dilogarithm identities related to conformal field theory, Lett. Math. Phys. **50** (1999) 213–228.
- [11] E. W. Weisstein, Unit square integrals. From *MathWorld*, a Wolfram Web resource, available at:
<http://mathworld.wolfram.com/UnitSquareIntegral.html>

Figures




Figure 1: Mapping the unit square in the xy -plane (left-hand side) onto the hyperbolic ‘quadrilateral’ in the uv -plane (right-hand side). The symmetry about the diagonal $v = u$, represented by the inclined dashed line, is exploited in the text. Note that the hyperbolic region formed in the uv -plane is bounded by the coordinate axes and the curves $v = f(u)$ and $v = g(u)$. Both these functions, as well as the expressions for the change of variables indicated in the central part, are stated elsewhere in the text. According to Eq. (4), α evaluates to $\sinh^{-1}(1) = \ln(1 + \sqrt{2}) \approx 0.88137$.