

Universal Elliptic Functions*

Yoshihiro Ônishi

The aim of this note is to discuss the power series expansion at $u = 0$ of the sigma function $\sigma(u)$ of the most general elliptic curve, namely, of

$$\mathcal{E} : y^2 + (\mu_1 x + \mu_3)y = x^3 + \mu_2 x^2 + \mu_4 x + \mu_6,$$

where μ_j are constants. Through out this note, $\boldsymbol{\mu}$ denotes the set consists of the coefficients μ_j s. We show, in particular, that the series expansion at $u = 0$ of the *square* of $\sigma(u)$ is Hurwitz integral over $\mathbb{Z}[\boldsymbol{\mu}] = \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_6]$, and $\sigma(u)$ itself is Hurwitz integral over $\mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6]$. Namely, any coefficient of the power series expansion of $\sigma(u)$ at $u = 0$ is of the form $c_n u^n/n!$ with $c_n \in \mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6]$. Although the author had thought that Weierstrass recursion in [6] implies directly Huriwitz integrality of $\sigma(u)$ at least if $\mu_1 = \mu_2 = \mu_3 = 0$, Victor Buchstaber pointed out we do not know how to prove that the prime 3 does not appear in the denominator of c_n above. On the other hand, the author had already written this paper because he had been interested in why c_n belongs $\mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6]$ and does not belong $\mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_6]$ in general. The method of this paper is completely different from that of Weierstrass, which is only an improvement of Nakayashiki's paper [4] and is aimed to be generalized for higher genus cases.

The fact that the power series expansion of the sigma function is not Hurwitz integral only at the prime 2 would relate with the result of Mazur-Tate [3].

In the last section, we give explicitly n -plication formula for the coordinate function of the curve \mathcal{E} .

Acknowledgement. Elena Bunkova read the draft of this paper carefully and corrected several mistakes. Buchstaber made the author realize the main result of this paper seems to be new. I deeply thank them two.

*Date: 2010.3.14

1 The Fundamental Differential Form

1.1 The most general elliptic curve

Let us consider the most general elliptic curve

$$(1.1) \quad \mathcal{E} : y^2 + (\mu_1 x + \mu_3)y = x^3 + \mu_2 x^2 + \mu_4 x + \mu_6.$$

In the sequel, we use notations

$$(1.2) \quad \begin{aligned} f(x, y) &= y^2 + (\mu_1 x + \mu_3)y - (x^3 + \mu_2 x^2 + \mu_4 x + \mu_6), \\ f_x(x, y) &= \frac{\partial}{\partial x} f(x, y) = \mu_1 y - (3x^2 + 2\mu_2 x + \mu_4), \\ f_y(x, y) &= \frac{\partial}{\partial y} f(x, y) = 2y + (\mu_1 x + \mu_3). \end{aligned}$$

We choose local parameter

$$(1.3) \quad t = -x/y$$

at ∞ on \mathcal{E} . We never use $x^{-1/2}$ as a local parameter. We call this t *arithmetic local parameter* of \mathcal{E} . We usually express by $\langle t \rangle$ the value determined by each value t . For instance, the coordinate x of \mathcal{E} is denoted by $x\langle t \rangle$. If we introduce also

$$(1.4) \quad s = 1/x,$$

the equation $f(x, y) = 0$ is rewritten as

$$(1.5) \quad s = (1 + \mu_2 s + \mu_4 s^2 + \mu_6 s^3)t^2 + (\mu_1 s + \mu_3 s^2)t.$$

Using this recursively, we have

$$(1.6) \quad \begin{aligned} s &= t^2 + \mu_1 t^3 + (\mu_1^2 + \mu_2)t^4 + (\mu_1^3 + 2\mu_2\mu_1 + \mu_3)t^5 + \\ &\quad (\mu_1^4 + 3\mu_2\mu_1^2 + 3\mu_3\mu_1 + \mu_2^2 + \mu_4)t^6 + \cdots . \end{aligned}$$

By (1.6), we see $x\langle t \rangle$ and $y\langle t \rangle$ are expressed as a power series in $\mathbb{Z}[\boldsymbol{\mu}][[t]]$ as

$$(1.7) \quad \begin{aligned} x\langle t \rangle &= t^{-2} - \mu_1 t^{-1} - \mu_2 - \mu_3 t - (\mu_3\mu_1 + \mu_4)t^2 - (\mu_3\mu_1^2 + \mu_4\mu_1 + \mu_2\mu_3)t^3 + \cdots , \\ y\langle t \rangle &= -t^{-3} + \mu_1 t^{-2} + \mu_2 t^{-1} + \mu_3 + (\mu_3\mu_1 + \mu_4)t + (\mu_3\mu_1^2 + \mu_4\mu_1 + \mu_2\mu_3)t^2 + \cdots . \end{aligned}$$

We choose

$$(1.8) \quad \omega_1(x, y) = \frac{dx}{f_y(x, y)} = \frac{dx}{2y + \mu_1 x + \mu_3}$$

as the base of the holomorphic 1-form (differentials of the 1st kind) on \mathcal{E} . Since

$$(1.9) \quad \begin{aligned} \omega_1(x, y) &= \frac{dx}{2y + \mu_1 x + \mu_3} = \frac{\frac{dx}{dt} dt}{2y + \mu_1 x + \mu_3} \in (1 + t \mathbb{Z}[\frac{1}{2}, \boldsymbol{\mu}][[t]])dt, \\ \omega_1(x, y) &= -\frac{dy}{f_x(x, y)} \in (1 + t \mathbb{Z}[\frac{1}{3}, \boldsymbol{\mu}][[t]])dt, \end{aligned}$$

we have

$$(1.10) \quad \begin{aligned} \omega_1(x, y) &= (1 + \mu_1 t + (\mu_2 + \mu_1^2)t^2 + (2\mu_1\mu_2 + 2\mu_3 + \mu_1^3)t^3 + \cdots)dt \\ &\in (1 + t \mathbb{Z}[\boldsymbol{\mu}][[t]])dt. \end{aligned}$$

1.2 The fundamental 2-form

We let consider

$$(1.11) \quad \Omega(x, y, z, w) = -\frac{y + w + \mu_1 z + \mu_3}{z - x} \omega_1(x, y) = -\frac{(y + w + \mu_1 z + \mu_3) dx}{(z - x)(2y + \mu_1 x + \mu_3)}.$$

This is a 1-form with respect to (x, y) that has simple pole at (z, w) and no other poles. Note that this is holomorphic at $(z, w + \mu_1 z + \mu_3)$. Indeed, the numerator becomes $(2w + \mu_1 z + \mu_3) = f_y(z, w)$ when $(x, y) = (z, w)$. If the value of the local parameter t gives the value x , we denote by t' the other value of the arithmetic local parameter which gives the same x coordinate $x\langle t \rangle$. Hence, $x\langle t \rangle = x\langle t' \rangle$. Then $y\langle t \rangle + y\langle t' \rangle = -(\mu_1 x\langle t \rangle + \mu_3)$. So that

$$(1.12) \quad \begin{aligned} t' &= -\frac{x\langle t' \rangle}{y\langle t' \rangle} = \frac{x\langle t \rangle}{y\langle t \rangle + \mu_1 x\langle t \rangle + \mu_3} \\ &= -t - \mu_1 t^2 - \mu_1^2 t^3 + (-\mu_1^3 - \mu_3) t^4 + (-\mu_1^4 - 3\mu_3 \mu_1) t^5 + \cdots \in t \mathbb{Z}[\mu_1, \mu_3][[t]]. \end{aligned}$$

The first equality of the above implies

$$(1.13) \quad \begin{aligned} tt' &= \frac{x\langle t \rangle^2}{(y\langle t \rangle + \mu_1 x\langle t \rangle + \mu_3)y\langle t \rangle} = \frac{x\langle t \rangle^2}{x\langle t \rangle^3 + \mu_2 x\langle t \rangle^2 + \mu_4 x\langle t \rangle + \mu_6} \\ &= \frac{1}{x\langle t \rangle(1 + \mu_2 \frac{1}{x\langle t \rangle} + \mu_4 \frac{1}{x\langle t \rangle^2} + \mu_6 \frac{1}{x\langle t \rangle^3})} = \frac{1}{x\langle t \rangle} (1 - \mu_2 \frac{1}{x\langle t \rangle} + \cdots). \end{aligned}$$

Finally, we see

$$(1.14) \quad \frac{1}{x\langle t \rangle} = tt' + \mu_2 (tt')^2 + \cdots \in tt' \mathbb{Z}[\boldsymbol{\mu}][[(tt')]].$$

Let us denote as $x_1 = x\langle t_1 \rangle$, $y_1 = y\langle t_1 \rangle$. Using Weierstrass preparation theorem, we define $p(t_1, t_2) \in (\mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]])^\times$ by

$$(1.15) \quad x_2^{-1} - x_1^{-1} = -(t_2 - t_1)(t_2' - t_1) p(t_1, t_2).$$

Then explicit calculation gives

$$(1.16) \quad \begin{aligned} p(t_1, t_2) &= 1 + \mu_1 t_1 + \mu_2 t_2^2 + (\mu_2 + \mu_1^2) t_1^2 + \mu_1 \mu_2 t_2^3 + \cdots \\ &\in x_1^{-1} / t_1^2 + t_2 \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]. \end{aligned}$$

The last one is shown by letting $t_2 = 0$. On the other hand, we have

$$(1.17) \quad \begin{aligned} y_1 + y_2 + \mu_1 x_2 + \mu_3 &= -\frac{x\langle t_1 \rangle}{t_1} + \frac{x\langle t_2' \rangle}{t_2'} \\ &= -\frac{x\langle t_1 \rangle}{t_1} + \frac{x\langle t_2 \rangle}{t_1} - \frac{x\langle t_2 \rangle}{t_1} + \frac{x\langle t_2' \rangle}{t_2'} \\ &= -\frac{1}{t_1} (x\langle t_1 \rangle - x\langle t_2 \rangle) - x\langle t_2 \rangle \left(\frac{1}{t_1} - \frac{1}{t_2'} \right) \end{aligned}$$

and

$$\begin{aligned}
& x_2 \left(\frac{1}{t_1} - \frac{1}{t_2'} \right) \frac{1}{x_2 - x_1} = \frac{-x_1^{-1}}{x_2^{-1} - x_1^{-1}} \left(\frac{1}{t_1} - \frac{1}{t_2'} \right) \\
&= \frac{-x_1^{-1}}{(t_2 - t_1)(t_2' - t_1) p(t_1, t_2)} \left(\frac{1}{t_1} - \frac{1}{t_2'} \right) \\
(1.18) \quad &= \frac{-x_1^{-1}}{(t_2 - t_1)(t_2' - t_1) (x_1^{-1}/t_1^2 + \text{"a series in } t_2 \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"})} \frac{t_2' - t_1}{t_1 t_2'} \\
&= \frac{-x_1^{-1}}{(t_2 - t_1) t_1 t_2' (x_1^{-1}/t_1^2 + \text{"a series in } t_2 \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"})} \\
&= \frac{t_1}{(t_2 - t_1) t_2} \cdot \frac{t_2}{t_2'} \cdot \frac{-x_1^{-1}/t_1^2}{(x_1^{-1}/t_1^2 + \text{"a series in } t_2 \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"})}.
\end{aligned}$$

Here we note that

$$(1.19) \quad t_2'/t_2 \in -1 + t_2 \mathbb{Z}[\boldsymbol{\mu}_1, \boldsymbol{\mu}_3][[t_2]].$$

At the last part of (1.18), since $x_1^{-1}/t_1^2 \in 1 + t_1 \mathbb{Z}[\boldsymbol{\mu}][[t_1]]$, we have

$$\begin{aligned}
(1.20) \quad & x_2 \left(\frac{1}{t_1} - \frac{1}{t_2'} \right) \frac{1}{x_2 - x_1} = - \left(\frac{1}{t_2 - t_1} - \frac{1}{t_2} \right) (\text{"a series in } 1 + t_2 \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"}) \\
&= + \frac{1}{t_2} - \frac{1}{t_2 - t_1} (\text{"a series in } 1 + t_2 \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"}) \\
&\quad + (\text{"a series in } \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"}).
\end{aligned}$$

Therefore,

$$\begin{aligned}
(1.21) \quad & \frac{y_1 + y_2 + \mu_1 x_2 + \mu_3}{x_2 - x_1} = \frac{1}{t_1} - \frac{x_2}{x_2 - x_1} \left(\frac{1}{t_1} - \frac{1}{t_2'} \right) \\
&= \frac{1}{t_1} - \frac{1}{t_2} + (\text{"a series in } \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"}) \\
&\quad + \frac{1}{t_2 - t_1} (\text{"a series in } 1 + t_2 \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]]\text{"}).
\end{aligned}$$

Defining $b\langle t_1, t_2 \rangle$ by

$$\begin{aligned}
(1.22) \quad & \left(\frac{y_1 + y_2 + \mu_1 x_2 + \mu_3}{x_2 - x_1} - \frac{1}{t_1} + \frac{1}{t_2} \right) \omega_1 \langle t_1 \rangle \\
&= \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]] \omega_1 \langle t_1 \rangle + \frac{1}{t_2 - t_1} b\langle t_1, t_2 \rangle dt_1 \\
& b\langle t_1, t_2 \rangle \in \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]],
\end{aligned}$$

since

$$\begin{aligned}
(1.23) \quad & \lim_{t_2 \rightarrow t_1} \frac{x_2 - x_1}{y_1 + y_2 + \mu_1 x_2 + \mu_3} \cdot \frac{1}{t_2 - t_1} = \frac{\frac{dx}{dt} \langle t_1 \rangle}{2y_1 + \mu_1 x_2 + \mu_3} \\
&= \omega_1 \langle t_1 \rangle / dt_1,
\end{aligned}$$

it must be

$$(1.24) \quad b\langle t_1, t_1 \rangle = 1.$$

Thus

$$(1.25) \quad b\langle t_1, t_2 \rangle \in 1 + (t_2 - t_1) \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]].$$

We summarize this fact as a theorem:

Theorem 1.26. *We have*

$$(1.27) \quad \left(\frac{y_1 + y_2 + \mu_1 x_2 + \mu_3}{x_2 - x_1} - \frac{1}{t_1} + \frac{1}{t_2} \right) \omega_1 \langle t_1 \rangle + \frac{dt_1}{t_1 - t_2} \in \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]] dt_1.$$

By using a computer, we have first several terms:

$$(1.28) \quad \begin{aligned} & \left(\frac{y + w + \mu_1 z + \mu_3}{z - x} - \frac{1}{t_1} + \frac{1}{t_2} \right) \omega_1(x, y) + \frac{dt_1}{t_1 - t_2} \\ &= \left(-\mu_2 t_1 - \mu_3 t_2 t_1 \right. \\ &\quad - (\mu_2 \mu_1 + 2\mu_3) t_1^2 \\ &\quad - (2\mu_3 \mu_1 + \mu_4) t_2 t_1^2 \\ &\quad - (\mu_3 \mu_1 + \mu_4) t_2^2 t_1 \\ &\quad - (\mu_2 \mu_1^2 + 4\mu_3 \mu_1 + \mu_2^2 + 2\mu_4) t_1^3 \\ &\quad - (\mu_3 \mu_1^2 + \mu_4 \mu_1 + \mu_2 \mu_3) t_2^3 t_1 \\ &\quad - (2\mu_3 \mu_1^2 + 2\mu_4 \mu_1 + \mu_2 \mu_3) t_2^2 t_1^2 \\ &\quad - (3\mu_3 \mu_1^2 + 2\mu_4 \mu_1 + 2\mu_2 \mu_3) t_2 t_1^3 \\ &\quad \left. - (\mu_2 \mu_1^3 + 6\mu_3 \mu_1^2 + 2\mu_2^2 \mu_1 + 4\mu_4 \mu_1 + 6\mu_2 \mu_3) t_1^4 - \dots \right) dt_1. \end{aligned}$$

By using Ω in (1.11), we consider a 2-form

$$(1.29) \quad \boldsymbol{\xi}(x, y; z, w) = \frac{d}{dx} \Omega(z, w; x, y) dx - \omega_1(z, w) \eta_1(x, y)$$

such that

$$(1.30) \quad \boldsymbol{\xi}(x, y; z, w) = \boldsymbol{\xi}(z, w; x, y)$$

with a differential of the 3rd kind η_1 that has a pole only at ∞ . Such an η_1 is determined modulo constant multiple of ω_1 . A solution is given by

$$(1.31) \quad \eta_1(x, y) = \frac{-x dx}{2y + \mu_1 x + \mu_3}.$$

The power series expansion with respect to t is given by

$$(1.32) \quad \begin{aligned} \eta_1(x, y) &= -t^{-2} - \mu_3 t - (\mu_4 + 2\mu_1 \mu_3) t^2 - (2\mu_1 \mu_4 + 2\mu_3 \mu_2 + 3\mu_1^2 \mu_3) t^3 - \dots \\ &\in -t^{-2} + t \mathbb{Z}[\boldsymbol{\mu}][[t]]. \end{aligned}$$

Under the situation, $\boldsymbol{\xi}$ in (1.29) is written as

$$(1.33) \quad \boldsymbol{\xi} = \frac{F(x, y, z, w) dx dz}{(x - z)^2 f_y(x, y) f_y(z, w)},$$

where

$$(1.34) \quad \begin{aligned} F(x, y; z, w) &= xz(x + z) + (\mu_1^2 + 2\mu_2)xz + \mu_1(z y + x w) \\ &\quad + (\mu_3 \mu_1 + \mu_4)(x + z) + 2y w + \mu_3(y + w) + \mu_3^2 + 2\mu_6. \end{aligned}$$

1.3 Legendre relation

Let us choose two closed paths α and β with their intersection being $\alpha \cdot \beta = -\beta \cdot \alpha = 1$ which generate the fundamental group of \mathcal{E} and let us define

$$(1.35) \quad \omega' = \int_{\alpha} \omega_1(x, y), \quad \omega'' = \int_{\beta} \omega_1(x, y), \quad \eta' = \int_{\alpha} \eta_1(x, y), \quad \eta'' = \int_{\beta} \eta_1(x, y).$$

Then we have Legendre relation

$$(1.36) \quad \omega''\eta' - \omega'\eta'' = 2\pi i,$$

where i is the imaginary unit.

2 The sigma function

2.1 Construction of the sigma function

Now, we let

$$(2.1) \quad R = \gcd \left(\text{rslt}_x(\text{rslt}_y(f, f_y), \text{rslt}_y(f, f_y)), \text{rslt}_y(\text{rslt}_x(f, f_y), \text{rslt}_x(f, f_y)) \right).$$

Then R is a squared element in $\mathbb{Z}[\mu]$. So we take a square root of it: $D = R^{1/2}$. More explicitly, if we define

$$(2.2) \quad \begin{aligned} b_2 &= \mu_1^2 + 4\mu_2, & b_4 &= 2\mu_4 + \mu_1\mu_3, & b_6 &= \mu_3^2 + 4\mu_6, \\ b_8 &= \mu_1^2\mu_6 + 4\mu_2\mu_6 - \mu_1\mu_3\mu_4 + \mu_2\mu_3^2 - \mu_4^2, \end{aligned}$$

then we let

$$(2.3) \quad D = -b_2^2b_8 - 8b_4^3 - 27b_6^2 + 9b_2b_4b_6.$$

Now let us define Weierstrass' *sigma function* by

$$(2.4) \quad \sigma(u) = D^{-1/8} \left(\frac{\pi}{\omega'} \right)^{1/2} \exp \left(-\frac{1}{2}u^2\eta'\omega'^{-1} \right) \vartheta \left[\begin{matrix} \frac{1}{2} \\ \frac{1}{2} \end{matrix} \right] (\omega'^{-1}u | \omega'^{-1}\omega'').$$

This function is characterized up to multiplicative constant as a $\text{SL}_2(\mathbb{Z})$ -invariant Jacobi form under the usual action.

For each $u \in \mathbb{C}$, there is unique pair of u' and $u'' \in \mathbb{R}$ determined by $u = u'\omega' + u''\omega''$. We use this notation convention also for each lattice point $\ell \in \Lambda$, and we write $\ell = \ell'\omega' + \ell''\omega''$. Moreover, for $u, v \in \mathbb{C}$, and $\ell \in \Lambda$, we let

$$(2.5) \quad L(u, v) = u(v'\eta' + v''\eta''), \quad \chi(\ell) = \exp \left(2\pi i \left(\frac{1}{2}\ell' - \frac{1}{2}\ell'' + \frac{1}{2}\ell'\ell'' \right) \right) (\in \{1, -1\}).$$

Lemma 2.6. *The function $\sigma(u)$ is an entire function and not depends of the choice of α and β . Therefore, it will be expanded at the origin in terms of μ_j s. Moreover, $\sigma(u)$ has poles of order 1 at each point of Λ and no pole elsewhere, and satisfies*

$$(2.7) \quad \sigma(u + \ell) = \chi(\ell)\sigma(u) \exp L(u + \frac{1}{2}\ell, \ell).$$

Proof. The claim that it does not depend of α and β is not so easy. Here, we refer [5], Chapter 9 and 10. The proof is using the transformation property of Dedekind η -function. See [1], pp.552–557, and [2], p.85 and pp.176–183. The zeroes of $\sigma(u)$ is given by calculating the integral around the boundary of the regular polygon associated to α and β after taking logarithm of (2.7). The equation (2.7) itself is shown by the translational relation for the theta series and Legendre relation (1.35). \square

In the proof of the first claim above, we know that there exists 8th root ε of 1 such that $\sigma(u) = \varepsilon u + O(u^2)$. We fix the 8th root in (2.4) as

$$(2.8) \quad \sigma(u) = u + O(u^2).$$

For an integral domain A with characteristic 0, and an indeterminate t , we denote by $A\langle\langle t \rangle\rangle$ the ring of the elements

$$(2.9) \quad \sum_{j=0}^{\infty} C_j \frac{t^j}{j!} \quad \text{with } C_j \in A.$$

We call such a series *Hurwitz integral* over A . We will finally have the power series expansion of $\sigma(u)$ at the origin as follows:

Theorem 2.10. *The power series expansion of the function $\sigma(u)^2$ around the origin belongs to $\mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_6]\langle\langle u \rangle\rangle$, and that of $\sigma(u)$ belongs to $\mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6]\langle\langle u \rangle\rangle$. Its first several terms are given by*

$$(2.11) \quad \begin{aligned} \sigma(u) = u &+ \left(\left(\frac{\mu_1}{2}\right)^2 + \mu_2\right) \frac{u^3}{3!} + \left(\left(\frac{\mu_1}{2}\right)^4 + 2\mu_2\left(\frac{\mu_1}{2}\right)^2 + \mu_3\mu_1 + \mu_2^2 + 2\mu_4\right) \frac{u^5}{5!} \\ &+ \left(\left(\frac{\mu_1}{2}\right)^6 + 3\mu_2\left(\frac{\mu_1}{2}\right)^4 + 6\mu_3\left(\frac{\mu_1}{2}\right)^3 + 3\mu_2^2\left(\frac{\mu_1}{2}\right)^2 + 6\mu_4\left(\frac{\mu_1}{2}\right)^2\right. \\ &\quad \left.+ 6\mu_3\mu_2\frac{\mu_1}{2} + \mu_2^3 + 6\mu_4\mu_2 + 6\mu_3^2 + 24\mu_6\right) \frac{u^7}{7!} + \dots \end{aligned}$$

The following is key relation through out this note:

Lemma 2.12. *The sigma function relates with the 2-form ξ by*

$$(2.13) \quad \frac{\sigma\left(\int_{\infty}^{(z_1, w_1)} \omega_1 - \int_{\infty}^{(x_1, y_1)} \omega_1\right) \sigma\left(\int_{\infty}^{(z, w)} \omega_1 - \int_{\infty}^{(x, y)} \omega_1\right)}{\sigma\left(\int_{\infty}^{(z, w)} \omega_1 - \int_{\infty}^{(x_1, y_1)} \omega_1\right) \sigma\left(\int_{\infty}^{(z_1, w_1)} \omega_1 - \int_{\infty}^{(x, y)} \omega_1\right)} = \exp\left(\int_{(z_1, w_1)}^{(z, w)} \int_{(x_1, y_1)}^{(x, y)} \xi\right),$$

where the integrals in the right hand side are given by jointing those in the left hand side.

Proof. Since we have the same factors in both sides when the point (x, y) or (z, w) varies through α or β , both sides must coincide up to multiplicative constant. If $(x, y) = (x_1, y_1)$, both sides are 1. Hence the multiplicative constant is 1 as a function of (z, w) . However, if $(z, w) = (z_1, w_1)$, both sides are 1 again. Therefore, the multiplicative constant is 1. \square

2.2 Solution to Jacobi's inversion problem

If we define Weierstrass \wp -function by

$$(2.14) \quad \wp(u) = -\frac{d^2}{du^2} \log \sigma(u),$$

expanding around the point (z, w)

$$(2.15) \quad \wp \left(\int_{\infty}^{(x,y)} \omega_1 - \int_{\infty}^{(z,w)} \omega_1 \right) = \frac{F(x, y, z, w)}{(x - z)^2},$$

which is obtained by taking 2nd derivative of the logarithm of (2.13), into power series of a local parameter corresponding the variable (z, w) around the point at infinity, we see that, if

$$(2.16) \quad u = \int_{\infty}^{(x,y)} \omega_1(x, y),$$

then

$$(2.17) \quad \wp(u) = x, \quad \wp'(u) = 2y + \mu_1 x + \mu_3.$$

First several terms of expansion of $\wp(u)$ is given by

$$(2.18) \quad \begin{aligned} \wp(u) = & \frac{1}{u^2} - \frac{1}{12}\mu_1^2 - \frac{1}{3}\mu_2 \\ & + \left(\frac{1}{240}\mu_1^4 + \frac{1}{30}\mu_2\mu_1^2 - \frac{1}{10}\mu_3\mu_1 + \frac{1}{15}\mu_2^2 - \frac{1}{5}\mu_4 \right) u^2 \\ & + \left(-\frac{1}{6048}\mu_1^6 - \frac{1}{504}\mu_2\mu_1^4 + \frac{1}{168}\mu_3\mu_1^3 + \left(-\frac{1}{126}\mu_2^2 + \frac{1}{84}\mu_4 \right) \mu_1^2 \right. \\ & \quad \left. + \frac{1}{42}\mu_2\mu_3\mu_1 - \frac{1}{28}\mu_3^2 - \frac{2}{189}\mu_2^3 + \frac{1}{21}\mu_4\mu_2 - \frac{1}{7}\mu_6 \right) u^4 \\ & + \left(\frac{1}{172800}\mu_1^8 + \frac{1}{10800}\mu_2\mu_1^6 - \frac{1}{3600}\mu_3\mu_1^5 + \left(\frac{1}{1800}\mu_2^2 - \frac{1}{1800}\mu_4 \right) \mu_1^4 \right. \\ & \quad \left. - \frac{1}{450}\mu_2\mu_3\mu_1^3 + \left(\frac{1}{300}\mu_3^2 + \frac{1}{675}\mu_2^3 - \frac{1}{225}\mu_4\mu_2 \right) \mu_1^2 - \frac{1}{225}\mu_2^2 + \frac{1}{75}\mu_4 \right) \mu_3\mu_1 \\ & \quad \left. + \left(\frac{1}{675}\mu_2^4 - \frac{2}{225}\mu_4\mu_2^2 + \frac{1}{75}\mu_4^2 \right) u^6 + O(u^8). \end{aligned}$$

2.3 Frobenius-Stickelberger formula

Lemma 2.19. (Frobenius-Stickelberger) *The following equality holds:*

$$(2.20) \quad \frac{\sigma(u+v)\sigma(u-v)}{\sigma(u)^2\sigma(v)^2} = x(u) - x(v).$$

Proof. By (2.7), the left hand side is a periodic function of both of u and v with respect Λ . On the other hand, since $\sigma(u)$ has poles of order 1 at each point in Λ , the divisors of both hand sides coincide. After expanding both sides into power series of u , they are of the form $1/u^2 + \dots$. Hence, the equality holds. \square

3 Hurwitz Integrality

Let A be an integral domain with characteristic 0. We denote by $A\langle\langle u \rangle\rangle$ is the ring consists of all *Hurwitz integral* series with respect to coefficient ring A .

Since

$$(3.1) \quad f(x\langle t \rangle, y) = (y - y\langle t \rangle)(y - y\langle t' \rangle).$$

for any t , we have $f_y(x\langle t \rangle, y) = (y - y\langle t \rangle) + (y - y\langle t' \rangle)$, so that

$$(3.2) \quad f_y(x\langle t \rangle, y\langle t \rangle) = y\langle t \rangle - y\langle t' \rangle.$$

This yields the following:

Lemma 3.3. *We have*

$$(3.4) \quad f_y(x\langle t \rangle, y\langle t \rangle) = \frac{1}{(tt')^3}(t-t')(t^2 + \text{“higher terms in } \mathbb{Z}[\boldsymbol{\mu}][[t]] \text{”}).$$

In this section, we prove Theorem 2.10. When

$$(3.5) \quad u = \int_{\infty}^{(x,y)} \omega_1,$$

by (1.10), we see

$$(3.6) \quad u = t + O(t^2) \quad \text{in } \mathbb{Z}[\boldsymbol{\mu}]\langle\langle t \rangle\rangle, \quad t = u + O(u^2) \quad \text{in } \mathbb{Z}[\boldsymbol{\mu}]\langle\langle u \rangle\rangle.$$

From (1.27), (1.29), (1.32), and (1.10), we have immediately, that

$$(3.7) \quad \boldsymbol{\xi}\langle t_1, t_2 \rangle - \frac{dt_1 dt_2}{(t_1 - t_2)^2} \in \mathbb{Z}[\boldsymbol{\mu}][t_1, t_2] dt_1 dt_2.$$

Explicitly, first several terms of this expansion are given by

$$(3.8) \quad \begin{aligned} \boldsymbol{\xi}\langle t_1, t_2 \rangle = & ((t_1 - t_2)^{-2} + \mu_3(t_1 + t_2) + (3\mu_3\mu_1 + 2\mu_4)t_1 t_2 \\ & + (2\mu_3\mu_1 + \mu_4)(t_1^2 + t_2^2) \\ & + (5\mu_3\mu_1^2 + 4\mu_4\mu_1 + 3\mu_2\mu_3)(t_1^2 t_2 + t_1 t_2^2) \\ & + (3\mu_3\mu_1^2 + 2\mu_4\mu_1 + 2\mu_2\mu_3)(t_1^3 + t_2^3) \\ & + (8\mu_3\mu_1^3 + 7\mu_4\mu_1^2 + 11\mu_2\mu_3\mu_1 + 3\mu_3^2 + 4\mu_4\mu_2 + 3\mu_6)t_1^2 t_2^2 \\ & + (7\mu_3\mu_1^3 + 6\mu_4\mu_1^2 + 10\mu_2\mu_3\mu_1 + 4\mu_3^2 + 4\mu_4\mu_2 + 4\mu_6)(t_1^3 t_2 + t_1 t_2^3) \\ & + (4\mu_3\mu_1^3 + 3\mu_4\mu_1^2 + 6\mu_2\mu_3\mu_1 + 3\mu_3^2 + 2\mu_4\mu_2 + 2\mu_6)(t_1^4 + t_2^4) \\ & + \dots) dt_1 dt_2, \end{aligned}$$

and

$$(3.9) \quad \begin{aligned} & \int_{t_1'}^{t_2'} \int_{t_1}^{t_2} \boldsymbol{\xi}\langle T_1, T_2 \rangle dT_1 dT_2 \\ & = -\log \left(-\frac{(t_2' - t_1)(t_2 - t_1')}{(t_2' - t_2)(t_1 - t_1')} \right) + \frac{\mu_3}{2} ((t_2'^2 - t_1'^2)(t_2 - t_1) + (t_2^2 - t_1^2)(t_2' - t_1')) \\ & \quad + \text{“a series divisible by } (t_1 - t_2)^2(t_1, t_2)^2 \text{ in } \mathbb{Z}[\boldsymbol{\mu}]\langle\langle t_1, t_2 \rangle\rangle \text{”} \end{aligned}$$

By plugging

$$(3.10) \quad \begin{aligned} (x, y) &= (x\langle t_2' \rangle, y\langle t_2' \rangle) = (x(-v), y(-v)) = (x(v), -y(v) - \mu_1 x(v) - \mu_3), \\ (z, w) &= (x\langle t_2 \rangle, y\langle t_2 \rangle) = (x(v), y(v)), \\ (x_1, y_1) &= (x\langle t_1' \rangle, y\langle t_1' \rangle) = (x(-u), y(-u)) = (x(u), -y(u) - \mu_1 x(u) - \mu_3), \\ (z_1, w_1) &= (x\langle t_1 \rangle, y\langle t_1 \rangle) = (x(u), y(u)), \end{aligned}$$

in (2.13), we have

$$(3.11) \quad \frac{\sigma(2u)\sigma(2v)}{\sigma(u+v)^2} = \exp \left(\int_{t_1}^{t_2} \int_{t_1'}^{t_2'} \boldsymbol{\xi}\langle T_1, T_2 \rangle dT_1 dT_2 \right)$$

Let u and v be analytic coordinates corresponding t_1 and t_2 , respectively, we have

$$\begin{aligned}
(x(u) - x(v))^2 &= \left(\frac{\sigma(u+v)\sigma(u-v)}{\sigma(u)^2\sigma(v)^2} \right)^2 \quad (\because (2.20)) \\
&= \frac{\sigma(u+v)^2}{\sigma(2u)\sigma(2v)} \frac{\sigma(2u)}{\sigma(u)^4} \frac{\sigma(2v)}{\sigma(v)^4} \sigma(u-v)^2 \\
&= \exp \left(- \int_{t_1'}^{t_2'} \int_{t_1}^{t_2} \boldsymbol{\xi} \langle T_1, T_2 \rangle dT_1 dT_2 \right) f_2 \langle t_1 \rangle f_2 \langle t_2 \rangle \sigma(u-v)^2 \\
(3.12) \qquad \qquad \qquad & \qquad \qquad \qquad (\because (2.13)) \\
&= \exp \left(- \log \frac{(t_2' - t_2)(t_1 - t_1')}{(t_2' - t_1)(t_2 - t_1')} \right. \\
&\quad \left. + \text{“a series in } \mathbb{Z}[\boldsymbol{\mu}] \langle \langle t_1, t_2 \rangle \rangle \text{”} \right) f_y \langle t_1 \rangle f_y \langle t_2 \rangle \sigma(u-v)^2 \\
&= \left(\frac{(t_2' - t_1)(t_2 - t_1')}{(t_2' - t_2)(t_1 - t_1')} \times \text{“a series of the form } 1 + \dots \text{ in } \mathbb{Z}[\boldsymbol{\mu}] \langle \langle t_1, t_2 \rangle \rangle \text{”} \right) \\
&\quad \times f_y \langle t_1 \rangle f_y \langle t_2 \rangle \sigma(u-v)^2.
\end{aligned}$$

We here recall (1.15):

$$\begin{aligned}
(3.13) \quad x \langle t_2 \rangle - x \langle t_1 \rangle &= \frac{(t_2' - t_1)(t_2 - t_1) p(t_1, t_2)}{x \langle t_1 \rangle^{-1} x \langle t_2 \rangle^{-1}}, \\
p(t_1, t_2) &= 1 + \mu_1 t_1 + \mu_2 t_2^2 + (\mu_2 + \mu_1^2) t_1^2 + \mu_1 \mu_2 t_2^3 + \dots
\end{aligned}$$

Exchanging t_1 and t_2 , we see

$$(3.14) \quad x \langle t_2 \rangle - x \langle t_1 \rangle = - \frac{x \langle t_2 \rangle^{-1} - x \langle t_1 \rangle^{-1}}{x \langle t_1 \rangle^{-1} x \langle t_2 \rangle^{-1}} = - \frac{(t_1' - t_2)(t_1 - t_2) p(t_2, t_1)}{x \langle t_1 \rangle^{-1} x \langle t_2 \rangle^{-1}}.$$

Dividing both sides of (2.20) by $u - v$ and (2.8) imply

$$(3.15) \quad \frac{\sigma(2u)}{\sigma(u)^4} = \frac{d}{du} x(u) = 1 / \frac{du}{dx} = f_2(x(u), y(u)).$$

Therefore,

$$\begin{aligned}
(3.16) \quad \frac{\sigma(2u)}{\sigma(u)^4} &= f_2 \langle t \rangle = f_2(x \langle t \rangle, y \langle t \rangle) = y \langle t \rangle - y \langle t' \rangle = - \frac{y \langle t' \rangle^{-1} - y \langle t \rangle^{-1}}{y \langle t \rangle^{-1} y \langle t' \rangle^{-1}} \\
&= - \frac{x \langle t \rangle}{t} + \frac{x \langle t \rangle}{t'} = (t - t') \frac{x \langle t \rangle}{tt'}.
\end{aligned}$$

Now we have arrived at the main result as follows:

Theorem 3.17. *Let u and v be analytic coordinates corresponding two values t_1 and t_2 of the arithmetic parameter in (1.3). Then the function $\sigma(u - v)^2$ is written as a product of formal power series:*

$$(3.18) \quad \sigma(u - v)^2 = (t_2 - t_1)^2 q(t_1)q(t_2) p(t_1, t_2) p(t_2, t_1) r(t_1, t_2),$$

where

$$(3.19) \quad \begin{aligned} p(t_1, t_2) &= \frac{x\langle t_2 \rangle^{-1} - x\langle t_1 \rangle^{-1}}{(t_2' - t_1)(t_2 - t_1)} = 1 + \mu_1 t_1 + \mu_2 t_2^2 + (\mu_2 + \mu_1^2) t_1^2 + \cdots \\ &\in 1 + (t_1, t_2) \mathbb{Z}[\boldsymbol{\mu}][[t_1, t_2]], \\ q(t) &= -x\langle t \rangle t t' = 1 - \mu_2 t^2 - \mu_2 \mu_1 t^3 - (\mu_2 \mu_1^2 + \mu_4) t^4 \\ &\quad - (\mu_2 \mu_1^3 + 2\mu_4 \mu_1 + \mu_2 \mu_3) t^5 + \cdots \in 1 + t \mathbb{Z}[\boldsymbol{\mu}][[t]], \\ r(t_1, t_2) &= \exp \left[\int_{t_1'}^{t_2'} \int_{t_1}^{t_2} \left(\boldsymbol{\xi}\langle t_1, t_2 \rangle - \frac{1}{(t_2 - t_1)^2} \right) \right] \\ &= 1 - \left(\frac{1}{12} \mu_1 \mu_3 + \frac{1}{6} \mu_4 \right) (t_1 - t_2)^4 - \left(\frac{1}{6} \mu_1^2 \mu_3 + \frac{1}{3} \mu_4 \mu_1 \right) (t_1 - t_2)^4 (t_1 + t_2) \\ &\quad + \left(-\left(\frac{1}{30} \mu_3^2 + \left(\frac{43}{180} \mu_1^3 + \frac{11}{90} \mu_2 \mu_1 \right) \mu_3 + \frac{43}{90} \mu_4 \mu_1^2 + \frac{11}{45} \mu_2 \mu_4 + \frac{2}{15} \mu_6 \right) (t_1^4 + t_2^4) \right. \\ &\quad + \left(\frac{2}{15} \mu_3^2 + \left(\frac{11}{90} \mu_1^3 + \frac{7}{45} \mu_2 \mu_1 \right) \mu_3 + \frac{11}{45} \mu_4 \mu_1^2 + \frac{14}{45} \mu_2 \mu_4 + \frac{8}{15} \mu_6 \right) t_1 t_2 (t_1^2 + t_2^2) \\ &\quad + \left(-\frac{1}{5} \mu_3^2 + \left(\frac{7}{30} \mu_1^3 - \frac{1}{15} \mu_2 \mu_1 \right) \mu_3 + \frac{7}{15} \mu_4 \mu_1^2 - \frac{2}{15} \mu_2 \mu_4 + \frac{1}{5} \mu_6 \right) t_1^2 t_2^2 \left. \right) (t_1 - t_2)^2 \\ &\quad + \cdots \in 1 + (t_1 - t_2)^2 (t_1, t_2)^2 \mathbb{Z}[\boldsymbol{\mu}]\langle\langle t_1, t_2 \rangle\rangle. \end{aligned}$$

Since $t_1 = u + \cdots \in \mathbb{Z}[\boldsymbol{\mu}]\langle\langle u \rangle\rangle$ and $t_2 = v + \cdots \in \mathbb{Z}[\boldsymbol{\mu}]\langle\langle v \rangle\rangle$, we have

$$(3.20) \quad \sigma(u - v)^2 \in (u - v)^2 \left(\text{“a series of the form } 1 + \cdots \text{ in } \mathbb{Z}[\boldsymbol{\mu}]\langle\langle u, v \rangle\rangle \text{”} \right).$$

What happens if we take a square root of $\sigma(u - v)^2$ is known by the following Lemma.

Lemma 3.21. *Let A be an integral domain with characteristic 0. Let $a_j \in A$ for $j \geq 1$ and*

$$(3.22) \quad h(z) = 1 + 2a_1 \frac{z}{1!} + 2a_2 \frac{z^2}{2!} + 2a_3 \frac{z^3}{3!} + \cdots,$$

be a power series with respect to an indeterminate z . Then a power series $\varphi(z)$ such that

$$(3.23) \quad h(z) = \varphi(z)^2$$

belongs to $A\langle\langle z \rangle\rangle$.

Proof. Since

$$(3.24) \quad \begin{aligned} &\left(1 + 2a_1 \frac{z}{1!} + 2a_2 \frac{z^2}{2!} + \cdots \right)^{-\frac{1}{2}} \\ &= 1 - \frac{1}{1!} \frac{1}{2} \left(2a_1 z + 2a_2 \frac{z^2}{2!} + 2a_3 \frac{z^3}{3!} + \cdots \right) + \frac{1}{2!} \frac{1}{2} \frac{3}{2} \left(2a_1 z + 2a_2 \frac{z^2}{2!} + 2a_3 \frac{z^3}{3!} + \cdots \right)^2 \\ &\quad - \frac{1}{3!} \frac{1}{2} \frac{3}{2} \frac{5}{2} \left(2a_1 z + 2a_2 \frac{z^2}{2!} + 2a_3 \frac{z^3}{3!} + \cdots \right)^3 + \cdots \\ &= 1 - \frac{1}{1!} \left(a_1 z + a_2 \frac{z^2}{2!} + a_3 \frac{z^3}{3!} + \cdots \right) + \frac{1}{2!} \cdot 1 \cdot 3 \left(a_1 z + a_2 \frac{z^2}{2!} + a_3 \frac{z^3}{3!} + \cdots \right)^2 \\ &\quad - \frac{1}{3!} \cdot 1 \cdot 3 \cdot 5 \left(a_1 z + a_2 \frac{z^2}{2!} + a_3 \frac{z^3}{3!} + \cdots \right)^3 + \cdots, \end{aligned}$$

the statement follows. \square

It is obvious that the expansion of $q(t)$ and $p(0, t)$ are of the form (3.22) for $A = \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_6] \langle\langle t \rangle\rangle$. However, the expansion of $p(t, 0)$ is of the form (3.22) only for $A = \mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6] \langle\langle t \rangle\rangle$. Since the denominators of the coefficients of $r(t)$ are come from the double integral in (3.19) with taking exponential, and the series of the double integral starts at a term of degree 4, we see that the expansion of $r(t)$ is also of the form (3.22) for $A = \mathbb{Z}[\mu_1, \mu_2, \mu_3, \mu_4, \mu_6] \langle\langle t \rangle\rangle$. Therefore, 3.21 implies that

$$(3.25) \quad \sigma(u) \in \mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6] \langle\langle t \rangle\rangle = \mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6] \langle\langle u \rangle\rangle.$$

If we compute this explicitly, we have

$$(3.26) \quad \begin{aligned} \sigma(u) &= t + \frac{1}{2}\mu_1 t^2 + \left(\frac{3}{2}\left(\frac{\mu_1}{2}\right)^2 + \frac{1}{2}\mu_2\right)t^3 \\ &\quad + \left(\frac{5}{2}\left(\frac{\mu_1}{2}\right)^3 + \frac{3}{2}\mu_2\left(\frac{\mu_1}{2}\right) + \frac{1}{2}\mu_3\right)t^4 \\ &\quad + \left(\frac{35}{8}\left(\frac{\mu_1}{2}\right)^4 + \frac{15}{4}\mu_2\left(\frac{\mu_1}{2}\right)^2 + \frac{29}{16}\mu_3\left(\frac{\mu_1}{2}\right) + \left(\frac{3}{8}\mu_2^2 + \frac{5}{12}\mu_4\right)\right)t^5 \\ &\quad + \left(\frac{63}{8}\left(\frac{\mu_1}{2}\right)^5 + \frac{35}{4}\mu_2\left(\frac{\mu_1}{2}\right)^3 + \frac{25}{3}\mu_3\left(\frac{\mu_1}{2}\right)^2 + \left(\frac{15}{8}\mu_2^2 + \frac{25}{12}\mu_4\right)\frac{\mu_1}{2} + \frac{5}{4}\mu_2\mu_3\right)t^6 \\ &\quad + \left(\frac{231}{16}\left(\frac{\mu_1}{2}\right)^6 + \frac{315}{16}\mu_2\left(\frac{\mu_1}{2}\right)^4 + \frac{8941}{360}\mu_3\left(\frac{\mu_1}{2}\right)^3 + \left(\frac{105}{16}\mu_2^2 + \frac{2641}{360}\mu_4\right)\left(\frac{\mu_1}{2}\right)^2 \right. \\ &\quad \left. + \frac{3091}{360}\mu_2\mu_3\left(\frac{\mu_1}{2}\right) + \frac{103}{120}\mu_3^2 + \frac{5}{16}\mu_2^3 + \frac{391}{360}\mu_4\mu_2 + \frac{13}{30}\mu_6\right)t^7 + O(t^8) \\ &= u + \left(\left(\frac{\mu_1}{2}\right)^2 + \mu_2\right)\frac{u^3}{3!} + \text{“higher terms in } \mathbb{Z}[\frac{\mu_1}{2}, \mu_2, \mu_3, \mu_4, \mu_6] \langle\langle u \rangle\rangle\text{”}. \end{aligned}$$

This result is no other than (2.11).

Remark 3.27. The author think that this result would be relate the result in [3].

4 n -plication formula

We mention here a small application. Using the expansion of the sigma function of the previous section, we can compute n -plication formula as follows.

If n is odd

$$(4.1) \quad \begin{aligned} \psi_n(u) := \frac{\sigma(nu)}{\sigma(u)^{n^2}} &= nx(u)^{\frac{n^2-1}{2}} + C_1 x(u)^{\frac{n^2-5}{2}} y(u) + C_2 x(u)^{\frac{n^2-3}{2}} \\ &+ C_3 x(u)^{\frac{n^2-7}{2}} y(u) + C_4 x(u)^{\frac{n^2-5}{2}} + \cdots + C_{n^2-1} \end{aligned}$$

is the n -plication polynomial, namely, the roots $(x(u), y(u))$ s of this polynomial are just the n -torsion points of \mathcal{E} . Comparing power series expansions of both sides with respect to u , we have explicit form of the coefficients C_j s. First several terms of them are given by

$$(4.2) \quad \begin{aligned} C_1 &= 0 \\ C_2 &= \frac{1}{24} n(n^2 - 1)\mu_1^2 + \frac{1}{6} n(n^2 - 1)\mu_2, \\ C_3 &= 0, \\ C_4 &= \frac{1}{1920} n(n^2 - 1)(n^2 - 9)\mu_1^4 + \frac{1}{240} n(n^2 - 1)(n^2 - 9)\mu_2\mu_1^2 \\ &+ \frac{1}{120} n(n^2 - 1)(n^2 + 6)\mu_3\mu_1 + \frac{1}{120} n(n^2 - 1)(n^2 - 9)\mu_2^2 + \frac{1}{60} n(n^2 - 1)(n^2 + 6)\mu_4, \\ C_5 &= 0, \\ C_6 &= \frac{1}{322560} n(n^2 - 1)(n^2 - 3^2)(n^2 - 5^2)\mu_1^6 + \frac{1}{26880} n(n^2 - 1)(n^2 - 3^2)(n^2 - 5^2)\mu_2\mu_1^4 \\ &+ \frac{1}{6720} n(n^2 - 1)(n^2 - 3^2)(n^2 + 10)\mu_3\mu_1^3 + \frac{1}{6720} n(n^2 - 1)(n^2 - 3^2)(n^2 - 5^2)\mu_2^2\mu_1^2 \\ &+ \frac{1}{3360} n(n^2 - 1)(n^2 - 3^2)(n^2 + 10)\mu_4\mu_1^2 + \frac{1}{1680} n(n^2 - 1)(n^2 - 3^2)(n^2 + 10)\mu_3\mu_2\mu_1 \\ &+ \frac{1}{5040} n(n^2 - 1)(n^2 - 3^2)(n^2 - 5^2)\mu_2^3 + \frac{1}{840} n(n^2 - 1)(n^2 - 3^2)(n^2 + 10)\mu_4\mu_2 \\ &+ \frac{1}{840} n(n^2 - 1)(n^4 + n^2 + 15)\mu_3^2 + \frac{1}{210} n(n^2 - 1)(n^4 + n^2 + 15)\mu_6. \end{aligned}$$

Note that these are elements of $\mathbb{Z}[\boldsymbol{\mu}]$ if n is odd.

If n is even

$$(4.3) \quad \begin{aligned} \psi_n(u) := \frac{\sigma(nu)}{\sigma(u)^{n^2}} &= nx(u)^{\frac{n^2-4}{2}} y(u) + C_1 x(u)^{\frac{n^2-2}{2}} + C_2 x(u)^{\frac{n^2-6}{2}} y(u) \\ &+ C_3 x(u)^{\frac{n^2-4}{2}} + C_4 x(u)^{\frac{n^2-8}{2}} y(u) + \cdots + C_{n^2-1} \end{aligned}$$

is the n -plication polynomial. The first several terms of this is given by

$$(4.4) \quad \begin{aligned} C_1 &= -\frac{1}{2} n\mu_1, \\ C_2 &= -\frac{1}{24} n(n^2 - 2^2)\mu_1^2 - \frac{1}{6} n(n^2 - 2^2)\mu_2, \\ C_3 &= -\frac{1}{48} n(n^2 - 2^2)\mu_1^3 - \frac{1}{12} n(n^2 - 2^2)\mu_2\mu_1 - \frac{1}{2} n\mu_3, \\ C_4 &= -\frac{1}{1920} n(n^2 - 2^2)(n^2 - 4^2)\mu_1^4 - \frac{1}{240} n(n^2 - 2^2)(n^2 - 4^2)\mu_2\mu_1^2 \\ &- \frac{1}{120} n(n^2 - 2^2)(n^2 + 9)\mu_3\mu_1 - \frac{1}{120} n(n^2 - 2^2)(n^2 - 4^2)\mu_2^2 \\ &- \frac{1}{60} n(n^2 - 2^2)(n^2 + 9)\mu_4, \\ C_5 &= -\frac{1}{3840} n(n^2 - 2^2)(n^2 - 4^2)\mu_1^5 - \frac{1}{480} n(n^2 - 2^2)(n^2 - 4^2)\mu_2\mu_1^3 \\ &- \frac{1}{240} n(n^2 - 2^2)(n^2 + 14)\mu_3\mu_1^2 - \frac{1}{240} n(n^2 - 2^2)(n^2 - 4^2)\mu_2^2\mu_1 \\ &- \frac{1}{120} n(n^2 - 2^2)(n^2 + 9)\mu_4\mu_1 - \frac{1}{12} n(n^2 - 2^2)\mu_3\mu_2. \end{aligned}$$

These are elements of $\mathbb{Z}[\boldsymbol{\mu}]$ if n is even.

References

- [1] H. F. Baker. *Abelian functions*. Cambridge Univ. Press, Cambridge, 1897.
- [2] J. Igusa. *Theta functions*. Springer-Verlag, Heidelberg, New York, 1972.
- [3] B. Mazur and J. Tate. The p -adic sigma function. *Duke Math. J.*, 62:663–688, 1991.
- [4] A. Nakayashiki. On algebraic expressions of sigma functions for (n, s) curves. <http://arxiv.org/abs/0803.2083>, 2007.
- [5] H. Rademacher. *Topics in analytic number theory, (Die Grundlehren der math. Wiss. Bd.169)*. Springer-Verlag, Berlin Heidelberg, New York, 1973.
- [6] K. Weierstrass. Zur theorie der elliptischen functionen. *Mathematische Werke Bd.2*, pages 245–255, 1894.