

INTRINSIC ENERGY IS A LOOP SCHUR FUNCTION

THOMAS LAM AND PAVLO PYLYAVSKYY

ABSTRACT. We give an explicit subtraction-free formula for the energy function in tensor products of Kirillov-Reshetikhin crystals for symmetric powers of the standard representation of $U'_q(\hat{\mathfrak{sl}}_n)$. The energy function is shown to be the tropicalization of a stretched staircase shape loop Schur function. The latter were introduced by the authors in the study of total positivity in loop groups.

1. INTRODUCTION

The intrinsic energy function plays an important role in the path model for affine highest weight crystals [7]. The energy function is also related to charge statistic of Lascoux-Schützenberger on semistandard tableaux (see [12]), which establishes a relation between one dimensional configuration sums arising in solvable lattice models and Kostka-Foulkes polynomials, cf. [2, 7, 8].

Let $B = B_1 \otimes \cdots \otimes B_m$ be a tensor product of $U'_q(\hat{\mathfrak{sl}}_n)$ Kirillov-Reshetikhin crystals, where each B_i is the crystal for a symmetric power of the standard representation. We identify B_i with the semistandard Young tableaux with row shape, filled with the numbers $1, 2, \dots, n$. Let $b = b_1 \otimes \cdots \otimes b_m \in B$, and write $x_i^{(r+i-1)}$ for the number of r 's in b_i . The upper index $(r+i-1)$ is to be considered as an element of $\mathbb{Z}/n\mathbb{Z}$. Our main result is the following formula for the intrinsic energy function \overline{D}_B of B .

Let $\delta_t = (t, t-1, \dots, 1)$ denote the staircase shape of side-length t .

Theorem 1.1. *We have*

$$\overline{D}_B(b) = \min_T \left\{ \sum_{(i,j) \in (n-1)\delta_{m-1}} x_{T(i,j)}^{(i-j)} \right\},$$

where the minimum is over all semistandard tableaux T of shape $(n-1)\delta_{m-1}$, and entries in $1, 2, \dots, m$.

In the physical interpretation, each b_i represents a particle, and the intrinsic energy function $\overline{D}_B(b)$ is defined as the sum of $\binom{m}{2}$ local energies of interactions of particles. Theorem 1.1 thus has the following interpretation: each tableau T encodes a way for m particles to interact *simultaneously*, and intrinsic energy is equal to the minimum of these.

In [10], motivated by the study of total positivity for loop groups, we introduced a generalization of the ring of symmetric functions, called *loop symmetric functions* and denoted LSym . In particular, we defined distinguished elements of LSym called loop Schur functions (see Section 2). It is shown in [10] that the algebra homomorphisms from

Date: November 1, 2021.

T.L. was supported by NSF grant DMS-0652641 and DMS-0901111, and by a Sloan Fellowship.

P.P. was supported by NSF grant DMS-0757165.

LSym to \mathbb{R} taking nonnegative values on (skew) loop Schur functions are in bijection with totally nonnegative elements of the formal loop group.

Recall that the tropicalization of a subtraction-free polynomial f , is obtained by replacing multiplication by addition, and replacing addition by taking minimums. Theorem 1.1 is equivalent to

Theorem 1.2. *The function \overline{D}_B is the tropicalization of the loop Schur function $s_{(n-1)\delta_{m-1}}^{(0)}$ in the variables $\{\mathbf{x}_i^{(s)}\}$.*

Theorem 1.2 is an immediate consequence of Theorems 2.5 and 3.2 below.

Theorems 1.1 and 1.2 are *canonical* in the sense that they correspond to the monomial expansion of a polynomial. That a piecewise-linear expression for $\overline{D}_B(b)$ exists is already clear from the literature. However, the fact a subtraction-free formula exists (or equivalently the rational version $\overline{D}_B(b)$ of energy is a polynomial with positive coefficients) is not apparent from the definition of $\overline{D}_B(b)$, even though the latter takes nonnegative values. (See also Remark 3.)

Example 1. Let $n = 2$ and $m = 3$. Then

$$\overline{D}_B(b) = \min(x_1^{(1)} + x_1^{(2)} + x_2^{(1)}, x_2^{(1)} + x_1^{(2)} + x_2^{(2)}, x_3^{(1)} + x_1^{(2)} + x_2^{(1)}, x_1^{(1)} + x_1^{(2)} + x_3^{(1)}, \\ x_2^{(1)} + x_1^{(2)} + x_3^{(1)}, x_3^{(1)} + x_1^{(2)} + x_3^{(2)}, x_2^{(1)} + x_2^{(2)} + x_3^{(1)}, x_3^{(1)} + x_2^{(2)} + x_3^{(1)})$$

corresponding to the following tableaux of shape $\delta_2 = (2, 1)$:

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|} \hline 1 & 3 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 3 \\ \hline 3 & \\ \hline \end{array}$$

We use in our calculations a birational analogue of the *combinatorial R-matrix*. It was previously studied by Kirillov [6] in the context of the Robinson-Schensted algorithm, by Noumi-Yamada [13, 17] in the context of discrete Painlevé systems, by Etingof [3] in the context Yang-Baxter equations, by Berenstein-Kazhdan [1] in the context of geometric crystals, and by the authors [10] in the context of total positivity of loop groups.

2. LOOP SYMMETRIC FUNCTIONS

Fix an integer $n > 1$ throughout.

2.1. Loop Schur functions. Let $(\mathbf{x}_i^{(r)})_{1 \leq i \leq m, r \in \mathbb{Z}/n\mathbb{Z}}$ be a rectangular array of variables. We recall from [10] the definition of the ring of *loop symmetric functions*¹ in the variables $\mathbf{x}_i^{(r)}$, denoted LSym_m . A detailed study of loop symmetric functions will appear in [11].

For $k \geq 1$ and $r \in \mathbb{Z}/n\mathbb{Z}$, define the *loop elementary symmetric functions* and *loop complete homogenous symmetric functions* by


$$e_k^{(r)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq m} \mathbf{x}_{i_1}^{(r)} \mathbf{x}_{i_2}^{(r+1)} \dots \mathbf{x}_{i_k}^{(r+k-1)} \\ h_k^{(r)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m) = \sum_{1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq m} \mathbf{x}_{i_1}^{(r)} \mathbf{x}_{i_2}^{(r-1)} \dots \mathbf{x}_{i_k}^{(r-k+1)}$$

¹In [10], there are two such rings: the ring of whirl loop symmetric functions, and the ring of curl loop symmetric functions. We use the former here. Furthermore, we only use finitely many variables here.

By convention, $e_k^{(r)} = h_k^{(r)} = 0$ for $k < 0$, and $e_0^{(r)} = h_0^{(r)} = 1$. Note that $e_k^{(r)} = 0$ for $k > m$. We call the upper index the *color*. When all n colors are identified, that is $\mathbf{x}_i^{(s)} = \mathbf{x}_i^{(s')}$ for all i and $s, s' \in \mathbb{Z}/n\mathbb{Z}$, these functions specialise to the usual elementary and complete homogenous symmetric functions [16]. We define LSym_m to be the ring generated by the $e_k^{(r)}$. Although it is not immediately obvious, the $h_k^{(r)}$ lie in LSym_m . In fact, both the $e_k^{(r)}$ and the $h_k^{(r)}$ are instances of distinguished elements of LSym_m called *loop Schur functions*.

A square $s = (i, j)$ in the i -th row and j -th column has *content* $c(s) = i - j$. We caution that our notion of content is the negative of the usual one. Let ρ/ν be a skew shape. Recall that a semistandard Young tableaux T with shape ρ/ν is a filling of each square $s \in \rho/\nu$ with an integer $T(s) \in \mathbb{Z}_{>0}$ so that the rows are weakly-increasing, and columns are increasing. For $r \in \mathbb{Z}/n\mathbb{Z}$, the r -weight \mathbf{x}^T of a tableaux T is given by $\mathbf{x}^T = \prod_{s \in \rho/\nu} \mathbf{x}_{T(s)}^{(c(s)+r)}$.

We shall draw our shapes and tableaux in English notation:



For $n = 3$ the 0-weight of the above tableaux is $(\mathbf{x}_1^{(1)})^2(\mathbf{x}_3^{(1)})^3\mathbf{x}_1^{(2)}\mathbf{x}_2^{(2)}\mathbf{x}_3^{(2)}\mathbf{x}_1^{(3)}\mathbf{x}_2^{(3)}(\mathbf{x}_4^{(3)})^2$. We define the *loop (skew) Schur function* by

$$s_{\rho/\nu}^{(r)}(\mathbf{x}) = \sum_T \mathbf{x}^T$$

where the summation is over all semistandard Young tableaux of (skew) shape ρ/ν . We have the following analog of the Jacobi-Trudi formula.

Theorem 2.1. [10, Theorem 7.6] *We have $s_{\lambda'/\mu'}^{(r)} = \det(e_{\lambda_i - \mu_j - i + j}^{(r-j+1+\mu_j)})$.*

Proposition 2.2. *For any $k > 0$ we have*

$$e_0^{(r)} h_k^{(r-1)} - e_1^{(r-1)} h_{k-1}^{(r-2)} + e_2^{(r-2)} h_{k-2}^{(r-3)} - \dots \pm e_k^{(r-k)} h_0^{(r-k-1)} = 0.$$

Proof. Let ρ_i denote the hook shape $(k - i, 1, \dots, 1)$ of size k . Then $e_i^{(r-i)} h_{k-i}^{(r-i-1)} = s_{\rho_{k-i}}^{(r-i-1)} + s_{\rho_{k-i+1}}^{(r-i)}$ is the sum of two loop Schur functions (one of which is zero if $i = 0$ or $i = k$). This holds since any two terms in $e_i^{(r-i)}$ and $h_{k-i}^{(r-i-1)}$, viewed as tableaux of column and row shapes, fit together to give a semistandard tableau of one of the two hook shapes, depending on the entry in their smallest boxes. It is clear that as we sum over all i , all the hook shape loop Schur functions cancel out. \square

2.2. Staircases. For $k \geq 1$ and $r \in \mathbb{Z}/n\mathbb{Z}$, define

$$\tau_k^{(r)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m) = \sum_{I=\{i_1 \leq i_2 \leq \dots \leq i_k\}} \mathbf{x}_{i_1}^{(r)} \mathbf{x}_{i_2}^{(r-1)} \dots \mathbf{x}_{i_k}^{(r-k+1)}$$

where the summation is over multisets $I \subset \{1, 2, 3, \dots, m\}$ such that no number occurs more than $n - 1$ times. Note that if $k > m(n - 1)$ we have $\tau_k^{(r)} = 0$. It can be shown that $\tau_k^{(r)}$ lies in LSym_m , but we shall not need it for what follows.

Lemma 2.3. *We have*

$$\tau_k^{(r)}(\mathbf{x}_1, \dots, \mathbf{x}_m) = \sum_{i=0}^{\infty} (-1)^i h_{k-in}^{(r)} e_i \left(\prod_{s \in \mathbb{Z}/n\mathbb{Z}} \mathbf{x}_1^{(s)}, \dots, \prod_{s \in \mathbb{Z}/n\mathbb{Z}} \mathbf{x}_m^{(s)} \right),$$

where the e_i in the above formula denotes the usual elementary symmetric function.

Proof. Let $\mathbf{x}_{i_1}^{(r)} \mathbf{x}_{i_2}^{(r-1)} \dots \mathbf{x}_{i_k}^{(r-k+1)}$ be a term in $h_k^{(r)}$. Let $J \subseteq I$ be the set of indexes which occur in $I = \{i_1 \leq i_2 \leq \dots \leq i_k\}$ more than $n-1$ times. Then the coefficient of this term on the right is equal to $\sum_{K \subseteq J} (-1)^{|K|}$. This is equal to 1 if $|J| = 0$ and to 0 otherwise. \square

Lemma 2.4. *For each k , we have*

$$\sum_{i=0}^{\infty} (-1)^i e_i^{(r-i)}(\mathbf{x}_1, \dots, \mathbf{x}_m) \tau_{k-i}^{(r-i-1)}(\mathbf{x}_1, \dots, \mathbf{x}_m) = 0.$$

Proof. Follows immediately from Proposition 2.2 and Lemma 2.3. \square

Define $\sigma_k^{(r)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m) = \sum_{i=0}^k \mathbf{x}_1^{(r)} \mathbf{x}_1^{(r-1)} \dots \mathbf{x}_1^{(r-i+1)} \tau_{k-i}^{(r-i)}(\mathbf{x}_2, \dots, \mathbf{x}_m)$.

Theorem 2.5. *For $m \geq 2$ and $r \in \mathbb{Z}/n\mathbb{Z}$, we have*

$$\begin{aligned} & s_{(n-1)\delta_{m-1}}^{(r)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m) \\ &= \sigma_{(n-1)(m-1)}^{(r)}(\mathbf{x}_1, \dots, \mathbf{x}_m) \sigma_{(n-1)(m-2)}^{(r+1)}(\mathbf{x}_2, \dots, \mathbf{x}_m) \dots \sigma_{(n-1)}^{(r+m-2)}(\mathbf{x}_{m-1}, \mathbf{x}_m). \end{aligned}$$

Remark 1. When all the colors are identified, that is, $\mathbf{x}_i^{(s)} = \mathbf{x}_i^{(s')}$ for all i and $s, s' \in \mathbb{Z}/n\mathbb{Z}$, Theorem 2.5 is a coarsening of a result of Jucis [5], see also [16, Ex. 7.30].

It is clear that Theorem 2.5 holds for $m = 2$, for then it states that $s_{n-1}^{(r)}(\mathbf{x}_1, \mathbf{x}_2) = \sigma_{n-1}^{(r)}(\mathbf{x}_1, \mathbf{x}_2)$. We shall prove Theorem 2.5 in Section 4.

3. AFFINE CRYSTALS

3.1. R -matrix. We shall use [15] as our main reference for affine crystals.

Recall that a *Kirillov-Reshetikhin crystal* of $U'_q(\hat{\mathfrak{sl}}_n)$ is the crystal graph corresponding to the highest weight module with highest weight proportional to one of the fundamental weights. An *affine crystal* is the tensor product of several Kirillov-Reshetikhin crystals. We shall restrict our attention to the set \mathfrak{C} of affine crystals that are tensor products of symmetric powers of the standard representation. Each element $b \in B$ of such a symmetric power can be identified with a single row semistandard tableau in the alphabet $1, \dots, n$.

If B_1, B_2 are Kirillov-Reshetikhin crystals, the *combinatorial R -matrix* is the unique isomorphism $R_{B_1, B_2} : B_1 \otimes B_2 \rightarrow B_2 \otimes B_1$ of affine crystals. It is known that the combinatorial R -matrices generate an action of S_m on $B_1 \otimes \dots \otimes B_m$.

The combinatorial R -matrix has a convenient interpretation [15] in terms of semistandard tableaux and the jeu de taquin algorithm [16]. Let $b_1 \otimes b_2$ be an element of $B_1 \otimes B_2$. Then $R_{B_1, B_2}(b_1 \otimes b_2) = c_1 \otimes c_2 \in B_2 \otimes B_1$ where c_1, c_2 are the unique pair of row shaped tableaux which jeu de taquin to the same tableau that b_1 and b_2 jeu de taquin to, as follows:

$$\begin{array}{|c|c|} \hline & 1 & 2 & 2 & 4 \\ \hline 1 & 3 & & & \\ \hline \end{array} \xrightarrow{R_{B_1, B_2}} \begin{array}{|c|c|c|c|} \hline & & 2 & 4 \\ \hline 1 & 1 & 2 & 3 \\ \hline \end{array} \text{ since both jeu de taquin to } \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 2 & 2 & 4 \\ \hline 3 & & & & \\ \hline \end{array}$$

We measure $\overline{H}(b_1, b_2) = 1$, $\overline{H}(b_2, b_3) = 2$. We also have seen above the result of applying R_{B_1, B_2} to the first two tensor factors, which allows us to measure $\overline{H}(s_1(b_2), b_3) = 2$. Therefore $\overline{D}_B(b) = 1 + 2 + 2 = 5$.

$$\begin{array}{|c|c|c|c|} \hline 1 & 2 & 2 & 4 \\ \hline 1 & 3 & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|c|} \hline & & 1 & 2 & 3 \\ \hline 1 & 2 & 2 & 4 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 2 & 4 & \\ \hline \end{array}$$

3.3. Product (summation) formula for intrinsic energy. In this section, we switch from piecewise linear functions to rational functions. The two worlds are connected via tropicalization: if f is a subtraction-free polynomial, we let $\text{trop}(f)$ denote the tropicalization of f , obtained by replacing addition by minimum, and multiplication by addition.

We are given a rectangular array of variables $\bar{x}_i^{(r)}$, $i = 1, \dots, m$, $r \in \mathbb{Z}/n\mathbb{Z}$, with columns $b_i = (\bar{x}_i^{(1)}, \bar{x}_i^{(2)}, \dots, \bar{x}_i^{(n)})$. It is very convenient to make the following change of variables: $\mathbf{x}_i^{(r)} = \bar{x}_i^{(r+1-i)}$. Define

$$\kappa_r(b_j, b_{j+1}) = \sum_{s=0}^{n-1} \left(\prod_{t=1}^s \mathbf{x}_{j+1}^{(r+t)} \prod_{t=s+1}^{n-1} \mathbf{x}_j^{(r+t)} \right),$$

so that $\text{trop}(\kappa_r(b_j, b_{j+1})) = \overline{\kappa}_{r-j+1}(b_j, b_{j+1})$.

In the variables $\mathbf{x}_i^{(r)}$, the birational R -matrix acts (see [17, Proposition 3.1]²) via algebra isomorphisms s_1, s_2, \dots, s_{m-1} of the field of rational functions in $\{\mathbf{x}_i^{(r)}\}$, given by

$$s_j(\mathbf{x}_j^{(r)}) = \frac{\mathbf{x}_{j+1}^{(r+1)} \kappa_{r+1}(b_j, b_{j+1})}{\kappa_r(b_j, b_{j+1})} \quad \text{and} \quad s_j(\mathbf{x}_{j+1}^{(r)}) = \frac{\mathbf{x}_j^{(r-1)} \kappa_{r-1}(b_j, b_{j+1})}{\kappa_r(b_j, b_{j+1})}$$

and $s_j(\mathbf{x}_k^{(r)}) = \mathbf{x}_k^{(r)}$ for $k \neq j, j+1$. We also have $\overline{H}(b_j \otimes b_{j+1}) = \kappa_j(b_j, b_{j+1})$, the rational analogue of the local coenergy. We let \overline{D}_B denote the rational analogue of the intrinsic energy function, so that $\text{trop}(\overline{D}_B) = \overline{D}_B$. The main result of this section is a product formula for \overline{D}_B . We remark that Kirillov [6] has also studied the rational functions \overline{H} and \overline{D}_B .

Lemma 3.1. *Suppose $1 \leq i < j \leq m$. Then*

$$\kappa_r(s_i s_{i+1} \cdots s_{j-2}(b_{j-1}), b_j) = \frac{\sigma_{(n-1)(j-i)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_j)}{\sigma_{(n-1)(j-i-1)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_{j-1})}$$

and

$$s_i s_{i+1} \cdots s_{j-1}(\mathbf{x}_j^{(r)}) = \frac{\mathbf{x}_i^{(r-j+i)} \sigma_{(n-1)(j-i)}^{(r-j+i-1)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_j)}{\sigma_{(n-1)(j-i)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_j)} = \frac{\sigma_{(n-1)(j-i)+1}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_j)}{\sigma_{(n-1)(j-i)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_j)}.$$

Proof. We prove the two statements in parallel by induction on $j-i$. For $j-i=1$ they coincide with the formulae for the κ_r and the R -action of s_i . By the induction assumption

$$s_i s_{i+1} \cdots s_{j-2}(\mathbf{x}_{j-1}^{(r)}) = \frac{\mathbf{x}_i^{(r-j+i+1)} \sigma_{(n-1)(j-i-1)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_{j-1})}{\sigma_{(n-1)(j-i-1)}^{(r-j+i+1)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_{j-1})}.$$

²Our variables $\bar{x}_i^{(r)}$ are nearly the same as Yamada's x_r^i , differing by a reversal of the orientation of the circle.

Therefore

$$\begin{aligned}
& \kappa_r(s_i s_{i+1} \cdots s_{j-2}(b_{j-1}), b_j) \\
&= \sum_{s=0}^{n-1} \left(\prod_{t=1}^s \mathbf{x}_j^{(r+t)} \prod_{t=s+1}^{n-1} s_i s_{i+1} \cdots s_{j-2}(\mathbf{x}_{j-1}^{(r+t)}) \right) \\
&= \sum_{s=0}^{n-1} \frac{\prod_{t=s-j+i+2}^{n-j+i} \mathbf{x}_i^{(r+t)} \sigma_{(n-1)(j-i-1)}^{(r+s-j+i+1)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_{j-1}) \prod_{t=1}^s \mathbf{x}_j^{(r+t)}}{\sigma_{(n-1)(j-i-1)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_{j-1})} \\
&= \frac{\sigma_{(n-1)(j-i)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_j)}{\sigma_{(n-1)(j-i-1)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_{j-1})}.
\end{aligned}$$

The last equality holds because in a term of $\sigma_{(n-1)(j-i)}^{(r-j+i)}(\mathbf{x}_i, \mathbf{x}_{i+1}, \dots, \mathbf{x}_j)$ the number of the $\mathbf{x}_j^{(t)}$ -s is at most $n-1$, while the number of the $\mathbf{x}_i^{(t)}$ -s and the $\mathbf{x}_j^{(t)}$ -s together should be at least $n-1$.

Now we can also prove the second claim, since

$$\begin{aligned}
s_i s_{i+1} \cdots s_{j-1}(\mathbf{x}_j^{(r)}) &= \frac{s_i s_{i+1} \cdots s_{j-2}(\mathbf{x}_{j-1}^{(r-1)}) \kappa_{r-1}(s_i s_{i+1} \cdots s_{j-2}(b_{j-1}), b_j)}{\kappa_r(s_i s_{i+1} \cdots s_{j-2}(b_{j-1}), b_j)} = \\
&= \frac{\mathbf{x}_i^{(r-j+i)} \sigma_{(n-1)(j-i-1)}^{(r-j+i-1)}(\mathbf{x}_i, \dots, \mathbf{x}_{j-1})}{\sigma_{(n-1)(j-i-1)}^{(r-j+i)}(\mathbf{x}_i, \dots, \mathbf{x}_{j-1})} \frac{\sigma_{(n-1)(j-i)}^{(r-j+i-1)}(\mathbf{x}_i, \dots, \mathbf{x}_j)}{\sigma_{(n-1)(j-i-1)}^{(r-j+i-1)}(\mathbf{x}_i, \dots, \mathbf{x}_{j-1})} \frac{\sigma_{(n-1)(j-i-1)}^{(r-j+i)}(\mathbf{x}_i, \dots, \mathbf{x}_{j-1})}{\sigma_{(n-1)(j-i)}^{(r-j+i)}(\mathbf{x}_i, \dots, \mathbf{x}_j)} \\
&= \frac{\mathbf{x}_i^{(r-j+i)} \sigma_{(n-1)(j-i)}^{(r-j+i-1)}(\mathbf{x}_i, \dots, \mathbf{x}_j)}{\sigma_{(n-1)(j-i)}^{(r-j+i)}(\mathbf{x}_i, \dots, \mathbf{x}_j)}.
\end{aligned}$$

□

Theorem 3.2. *We have*

$$\bar{\mathbb{D}}_B(b) = \sigma_{(n-1)(m-1)}^{(n)}(\mathbf{x}_1, \dots, \mathbf{x}_m) \sigma_{(n-1)(m-2)}^{(1)}(\mathbf{x}_2, \dots, \mathbf{x}_m) \cdots \sigma_{(n-1)}^{(m-2)}(\mathbf{x}_{m-1}, \mathbf{x}_m).$$

Proof. The result follows from Lemma 3.1 and (1). □

Remark 2. Comparing [12, Theorem 4.2] with Lemma 3.1 and Theorem 3.2 one can see that the tropicalization of the $\sigma_{(n-1)(m-1-i)}^{(i)}(\mathbf{x}_{i+1}, \dots, \mathbf{x}_m)$ is essentially the index $\text{ind}(m-i)$ in the index decomposition of charge in [12].

Remark 3. Theorem 3.2 gives the irreducible factorization of $\bar{\mathbb{D}}_B(b)$ (whereas Theorem 1.2 gives the monomial expansion). To see this, one first notes that $\sigma_{(n-1)}^{(r)}(\mathbf{x}_1, \mathbf{x}_2)$ has a unique monomial which involves $\mathbf{x}_2^{(r)}$, from which one deduces the irreducibility. Now suppose that $\sigma_{(n-1)(m-1)}^{(r)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m)$ factorizes non-trivially as the product fg . We may write $f = a \mathbf{x}_m^{(r-(n-1)(m-2))} + b$ and $g = c$ as polynomials in $\mathbf{x}_m^{(r-(n-1)(m-2))}$, where a, b, c do not involve $\mathbf{x}_m^{(r-(n-1)(m-2))}$. One verifies that none of the variables $\mathbf{x}_m^{(s)}$ divide $\sigma_{(n-1)(m-1)}^{(r)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m)$ and every monomial which contains $\mathbf{x}_m^{(r-(n-1)(m-2))}$ is divisible by the product $\mathbf{x}_m^{(r-(n-1)(m-2))} \mathbf{x}_m^{(r-(n-1)(m-2)-1)} \cdots \mathbf{x}_m^{(r-(n-1)(m-1)+1)}$. Thus we have

Proof. Since $\det(B_{i,m}) = s_{(n-1)\delta_m/(i-1)}^{(r-1)}$ is never the zero polynomial in the $e_i^{(s)}$, we deduce that considered as a matrix with coefficients in the field $\text{Frac}(\mathbf{x}_i^{(s)})$, the matrix B_m has maximal rank. There is thus, up to scaling, a unique solution to the equation $B_m \cdot v = 0$. By expanding the determinant of the matrix obtained from B_m by repeating a row, it is easy to see that $v = (\det(B_{1,m}), -\det(B_{2,m}), \dots, \pm \det(B_{na+n,m}))$ is a solution. But by Proposition 4.2 so is \mathcal{T} . Thus the two vectors are proportional, and it remains to check that the scaling coefficient is $\det(A_m)$. We have $\det(B_{1,m}) = e_m^{(r-1)} \dots e_m^{(r-n+1)} \det(A_m)$, while at the same time $\tau_{(n-1)m}^{(r-1)} = e_m^{(r-1)} \dots e_m^{(r-n+1)}$, and the statement follows. \square

Proof of Theorem 2.5. We have already verified the case $m = 2$, so we suppose that $m > 2$, and by induction on m that

$$\begin{aligned} & s_{(n-1)\delta_{m-1}}^{(s+1)}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{m+1}) \\ &= \sigma_{(n-1)(m-1)}^{(s+1)}(\mathbf{x}_2, \dots, \mathbf{x}_{m+1}) \sigma_{(n-1)(m-2)}^{(s+2)}(\mathbf{x}_3, \dots, \mathbf{x}_{m+1}) \cdots \sigma_{(n-1)}^{(s+m-1)}(\mathbf{x}_m, \mathbf{x}_{m+1}) \end{aligned}$$

for every $s \in \mathbb{Z}/n\mathbb{Z}$. We calculate that

$$\begin{aligned} & s_{(n-1)\delta_m}^{(r-1)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{m+1}) \\ &= \sum_{i=0}^{(n-1)m} \mathbf{x}_1^{(r-1)} \mathbf{x}_1^{(r-2)} \cdots \mathbf{x}_1^{(r-i)} s_{(n-1)\delta_m/(i)}^{(r-1)}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{m+1}) \\ &= \sum_{i=0}^{(n-1)m} \mathbf{x}_1^{(r-1)} \mathbf{x}_1^{(r-2)} \cdots \mathbf{x}_1^{(r-i)} \det(B_{i,m})(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{m+1}) \\ &= \left(\sum_{i=0}^{(n-1)m} \mathbf{x}_1^{(r-1)} \mathbf{x}_1^{(r-2)} \cdots \mathbf{x}_1^{(r-i)} \tau_{(n-1)m-i+1}^{(r-i-1)}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{m+1}) \right) s_{(n-1)\delta_{m-1}}^{(r)}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{m+1}) \\ &= \sigma_{(n-1)m}^{(r-1)}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{m+1}) s_{(n-1)\delta_{m-1}}^{(r)}(\mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_{m+1}) \end{aligned}$$

where in the first equality we used the tableau definition of $s_{(n-1)\delta_m}^{(r)}$, and in the penultimate equality we used Proposition 4.3. \square

Remark 4. It is clear from the theory developed in [10] that loop Schur functions are invariants of the action of the symmetric group S_m via the birational R -action. Thus we have demonstrated directly that the energy function is an invariant of this action. This property is not obvious from the definition we use.

REFERENCES

- [1] A. BERENSTEIN AND D. KAZHDAN: Geometric and unipotent crystals. II. From unipotent bicrystals to crystal bases. *Quantum groups*, 13–88, *Contemp. Math.*, 433, Amer. Math. Soc., Providence, RI, 2007.
- [2] E. DATE, M. JIMBO, A. KUNIBA, T. MIWA, AND M. OKADO: One dimensional configuration sums in vertex models and affine Lie algebra characters. *LMP*, **17**, 69–77, 1989.
- [3] P. ETINGOF: Geometric crystals and set-theoretical solutions to the quantum Yang-Baxter equation. *Comm. Algebra* **31** (2003), no. 4, 1961–1973.

- [4] G. HATAYAMA, K. HIKAMI, R. INOUE, A. KUNIBA, T. TAKAGI, AND T. TOKIHIRO: The $A_M^{(1)}$ automata related to crystals of symmetric tensors. (English summary) *J. Math. Phys.* **42** (2001), no. 1, 274–308.
- [5] A.-A.A. JUCIS: Tournaments and generalized Young tableaux. (Russian) *Mat. Zametki* **27** (1980), no. 3, 353–359, 492.
- [6] A. KIRILLOV: Introduction to tropical combinatorics. (English summary) *Physics and combinatorics, 2000 (Nagoya)*, 82–150, World Sci. Publ., River Edge, NJ, 2001.
- [7] S.-J. KANG, M. KASHIWARA, K.C. MISRA, T. MIWA, T. NAKASHIMA, AND A. NAKAYASHI: Affine crystals and vertex models, in *Infinite analysis Part A (Kyoto 1991)*, 449–484, *Adv. Ser. Math. Phys.*, **16**, World Sci. Publishing, River Edge, NJ, 1992.
- [8] S.-J. KANG, M. KASHIWARA, K.C. MISRA, T. MIWA, T. NAKASHIMA, AND A. NAKAYASHI: Perfect crystals of quantum affine Lie algebras. *Duke Math. J.*, **68**, no. 3, 499–607, 1992.
- [9] A. KIRILLOV, A. KUNIBA, AND T. NAKANISHI: Skew Young diagram method in spectral decomposition of integrable lattice models. II. Higher levels. *Nuclear Phys. B* **529** (1998), no. 3, 611–638.
- [10] T. LAM AND P. PYLYAVSKYY: Total positivity in loop groups I: whirls and curls, *Adv. in Math.*, to appear.
- [11] T. LAM AND P. PYLYAVSKYY: Loop symmetric functions, in preparation.
- [12] A. NAKAYASHIKI AND Y. YAMADA: Kostka polynomials and energy functions in solvable lattice models. *Selecta Mathematica* **3** (1997), 547–600.
- [13] M. NOUMI AND Y. YAMADA: Affine Weyl group symmetry in Painlevé hierarchies. *Painlevé systems, hypergeometric systems and asymptotic analysis (Kyoto, 1999)*.
- [14] M. OKADO, A. SCHILLING, AND M. SHIMOZONO: Virtual crystals and fermionic formulas of type $D_{n+1}^{(2)}$, $A_{2n}^{(2)}$, and $C_n^{(1)}$. *Represent. Theory* **7** (2003), 101–163.
- [15] M. SHIMOZONO: Crystals for dummies, available at <http://www.aimath.org/WWN/kostka/crysdumb.pdf>.
- [16] R. STANLEY: *Enumerative Combinatorics, Volume 2*, Cambridge Studies in Advanced Mathematics 62, Cambridge University Press, 2001.
- [17] Y. YAMADA: A birational representation of Weyl group, combinatorial R -matrix and discrete Toda equation. *Physics and combinatorics, 2000 (Nagoya)*, 305–319, World Sci. Publ., River Edge, NJ, 2001.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109 USA.

E-mail address: tfylam@umich.edu

URL: <http://www.math.lsa.umich.edu/~tfylam>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109 USA.

E-mail address: pavlo@umich.edu

URL: <http://sites.google.com/site/pylyavskyy/>