

A REFINED BIJECTION BETWEEN ALTERNATING PERMUTATIONS AND 0-1-2 INCREASING TREES

HEESUNG SHIN

ABSTRACT. We construct a refined bijection φ between alternating permutations and 0-1-2 increasing trees with degree at most 2. It satisfies that the first element of alternating permutation π is equal to the first vertex in $\varphi(\pi)$ in the postorder.

1. INTRODUCTION

Let \mathcal{A}_n be the set of *alternating permutations* $\pi = \pi_1\pi_2\dots\pi_n$ on $[n] := \{1, 2, \dots, n\}$ satisfying

$$\pi_1 > \pi_2 < \pi_3 > \pi_4 < \dots.$$

Let $\mathcal{A}_{n,k} = \{\pi \in \mathcal{A}_n : \pi_1 = k\}$. Let \mathcal{T}_n be the set of *0-1-2 increasing trees* T on $[n]$ with degree at most 2. Given a increasing tree T , the unique path towards a smallest child of each vertex from the root 1 is called the *main chain* of T and we denote the leaf at the end of the main chain by $p(T)$. Let $\mathcal{T}_{n,k} = \{T \in \mathcal{T}_n : p(T) = k\}$.

It is important that \mathcal{A}_n and \mathcal{T}_n are equinumerous (see [Foa72, FS74], and also the bijection in [FS73, Don75]). Also $\mathcal{A}_{n,k}$ and $\mathcal{T}_{n,k}$ are equinumerous [Pou82, KPP94]. However, no one has managed to establish explicit bijections between $\mathcal{A}_{n,k}$ and $\mathcal{T}_{n,k}$, as mentioned in the paper of Kuznetsov, Pak and Postnikov [KPP94], who asked also for a combinatorial explanation of the identity


$$|\mathcal{T}_{n,k}| = |\mathcal{T}_{n,k-1}| + |\mathcal{T}_{n-1,n-k+1}|$$

in the model of 0-1-2 increasing trees. The aim of this paper is to give a bijection between $\mathcal{A}_{n,k}$ and $\mathcal{T}_{n,k}$. To the best knowledge of the author, this is the first bijection between these two sets.

2. A BIJECTION BETWEEN $\mathcal{A}_{n,k}$ AND $\mathcal{T}_{n,k}$

For $n = 1$ or 2 , since $|\mathcal{A}_n| = |\mathcal{T}_n| = 1$, we can define trivially $\varphi : \mathcal{A}_{n,k} \rightarrow \mathcal{T}_{n,k}$. For $n \geq 3$, given $\pi \in \mathcal{A}_{n,k}$ ($k = \pi_1$), we define the mapping $\varphi : \mathcal{A}_{n,k} \rightarrow \mathcal{T}_{n,k}$ recursively as follows:

- (a) If $\pi_2 = k - 1$, then define $\pi' \in \mathcal{A}_{n-2,i-2}$ by deleting $k - 1$ and k from π and relabeling by $[n - 2]$ where $i > k$. We get $T' = \varphi(\pi') \in \mathcal{T}_{n-2,i-2}$. Relabel T' by $\{1, \dots, k - 2, k + 1, \dots, n\}$, denoted by T'' . Let m be the minimal vertex greater than k in the main chain of T'' and j the parent of m in T'' . Then insert a vertex $k - 1$ in the middle of the edge (j, m) and add the edge $(k, k - 1)$.




We get the tree $T = \varphi(\pi) \in \mathcal{T}_{n,k}$.

π	1	213	312	21534	31524	41523
π_1	1	2	3	2	3	4
$\text{inv}(\pi)$	0	1	2	3	4	5
$31-2(\pi)$	0	0	1	1	2	3
$\varphi(\pi)$						
$p(\varphi(\pi))$	1	2	3	2	3	4
π	51423	5471623	6471523	548691723	648591723	748591623
π_1	5	5	6	5	6	7
$\text{inv}(\pi)$	6	13	14	21	22	23
$31-2(\pi)$	4	4	5	5	6	7
$\varphi(\pi)$						
$p(\varphi(\pi))$	5	5	6	5	6	7


FIGURE 1. Constructing a tree $\varphi(748591623)$ by a recursive algorithm

(b) If $\pi_2 < k - 1$, then define $\pi' = (k - 1 \ k)\pi \in \mathcal{A}_{n, k-1}$ (exchange $k - 1$ and k in π). We get $T' = \varphi(\pi') \in \mathcal{T}_{n, k-1}$.

(1) If k is a sibling of $k - 1$ in T' , then we get the tree $T = \varphi(\pi) \in \mathcal{T}_{n, k}$ modifying as follows:



(2) If k is a not sibling of $k - 1$ in T' , then we get the tree $T = \varphi(\pi) \in \mathcal{T}_{n, k}$ exchanging the labels $k - 1$ and k in T' .




Theorem 1. For all $n \geq 1$ and $k \in [n]$ The mapping φ is a bijection between $\mathcal{A}_{n, k}$ and $\mathcal{T}_{n, k}$ satisfying

$$\pi_1 = p(\varphi(\pi)).$$


Proof. It is sufficient to construct the inverse mapping of φ . Given $T \in \mathcal{T}_{n,k}$ ($k = p(T)$), we define $\pi = \varphi^{-1}(T)$ recursively as follows:

- (A) If $k-1$ is a parent of k in T , then let $m(> k)$ be the another child of $k-1$ ($m = \infty$ if $k-1$ has only child k) and $s(> k)$ be a sibling of $k-1$ ($s = \infty$ if $k-1$ has no sibling).
- (1) If $m < \infty$ and $m < s$ (Case a), then define T' by deleting vertex $k-1$ and k and their adjacent edges from T and adding new edge (m, j) where j is a parent of $k-1$ in T .




We get $\pi' = \varphi^{-1}(T'') \in \mathcal{A}_{n-2, i-2}$ where $T'' \in \mathcal{A}_{n-2, i-2}$ is induced from T' relabeling by $[n-2]$, where $i > k$. Relabeling π' by $\{1, \dots, k-2, k+1, \dots, n\}$, denoted by π'' , and define $\pi = (k, k-1, \pi'') \in \mathcal{A}_{n,k}$ where $\pi'_1 = i$.

- (2) If $m = \infty$ or $m > s$ (Case b1), then we get the tree $T' \in \mathcal{T}_{n, k-1}$ modifying as follows:



Define $\pi' = \varphi^{-1}(T') \in \mathcal{A}_{n, k-1}$ and $\pi = (k-1, k)\pi' \in \mathcal{A}_{n,k}$ (exchange $k-1$ and k in π').

- (B) If $k-1$ is not a parent of k in T (Case b2), then we get the tree $T' \in \mathcal{T}_{n,k}$ exchanging the labels $k-1$ and k in T .



Define $\pi' = \varphi^{-1}(T') \in \mathcal{A}_{n, k-1}$ and $\pi = (k-1, k)\pi' \in \mathcal{A}_{n,k}$ (exchange $k-1$ and k in π').

□

Remark. By considering a description of φ , given a elements $\pi \in \mathcal{A}_{n,k}$, π in Case (a) can be induced from $\cup_{i=k+1}^n \mathcal{A}_{n-2, i-2}$ and π in Case (b) from $\mathcal{A}_{n, k-1}$. It yields that two following sets are *isomorphic*

$$\mathcal{A}_{n,k} \simeq \mathcal{A}_{n, k-1} \cup (\cup_{i=k+1}^n \mathcal{A}_{n-2, i-2}),$$

for $2 \leq k \leq n$. We get the recurrence relation

$$a_{n,k} = a_{n,k-1} + \sum_{i=k+1}^n a_{n-2,i-2},$$

where $a_{n,k}$ is the cardinality of the set $\mathcal{A}_{n,k}$. Indeed, we are able to generalize this recurrence relation. Let $a_{n,k}(q, p) = \sum_{\sigma \in \mathcal{A}_{n,k}} q^{\text{inv}(\sigma)} p^{31-2(\sigma)}$. We have also

$$a_{n,k}(q, p) = q p a_{n,k-1}(q, p) + q^{2k-3} \sum_{i=k+1}^n a_{n-2,i-2}(q, p),$$

for $2 \leq k \leq n$.

Similarly, by considering a description of φ^{-1} , given an element $T \in \mathcal{T}_{n,k}$, T in Case (A1) can be induced from $\cup_{i=k+1}^n \mathcal{T}_{n-2,i-2}$ and T in Case (A2) and Case (B) from $\mathcal{T}_{n,k-1}$. It yields






$$\mathcal{T}_{n,k} \simeq \mathcal{T}_{n,k-1} \cup (\cup_{i=k+1}^n \mathcal{T}_{n-2,i-2}).$$

We get the recurrence relation

$$t_{n,k} = t_{n,k-1} + \sum_{i=k+1}^n t_{n-2,i-2},$$

where $t_{n,k}$ is the cardinality of the set $\mathcal{T}_{n,k}$.

Example. The following is the table of φ on \mathcal{A}_4 .

π	2143	3142	3241	4132	4231
π_1	2	3	3	4	4
$\text{inv}(\pi)$	2	3	3	4	4
$31-2(\pi)$	0	1	0	2	1
$\varphi(\pi)$					
$p(\varphi(\pi))$	2	3	3	4	4

REFERENCES

- [Don75] R. Donaghey, *Alternating permutations and binary increasing trees*, J. Combinatorial Theory Ser. A **18** (1975), 141–148.
- [Foa72] D. Foata, *Groupes de réarrangements et nombres d'Euler*, C. R. Acad. Sci. Paris Sér. A-B **275** (1972), A1147–A1150.
- [FS73] D. Foata and M.-P. Schützenberger, *Nombres d'Euler et permutations alternantes*, A survey of combinatorial theory (Proc. Internat. Sympos., Colorado State Univ., Fort Collins, Colo., 1971), North-Holland, Amsterdam, 1973, pp. 173–187.
- [FS74] D. Foata and V. Strehl, *Rearrangements of the symmetric group and enumerative properties of the tangent and secant numbers*, Math. Z. **137** (1974), 257–264.
- [KPP94] A. G. Kuznetsov, I. M. Pak, and A. E. Postnikov, *Increasing trees and alternating permutations*, Uspekhi Mat. Nauk **49** (1994), no. 6(300), 79–110.
- [Pou82] C. Poupard, *De nouvelles significations énumératives des nombres d'Entringer*, Discrete Math. **38** (1982), no. 2-3, 265–271.

(Heesung Shin) UNIVERSITÉ DE LYON; UNIVERSITÉ LYON 1; INSTITUT CAMILLE JORDAN; UMR 5208 DU CNRS; 43, BOULEVARD DU 11 NOVEMBRE 1918, F-69622 VILLEURBANNE CEDEX, FRANCE
E-mail address: hshin@math.univ-lyon1.fr