ADDENDUM TO: "AN EXPANSION OF THE JONES REPRESENTATION OF GENUS 2 AND THE TORELLI GROUP"

YASUSHI KASAHARA

ABSTRACT. We observe that the determinant of the representation provides a little restriction for the structure of the graded quotients $\mathcal{G}_k^{\mathbb{Q}}\mathcal{F}$ introduced in both [2] and [3] that any one of them does not contain the trivial 1-dimensional summand.

In our previous papers [2] and [3], we studied the Jones representation of genus 2

 $\rho: \mathcal{M}_2 \to \mathrm{GL}(5, \mathbb{Z}[t, t^{-1}])$

where \mathcal{M}_2 denotes the mapping class group of an closed oriented surface of genus 2. We considered certain expansions of the involved parameter starting with the substitutions $t = -e^h$ in the former paper [2], and $t = e^h$ in the latter one [3], which induce two *different* descending central filtrations, both denoted by the same symbol $\{\mathcal{F}_k\}_{k\geq 1}$, of the Torelli subgroup $\mathcal{I}_2 \subset \mathcal{M}_2$. In each case, the associated graded quotients $\mathcal{G}_k \mathcal{F} = \mathcal{F}_k / \mathcal{F}_{k+1}$ are free abelian groups of finite rank, and their scalar extensions $\mathcal{G}_k^{\mathbb{Q}} \mathcal{F} = \mathcal{G}_k \mathcal{F} \otimes \mathbb{Q}$ have natural structures of modules over $\operatorname{Sp}(4, \mathbb{Q})$ and the symmetric group \mathfrak{S}_6 of degree 6, respectively, and the direct sum $\bigoplus_{k\geq 1} \mathcal{G}_k^{\mathbb{Q}} \mathcal{F}$ forms a graded Lie algebra. We determined, in [2] and [3], respectively, the Lie subalgebra generated by the homogeneous component of degree 1. However, the structure of the graded quotients beyond this Lie subalgebra has remained unknown for the both cases.

In this note, we notify that the determinant of the representation ρ provides a little restriction for the structure of each graded quotient for the both cases that it does not contain the trivial 1-dimensional summand, which we had overlooked previously. While our argument is quite elementary, it is similar to that for the Morita trace for the images of the Johnson homomorphisms [4].

Date: April 19, 2010.

²⁰⁰⁰ Mathematics Subject Classification. Primary 57N05; Secondary 20F38, 20C08, 20F40.

Y. KASAHARA

From now on, we freely use the notations in our previous papers [2] and [3], which are common, as well as our argument below, except for the initial substitutions $t = -e^h$ and $t = e^h$, respectively. We begin our observation with:

Fact. The determinant of the Jones representation ρ lies in $\{\pm 1\}$.

This fact follows from the construction by Jones [1], and in fact holds for all the representations of the mapping class groups of punctured spheres constructed in [1] from rectangular Young diagrams. For the genus 2 case, however, one can verify the fact easily by using the explicit matrix form given in [1] or [2].

It is well-known that the abelianization of \mathcal{M}_2 is $\mathbb{Z}/10\mathbb{Z}$, and one can easily see that the image of the Torelli subgroup \mathcal{I}_2 in the abelianization is $\mathbb{Z}/5\mathbb{Z}$. Therefore, the above fact immediately implies that the determinant of ρ is identically 1 on \mathcal{I}_2 . Hence, for each $k \geq 1$, we have

$$\det \varphi^{(k)}(\mathcal{F}_k) \equiv 1 \mod h^{k+1}$$

On the other hand, for any $x \in \mathcal{F}_k$, $\varphi^{(k)}(x)$ has the form

$$\varphi^{(k)}(x) = I + h^k \cdot \Delta_k(x) \mod R[k+1]$$

where I denotes the identity matrix, and Δ_k is a homomorphism of \mathcal{F}_k , with kernel \mathcal{F}_{k+1} , into the additive group $M(5, \mathbb{Q})$ of the matrix algebra of degree 5 over \mathbb{Q} . Considering $M(5, \mathbb{Q})$ as an \mathcal{M}_2 -module by $\varphi^{(0)} \otimes$ $(\varphi^{(0)})^*$, Δ_k is also \mathcal{M}_2 -equivariant. We remark that the homomorphism $h^k \cdot \Delta_k$, with its target considered as $\mathcal{G}_k R = R[k]/R[k+1]$, is denoted by δ_k in our previous papers. Essential for this note is the following:

Lemma. For any $k \geq 1$ and $x \in \mathcal{F}_k$,

$$\det \varphi^{(k)}(x) = 1 + h^k \cdot \operatorname{trace} \Delta_k(x) \mod h^{k+1}$$

where trace denotes the usual matrix trace.

Proof. Writing $I + h^k \cdot \Delta_k(x) = (a_{i,j})$, consider the definition for determinant

$$\det \varphi^{(k)}(x) = \sum_{\sigma \in \mathfrak{S}_5} \operatorname{sgn} \sigma \prod_{i=1}^5 a_{i,\sigma(i)}.$$

It is easy to see that each summand except for $\sigma = 1$ has at least two factors from the off diagonal, and hence is a multiple of h^{2k} , which vanishes modulo h^{k+1} . Therefore, det $\varphi^{(k)}(x)$ is simply the product of the diagonal entries, which is equal to the right-hand side in the lemma.

Therefore, we see that $\Delta_k(\mathcal{F}_k)$ satisfies trace $\equiv 0$. For each of the cases [2] and [3], the \mathcal{M}_2 -module $M(5, \mathbb{Q})$, which is identical to $\mathcal{G}_k R$, is in fact an Sp(4, \mathbb{Q})-module and an \mathfrak{S}_6 -module, respectively, and its irreducible decomposition is explicitly given by

$$[2, (4.1)] \qquad \qquad \Gamma_{0,2} \oplus \Gamma_{2,0} \oplus \Gamma_{0,0}$$

and

$$[3, (3.4)] \qquad \qquad [6] \oplus [4, 2] \oplus [2^3] \oplus [3, 1^3]$$

respectively.

Among the summands in each case, the trivial 1-dimensional one, which corresponds to $\Gamma_{0,0}$ and [6], respectively, can be identified with the center of $M(5, \mathbb{Q})$, and therefore has *nonzero* trace. This shows:

Proposition. The trivial 1-dimensional summand does not appear in $\mathcal{G}_k^{\mathbb{Q}}\mathcal{F}$ for any $k \geq 1$.

We remark that all the other summands of $M(5, \mathbb{Q})$ are all trace free since we already know by [2, Theorem 1.3] and [3, Theorem 2.2], respectively, that any non-trivial summand of $M(5, \mathbb{Q})$ appears in some homogeneous component of the Lie subalgebra generated by $\mathcal{G}_1^{\mathbb{Q}}\mathcal{F}$.

Acknowledgement. The author is grateful to Professor Shigeyuki Morita for asking about the determinant of the Jones representation for over ten years intermittently.

References

- V. F. R. Jones, Hecke algebra representations of braid groups and link polynomials, Ann. of Math. 126 (1987), 335–388.
- Y. Kasahara, An expansion of the Jones representation of genus 2 and the Torelli group, Algebr. Geom. Topol. 1 (2001), 39–55 (electronic).
- 3. _____, An expansion of the Jones representation of genus 2 and the Torelli group. II, J. Knot Theory Ramifications 13 (2004), no. 2, 297–306.
- S. Morita, Abelian quotients of subgroups of the mapping class group of surfaces, Duke Math. J. 70 (1993), no. 3, 699–726.

DEPARTMENT OF MATHEMATICS, KOCHI UNIVERSITY OF TECHNOLOGY, TOSAYA-MADA, KAMI CITY, KOCHI, 782-8502 JAPAN

E-mail address: kasahara.yasushi@kochi-tech.ac.jp