Limit sets and commensurability of Kleinian groups

WEN-YUAN YANG[™] and YUE-PING JIANG

Abstract

In this paper, we obtain several results on the commensurability of two Kleinian groups and their limit sets. We prove that two finitely generated subgroups G_1 and G_2 of an infinite co-volume Kleinian group $G \subset \text{Isom}(\mathbf{H}^3)$ having $\Lambda(G_1) = \Lambda(G_2)$ are commensurable. In particular, it is proved that any finitely generated subgroup H of a Kleinian group $G \subset \text{Isom}(\mathbf{H}^3)$ with $\Lambda(H) = \Lambda(G)$ is of finite index if and only if H is not a virtually fiber subgroup.

2000 Mathematics subject classification. primary 30F40. Keywords and phrases: Commensurability, Geometrically finite, Limit set, Hyperbolic element.

1. Introduction

Two groups G_1 and G_2 are commensurable if their intersection $G_1 \cap G_2$ is of finite index in both G_1 and G_2 . In this paper, we investigate the following question asked by J. Anderson [5]: namely, if $G_1, G_2 \subset \text{Isom}(\mathbf{H}^n)$ are finitely generated and discrete, does $Ax(G_1) = Ax(G_2)$ imply that G_1 and G_2 are commensurable? Here we use Ax(G)to denote the set of axes of the hyperbolic elements of $G \subset \text{Isom}(\mathbf{H}^n)$.

The question has been discussed by several authors. In 1990, G. Mess [11] showed that if G_1 and G_2 are non-elementary finitely generated Fuchsian groups having the same nonempty set of simple axes, then G_1 and G_2 are commensurable. Using some technical results on arithmetic Kleinian groups, D. Long and A. Reid [9] gave an affirmative answer to this question in the case where G_1 and G_2 are arithmetic Kleinian groups. Note that all the confirmed cases for the question are geometrically finite groups. So it is natural to ask if the question is true with the assumption that G_1 and G_2 are geometrically finite. Recently, P. Susskind [13] constructed two geometrically finite Kleinian groups in Isom(\mathbf{H}^n) (for $n \ge 4$) having the same action on some invariant 2-hyperbolic plane but whose intersection is infinitely generated. So this implies that these two geometrically finite groups are not commensurable although they have the same axes set. But it is worth pointing that these two geometrically finite Kleinian groups generate a non-discrete group. This example suggests that some additional conditions need to be imposed to eliminate such 'bad' groups.

The first author is supported by the China-funded Postgraduates Studying Aboard Program for Building Top University. The second author is supported by National Natural Science Foundation of China (No. 10671059) and Doctorate Foundation of the Ministry of Education of China (No. 20060532023).

In higher dimensions, we have the following consequence of P. Susskind and G. Swarup's results [14] on the limit set of the intersection of two geometrically finite Kleinian groups.

PROPOSITION 1.1. Let G_1 and G_2 be two non-elementary geometrically finite subgroups of a Kleinian group $G \subset \text{Isom}(\mathbf{H}^n)$. Then G_1 and G_2 are commensurable if and only if the limit sets $\Lambda(G_1)$ and $\Lambda(G_2)$ are equal. In particular, $\Lambda(G_1) = \Lambda(G_2)$ if and only if $\operatorname{Ax}(G_1) = \operatorname{Ax}(G_2) = \operatorname{Ax}(G_1 \cap G_2)$.

In several cases, the condition that both subgroups lie in a larger discrete group can be dropped.

COROLLARY 1.2. Let $G_1, G_2 \subset \text{Isom}(\mathbf{H}^n)$ be two non-elementary geometrically finite Kleinian groups of the second kind leaving no m hyperbolic planes invariant for m < n - 1. Then G_1 and G_2 are commensurable if and only if $\Lambda(G_1) = \Lambda(G_2)$.

In dimension 3, we can refine the analysis of the limit sets using Anderson's results [3] to get the following result in the essence of the recent solution of the Tameness Conjecture (see [1] and [6]) which states that all finitely generated Kleinian groups in $Isom(\mathbf{H}^3)$ are topologically tame.

THEOREM 1.3. Let G_1 , G_2 be two non-elementary finitely generated subgroups of an infinite co-volume Kleinian group $G \subset \text{Isom}(\mathbf{H}^3)$. Then G_1 and G_2 are commensurable if and only if $\Lambda(G_1) = \Lambda(G_2)$. In particular, $\Lambda(G_1) = \Lambda(G_2)$ if and only if $\operatorname{Ax}(G_1) = \operatorname{Ax}(G_2) = \operatorname{Ax}(G_1 \cap G_2)$

Similarly, in the following case we are able to remove the ambient discrete group.

COROLLARY 1.4. Let $G_1, G_2 \subset \text{Isom}(\mathbf{H}^3)$ be two non-elementary finitely generated Kleinian groups of the second kind whose limit sets are not circles. Then G_1 and G_2 are commensurable if and only if $\Lambda(G_1) = \Lambda(G_2)$.

In fact, under the hypotheses of the above results, the condition of having the same limit sets of two Kleinian subgroups exactly implies having the same axes sets. But Anderson's original formulation of the question is only to suppose that two Kleinian groups have the same axes set. So it is interesting to explore whether there is some essential difference between the limit set and axes set. The following theorem is a result in this direction, suggesting that the 'same axes set' condition is necessary in general for Anderson's question.

THEOREM 1.5. Let H be a non-elementary finitely generated subgroup of a Kleinian group $G \subset \text{Isom}(\mathbf{H}^3)$. Suppose that $\Lambda(H) = \Lambda(G)$. Then [G : H] is finite if and only if G is not virtually fibered over H. In particular, [G : H] is finite if and only if Ax(H) = Ax(G).

Remark 1.6. We remark that the case of H being geometrically finite is proved by P. Susskind and G. Swarup [14, Theorem 1]. Theorem 1.5 actually proves the special case of Anderson's question where G_1 is a subgroup of G_2 .

The paper is organized as follows. In section 2, we collect some results on the limit set of the intersection of two Kleinian groups and prove some useful lemmas for later use. In section 3, we prove Theorems 1.3 and 1.5.

2. Preliminaries

Let \mathbf{B}^n denote the closed ball $\mathbf{H}^n \cup \mathbf{S}^{n-1}$, whose boundary \mathbf{S}^{n-1} is identified via stereographic projection with $\overline{\mathbf{R}}^{n-1} = \mathbf{R}^{n-1} \cup \infty$. Let $\operatorname{Isom}(\mathbf{H}^n)$ be the full group of isometries of \mathbf{H}^n and let $G \subset \operatorname{Isom}(\mathbf{H}^n)$ be a *Kleinian* group; that is, a discrete subgroup of $\operatorname{Isom}(\mathbf{H}^n)$. Then G acts discontinuously on \mathbf{H}^n if and only if G is discrete. Furthermore, G acts on \mathbf{S}^{n-1} as a group of conformal homeomorphisms. The *set of discontinuity* $\Omega(G)$ of G is the subset of \mathbf{S}^{n-1} on which G acts discontinuously; the *limit set* $\Lambda(G)$ is the complement of $\Omega(G)$ in \mathbf{S}^{n-1} . A Kleinian group is said to be of the *second kind* if $\Omega(G)$ is nonempty otherwise it is said to be of the *first kind*.

The elements of $\text{Isom}(\mathbf{H}^n)$ are classified in terms of their fixed point sets. An element $g \neq \text{id}$ in $\text{Isom}(\mathbf{H}^n)$ is *elliptic* if it has a fixed point in \mathbf{H}^n , *parabolic* if it has exactly one fixed point which lies in \mathbf{S}^{n-1} , *hyperbolic* if it has exactly two fixed points which lie in \mathbf{S}^{n-1} . The unique geodesic joining the two fixed points of the hyperbolic element g, which is invariant under g, is called the *axis* of the hyperbolic element and is denoted by ax(g). The limit set $\Lambda(G)$ is the closure of the set of fixed points of hyperbolic and parabolic elements of G. A Kleinian group whose limit set contains fewer than three points is called *elementary* and is otherwise called *non-elementary*.

For a non-elementary Kleinian group *G*, define $\overline{C}(G)$ to be the smallest nonempty convex set in \mathbf{H}^n which is invariant under the action of *G*; this is the *convex hull* of $\Lambda(G)$. The quotient $C(G) = \widetilde{C}(G)/G$ is the *convex core* of $M = \mathbf{H}^n/G$. The group *G* is *geometrically finite* if the convex core C(G) has finite volume.

By Margulis's lemma, it is known that there is a positive constant ϵ_0 such that for any Kleinian group $G \in \text{Isom}(\mathbf{H}^n)$ and $\epsilon < \epsilon_0$, the part of \mathbf{H}^n/G where the injectivity radius is less than ϵ is a disjoint union of tubular neighbourhoods of closed geodesics, whose lengths are less than 2ϵ , and cusp neighbourhoods. In dimensions 2 and 3, these cusp neighbourhoods can be taken to be disjoint quotients of horoballs by the corresponding parabolic subgroup. This set of disjoint horoballs is called a *precisely invariant system of horoballs* for *G*. In dimension 3, it is often helpful to identify the infinity boundary \mathbf{S}^2 of \mathbf{H}^3 with the extended complex plane $\overline{\mathbf{C}}$. In particular, the fixed point of a rank 1 parabolic subgroup *J* of *G* is called *doubly cusped* if there are two disjoint circular discs $B_1, B_2 \subset \overline{\mathbf{C}}$ such that $B_1 \cup B_2$ is precisely invariant under *J* in *G*. In this case, the parabolic elements of *J* are also called doubly cusped.

In dimension 3, we call a Kleinian group *G* topologically tame if the manifold $M = \mathbf{H}^3/G$ is homeomorphic to the interior of a compact 3-manifold. Denote by M^c the complement of these cusp neighbourhoods. Using the relative core theorem of McCullough [10], there exists a compact submanifold *N* of M^c such that the inclusion of *N* in M^c is a homotopy equivalence, every torus component of $\partial(M^c)$ lies in *N*, and *N* meets each annular component of $\partial(M^c)$ in an annulus. Call such an

N a relative compact core for *M*. The components of $\partial(N) - \partial(M^c)$ are the relative boundary components of *N*. The ends of M^c are in one-to-one correspondence with the components of $M^c - N$. An end *E* of M^c is geometrically finite if it has a neighbourhood disjoint from *C*(*G*). Otherwise, *E* is geometrically infinite.

A Kleinian group $G \subset \text{Isom}(\mathbf{H}^3)$ is *virtually fibered* over a subgroup H if there are finite index subgroups G^0 of G and H^0 of H such that \mathbf{H}^3/G^0 has finite volume and fibers over the circle with the fiber subgroup H^0 . Note that H^0 is then a normal subgroup of G^0 , and so $\Lambda(H^0) = \Lambda(G^0) = \mathbf{S}^2$.

In order to analyze the geometry of a geometrically infinite Kleinian group, we will use Canary's covering theorem, which generalizes a theorem of Thurston [15]. Note that the Tameness Theorem ([1] and [6]) states that all finitely generated Kleinian groups in Isom(\mathbf{H}^3) are topologically tame.

THEOREM 2.1 ([7, The Covering Theorem]). Let G be a torsion free Kleinian group in Isom(\mathbf{H}^3) and let H be a non-elementary finitely generated subgroup of G. Let $N = \mathbf{H}^3/G$, let $M = \mathbf{H}^3/H$, and let $p : M \to N$ be the covering map. If M has a geometrically infinite end E, then either G is virtually fibered over H or E has a neighbourhood U such that p is finite-to-one on U.

Now we list several results on the limit set of the intersection of two Kleinian groups, which describe $\Lambda(G_1 \cap G_2)$ in terms of $\Lambda(G_1)$ and $\Lambda(G_2)$, where G_1 and G_2 are subgroups of a Kleinian group G. Here we only collect the results used in this paper and state them in an appropriate form for our purpose. See [4] for a useful survey and the bibliography therein for the results in full details.

THEOREM 2.2 ([14, Theorem 3]). Let G_1 , G_2 be two geometrically finite subgroups of Kleinian group $G \subset \text{Isom}(\mathbf{H}^n)$. Then $\Lambda(G_1) \cap \Lambda(G_2) = \Lambda(G_1 \cap G_2) \cup P$ where P consists of some parabolic fixed points of G_1 and G_2 .

PROPOSITION 2.3 ([14, Corollary 1]). Let *H* be geometrically finite and *j* be a hyperbolic element with a fixed point in $\Lambda(H)$. If $\langle H, j \rangle$ is discrete, then $j^n \in H$ for some n > 0.

Based on the above results, we get the following lemma characterizing the relationship between limit sets and axes sets.

LEMMA 2.4. Let G_1 , G_2 be two geometrically finite subgroups of Kleinian group $G \subset \text{Isom}(\mathbf{H}^n)$. Then $\Lambda(G_1) = \Lambda(G_2)$ if and only if $\text{Ax}(G_1) = \text{Ax}(G_2) = \text{Ax}(G_1 \cap G_2)$

PROOF. If G_1 and G_2 are geometrically finite subgroups of a Kleinian group G, then $G_1 \cap G_2$ is again geometrically finite ([14, Theorem 4]). It is well known that a hyperbolic element cannot share one fixed point with a parabolic element in a discrete group. Thus by applying Theorem 2.2 to $\Lambda(G_1) \cap \Lambda(G_2)$, we can conclude that any hyperbolic element $h \in G_i$ has at least one fixed point in $\Lambda(G_1 \cap G_2)$ for i = 1, 2. Now by Proposition 2.3, we have $h^j \in G_1 \cap G_2$ for some large integer j > 0. This implies the axis ax(h) of h belongs to $Ax(G_1 \cap G_2)$. Therefore we have $Ax(G_1) = Ax(G_2) = Ax(G_1 \cap G_2)$ and it also follows that P is the empty set.

The other direction is easy to see using the fact that the set of fixed points of hyperbolic elements of *G* is dense in $\Lambda(G)$.

In dimension 3, J. Anderson [3] carried out a more careful analysis on the limit set of the intersection of two topologically tame Kleinian groups. Combined with the recent solution of the Tameness Conjecture ([1] and [6]), we have the following theorem.

THEOREM 2.5 ([3, Theorem C]). Let $G \subset \text{Isom}(\mathbf{H}^3)$ be a Kleinian group, and let G_1 and G_2 be non-elementary finitely generated subgroups of G, then $\Lambda(G_1) \cap \Lambda(G_2) = \Lambda(G_1 \cap G_2) \cup P$ where P is empty or consists of some parabolic fixed points of G_1 and G_2 .

PROPOSITION 2.6 ([3, Theorem A]). Let *H* be finitely generated Kleinian group and *j* be a hyperbolic element with a fixed point in $\Lambda(H)$. If $\langle H, j \rangle$ is discrete, then either $\langle H, j \rangle$ is virtually fibered over *H* or $j^n \in H$ for some n > 0.

Similarly, we obtain the following lemma.

LEMMA 2.7. Let G_1 and G_2 be two non-elementary finitely generated subgroups of an infinite co-volume Kleinian group $G \subset \text{Isom}(\mathbf{H}^3)$. Then $\Lambda(G_1) = \Lambda(G_2)$ if and only if $\text{Ax}(G_1) = \text{Ax}(G_2) = \text{Ax}(G_1 \cap G_2)$

PROOF. Observe that our hypothesis '*G* is an infinite co-volume Kleinian group' implies that *G* is not virtually fibered over G_1 . Since the intersection of any pair of finitely generated subgroups of a Kleinian group is finitely generated (see [2]), we see that $G_1 \cap G_2$ is finitely generated. Using Theorem 2.5 and Proposition 2.6, we can argue exactly as in Lemma 2.4 to obtain $Ax(G_1) = Ax(G_2) = Ax(G_1 \cap G_2)$, if we suppose $\Lambda(G_1) = \Lambda(G_2)$.

REMARK 2.8. The condition of G being infinite co-volume cannot be dropped, as will be seen in the proof of Theorem 1.5. Namely, the Kleinian group G fibering over H has a different axes set from that of its fiber subgroup H.

In the following two lemmas, we give some useful properties about the same axes sets of two Kleinian groups.

LEMMA 2.9. Let G be a non-elementary finitely generated Kleinian group and H a subgroup of finite index in G. Then Ax(G) = Ax(H).

PROOF. It is obvious that $Ax(H) \subset Ax(G)$. Conversely, since [G : H] is finite, for any hyperbolic element g with axis $ax(g) \in Ax(G)$, there are two integers i and j such that $g^i H = g^j H$ and thus $g^{i-j} \in H$. It follows that $ax(g) \in Ax(H)$. The proof is complete.

REMARK 2.10. In fact, our Theorem 1.5 proves that the converse of the above Lemma is also true when *H* is a finitely generated subgroup of $G \subset \text{Isom}(\mathbf{H}^3)$.

LEMMA 2.11. Let G be a non-elementary finitely generated, torsion free Kleinian group and H be a subgroup of G. Suppose that Ax(G) = Ax(H). Then for every hyperbolic element $g \in G$, $g^n \in H$ for some n > 0.

PROOF. For any hyperbolic element $g \in G$, we can choose a hyperbolic element h from H such that ax(g) = ax(h) by the hypothesis Ax(G) = Ax(H). It follows that the subgroup $\langle g, h \rangle$ is elementary and torsion free. By the characterization of elementary Kleinian groups it follows that $\langle g, h \rangle$ is actually a cyclic subgroup $\langle f \rangle$ of G. Thus we can write $g = f^m$ and $h = f^n$ for two appropriate integers m, n. Now we have found the integer n such that $g^n = h^m \in H$, which proves the lemma.

3. Proofs

PROOF OF PROPOSITION 1.1. Recall G_1 and G_2 are commensurable if the intersection $G_1 \cap G_2$ is of finite index in both G_1 and G_2 . By Lemma 2.9, we have $Ax(G_1) = Ax(G_2)$ and thus $\Lambda(G_1) = \Lambda(G_2)$, if G_1 and G_2 are commensurable. So it remains to prove the converse.

If $\Lambda(G_1) = \Lambda(G_2)$, we have $\operatorname{Ax}(G_1) = \operatorname{Ax}(G_2) = \operatorname{Ax}(G_1 \cap G_2)$ by Lemma 2.4. Therefore it follows that $\Lambda(G_1) = \Lambda(G_2) = \Lambda(G_1 \cap G_2)$, since the set of fixed points of hyperbolic elements of *G* is dense in $\Lambda(G)$. Now we can conclude that $G_1 \cap G_2$ is of finite index in both G_1 and G_2 , by using Theorem 1 in [14] which states that any geometrically finite subgroup *H* of a Kleinian group *G* is of finite index in *G* if $\Lambda(H) = \Lambda(G)$.

The second assertion is just Lemma 2.4. This completes the proof.

PROOF OF COROLLARY 1.2. It is well known that the stabilizer in Isom(\mathbf{H}^n) of the limit set of a non-elementary Kleinian group *G* of the second kind leaving no *m* hyperbolic planes invariant for m < n - 1, is itself a Kleinian group. See for example Greenberg [8], where the discreteness of the stabilizer of that limit set is proved.

Thus $Ax(G_1) = Ax(G_2)$ implies that G_1 and G_2 together lie in a common Kleinian group, which is the stabilizer of the common limit set of G_1 and G_2 . Thus Proposition 1.1 completes the proof.

PROOF OF THEOREM 1.3. Observe that the fundamental domain of the subgroup *H* is the union of translates of the fundamental domain of *G* by left *H*-coset representatives in *G*. So the subgroup $\langle G_1, G_2 \rangle$ also has infinite co-volume, and we can assume that *G* is finitely generated by replacing *G* by $\langle G_1, G_2 \rangle$. As the conclusion is easily seen to be unaffected by passage to a finite index subgroup, we may use Selberg's Lemma to pass to a finite index, torsion free subgroup of *G*. Hence, without loss of generality, we may assume that *G* is finitely generated and torsion free.

By Proposition 1.1, the conclusion is trivial if G is geometrically finite. So we suppose that G is geometrically infinite. Then $M = \mathbf{H}^3/G$ has infinite volume. Let C be a compact core for M. Since M has infinite volume, ∂C contains a surface of

genus at least two. Then using Thurston's geometrization theorem for Haken threemanifolds (see [12]), there exists a geometrically finite Kleinian group with non-empty discontinuity domain, which is isomorphic to G.

Now our task is to give an algebraic characterization of the limit set of $G_1 \cap G_2$ in G_i such that the relationships between $\Lambda(G_1 \cap G_2)$ and $\Lambda(G_i)$ can be passed to the ones between target isomorphic groups under the above isomorphism of G. Then the conclusion of Theorem 1.3 follows from Proposition 1.1. We claim that for every element $g \in G_1$, there exists an integer k such that $g^k \in G_1 \cap G_2$.

Firstly, by Lemma 2.7, we obtain that $Ax(G_1) = Ax(G_2) = Ax(G_1 \cap G_2)$. So for any hyperbolic element $g \in G_1$, the integer k obtained in Lemma 2.11 is such that $g^k \in G_1 \cap G_2$. Now we consider the remaining parabolic elements. Theorem B of [3] says that if no nontrivial power of a parabolic element $h \in G_1$ lies in $G_1 \cap G_2$, then there exists a doubly cusped parabolic element $f \in G_1 \cap G_2$ with the same fixed point ξ as h. Normalizing their fixed point ξ to ∞ , we can suppose that f(z) = z + 1and $h(z) = z + \tau$, where $\text{Im}(\tau) \neq 0$. Since f is doubly cusped in $G_1 \cap G_2$, then $\Lambda(G_1 \cap G_2) \subset \{z : |\text{Im}(z)| < c\}$, for some constant c. But on the other hand, $\Lambda(G_1 \cap G_2)$ is also kept invariant under h, which contradicts the fact that $\Lambda(G_1 \cap G_2)$ is invariant under f. Therefore the claim is proved for all elements including parabolic elements. A similar claim holds for $G_1 \cap G_2$ and G_2 .

Under the isomorphism, using the above claims, we can conclude that the limit set of the (isomorphic) image of $G_1 \cap G_2$ is equal to those of the (isomorphic) images of G_1 and G_2 . The proof is complete as a consequence of Proposition 1.1.

REMARK 3.1. Theorem 1.3 can be thought as a geometric version of Lemma 5.4 in [3], which uses an algebraic assumption on the limit sets of the groups involved.

PROOF OF COROLLARY 1.4. This is proved similarly to Corollary 1.2.

PROOF OF THEOREM 1.5. In view of Lemma 2.9, we may assume, without loss of generality, that H is finitely generated and torsion free by using Selberg's Lemma to pass to a finite index, torsion free subgroup of H.

If *H* is geometrically finite, then the conclusion follows from a result of P. Susskind and G. Swarup [14], which states that a non-elementary geometrically finite subgroup sharing the same limit set with the ambient discrete group is of finite index. So next we suppose that *H* is geometrically infinite. Then there exist finitely many geometrically infinite ends E_i for the manifold $N := \mathbf{H}^3/H$.

We first claim that Ax(H) = Ax(G) implies that H cannot be a virtually fiber subgroup of G. Otherwise, by taking finite index subgroups of G and H, we can suppose H is normal in G. Then it follows that every element of the quotient group G/H has finite order by Lemma 2.11. Thus G/H could not be isomorphic to \mathbb{Z} . This is a contradiction. So H is not a virtually fiber subgroup of G.

Using the Covering Theorem 2.1, we know that for each geometrically infinite end E_i , there exists a neighbourhood U_i of E_i such that the covering map $\mathcal{P}: N \to M := \mathbf{H}^3/G$ is finite to one on U_i .

Now we argue by way of contradiction. Let $Q_N : \mathbf{H}^3 \to N$ and let $Q_M : \mathbf{H}^3 \to M$ be the covering maps and notice that $Q_M = \mathcal{P} \circ Q_N$. Suppose that [G : H] is infinite. This implies that \mathcal{P} is an infinite covering map. By the definition of a geometrically infinite end, we can take a point *z* from the neighbourhood U_1 of a geometrically infinite end E_1 such that *z* also lies in the convex core of *N*. By lifting the point $\mathcal{P}(z) \in M$ to \mathbf{H}^3 , it is easy to see that the infinite set $\widetilde{S} := Q_M^{-1}(\mathcal{P}(z))$ lies in the common convex hull $\widetilde{C}(H) = \widetilde{C}(G) \subset \mathbf{H}^3$, by observing that $\widetilde{C}(G)$ is invariant under *G* and the preimage $Q_N^{-1}(z) \subset \widetilde{S}$ lies in $\widetilde{C}(H)$. Since \mathcal{P} is an infinite covering map, the set $S := \mathcal{P}^{-1}(\mathcal{P}(z))$ is infinite. By considering $Q_M = \mathcal{P} \circ Q_N$, it follows that $S = Q_N(\widetilde{S}) \subset N$ and thus *S* lies in the convex core of *N*, because $\widetilde{S} \subset \widetilde{C}(H)$.

We claim that we can take a smaller invariant horoball system for H such that infinitely many points of S lie outside all cusp ends of N. Otherwise, we can suppose that infinitely many points of S are contained inside a cusp end E_c of N, since there are only finitely many cusp ends for N. Thus infinitely many points of $Q_N^{-1}(S)$ lie in the corresponding horoball B for the end E_c . Normalizing the parabolic fixed point for E_c to ∞ in the upper half space model of \mathbf{H}^3 , the horoball B at ∞ is precisely invariant under the stabilizer of ∞ in H. On the other hand, we have that infinitely many points of $Q_N^{-1}(S)$ have the same height, since the covering map \mathcal{P} maps $S \subset N$ to a single point on M, and the horoball B is also precisely invariant under the stabilizer of ∞ in G, which is a Euclidean group preserving the height of points in the horoball B. Then we can take a smaller horoball for E_c such that these infinitely many points of $Q_N^{-1}(S)$ lie outside the horoball.

Continuing the above process for all cusp ends of N, we can get a new invariant horoball system such that infinitely points of S lie outside these cusp ends of N.

Since *S* projects to a single point on *M*, we can conclude that *S* cannot lie in any compact subset of the convex core of *N*. Thus by the above second claim, there exist infinitely many points of *S* which can only lie in geometrically infinite ends of *N*. This is a contradiction to the Covering Theorem 2.1, which states that the covering map \mathcal{P} restricted to each geometrically infinite end is finite-to-one, if *H* is not a virtually fiber subgroup of *G*.

Acknowledgements

The first author would like to sincerely thank Prof. Jim Anderson for helpful communications and Prof. Leonid Potyagailo for many helpful comments and discussions about the original version. The authors would like to express their gratitude to the referee for his/her many helpful comments, which substantially improved the exposition of the paper.

References

- [1] I. Agol, Tameness of hyperbolic 3-manifolds, preprint.
- [2] J. Anderson, On the finitely generated intersection property for Kleinian groups, Complex Variables Theory Appl. 17 (1991), 111–112.

- [3] J. Anderson, Intersections of topologically tame subgroups of Kleinian groups, J. Anal. Math. 65 (1995), 77–94.
- [4] J. Anderson, The limit set intersection theorem for finitely generated Kleinian groups, Math. Res. Lett. 3 (1996), 675–692.
- [5] M. Bestvina, Questions in geometric group Theory, http://www.math.utah.edu/~bestvina.
- [6] D. Calegari and D. Gabai, Shrinkwrapping and the taming of hyperbolic 3-manifolds, J. Amer. Math. Soc. 19 (2006), 385–446.
- [7] R. Canary, A covering theorem for hyperbolic 3-manifolds and its applications, Topology, 35(1996), 751–778.
- [8] L. Greenberg, Commensurable groups of Moebius transformations, in: Discontinuous groups and Riemann surfaces, ed. by L. Greenberg, Annals of Mathematics Studies 79, Princeton University Press, Princeton, 1974, 227–237.
- [9] D. Long and A. Reid, On Fuchsian groups with the same set of axes, Bull. London Math. Soc. 30 (1998), 533–538.
- [10] D. McCullough, Compact submanifolds of 3-manifolds with boundary, Quart. J. Math. Oxford 37 (1986), 299–307.
- [11] G. Mess, Fuchsian groups with the same simple axes, Preprint (1990).
- [12] J. Morgan, On Thurston's uniformization theorem for 3-dimensional manifolds, in The Smith Conjecture, ed. by H. Bass and J. Morgan, Academic Press, 1984, pp. 37–125.
- [13] P. Susskind, An infinitely generated intersection of the hyperbolic group, Proc. Amer. Math. Soc. 129 (2001), 2643–2646.
- [14] P. Susskind and G. Swarup, Limit sets of geometrically finite hyperbolic groups, Amer. J. Math. 114 (1992), 233–250.
- [15] W. Thurston, The geometry and topology of 3-manifolds, Lecture Notes, Princeton Mathematics Department.

Wen-yuan Yang, College of Mathematics and Econometrics, Hunan University, Changsha, Hunan 410082 People's Republic of China

U.F.R. de Mathematiques, Universite de Lille 1,59655 Villneuve D'Ascq Cedex, France

e-mail: wyang@math.univ-lille1.fr

Yue-ping Jiang, College of Mathematics and Econometrics, Hunan University, Changsha, Hunan 410082 People's Republic of China e-mail: ypjiang731@163.com