#### Examples of rigid and flexible Seifert fibred cone-manifolds

Alexander Kolpakov\*

#### Abstract

The present paper gives an example of a rigid spherical cone-manifold and that of a flexible one which are both Seifert fibred.

**Keywords** cone-manifold, rigidity, flexibility, Seifert fibration **MSC (2010)** 53A35, 57R18, 57M25

#### 1 Introduction

The theory of three-dimensional orbifolds and cone-manifolds attracts attention of many mathematicians since the original work of Thurston [29]. An introduction to the theory of orbifolds could be found in [29, Ch.13]. For a basic introduction to the geometry of three-dimensional cone-manifolds and cone-surfaces we refer the reader to [6]. The main motivation for studying three-dimensional cone-manifolds comes from Thurston's approach to geometrization of three-orbifolds: three-dimensional cone-manifolds provide a way to deform geometric orbifold structures. The orbifold theorem has been proven in full generality by M. Boileau, B. Leeb and J. Porti, see [1, 2]. One of the main questions in the theory of three-dimensional cone-manifolds is the rigidity problem. First, the rigidity property was discovered for hyperbolic manifolds (so-called Mostow-Prasad rigidity, see [19, 24]). After that, the global rigidity property for hyperbolic three-dimensional cone-manifolds with singular locus a link and cone angles less than  $\pi$  was proven by S. Kojima [16]. The key result that implies global rigidity is due to Hodgson and Kerckhoff [13], who showed the local rigidity of hyperbolic cone manifolds with singularity of link or knot type and cone angles less than  $2\pi$ . The de Rham rigidity for spherical orbifolds was established in [26, 27]. Detailed analysis of the rigidity property for three-dimensional cone-manifolds

<sup>\*</sup>Supported by the Schweizerischer National fonds SNF no. 200020-121506/1 and no. 200021-131967/1

was carried out in [31, 32] for hyperbolic and spherical cone-manifolds with singularity a trivalent graph and cone angles less than  $\pi$ .

Recently, the local rigidity for hyperbolic cone-manifolds with cone angles less than  $2\pi$  was proven in [18, 33]. However, examples of infinitesimally flexible hyperbolic cone-manifolds had already been given in [5]. For other examples of flexible cone-manifolds one may refer to [15, 21, 28].

The theorem of [32] concerning the global rigidity for spherical threedimensional cone-manifolds was proven under the condition of being not Seifert fibred. Recall that due to [22] a cone-manifold is Seifert fibred if its underlying space carries a Seifert fibration such that components of the singular stratum are leafs of the fibration. In particular, if its singular stratum is represented by a link, then the complement is a Seifert fibred three-manifold. All Seifert fibred link complements in the three-sphere are described by [4]. In the present paper, we give an explicit example of a rigid spherical cone-manifold and a flexible one which are both Seifert fibred. The singular locus for each of these cone-manifolds is a link and the underlying space is the three-sphere  $\mathbb{S}^3$ . The rigid cone-manifold given in the paper has cone-angles of both kinds, less or greater than  $\pi$ . The flexible one has cone-angles strictly greater than  $\pi$ . Deformation of its geometric structure comes essentially from those of the base cone-surface. However, hyperbolic orbifolds, which are Seifert fibred over a disc, are rigid. Their geometric structure degenerates to the minimal-perimeter hyperbolic polygon, as shown in [23]. These are uniquely determined by cone angles.

The paper is organised as follows: first, we recall some common facts concerning spherical geometry. In the second section, the geometry of the Hopf fibration is considered and a number of lemmas are proven. After that, we construct two explicit examples of Seifert fibred cone-manifolds. The first one is a globally rigid cone-manifold and its moduli space is parametrised by its cone angles only. The second one is a flexible Seifert fibred cone-manifold. This means that we can deform its metric while keeping its cone angles fixed. Rigorously speaking, the following assertion is proven: the given cone-manifold has a one-parameter family of distinct spherical cone metrics with the same cone angles.

**Acknowledgement**. The author is grateful to Prof. J. Porti (Universitat Autònoma de Barcelona) and Prof. J.-M. Schlenker (Institut de Mathématiques de Toulouse) for their valuable comments on the paper and discussion of the subject.

## 2 Spherical geometry

Below we present several common facts concerning spherical geometry in dimension two and three.

Let us identify a point p = (w, x, y, z) of the three-dimensional sphere

$$\mathbb{S}^3 = \{(w, x, y, z) \in \mathbb{R}^4 | w^2 + x^2 + y^2 + z^2 = 1\}$$

with an  $SU_2(\mathbb{C})$  matrix of the form

$$P = \left( \begin{array}{cc} w + ix & y + iz \\ -y + iz & w - ix \end{array} \right).$$

Then, replace the group Isom<sup>+</sup>  $\mathbb{S}^3 \cong SO_4(\mathbb{R})$  of orientation preserving isometries with its two-fold covering  $SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$ . Finally, define the action of  $\langle A, B \rangle \in SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$  on  $P \in SU_2(\mathbb{C})$  by

$$\langle A, B \rangle : P \longmapsto A^t P \overline{B}.$$

Thus, we define the action of  $SO_4(\mathbb{R}) \cong SU_2(\mathbb{C}) \times SU_2(\mathbb{C})/\{\pm id\}$  on the three-sphere  $\mathbb{S}^3$ .

By assuming w=0, we obtain the two-dimensional sphere

$$\mathbb{S}^2 = \{(x, y, z) \in \mathbb{R}^3 | x^2 + y^2 + z^2 = 1\}.$$

Let us identify a point (x, y, z) of  $\mathbb{S}^2$  with the matrix

$$Q = \begin{pmatrix} ix & y + iz \\ -y + iz & -ix \end{pmatrix},$$

which represents a pure imaginary unit quaternion  $Q \in \mathbf{H}$ .

Instead of Isom<sup>+</sup>  $\mathbb{S}^2 \cong SO_3(\mathbb{R})$  we use its two-fold covering  $SU_2(\mathbb{C})$  acting by

$$A: q \longmapsto A^t q \overline{A}$$

for every  $A \in SU_2(\mathbb{C})$  and every  $q \in \mathbb{S}^2$ .

Equip each  $\mathbb{S}^3$  and  $\mathbb{S}^2$  with an intrinsic metric of constant sectional curvature +1. We call the distance between two points P and Q of  $\mathbb{S}^n$  (n=2,3) a real number d(P,Q) uniquely defined by the conditions

$$0 \le d(P, Q) \le \pi,$$

$$\cos d(P,Q) = \frac{1}{2} \operatorname{tr} P^t \overline{Q}.$$

The next step is to describe spherical geodesic lines in  $\mathbb{S}^n$ . Let us recall the following theorem [25, Theorem 2.1.5].

**Theorem 1** A function  $\lambda : \mathbb{R} \to \mathbb{S}^n$  is a geodesic line if and only if there are orthogonal vectors x, y in  $\mathbb{S}^n$  such that

$$\lambda(t) = (\cos t)x + (\sin t)y.$$

Taking into account the preceding discussion, we may reformulate the statement above.

**Lemma 1** Every geodesic line (a great circle) in  $\mathbb{S}^3$  (respectively,  $\mathbb{S}^2$ ) could be represented in the form

$$C(t) = P\cos t + Q\sin t,$$

where  $P, Q \in SU_2(\mathbb{C})$  (respectively  $P, Q \in \mathbf{H}$ ) satisfy orthogonality condition

$$\cos d(P,Q) = 0.$$

By virtue of this lemma, one may regard P as the starting point of the curve C(t) and Q as the velocity vector at P, since C(0) = P,  $\dot{C}(0) = \frac{d}{dt} C(t)|_{t=0} = Q$  and  $d(C(0), \dot{C}(0)) = \frac{\pi}{2}$  (the latter holds up to a change of the parameter sign).

Given two geodesic lines  $C_1(t)$  and  $C_2(t)$ , define their common perpendicular  $C_{12}(t)$  as a geodesic line such that there exist  $0 \le t_1, t_2 \le 2\pi, \ 0 \le \delta \le \pi$  with the following properties:

$$C_{12}(0) = C_1(t_1), C_{12}(\delta) = C_2(t_2),$$
  
$$d(\dot{C}_{12}(0), \dot{C}_1(t_1)) = d(\dot{C}_{12}(\delta), \dot{C}_2(t_2)) = \frac{\pi}{2}.$$

We call  $\delta$  the distance between the geodesics  $C_1(t)$  and  $C_2(t)$ . Note, that for an arbitrary pair of geodesics their common perpendicular should not be unique.

For an additional explanation of spherical geometry we refer the reader to [25] and [31, Chapter 6.4.2].

# 3 Links arising from the Hopf fibration

The present section is devoted to the construction of a family of links  $\mathcal{H}_n$   $(n \geq 2)$  which we shall use later. These links have a nice property – each of them is formed by  $n \geq 2$  fibres of the Hopf fibration. Recall that the Hopf map  $h: \mathbb{S}^3 \xrightarrow{\mathbb{S}^1} \mathbb{S}^2$  has geometric nature [14, p. 654]. Our aim is to prove a number of lemmas concerning the geometry of the Hopf fibration in more detail.

#### 3.1 Links $\mathcal{H}_n$ as fibres of the Hopf fibration

The Hopf map h is defined as follows [14]: for every point  $(w, x, y, z) \in \mathbb{S}^3$  let its image on  $\mathbb{S}^2$  be

$$h(w, x, y, z) = (2(xz + wy), 2(yz - wx), 1 - 2(x^2 + y^2)).$$

The fibre  $h^{-1}(a,b,c)$  over the point  $(a,b,c)\in\mathbb{S}^2$  is a geodesic line in  $\mathbb{S}^3$  of the form

$$C(t) = \frac{1}{\sqrt{2(1+c)}} \left( (1+c, -b, a, 0) \cos t + (0, a, b, 1+c) \sin t \right).$$

The exceptional point (0,0,-1) has the fibre  $(0,\cos t, -\sin t, 0)$ . The line C(t) is a great circle of  $\mathbb{S}^3$  and can be rewritten in the matrix form

$$C(t) = P(a, b, c)\cos t + Q(a, b, c)\sin t,$$

where

$$P(a,b,c) = \frac{1}{\sqrt{2(1+c)}} \begin{pmatrix} (1+c) - ib & a \\ -a & (1+c) + ib \end{pmatrix},$$
$$Q(a,b,c) = P(a,b,c) \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

We call

$$F(t) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \cos t + \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \sin t$$

the generic fibre  $h^{-1}(0,0,1)$ . Moreover, every fibre  $h^{-1}(a,b,c)$  can be described as a circle C(t) = P(a,b,c) F(t). Note, that P(a,b,c) is an  $SU_2(\mathbb{C})$  matrix. Thus C(t) could be obtained from F(t) by means of the isometry  $\langle P(a,b,c)^t, \mathrm{id} \rangle$ . For the exceptional point  $(0,0,-1) \in \mathbb{S}^2$ , we set

$$P(0,0,-1) = \left(\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right).$$

It is known, that every pair of distinct fibres of the Hopf fibration represents simply linked circles in  $\mathbb{S}^3$  (the Hopf link). Thus, n fibres form a link  $\mathcal{H}_n$  whose every two components form the Hopf link. One can obtain it by drawing n straight vertical lines on a cylinder and identifying its ends by a rotation through the angle of  $2\pi$ . Hence  $\mathcal{H}_n$  is an (n,n) torus link.

Another remark is that the  $\mathcal{H}_n$  link could be arranged around a point in order to reveal its n-th order symmetry, as depicted in Fig. 1. This fact allows us to consider n-fold branched coverings of the corresponding conemanifolds with singular locus  $\mathcal{H}_n$  that appear in Section 4.




Figure 1: n-fold branched covering of (2, 2n) torus link by  $\mathcal{H}_n$ 

#### 3.2 Geometry of the Hopf fibration

Here and below we use the polar coordinate system  $(\psi, \theta)$  on  $\mathbb{S}^2$  instead of the Cartesian one. Suppose

$$a = \cos \psi \sin \theta, \ b = \sin \psi \sin \theta, \ c = \cos \theta,$$
  $0 \le \psi \le 2\pi, \ 0 \le \theta \le \pi$ 

and let

$$M(\psi, \theta) = P(a, b, c) = \begin{pmatrix} \cos \frac{\theta}{2} - i \sin \psi \sin \frac{\theta}{2} & \cos \psi \sin \frac{\theta}{2} \\ -\cos \psi \sin \frac{\theta}{2} & \cos \frac{\theta}{2} + i \sin \psi \sin \frac{\theta}{2} \end{pmatrix}.$$

A rotation of  $\mathbb{S}^3$  about the generic fibre F(t) through angle  $\omega$  has the form  $\langle R(\omega), R(\omega) \rangle$ , where

$$R(\omega) = \begin{pmatrix} \cos\frac{\omega}{2} & i\sin\frac{\omega}{2} \\ i\sin\frac{\omega}{2} & \cos\frac{\omega}{2} \end{pmatrix}.$$

The image of F(t) under the Hopf map h is (0,0) w.r.t. the polar coordinates. The following lemma shows how to obtain a rotation about the pre-image  $h^{-1}(\psi,\theta)$  of an arbitrary point  $(\psi,\theta)$ .

**Lemma 2** A rotation through angle  $\omega$  about an axis C(t) in  $\mathbb{S}^3$  which is the pre-image of a point  $(\psi, \theta) \in \mathbb{S}^2$  with respect to the Hopf map is

$$\langle \overline{M(\psi,\theta)} R(\omega) M(\psi,\theta)^t, R(\omega) \rangle.$$

**Proof.** Since we have that  $C(t) = M(\psi, \theta)F(t)$  and  $R(\omega)^t F(t)\overline{R(\omega)} = F(t)$  for every  $0 \le t \le 2\pi$ , then

$$\begin{split} \left(\overline{M(\psi,\theta)}R(\omega)M(\psi,\theta)^t\right)^tC(t)\overline{R(\omega)} &= M(\psi,\theta)R(\omega)^tF(t)\overline{R(\omega)} = \\ &= M(\psi,\theta)F(t) = C(t) \end{split}$$

by a straightforward computation. Here we use the fact that  $M(\psi, \theta) \in SU_2(\mathbb{C})$ , and so  $\overline{M(\psi, \theta)^t}M(\psi, \theta) = \mathrm{id}$ .  $\square$ Another remarkable property of the Hopf fibration is discussed below.

**Lemma 3** Every two fibres  $C_1(t)$  and  $C_2(t)$  of the Hopf fibration are equidistant geodesic lines (great circles) in  $\mathbb{S}^3$ . If  $C_i(t)$ ,  $i \in \{1, 2\}$  are pre-images of the points  $\widehat{C}_i \in \mathbb{S}^2$ , then the length  $\delta$  of the common perpendicular for  $C_1(t)$  and  $C_2(t)$  equals  $\frac{1}{2}d(\widehat{C}_1, \widehat{C}_2)$ .

**Proof.** The proof follows from the fact that the Hopf fibration is a Riemannian submersion between  $\mathbb{S}^3$  and  $\mathbb{S}^2_{\frac{1}{2}} = \{(x,y,z) \in \mathbb{R}^3 | x^2 + y^2 + z^2 = \frac{1}{4} \}$  with their standard Riemannian metrics of sectional curvature +1 and +4 respectively, see Proposition 1.1 and Proposition 1.2 of [9].  $\square$  Every rotation about a fibre of the Hopf fibration induces a rotation about a point of its base.

**Lemma 4** Given a rotation  $\langle A, B \rangle \in SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$  about a fibre C(t) of the Hopf fibration, the transformation  $A \in SU_2(\mathbb{C})$  induces a rotation of  $\mathbb{S}^2$  about the point to which C(t) projects under the Hopf map.

**Proof.** Rotation about the fibre  $C(t) = M(\psi, \theta)F(t)$  which projects to the point  $(\psi, \theta) \in \mathbb{S}^2$  has the form

$$\langle A, B \rangle = \langle \overline{M(\psi, \theta)} R(\omega) M(\psi, \theta)^t, R(\omega) \rangle.$$

Observe that the rotation  $\langle R(\omega), R(\omega) \rangle$  fixes the geodesic F(t) in  $\mathbb{S}^3$  and  $R(\omega)$  fixes the point  $\widehat{F} = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$  in  $\mathbb{S}^2$ . Thus  $A \in SU_2(\mathbb{C})$  fixes the point  $\widehat{C} = M(\psi, \theta) \widehat{F} M(\psi, \theta)^{\overline{t}}$ . By a straightforward computation, we obtain that

$$\widehat{C} = \left( \begin{array}{cc} i\cos\psi\sin\theta & \sin\theta\sin\psi + i\cos\theta \\ -\sin\theta\sin\psi + i\cos\theta & -i\cos\psi\sin\theta \end{array} \right).$$

The point  $\widehat{C} \in \mathbb{S}^2$  corresponds to  $(\psi, \theta)$  w.r.t. the polar coordinates.  $\square$ 

## 4 Examples of rigidity and flexibility

In this section we work out two principal examples of Seifert fibred conemanifolds: the first represents a rigid cone-manifold, the second one is flexible.

## 4.1 Case of rigidity: the cone-manifold $\mathcal{H}_3(\alpha, \beta, \gamma)$

Let  $\mathcal{H}_3(\alpha, \beta, \gamma)$  denote a three-dimensional cone-manifold with underlying space the sphere  $\mathbb{S}^3$  and singular locus formed by the link  $\mathcal{H}_3$  with cone angles  $\alpha$ ,  $\beta$  and  $\gamma$  along its components. The remaining discussion is devoted to the proof of

**Theorem 2** The cone-manifold  $\mathcal{H}_3(\alpha, \beta, \gamma)$  admits a spherical structure if the following inequalities are satisfied:

$$2\pi - \gamma < \alpha + \beta < 2\pi + \gamma,$$

$$-2\pi + \gamma < \alpha - \beta < 2\pi - \gamma.$$

The spherical structure on  $\mathcal{H}_3(\alpha, \beta, \gamma)$  is unique (i.e.  $\mathcal{H}_3(\alpha, \beta, \gamma)$  is globally rigid).

The lengths  $\ell_{\alpha}$ ,  $\ell_{\beta}$ ,  $\ell_{\gamma}$  of its singular strata are pairwise equal and the following formula holds:

$$\ell_{\alpha} = \ell_{\beta} = \ell_{\gamma} = \frac{\alpha + \beta + \gamma}{2} - \pi.$$

The volume of  $\mathcal{H}_3(\alpha, \beta, \gamma)$  equals

Vol 
$$\mathcal{H}_3(\alpha, \beta, \gamma) = \frac{1}{2} \left( \frac{\alpha + \beta + \gamma}{2} - \pi \right)^2$$
.

**Proof.** First, we construct a holonomy map for  $\mathcal{H}_3(\alpha, \beta, \gamma)$ . By applying Wirtinger's algorithm, one obtains the following fundamental group presentation for the link  $\mathcal{H}_3$ :

$$\Gamma = \pi_1(\mathbb{S}^3 \setminus \mathcal{H}_3) = \langle a, b, c, h | acb = bac = cba = h, h \in Z(\Gamma) \rangle,$$

that is a central extension by h of the thrice-punctured sphere group

$$\Gamma_0 = \pi_1(\mathbb{S}^2 \setminus \{3 \text{ points}\}) = \langle a, b, c | acb = bac = cba = \text{id} \rangle.$$

Consider a holonomy map

$$\rho: \Gamma \longmapsto \operatorname{Isom}^+ \mathbb{S}^3 \cong SO_4(\mathbb{R}).$$




Figure 2: The link  $\mathcal{H}_3$ 

Let  $\widetilde{\rho}$  denote its lift to  $SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$ , which is a two-fold covering of  $SO_4(\mathbb{R})$  (see [7]):

$$\widetilde{\rho} = \langle \widetilde{\rho}_1, \widetilde{\rho}_2 \rangle : \Gamma \longmapsto SU_2(\mathbb{C}) \times SU_2(\mathbb{C}).$$

Let us note, that if holonomy images of any two generators of  $\Gamma$  commute, then the whole homomorphic image  $\widetilde{\rho}(\Gamma)$  is abelian. Thus, for a representation  $\widetilde{\rho}$  we have that the following three cases, up to a suitable conjugation, are possible:

- (i)  $\widetilde{\rho} = (\widetilde{\rho}_1, \widetilde{\rho}_2) : \Gamma \to SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$ , both  $\widetilde{\rho}_1$  and  $\widetilde{\rho}_2$  are non-abelian,
- (ii)  $\widetilde{\rho}:\Gamma\to\mathbb{S}^1\times\mathbb{S}^1,$  an abelian representation,
- (iii)  $\widetilde{\rho} = (\widetilde{\rho}_1, \widetilde{\rho}_2) : \Gamma \to SU_2(\mathbb{C}) \times \mathbb{S}^1$ , where  $\widetilde{\rho}_1$  is non-abelian.

For case (i), let us first suppose that  $\tilde{\rho}(h)$  is non-trivial. Since the holonomy images of the meridians a, b and c have to commute with the holonomy image of h, they are simultaneously diagonalisable. We arrive at case (ii). If  $\tilde{\rho}(h)$  is trivial, then we have two non-abelian representations  $\tilde{\rho}_i: \Gamma_0 \to SU_2(\mathbb{C})$ . Since the holonomy images of the meridians correspond to rotations along geodesic lines in  $\mathbb{S}^3$ , it follows by [2, Lemma 9.2] that  $\operatorname{tr} \tilde{\rho}_1(x) = \operatorname{tr} \tilde{\rho}_2(x)$  for  $x \in \{a, b, c\}$ . The base space of the fibred cone-manifold  $\mathcal{H}_3(\alpha, \beta, \gamma)$  is a turnover  $\mathbb{S}^2(\alpha, \beta, \gamma)$ , with  $\alpha, \beta, \gamma$  cone angles. Then, by [10, Lemma 4.1], up to a conjugation,  $\tilde{\rho} = (\tilde{\rho}_1, \tilde{\rho}_1)$ . The representation  $\rho: \Gamma \to SO(4)$  is conjugate into SO(3) and the holonomy images of the meridians have a common fixed point in  $\mathbb{S}^3$ . Thus, their axis intersect, which does not

correspond to a non-degenerate spherical structure on the cone-manifold  $\mathcal{H}_3(\alpha, \beta, \gamma)$ .

For case (ii), up to a suitable conjugation, the representation  $\tilde{\rho}$  preserves the Hopf fibration. Thus, by Lemma 4, it descends to an abelian representation of  $\Gamma_0$ , which cannot be a holonomy of a non-degenerate spherical structure on the base of the fibration.

Finally, case (iii) is left. By [2, Lemma 9.2], one has

$$\widetilde{\rho}(a) = \langle m_a^t R(\alpha) \overline{m_a}, R(\alpha) \rangle,$$

$$\widetilde{\rho}(b) = \langle m_b^t R(\beta) \overline{m_b}, R(\beta) \rangle,$$

$$\widetilde{\rho}(c) = \langle m_c^t R(\gamma) \overline{m_c}, R(\gamma) \rangle$$

for  $m_a$ ,  $m_b$ ,  $m_c \in SU_2(\mathbb{C})$ .

Note, that every matrix  $m \in SU_2(\mathbb{C})$  is of the form  $m = R(\tau)M(\psi,\theta)$  for suitable  $0 \le \psi \le \pi$ ,  $0 \le \theta, \tau \le 2\pi$ . Then we obtain that the image of every meridian in  $\Gamma = \pi_1(\mathbb{S}^3 \setminus \mathcal{H}_3)$  has the form

$$\langle m^t R(\omega) \overline{m}, R(\omega) \rangle = \langle M^t(\psi, \theta) R^t(\tau) R(\omega) \overline{R(\tau)} \overline{M(\psi, \tau)}, R(\omega) \rangle = \langle M^t(\psi, \theta) R(\omega) \overline{M(\psi, \theta)}, R(\omega) \rangle,$$

since  $R(\omega)$  and  $R(\tau)$  commute. Hence, Lemma 2 implies that every meridian is mapped by  $\widetilde{\rho}$  to a rotation about an appropriate fibre of the Hopf fibration. By Propositions 2.1 and 2.2 of [9], the holonomy preserves the fibration structure.

Let  $A = \widetilde{\rho}(a)$ ,  $B = \widetilde{\rho}(b)$ ,  $C = \widetilde{\rho}(c)$  be holonomy images of the generators a, b, c for  $\Gamma = \pi_1(\mathbb{S}^3 \setminus \mathcal{H}_3)$ .

After a suitable conjugation in  $SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$ , we obtain

$$A = \langle A_l, A_r \rangle = \langle R(\alpha), R(\alpha) \rangle ,$$

$$B = \langle B_l, B_r \rangle = \left\langle \overline{M(0, \phi)} R(\beta) M(0, \phi)^t, R(\beta) \right\rangle ,$$

$$C = \langle C_l, C_r \rangle = \left\langle \overline{M(\psi, \theta)} R(\gamma) M(\psi, \theta)^t, R(\gamma) \right\rangle .$$

In order for the holonomy map  $\widetilde{\rho}$  to be a homomorphism, the following relations should hold:

$$A_lC_lB_l = B_lA_lC_l = C_lB_lA_l,$$
  

$$A_rC_rB_r = B_rA_rC_r = C_rB_rA_r.$$

The latter of them are satisfied by the construction of  $\tilde{\rho}: \Gamma \to SU_2(\mathbb{C}) \times \mathbb{S}^1$ . Let us consider the former relations. By Lemma 4, the elements  $A_l$ ,  $B_l$  and  $C_l$  are rotations of  $\mathbb{S}^2$  about the points  $\hat{F}_a = (0,0)$ ,  $\hat{F}_b = (0,\phi)$  and  $\hat{F}_c = (\psi,\theta)$ , respectively. Since  $\hat{F}_a$ ,  $\hat{F}_b$ ,  $\hat{F}_c$  form a triangle on  $\mathbb{S}^2$  and the base space of  $\mathcal{H}_3(\alpha,\beta,\gamma)$  is a turnover with  $\alpha$ ,  $\beta$ ,  $\gamma$  cone angles, one may expect the following

**Lemma 5** The points  $\widehat{F}_a=(0,0)$ ,  $\widehat{F}_b=(0,\phi)$  and  $\widehat{F}_c=(\psi,\theta)$  form a triangle with angles  $\frac{\alpha}{2}$ ,  $\frac{\beta}{2}$  and  $\frac{\gamma}{2}$  at the corresponding vertices.

**Proof.** By a straightforward computation, we obtain that

$$A_l C_l B_l - B_l A_l C_l = \begin{pmatrix} iR_1 & R_2 + iR_3 \\ -R_2 + iR_3 & -iR_1 \end{pmatrix},$$

$$C_l B_l A_l - B_l A_l C_l = \begin{pmatrix} iR_4 & R_5 + iR_3 \\ -R_5 + iR_3 & -iR_4 \end{pmatrix},$$

where

$$R_1 = 2\sin\frac{\beta}{2}\sin\frac{\gamma}{2}\sin\theta\cos\phi\sin\left(\frac{\alpha}{2} - \psi\right),\,$$

$$R_2 = 2\sin\frac{\beta}{2}\left(\cos\frac{\gamma}{2}\sin\frac{\alpha}{2}\sin\phi + \sin\frac{\gamma}{2}\left(-\cos\phi\cos\left(\frac{\alpha}{2} - \psi\right)\sin\theta + \cos\frac{\alpha}{2}\cos\theta\sin\phi\right)\right),$$

$$R_3 = -2\sin\frac{\beta}{2}\sin\frac{\gamma}{2}\sin\theta\sin\phi\sin\left(\frac{\alpha}{2} - \psi\right),\,$$

$$R_4 = 2\sin\frac{\gamma}{2}\left(\cos\theta\sin\frac{\alpha}{2}\sin\frac{\beta}{2}\sin\phi - \left(\cos\frac{\beta}{2}\sin\frac{\alpha}{2} + \cos\frac{\alpha}{2}\sin\frac{\beta}{2}\cos\phi\right)\sin\theta\sin\psi\right),\,$$

$$R_5 = 2\sin\frac{\gamma}{2} \left(\cos\frac{\beta}{2}\cos\psi\sin\frac{\alpha}{2}\sin\theta + \cos\frac{\alpha}{2}\sin\frac{\beta}{2}(\cos\phi\cos\psi\sin\theta - \cos\theta\sin\phi)\right).$$

In order to determine the parameters  $\phi$ ,  $\psi$  and  $\theta$ , one can proceed as follows: these are determined by the system of equations  $R_k = 0, k \in \{1, ..., 5\}$ 

under the restrictions  $0 < \alpha, \beta, \gamma < 2\pi$  and  $0 < \psi \le 2\pi$ ,  $0 < \theta \le \pi$ . Thus, the common solutions to  $R_1$  and  $R_3$  are  $\psi = \frac{\alpha}{2}$  and  $\psi = \frac{\alpha}{2} \pm \pi$ . We claim that the cone angles in the base space of  $\mathcal{H}_3(\alpha, \beta, \gamma)$  and along its fibres are the same, and choose  $\psi = \frac{\alpha}{2}$ .

Taking into account that  $0 < \alpha, \beta, \gamma < 2\pi$  (this implies that the sine functions of half cone angles are non-zero), turn the set of relations  $R_k$ ,  $k \in \{1, \ldots, 5\}$  into a new one:

$$\widetilde{R}_1 = -\cos\phi\sin\frac{\gamma}{2}\sin\theta + \left(\sin\frac{\alpha}{2}\cos\frac{\gamma}{2} + \cos\frac{\alpha}{2}\sin\frac{\gamma}{2}\cos\theta\right)\sin\phi,$$

$$\widetilde{R}_2 = -\cos\theta\sin\frac{\beta}{2}\sin\phi + \left(\sin\frac{\alpha}{2}\cos\frac{\beta}{2} + \cos\frac{\alpha}{2}\sin\frac{\beta}{2}\cos\phi\right)\sin\theta.$$

Note, that the conditions of Theorem 2 concerning cone angles are exactly the existence conditions for a spherical triangle with angles  $\frac{\alpha}{2}$ ,  $\frac{\beta}{2}$  and  $\frac{\gamma}{2}$ . For the latter, the following trigonometric identities (spherical cosine and sine rules) are satisfied [25, Theorems 2.5.2 and 2.5.4]:

$$\cos \phi = \frac{\cos \frac{\gamma}{2} + \cos \frac{\alpha}{2} \cos \frac{\beta}{2}}{\sin \frac{\alpha}{2} \sin \frac{\beta}{2}},$$
$$\cos \theta = \frac{\cos \frac{\beta}{2} + \cos \frac{\alpha}{2} \cos \frac{\gamma}{2}}{\sin \frac{\alpha}{2} \sin \frac{\gamma}{2}},$$

$$\frac{\sin\phi}{\sin\frac{\gamma}{2}} = \frac{\sin\theta}{\sin\frac{\beta}{2}}.$$

These identities state that the points  $\widehat{F}_a$ ,  $\widehat{F}_b$  and  $\widehat{F}_c$  form a triangle on  $\mathbb{S}^2$  with angles  $\frac{\alpha}{2}$ ,  $\frac{\beta}{2}$  and  $\frac{\gamma}{2}$  at the corresponding vertices. Its double provides the base turnover with cone angles  $\alpha$ ,  $\beta$  and  $\gamma$  for the fibred cone-manifold  $\mathcal{H}_3(\alpha, \beta, \gamma)$ .

On substituting the expressions for  $\cos \phi$  and  $\cos \psi$  above in the relations  $\widetilde{R}_k$ ,  $k \in \{1,2\}$  and taking into account the sine rule, one obtains that  $\widetilde{R}_k = 0$ ,  $k \in \{1,2\}$ . The lemma is proven.  $\square$ 

Let  $\mathcal S$  denote the domain of cone angles indicated in the statement of the theorem:

$$S = \left\{ \overrightarrow{\alpha} = (\alpha, \beta, \gamma) \middle| \begin{array}{c} 2\pi - \gamma < \alpha + \beta < 2\pi + \gamma \\ -2\pi + \gamma < \alpha - \beta < 2\pi - \gamma \end{array} \right\}.$$

Let  $\mathcal{S}^*$  denote the subset of  $\mathcal{S}$ , such that for every triple of cone angles  $\overrightarrow{\alpha} = (\alpha, \beta, \gamma) \in \mathcal{S}^*$  there exists a spherical structure on  $\mathcal{H}_3(\overrightarrow{\alpha})$ . Our next step is to show that  $\mathcal{S}^*$  coincides with  $\mathcal{S}$ .

The set  $\mathcal{S}^*$  is non-empty. From [8], it follows that  $\mathcal{H}_3(\pi, \pi, \pi)$  has a spherical structure. The orbifold  $\mathcal{H}_3(\pi, \pi, \pi)$  is Seifert fibred and its base is a turnover with cone angles equal to  $\pi$ . Thus, the point  $(\pi, \pi, \pi) \in \mathcal{S}$  belongs to  $\mathcal{S}^*$ . The set  $\mathcal{S}^*$  is open, because a deformation of the holonomy induces a deformation of the structure [20].

In order to prove that the set  $\mathcal{S}^*$  is closed, we consider a sequence  $\overrightarrow{\alpha}_n = (\alpha_n, \beta_n, \gamma_n)$  in  $\mathcal{S}^*$  converging to  $\overrightarrow{\alpha}_{\infty} = (\alpha_{\infty}, \beta_{\infty}, \gamma_{\infty})$  in  $\mathcal{S}$ . Since every spherical cone-manifold with cone angles  $\leq 2\pi$  is an Alexandrov space with curvature  $\geq 1$  [3], we obtain that the diameter of  $\mathcal{H}_3(\overrightarrow{\alpha}_n)$  is bounded above: diam  $\mathcal{H}_3(\overrightarrow{\alpha}_n) \leq \pi$ .

Let dist  $\mathcal{H}_3(\overrightarrow{\alpha}_n)$  denote the minimum of the mutual distances between the axis of rotations A, B and C. Since  $\overrightarrow{\alpha}_{\infty} \in \mathcal{S}$ , we have by Lemma 5 that the turnover  $\mathbb{S}^2(\overrightarrow{\alpha}_{\infty})$  is non-degenerate. By making use of Lemma 3, one obtains that (restricting to a subsequence, if needed) for every  $\overrightarrow{\alpha}_n \in \mathcal{S}$ ,  $n = 1, 2, \ldots$  the function dist  $\mathcal{H}_3(\overrightarrow{\alpha}_n)$  is uniformly bounded below away from zero:

dist 
$$\mathcal{H}_3(\overrightarrow{\alpha}_n) \ge d_0 > 0, \ n = 1, 2, \dots$$

Then we use the following facts [3]:

- 1. The Gromov-Hausdorff limit of Alexandrov spaces with curvature  $\geq 1$ , dimension = 3 and bounded diameter is an Alexandrov space with curvature  $\geq 1$  and dimension  $\leq 3$ ,
- 2. Dimension of an Alexandrov space with curvature  $\geq 1$  holds the same at every point (the word "dimension" means Hausdorff or topological dimension, which are equal in the case of curvature  $\geq 1$ ).

Since dist  $\mathcal{H}_3(\overrightarrow{\alpha}_n) \geq d_0 > 0$ , the sequence  $\mathcal{H}_3(\overrightarrow{\alpha}_n)$  does not collapse. Thus, the cone-manifold  $\mathcal{H}_3(\overrightarrow{\alpha}_{\infty})$  has a non-degenerate spherical structure and  $\overrightarrow{\alpha}_{\infty} \in \mathcal{S}^*$ .

The subset  $S^* \subset S$  is non-empty, as well as both closed and open. This implies  $S^* = S$ .

Finally, we claim the following fact concerning the geometric characteristics of  $\mathcal{H}_3(\alpha, \beta, \gamma)$  cone-manifold:

**Lemma 6** Let  $\ell_{\alpha}$ ,  $\ell_{\beta}$ ,  $\ell_{\gamma}$  denote the lengths of the singular strata for  $\mathcal{H}_3(\alpha, \beta, \gamma)$  cone-manifold with cone angles  $\alpha$ ,  $\beta$  and  $\gamma$ . Then

$$\ell_{\alpha} = \ell_{\beta} = \ell_{\gamma} = \frac{\alpha + \beta + \gamma}{2} - \pi.$$

The volume of  $\mathcal{H}_3(\alpha, \beta, \gamma)$  is

Vol 
$$\mathcal{H}_3(\alpha, \beta, \gamma) = \frac{1}{2} \left( \frac{\alpha + \beta + \gamma}{2} - \pi \right)^2$$
.

**Proof.** Let us calculate the geometric parameters explicitly, using the holonomy map defined above. First, we introduce two notions suitable for the further discussion. Given an element  $M = \langle M_l, M_r \rangle \in SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$ , one may assume that the pair of matrices  $\langle M_l, M_r \rangle$  is conjugated, by means of a certain element  $\langle C_l, C_r \rangle \in SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$ , to the pair of diagonal matrices

$$\left\langle \left(\begin{array}{cc} e^{i\gamma} & 0\\ 0 & e^{-i\gamma} \end{array}\right), \left(\begin{array}{cc} e^{i\varphi} & 0\\ 0 & e^{-i\varphi} \end{array}\right) \right\rangle$$

with  $0 \le \gamma, \varphi \le \pi$ .

Then call the translation length of M the quantity  $\delta(M) := \varphi - \gamma$  and call the "jump" of M the quantity  $\nu(M) := \varphi + \gamma$ , see [11] and [31, Ch.6.4.2]. We suppose that  $\varphi > \gamma$ , otherwise changing  $\gamma$ ,  $\varphi$  for  $2\pi - \gamma$  and  $\pi - \varphi$  makes the considered tuple to have the desired form.

Recall that the representation of  $\Gamma = \pi_1(\mathbb{S}^3 \setminus \mathcal{H}_3)$  is

$$\Gamma = \langle a, b, c, h | acb = bac = cba = h, h \in Z(\Gamma) \rangle,$$

where a, b, c are meridians and h is a longitudinal loop that represents a fibre. Denote by H the image of h under the holonomy map  $\widetilde{\rho}$ . Then we obtain

$$\ell_{\alpha} = \ell_{\beta} = \ell_{\gamma} = \delta(H).$$

Since  $A = \widetilde{\rho}(a)$  and  $H = \widetilde{\rho}(h)$  commute, there exists an element  $C = \langle C_l, C_r \rangle$  of  $SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$  such that

$$CAC^{-1} = \left\langle \left( \begin{array}{cc} e^{i\frac{\alpha}{2}} & 0\\ 0 & e^{-i\frac{\alpha}{2}} \end{array} \right), \left( \begin{array}{cc} e^{i\frac{\alpha}{2}} & 0\\ 0 & e^{-i\frac{\alpha}{2}} \end{array} \right) \right\rangle,$$

$$CHC^{-1} = \left\langle \left( \begin{array}{cc} e^{i\gamma(H)} & 0 \\ 0 & e^{-i\gamma(H)} \end{array} \right), \left( \begin{array}{cc} e^{i\varphi(H)} & 0 \\ 0 & e^{-i\varphi(H)} \end{array} \right) \right\rangle.$$

By a straightforward computation similar to that in Lemma 5, one obtains

$$2\cos\gamma(H) = \operatorname{tr} H_I = \operatorname{tr} A_I C_I B_I = \operatorname{tr}(-\operatorname{id}) = 2\cos\pi$$

and

$$2\cos\varphi(H) = \operatorname{tr} H_r = \operatorname{tr} A_r C_r B_r = 2\cos\frac{\alpha + \beta + \gamma}{2}.$$

From the foregoing discussion, the singular stratum's length is

$$\ell_{\alpha} = \delta(H) = \frac{\alpha + \beta + \gamma}{2} - \pi.$$

An analogous equality holds for  $\ell_{\beta}$  and  $\ell_{\gamma}$ . By the Schläfli formula [12], the following relation holds:

2 dVol 
$$\mathcal{H}_3(\alpha, \beta, \gamma) = \ell_{\alpha} d\alpha + \ell_{\beta} d\beta + \ell_{\gamma} d\gamma$$
.

Solving this differential equality, we obtain that

$$\operatorname{Vol} \mathcal{H}_{3}(\alpha, \beta, \gamma) = \frac{1}{2} \left( \frac{\alpha + \beta + \gamma}{2} - \pi \right)^{2} + \operatorname{Vol}_{0},$$

where Vol<sub>0</sub> is an arbitrary constant. Since the geometric structure on the base space of the fibration (consequently, on the whole  $\mathcal{H}_3(\alpha, \beta, \gamma)$  conemanifold) degenerates when  $\alpha + \beta + \gamma \longrightarrow 2\pi$ , the equality Vol<sub>0</sub> = 0 follows from the volume function continuity.  $\square$ 

Consider a holonomy  $\tilde{\rho} = \langle \tilde{\rho}_1, \tilde{\rho}_2 \rangle$ :  $\Gamma = \pi_1(\mathbb{S}^3 \setminus \mathcal{H}_3) \to SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$  for  $\mathcal{H}_3(\alpha, \beta, \gamma)$  cone-manifold. As we already know from the preceding discussion, one has  $\tilde{\rho}: \Gamma \to SU_2(\mathbb{C}) \times \mathbb{S}^1$  essentially, and  $\tilde{\rho}_1$  determines  $\tilde{\rho}_2$  up to a conjugation by means of the equality  $\operatorname{tr} \tilde{\rho}_1(m) = \operatorname{tr} \tilde{\rho}_2(m)$  for meridians in  $\Gamma$ . So any deformation of  $\tilde{\rho}$  is a deformation of  $\tilde{\rho}_1$ . In the case of  $\mathcal{H}_3(\alpha, \beta, \gamma)$ , the map  $\tilde{\rho}_1$  is a non-abelian representation of the base turnover group. Spherical turnover is rigid, that means  $\tilde{\rho}_1$  is determined only by the corresponding cone angles. Thus  $\mathcal{H}_3(\alpha, \beta, \gamma)$  is locally rigid.

The global rigidity follows from the fact that every  $\mathcal{H}_3(\alpha, \beta, \gamma)$  cone-manifold could be deformed to the orbifold  $\mathcal{H}_3(\pi, \pi, \pi)$  by a continuous path through locally rigid structures. This assertion holds since  $\mathcal{S}^*$  contains the point  $(\pi, \pi, \pi)$  and  $\mathcal{S}^*$  is convex. The global rigidity of  $\mathcal{H}_3(\pi, \pi, \pi)$  spherical orbifold follows from [26, 27] and implies the global rigidity of  $\mathcal{H}_3(\alpha, \beta, \gamma)$  by means of deforming the orbifold structure backwards to the considered conemanifold one.  $\square$ 

### 4.2 Case of flexibility: the cone-manifold $\mathcal{H}_4(\alpha)$

Let  $\mathcal{H}_4(\alpha)$  denote a three-dimensional cone-manifold with underlying space the sphere  $\mathbb{S}^3$  and singular locus formed by the link  $\mathcal{H}_4$  with cone angle  $\alpha$ along all its components.

The following theorem provides an example of a flexible cone-manifold which is Seifert fibred.




Figure 3: The link  $\mathcal{H}_4$ 

**Theorem 3** The cone-manifold  $\mathcal{H}_4(\alpha)$  admits a spherical structure if

$$\pi < \alpha < 2\pi$$
.

This structure is not unique (i.e.  $\mathcal{H}_4(\alpha)$  is not globally, nor locally rigid). The deformation space contains an open interval, that provides a one-parameter family of distinct spherical cone-metrics on  $\mathbb{S}^3$ . The length of each singular stratum is

$$\ell = 2(\alpha - \pi).$$

The volume of  $\mathcal{H}_4(\alpha)$  equals

$$Vol \mathcal{H}_4(\alpha) = 2(\alpha - \pi)^2.$$

**Proof.** The following lemma precedes the proof of the theorem.

**Lemma 7** Given a quadrangle Q on  $\mathbb{S}^2$  with three right angles and one angle  $\frac{\alpha}{2}$  (see Fig. 4), the following statements hold:

- 1. The quadrangle Q exists if  $\pi < \alpha < 2\pi$ ,
- 2.  $\sin \ell_1 \sin \ell_2 = -\cos \frac{\alpha}{2}$ ,
- 3.  $\cos \phi = \frac{\cos \ell_1 \cos \ell_2}{\sin \frac{\alpha}{2}}$ ,

4.  $\cos \psi = \tan \ell_1 \cot \phi$ ,

5. 
$$0 \le \ell_1, \, \ell_2, \, \phi, \, \psi \le \frac{\pi}{2}$$
.

**Proof.** We refer the reader to  $[30, \S 3.2]$  for a detailed proof of the statements above.  $\square$ 




Figure 4: The quadrangle Q

Given a quadrangle Q from Lemma 7 (so-called Saccheri's quadrangle) one can construct another one, depicted in Fig.5, by reflecting Q in its sides incident to the vertex O. We may regard O to be the point  $(0,0) \in \mathbb{S}^2$ . Thus, the fibres over the corresponding vertices are

$$F_a(t) = M(\psi, \phi) F(t),$$
  

$$F_b(t) = M(\pi - \psi, \phi) F(t),$$
  

$$F_c(t) = M(\pi + \psi, \phi) F(t),$$
  

$$F_d(t) = M(2\pi - \psi, \phi) F(t).$$

Let  $A = \langle A_l, A_r \rangle$ ,  $B = \langle B_l, B_r \rangle$ ,  $C = \langle C_l, C_r \rangle$ ,  $D = \langle D_l, D_r \rangle$  denote the respective rotations through angle  $\alpha$  about the axis  $F_a$ ,  $F_b$ ,  $F_c$  and  $F_d$ . From Lemma 2, one obtains

$$A_{l} = \overline{M(\psi, \phi)} R(\alpha) M(\psi, \phi)^{t}, A_{r} = R(\alpha);$$

$$B_{l} = \overline{M(\pi - \psi, \phi)} R(\alpha) M(\pi - \psi, \phi)^{t}, B_{r} = R(\alpha);$$

$$C_{l} = \overline{M(\pi + \psi, \phi)} R(\alpha) M(\pi + \psi, \phi)^{t}, B_{r} = R(\alpha);$$

$$D_{l} = \overline{M(2\pi - \psi, \phi)} R(\alpha) M(2\pi - \psi, \phi)^{t}, D_{r} = R(\alpha).$$

We assume that  $\ell_1$ ,  $\ell_2$ ,  $\phi$  and  $\psi$  satisfy the identities of Lemma 7.




Figure 5: The base quadrangle P for  $\mathcal{H}_4(\alpha)$ 

The fundamental group of  $\pi_1(\mathbb{S}^3 \setminus \mathcal{H}_4)$  has the presentation

$$\Gamma = \pi_1(\mathbb{S}^3 \setminus \mathcal{H}_4) = \langle a, b, c, d, h | adcb = badc = cbad = dcba = h, h \in Z(\Gamma) \rangle.$$

Let us construct a lift of the holonomy map  $\widetilde{\rho}:\Gamma\to SU_2(\mathbb{C})\times SU_2(\mathbb{C})$  as follows:

$$\widetilde{\rho}(a) = A, \ \widetilde{\rho}(b) = B, \ \widetilde{\rho}(c) = C, \ \widetilde{\rho}(d) = D.$$

Here we choose  $\widetilde{\rho}: \Gamma \to SU_2(\mathbb{C}) \times \mathbb{S}^1$  by the same reason as in Theorem 2. In order to show that the map  $\widetilde{\rho}$  is a homomorphism, one has to check whether the following relations are satisfied:

$$A_l D_l C_l B_l = B_l A_l D_l C_l = C_l B_l A_l D_l = D_l C_l B_l A_l,$$
  

$$A_r D_r C_r B_r = B_r A_r D_r C_r = C_r B_r A_r D_r = D_r C_r B_r A_r.$$

The latter relations hold in view of the fact that the matrices  $A_r$ ,  $B_r$ ,  $C_r$  and  $D_r$  pairwise commute. Then, we show that the following equality holds:

$$A_l D_l C_l B_l = id.$$

To do this, split the quadrangle P into two triangles by drawing a geodesic line from B to D. Since  $A_l$ ,  $B_l$ ,  $C_l$  and  $D_l$  are rotations about the vertices of the quadrangle depicted in Fig. 6, let us decompose the rotations  $B_l = B'_l B''_l$  and  $D_l = D'_l D''_l$  into the products of rotations  $B'_l$ ,  $B''_l$  through angles  $\beta_1$ ,




Figure 6: Section of P by the line joining vertices B and D

 $\beta_2$  and the rotations  $D'_l$ ,  $D''_l$  through angles  $\delta_1$  and  $\delta_2$ , respectively. The following equalities hold:  $\beta_1 + \beta_2 = \frac{\alpha}{2}$  and  $\delta_1 + \delta_2 = \frac{\alpha}{2}$ . Thus, the triples  $D''_l$ ,  $C_l$ ,  $B'_l$  and  $A_l$ ,  $D'_l$ ,  $B''_l$  consist of rotations about the vertices of two disjoint triangles depicted in Fig. 6. Similar to the computation of Lemma 6, we have

$$D_l''C_lB_l' = -\mathrm{id}$$

and

$$A_l D_l' B_l'' = -\mathrm{id}.$$

From the identities above, it follows that

$$A_l D_l C_l B_l = A_l D_l' D_l'' C_l B_l' B_l'' = -A_l D_l' B_l'' = id.$$

The statement holds under a cyclic permutation of the factors. Thus,

$$A_l D_l C_l B_l = B_l A_l D_l C_l = C_l B_l A_l D_l = D_l C_l B_l A_l = id.$$

Below we shall consider the side-length  $\ell_1$  as a parameter. Let  $\ell_1 := \tau$ . Then by Lemma 7 one has that  $\sin \ell_2 = -\frac{\cos \frac{\alpha}{2}}{\sin \tau}$  and  $\ell_2 := \ell_2(\tau)$  is a well-defined continuous function of  $\tau$ . The quadrangle P depends on the parameter  $\tau$  continuously while keeping the angles in its vertices equal to  $\frac{\alpha}{2}$ .

Let  $\mathcal{H}_4(\alpha;\tau)$  denote a three-dimensional cone-manifold with underlying space the sphere  $\mathbb{S}^3$  and singular locus the link  $\mathcal{H}_4$  with cone angle  $\alpha$  along its components. Furthermore, its holonomy map is determined by the quadrangle P described above (see Fig. 5) depending on the parameter  $\tau$ . This means that the double of P forms a "pillowcase" cone-surface with all cone angles equal to  $\alpha$ , which is the base space for the fibred cone-manifold  $\mathcal{H}_4(\alpha;\tau)$ .

Let  $\mathbb{L}_n(\alpha,\beta)$  be a cone-manifold with underlying space the sphere  $\mathbb{S}^3$  and singular locus a torus link of the type (2,2n) with cone angles  $\alpha$  and  $\beta$  along its components. Torus links of the type (2,2n) are two-bridge links. The corresponding cone-manifolds were previously considered in [17, 22]. Since the cone-manifold  $\mathcal{H}_4(\alpha)$  forms a 4-fold branched covering of the conemanifold  $\mathbb{L}_4(\alpha,\frac{\pi}{2})$ , from [17, Theorem 2] we obtain that  $\mathcal{H}_4(\alpha)$  has a spherical structure if  $\pi < \alpha < 2\pi$ . The length of each singular stratum equals to  $\ell = 2(\alpha - \pi)$  and the volume is  $\operatorname{Vol} \mathcal{H}_4(\alpha) = 2(\alpha - \pi)^2$ .

Under the assumption that  $\ell_1 = \ell_2$ , the base quadrangle depicted in Fig. 5 appears to have a four order symmetry. Moreover, by making use of Lemma 7, one may derive the following equalities:  $\psi = \frac{\pi}{4}$ ,  $\cos \phi = \cot \frac{\alpha}{4}$ . The general formulas for the holonomy of  $\mathcal{H}_4(\alpha)$  cone-manifold derived above subject to the condition  $\ell_1 = \ell_2$  (equivalently, the cone-manifold  $\mathcal{H}_4(\alpha)$  has a four order symmetry) give the holonomy map induced by the covering. Thus  $\mathcal{H}_4(\alpha) \cong \mathcal{H}_4(\alpha; \arccos(\sqrt{2}\cos\frac{\alpha}{4}))$  is a spherical cone-manifold.

We claim that one can vary the parameter  $\tau$  in certain ranges while keeping spherical structure on  $\mathcal{H}_4(\alpha;\tau)$  non-degenerate.

**Lemma 8** If  $\tau$  varies over  $(\frac{\alpha-\pi}{2}, \frac{\pi}{2})$ , the cone-manifold  $\mathcal{H}_4(\alpha; \tau)$  has a non-degenerate spherical structure.

**Proof.** The proof has much in common with the proof of the spherical structure existence on  $\mathcal{H}_3(\alpha, \beta, \gamma)$  cone-manifold given in Theorem 2. Let us express the identities of Lemma 7 in terms of the parameter  $\ell_1 := \tau$ . We obtain

$$\cos \phi = \cos \tau \sqrt{1 - \cot^2 \frac{\alpha}{2} \cot^2 \tau},$$

$$\cos \psi = \sqrt{\frac{1 - \cot^2 \frac{\alpha}{2} \cot^2 \tau}{1 + \cot^2 \frac{\alpha}{2} \cot^4 \tau}},$$

$$\sin \ell_2 = -\frac{\cos \frac{\alpha}{2}}{\sin \tau}.$$

Since Lemma 7 states that  $0 \le \phi$ ,  $\psi$ ,  $\ell_2 \le \frac{\pi}{2}$ , the functions  $\phi := \phi(\tau)$ ,  $\psi := \psi(\tau)$ ,  $\ell_2 := \ell_2(\tau)$  are well-defined and depend continuously on  $\tau$ . Moreover, the following relations hold:

$$\cos b_1 = \frac{\cos \phi}{\cos \ell_2} = \cos \tau \sqrt{\frac{\sin^2 \tau - \cot^2 \frac{\alpha}{2} \cos^2 \tau}{\sin^2 \tau - \cos^2 \frac{\alpha}{2}}},$$
$$\cos b_2 = \frac{\cos \phi}{\cos \tau} = \sqrt{1 - \cot^2 \frac{\alpha}{2} \cot^2 \tau}.$$

If one sets the centre O of the quadrangle P to  $(0,0) \in \mathbb{S}^2$ , the whole quadrangle is situated in the upper hemisphere provided  $\phi < \frac{\pi}{2}$ . From the fact that  $\cos b_1 \geq \cos \phi$  and  $\cos b_2 \geq \cos \phi$ , it follows  $b_1, b_2 \leq \phi$ . Thus  $b_1, b_2 \leq \frac{\pi}{2}$  and the functions  $b_1 := b_1(\tau), b_2 := b_2(\tau)$  are well-defined and continuous with respect to  $\tau$ .

Observe that if the condition  $\frac{\alpha-\pi}{2} < \tau < \frac{\pi}{2}$  is satisfied, then the required inequality  $\phi < \frac{\pi}{2}$  holds.

Let  $S_{\alpha}^*$  denote the subset of  $S_{\alpha} = \{\tau | \frac{\alpha - \pi}{2} < \tau < \frac{\pi}{2}\}$  that consists of the points  $\tau \in S_{\alpha}$  such that the cone-manifold  $\mathcal{H}_4(\alpha; \tau)$  has a non-degenerate spherical structure. We show  $S_{\alpha}^* = S_{\alpha}$  by means of the fact that  $S_{\alpha}^*$  is both open and closed non-empty subset of  $S_{\alpha}$ .

As noticed above,  $\tau = \arccos(\sqrt{2}\cos\frac{\alpha}{4})$  belongs to  $\mathcal{S}_{\alpha}^*$ . Hence the set  $\mathcal{S}_{\alpha}^*$  is non-empty.

The set  $\mathcal{S}_{\alpha}^{*}$  is open by the fact that a deformation of the holonomy implies a deformation of the structure [20]. To prove that  $\mathcal{S}_{\alpha}^{*}$  is closed, consider a sequence  $\tau_{n}$  converging in  $\mathcal{S}_{\alpha}^{*}$  to  $\tau_{\infty} \in \mathcal{S}_{\alpha}$ .

The lengths of common perpendiculars between the axis of rotations A, B, C and D defined above equal respectively  $b_1$ ,  $b_2$  and  $\phi$ .

Since  $\tau_{\infty}$  corresponds to a non-degenerated quadrangle, every cone-manifold  $\mathcal{H}_4(\alpha;\tau_n)$  has the quantities  $b_1(\tau_n)$ ,  $b_2(\tau_n)$  and  $\phi(\tau_n)$  uniformly bounded below away from zero. By the arguments similar to those of Theorem 2, we obtain that  $\mathcal{H}_4(\alpha;\tau_{\infty})$  is a non-degenerate spherical cone-manifold. Thus  $\tau_{\infty}$  belongs to  $\mathcal{S}_{\alpha}^*$ . Hence  $\mathcal{S}_{\alpha}^*$  is closed.

Finally, we obtain that  $S_{\alpha}^* = S_{\alpha}$ . Thus, while  $\tau$  varies over  $(\frac{\alpha - \pi}{2}, \frac{\pi}{2})$  the cone-manifold  $\mathcal{H}_4(\alpha; \tau)$  does not collapse.  $\square$ 

The following lemma shows that the interval  $(\frac{\alpha-\pi}{2}, \frac{\pi}{2})$  represents a part of the deformation space for possible spherical structures on  $\mathcal{H}_4(\alpha; \tau)$ .

**Lemma 9** The cone-manifolds  $\mathcal{H}_4(\alpha; \tau_1)$  and  $\mathcal{H}_4(\alpha; \tau_2)$  with  $\pi < \alpha < 2\pi$  and  $\frac{\alpha - \pi}{2} < \tau_1, \tau_2 < \frac{\pi}{2}$  are not isometric if  $\tau_1 \neq \tau_2$ .

**Proof.** If the cone-manifolds  $\mathcal{H}_4(\alpha; \tau_1)$  and  $\mathcal{H}_4(\alpha; \tau_2)$  were isometric, then their holonomy maps  $\widetilde{\rho}_i$ , i=1,2 would be conjugated representations of  $\Gamma = \pi_1(\mathbb{S}^3 \setminus \mathcal{H}_4)$  into  $SU_2(\mathbb{C}) \times SU_2(\mathbb{C})$ . Then the mutual distances between the axis of rotations  $A_i$ ,  $B_i$ ,  $C_i$  and  $D_i$ , i=1,2, coming from the holonomy maps  $\widetilde{\rho}_1$  and  $\widetilde{\rho}_2$  would be equal for the corresponding pairs. From Lemma 3, it follows that the common perpendicular length for the given fibres  $C_1$  and  $C_2$  is half the distance between the images of  $C_1$  and  $C_2$  under the Hopf map. By applying Lemmas 3 and 8 to the base quadrangle P of  $\mathcal{H}_4(\alpha; \tau_i)$ , i=1,2 one makes sure that the inequality  $\tau_1 \neq \tau_2$  implies the inequality for the lengths of corresponding common perpendiculars.  $\square$ 

Note, that by the Schläfli formula the volume of  $\mathcal{H}_4(\alpha)$  remains the same under any deformation preserving cone angles. Then the formulas for the volume and the singular stratum length follow from the covering properties of  $\mathcal{H}_4(\alpha) \xrightarrow{4:1} \mathbb{L}_4(\alpha, \frac{\pi}{2})$  and Theorem 2 of [17]. Thus, Theorem 3 is proven.  $\square$ 

## References

- [1] BOILEAU, M., LEEB, B., PORTI, J. "Uniformization of small 3-orbifolds". C.R. Acad. Sci. Paris Se'r. I Math. 332(1), 57-62 (2001).
- [2] Boileau, M., Leeb, B., Porti, J. "Geometrization of 3-dimensional orbifolds". Ann. Math. **162**(1), 195-250 (2005).
- [3] Burago, Yu., Gromov, M., Perelman, G. "A. D. Aleksandrov spaces with curvature bounded below". Russian Math. Surveys 47, 1-58 (1992).
- [4] BURDE, G., MURASUGI, K. "Links and Seifert fiber spaces". Duke Math. J. 37(1), 89-93 (1970).
- [5] Casson, A. "An example of weak non-rigidity for cone manifolds with vertices". Talk at the Third MSJ regional workshop, Tokyo, 1998.
- [6] COOPER, D., HODGSON, C., KERCKHOFF, S. "Three-dimensional orbifolds and cone-manifolds", Postface by S. Kojima. Tokyo: Mathematical Society of Japan, 2000. (MSJ Memoirs; 5)
- [7] Culler, M. "Lifting representations to covering groups". Adv. Math. **59**(1), 64-70 (1986).
- [8] Dunbar, W.D. "Geometric orbifolds". Rev. Mat. Univ. Complut. Madrid 1, 67-99 (1988).
- [9] GLUCK, H., ZILLER, W. "The geometry of the Hopf fibrations". L'Enseign. Math. **32**, 173-198 (1986).
- [10] GOLDMAN, W. "Ergodic Theory on Moduli Spaces". Ann. Math. **146**(3), 475-507 (1997).
- [11] HILDEN, H.M., LOZANO, M.T., MONTESINOS-AMILIBIA, J.-M. "Volumes and Chern-Simons invariants of cyclic coverings over rational knots". Proceedings of the 37-th Taniguchi Symposium on Topology and Teichmuller Spaces held in Finland, July 1995, ed. by Sadayoshi Kojima et al. (1996), 31-35.

- [12] Hodgson, C. "Degeneration and regeneration of Hyperbolic Structures on Three-Manifolds". Princeton: Thesis, 1986.
- [13] Hodgson, C., Kerckhoff, S. "Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery". J. Differential Geom. 48(1), 1-59 (1998).
- [14] HOPF, H. "Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche". Math. Ann. **104**, 637-665 (1931).
- [15] IZMESTIEV, I. "Examples of infinitesimally flexible 3-dimensional hyperbolic cone-manifolds". J. Math. Soc. Japan 63(2), 581-598 (2011); arXiv:0910.2876.
- [16] KOJIMA, S. "Deformations of hyperbolic 3-cone-manifolds". J. Diff. Geom. 49(3), 469-516 (1998).
- [17] KOLPAKOV, A.A., MEDNYKH, A.D. "Spherical structures on torus knots and links". Siberian Math. J. **50**(5), 856-866 (2009).
- [18] Montcouquiol, G. "Deformation of hyperbolic convex polyhedra and 3-cone-manifolds". arXiv:0903.4743.
- [19] Mostow, G.D. "Quasi-conformal mappings in n-space and the rigidity of hyperbolic space forms". Inst. Hautes Etudes Sci. Publ. Math. 34, 53-104 (1968).
- [20] PORTI, J. "Regenerating hyperbolic and spherical cone structures from Euclidean ones". Topology **37**(2), 365-392 (1998).
- [21] PORTI, J. "Regenerating hyperbolic cone structures from Nil". Geom. Topol. 6, 815-852 (2002).
- [22] PORTI, J. "Spherical cone structures on 2-bridge knots and links". Kobe J. Math. 21(1-2), 61-70 (2004).
- [23] PORTI, J. "Regenerating hyperbolic cone 3-manifolds from dimension 2". arXiv:1003.2494.
- [24] Prasad, G. "Strong rigidity of **Q**-rank 1 lattices". Invent. Math. **21**, 255-286 (1973).
- [25] RATCLIFFE, J. "Foundations of hyperbolic manifolds". New York: Springer-Verlag, 1994. (Graduate Texts in Math.; 149).

- [26] DE RHAM, G. "Reidemeister's torsion invariant and rotations of  $S^n$ "; in Differential Analysis, Bombay Colloq., Oxford Univ. Press, London, 1964, 27-36.
- [27] ROTHENBERG, M. "Torsion invariants and finite transformation groups". Proceedings of Symposia in Pure Math. **32**, 267-311 (1978).
- [28] SCHLENKER, J.-M. "Dihedral angles of convex polyhedra". Discrete Comput. Geom. 23, 409-417 (2000).
- [29] Thurston, W.P. "Geometry and topology of three-manifolds". Princeton Univ., 1979. (Princeton University Lecture Notes)
- [30] VINBERG E.B., ED. "Geometry II. Spaces of Constant Curvature". New York: Springer-Verlag, 1993. (Encyclopaedia of Mathematical Sciences; 29)
- [31] Weiss, H. "Local rigidity of 3-dimensional cone-manifolds". J. Diff. Geom. **71**(3), 437-506 (2005).
- [32] Weiss, H. "Global rigidity of 3-dimensional cone-manifolds". J. Diff. Geom. **76**(3), 495-523 (2007).
- [33] Weiss, H. "The deformation theory of hyperbolic cone-3-manifolds with cone-angles less than  $2\pi$ ". arXiv:0904.4568.

Alexander Kolpakov Department of Mathematics University of Fribourg chemin du Musée 23 CH-1700 Fribourg, Switzerland kolpakov.alexander@gmail.com