EPIMORPHISMS BETWEEN 2-BRIDGE LINK GROUPS: HOMOTOPICALLY TRIVIAL SIMPLE LOOPS ON 2-BRIDGE SPHERES

DONGHI LEE AND MAKOTO SAKUMA

ABSTRACT. We give a complete characterization of those essential simple loops on 2-bridge spheres of 2-bridge links which are null-homotopic in the link complements. By using this result, we describe all upper-meridianpair-preserving epimorphisms between 2-bridge link groups.

CONTENTS

1.	Introduction	1
2.	Main result	3
3.	Presentations of 2-bridge link groups	7
4.	Sequences associated with 2-bridge links	10
5.	Small cancellation conditions for 2-bridge link groups	22
6.	Van Kampen diagrams over 2-bridge link groups	28
7.	Proof of Main Theorem 2.3	32
8.	Relation with a question by Minsky	36
References		37

1. INTRODUCTION

For a knot or a link, K, in S^3 , the fundamental group $\pi_1(S^3 - K)$ of the complement is called the *knot group* or the *link group* of K, and is denoted by G(K). For prime knots, the knot groups are complete invariants for the knot types (see [8]). Moreover we have a partial order on the set of prime

²⁰¹⁰ Mathematics Subject Classification. Primary 57M25, 20F06

The first author was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology(2009-0065798). The second author was supported by JSPS Grants-in-Aid 18340018 and 21654011, and was partially supported by JSPS Core-to-Core Program 18005.

knots, by setting $\tilde{K} \geq K$ if there is an epimorphism $G(\tilde{K}) \rightarrow G(K)$ (see, for example, [32, Proposition 3.2]). Epimorphisms among link groups have received considerable attention and they have been studied in various places in the literature (see [1, 2, 5, 6, 10, 11, 15, 16, 17, 25, 27, 30, 32, 33, 34, 35, 36] and references therein).

In [25], a systematic construction of epimorphisms between 2-bridge link groups was given. The construction is based on a systematic construction of essential simple loops on 2-bridge spheres of 2-bridge links which are null-homotopic in the link complements. Thus the following question naturally arises (see [25, Question 9.1(2)]).

Question 1. Let K be a 2-bridge link, and let S be a 4-times punctured sphere in $S^3 - K$ determined by a 2-bridge sphere. Then which essential simple loops on S are null-homotopic in $S^3 - K$?

It should be noted that each 2-bridge link admits a unique 2-bridge sphere up to isotopy (see [31]), and hence the 4-times punctured sphere S in the above problem is unique up to isotopy.

In this paper, we give a complete answer to the above question (Main Theorem 2.3). In fact, we show that those essential simple loops on S constructed in [25, Corollary 4.7] are the only essential simple loops on S which are nullhomotopic in the 2-bridge link complement. This enables us to describe all epimorphisms between 2-bridge link groups which map the upper meridian pair of the source group to the upper meridian pair of the target group (Main Theorem 2.4). In fact, this theorem says that any such epimorphism is equivalent to that constructed in [25, Theorem 1.1].

To the authors' knowledge, every known pair of 2-bridge knots (\tilde{K}, K) with $\tilde{K} \geq K$ belongs to the list in [25, Theorem 1.1]. Kitano and Suzuki and their coworkers verified this for 2-bridge knots up to 11-crossings in [10, 15, 16]. Gonzaléz-Acũna and Ramírez [6] determined the 2-bridge knots whose knot groups have epimorphisms to the (2, p) torus knot group, and their result implies that every such 2-bridge knot group is isomorphic to one constructed in [25, Theorem 1.1]. In their recent work [1], Boileau, Boyer, Reid and Wang proved Simon's conjecture (see [14, Problem 1.12(D)]) for 2-bridge knot groups, namely they have shown that each 2-bridge knot group surjects onto only finitely many distinct knot groups. To be more precise, they have shown that if a 2-bridge knot group G(K) surjects onto a non-trivial knot group G(K'), then K' is a 2-bridge knot and the epimorphism is induced by a map between the knot complements of non-zero degree. The last condition is satisfied for all epimorphisms in [25, Theorem 1.1]. In fact, they are induced by a very

nice map $(S^3, K) \to (S^3, K')$, called a branched-fold map [25, Theorem 1.2]. Thus it would be natural to expect that any epimorphism between 2-bridge knot groups is equivalent to one in [25, Theorem 1.1]. In fact, some evidence for this conjecture was provided recently by Hoste and Shanahan [11].

Question 1 can be regarded as a special case of the more general question that, for a given link L and a bridge sphere F for L, which essential simple loops on F are null-homotopic in $S^3 - L$. The latter question in turn can be regarded as a variation of the question that, for a given 3-manifold M and its Heegaard surface F, which essential simple loops on F are null-homotopic in M. In [7, Question 5.4], Minsky refined this to a certain question which generalizes Question 1. Thus our result may be regarded as an answer to a special variation of Minsky's question (see Section 8).

The authors would like to thank Norbert A'Campo, Hirotaka Akiyoshi, Brian Bowditch, Danny Calegari, Max Forester, Koji Fujiwara, Yair Minsky, Ser Peow Tan and Caroline Series for stimulating conversations. They also thank the referee for his/her careful reading of the manuscript.

2. Main result

Consider the discrete group, H, of isometries of the Euclidean plane \mathbb{R}^2 generated by the π -rotations around the points in the lattice \mathbb{Z}^2 . Set $(\mathbf{S}^2, \mathbf{P}) := (\mathbb{R}^2, \mathbb{Z}^2)/H$ and call it the *Conway sphere*. Then \mathbf{S}^2 is homeomorphic to the 2-sphere, and \mathbf{P} consists of four points in \mathbf{S}^2 . We also call \mathbf{S}^2 the Conway sphere. Let $\mathbf{S} := \mathbf{S}^2 - \mathbf{P}$ be the complementary 4-times punctured sphere. For each $r \in \hat{\mathbb{Q}} := \mathbb{Q} \cup \{\infty\}$, let α_r be the simple loop in \mathbf{S} obtained as the projection of a line in $\mathbb{R}^2 - \mathbb{Z}^2$ of slope r. Then α_r is *essential* in \mathbf{S} , i.e., it does not bound a disk in \mathbf{S} and is not homotopic to a loop around a puncture. Conversely, any essential simple loop in \mathbf{S} is isotopic to α_r for a unique $r \in \hat{\mathbb{Q}}$. Then r is called the *slope* of the simple loop. Similarly, any simple arc δ in \mathbf{S}^2 joining two different points in \mathbf{P} such that $\delta \cap \mathbf{P} = \partial \delta$ is isotopic to the image of a line in \mathbb{R}^2 of some slope $r \in \hat{\mathbb{Q}}$ which intersects \mathbb{Z}^2 . We call r the *slope* of δ .

A trivial tangle is a pair (B^3, t) , where B^3 is a 3-ball and t is a union of two arcs properly embedded in B^3 which is parallel to a union of two mutually disjoint arcs in ∂B^3 . Let τ be the simple unknotted arc in B^3 joining the two components of t as illustrated in Figure 1. We call it the core tunnel of the trivial tangle. Pick a base point x_0 in int τ , and let (μ_1, μ_2) be the generating pair of the fundamental group $\pi_1(B^3 - t, x_0)$ each of which is represented by a based loop consisting of a small peripheral simple loop around a component of t and a subarc of τ joining the circle to x_0 . For any base point $x \in B^3 - t$, the generating pair of $\pi_1(B^3 - t, x)$ corresponding to the generating pair (μ_1, μ_2) of $\pi_1(B^3 - t, x_0)$ via a path joining x to x_0 is denoted by the same symbol. The pair (μ_1, μ_2) is unique up to (i) reversal of the order, (ii) replacement of one of the members with its inverse, and (iii) simultaneous conjugation. We call the equivalence class of (μ_1, μ_2) the *meridian pair* of the fundamental group $\pi_1(B^3 - t)$.



FIGURE 1. A trivial tangle

By a rational tangle, we mean a trivial tangle (B^3, t) which is endowed with a homeomorphism from $\partial(B^3, t)$ to $(\mathbf{S}^2, \mathbf{P})$. Through the homeomorphism we identify the boundary of a rational tangle with the Conway sphere. Thus the slope of an essential simple loop in $\partial B^3 - t$ is defined. We define the *slope* of a rational tangle to be the slope of an essential loop on $\partial B^3 - t$ which bounds a disk in B^3 separating the components of t. (Such a loop is unique up to isotopy on $\partial B^3 - t$ and is called a *meridian* of the rational tangle.) We denote a rational tangle of slope r by $(B^3, t(r))$. By van Kampen's theorem, the fundamental group $\pi_1(B^3 - t(r))$ is identified with the quotient $\pi_1(\mathbf{S})/\langle \langle \alpha_r \rangle \rangle$, where $\langle \langle \alpha_r \rangle \rangle$ denotes the normal closure.

For each $r \in \hat{\mathbb{Q}}$, the 2-bridge link K(r) of slope r is defined to be the sum of the rational tangles of slopes ∞ and r, namely, $(S^3, K(r))$ is obtained from $(B^3, t(\infty))$ and $(B^3, t(r))$ by identifying their boundaries through the identity map on the Conway sphere (S^2, P) . (Recall that the boundaries of rational tangles are identified with the Conway sphere.) K(r) has one or two components according as the denominator of r is odd or even. We call $(B^3, t(\infty))$ and $(B^3, t(r))$, respectively, the upper tangle and lower tangle of the 2-bridge link. The 2-bridge links are classified by the following theorem of Schubert [31] (cf. [3, 13]). **Theorem 2.1** (Schubert). Two 2-bridge links K(q/p) and K(q'/p') are equivalent (i.e., there is a homeomorphism from S^3 to itself sending K(q/p) to K(q'/p')), if and only if the following conditions hold.

- (1) p = p'.
- (2) Either $q \equiv \pm q' \pmod{p}$ or $qq' \equiv \pm 1 \pmod{p}$.

Let \mathcal{D} be the *Farey tessellation*, that is, the tessellation of the upper half space \mathbb{H}^2 by ideal triangles which are obtained from the ideal triangle with the ideal vertices $0, 1, \infty \in \mathbb{Q}$ by repeated reflection in the edges. Then \mathbb{Q} is identified with the set of the ideal vertices of \mathcal{D} . For each $r \in \hat{\mathbb{Q}}$, let Γ_r be the group of automorphisms of $\mathcal D$ generated by reflections in the edges of $\mathcal D$ with an endpoint r. It should be noted that Γ_r is isomorphic to the infinite dihedral group and the region bounded by two adjacent edges of \mathcal{D} with an endpoint r is a fundamental domain for the action of Γ_r on \mathbb{H}^2 , by virtue of Poincare's fundamental polyhedron theorem (see, for example, [26]). Let Γ_r be the group generated by Γ_r and Γ_{∞} . When $r \in \mathbb{Q} - \mathbb{Z}$, Γ_r is equal to the free product $\Gamma_r * \Gamma_\infty$, having a fundamental domain shown in Figure 2. Otherwise, $\tilde{\Gamma}_r$ is the group generated by reflections in the edges of \mathcal{D} or Γ_{∞} according as $r \in \mathbb{Z}$ or $r = \infty$. It should be noted that Theorem 2.1 says that two 2-bridge links K(r) and K(r') are equivalent if and only if there is an automorphism of \mathcal{D} which sends $\{\infty, r\}$ to $\{\infty, r'\}$. Thus the conjugacy class of the group Γ_r in the automorphism group of \mathcal{D} is uniquely determined by the link K(r).

We recall the following fact ([25, Proposition 4.6 and Corollary 4.7]) which describes the role of $\hat{\Gamma}_r$ in the study of 2-bridge link groups.

Proposition 2.2. For every 2-bridge link K(r), the following holds. If two elements s and s' of $\hat{\mathbb{Q}}$ lie in the same $\hat{\Gamma}_r$ -orbit, then α_s and $\alpha_{s'}$ are homotopic in $S^3 - K(r)$. In particular, if s belongs to the orbit of ∞ or r by $\hat{\Gamma}_r$, then α_s is null-homotopic in $S^3 - K(r)$.

Our main theorem says that the converse to the last statement in the above proposition is valid.

Main Theorem 2.3. The loop α_s is null-homotopic in $S^3 - K(r)$ if and only if s belongs to the $\hat{\Gamma}_r$ -orbit of ∞ or r.

This theorem may be paraphrased as follows, with a detailed reason explained in Section 3.

Main Theorem 2.4. There is an upper-meridian-pair-preserving epimorphism from G(K(s)) to G(K(r)) if and only if s or s+1 belongs to the $\hat{\Gamma}_r$ -orbit of r or ∞ .



FIGURE 2. A fundamental domain of $\overline{\Gamma}_r$ in the Farey tessellation (the shaded domain) for $r = 5/17 = \frac{1}{3 + \frac{1}{2 + \frac{1}{2}}} =: [3, 2, 2].$

Since the if part is [25, Theorem 1.1], the heart of this theorem is the only if part.

The remainder of this paper is organized as follows. In Section 3, we introduce the so-called upper presentation $G(K(r)) = \langle a, b | u_r \rangle$ of a 2-bridge link group, where $\{a, b\}$ is the upper meridian pair of K(r). This upper presentation of a 2-bridge link group will be used throughout this paper. In Section 4, we define two sequences S(r) and T(r) of slope r and two cyclic sequences CS(r) and CT(r) of slope r all of which arise from the single relator u_r of the presentation $G(K(r)) = \langle a, b | u_r \rangle$, and observe several important properties of these sequences so that we can adopt, in the succeeding sections, small cancellation theory which is one of the geometric techniques in combinatorial group theory. In Section 5, we show that the presentation $G(K(r)) = \langle a, b | u_r \rangle$, where 0 < r < 1, satisfies small cancellation conditions C(4) and T(4). In Section 6, by applying the Curvature Formula of Lyndon and Schupp (see [24]) to van Kampen diagrams over $G(K(r)) = \langle a, b | u_r \rangle$, we obtain that if α_s is null-homotopic in $S^3 - K(r)$, where 0 < r < 1, then the cyclic word (u_s) contains some particular part of the the cyclic word $(u_r^{\pm 1})$. In Section 7, we prove the only if part of Main Theorem 2.3 by showing that if a rational number s belongs to a natural fundamental domain of the action of Γ_r on the domain of discontinuity of $\hat{\Gamma}_r$, then α_s is not null-homotopic in $S^3 - K(r)$. In

the final section, Section 8, we describe the relation of Main Theorem 2.3 with the question raised by Minsky in [7, Question 5.4].

3. Presentations of 2-bridge link groups

In this section, we introduce the upper presentation of a 2-bridge link group which we shall use throughout this paper. By van Kampen's theorem, the link group $G(K(r)) = \pi_1(S^3 - K(r))$ is identified with $\pi_1(\mathbf{S})/\langle\langle \alpha_{\infty}, \alpha_r \rangle\rangle$. We call the image in the link group of the meridian pair of the fundamental group $\pi_1(B^3 - t(\infty))$ (resp. $\pi_1(B^3 - t(r))$ the upper meridian pair (resp. lower meridian pair). The link group is regarded as the quotient of the rank 2 free group, $\pi_1(B^3 - t(\infty)) \cong \pi_1(\mathbf{S})/\langle\langle \alpha_{\infty} \rangle\rangle$, by the normal closure of α_r . This gives a one-relator presentation of the link group, which is called the upper presentation (see [4]).



FIGURE 3. $\pi_1(B^3 - t(\infty), x_0) = F(a, b)$, where a and b are represented by μ_1 and μ_2 , respectively.

To find the upper presentation of G(K(r)) explicitly, let a and b, respectively, be the elements of $\pi_1(B^3 - t(\infty), x_0)$ represented by the oriented loops μ_1 and μ_2 based on x_0 as illustrated in Figure 3. Then $\{a, b\}$ forms the meridian pair of $\pi_1(B^3 - t(\infty))$, which is identified with the free group F(a, b). Note that μ_i intersects the disk, δ_i , in B^3 bounded by a component of $t(\infty)$ and the essential arc, γ_i , on $\partial(B^3, t(\infty)) = (\mathbf{S}^2, \mathbf{P})$ of slope 1/0, in Figure 3. Obtain a word u_r in $\{a, b\}$ by reading the intersection of the (suitably oriented) loop α_r with $\gamma_1 \cup \gamma_2$, where a positive intersection with γ_1 (resp. γ_2) corresponds to a(resp. b). Then the cyclic word (u_r) represents the free homotopy class of α_r (see Section 4 for the precise definition of a cyclic word). It then follows that

$$G(K(r)) = \pi_1(S^3 - K(r)) \cong \pi_1(B^3 - t(\infty)) / \langle \langle \alpha_r \rangle \rangle$$
$$\cong F(a, b) / \langle \langle u_r \rangle \rangle \cong \langle a, b | u_r \rangle.$$

If $r \neq \infty$, then α_r intersects γ_1 and γ_2 alternately, and hence a and b appear in (u_r) alternately. It is known that there is a nice formula to find u_r as follows (see [28, Proposition 1]).

Lemma 3.1. Let p and q be relatively prime positive integers such that $p \ge 1$. For $1 \le i \le p - 1$, let

$$\epsilon_i = (-1)^{\lfloor iq/p \rfloor}.$$

where |x| is the greatest integer not exceeding x.

(1) If p is odd, then

$$u_{q/p} = a\hat{u}_{q/p}b^{(-1)^q}\hat{u}_{q/p}^{-1},$$

where $\hat{u}_{q/p} = b^{\epsilon_1} a^{\epsilon_2} \cdots b^{\epsilon_{p-2}} a^{\epsilon_{p-1}}$. (2) If p is even, then $u_{q/p} = a \hat{u}_{q/p} a^{-1} \hat{u}_{q/p}^{-1}$,

where $\hat{u}_{q/p} = b^{\epsilon_1} a^{\epsilon_2} \cdots a^{\epsilon_{p-2}} b^{\epsilon_{p-1}}$.

Remark 1. (1) The word $\hat{u}_{q/p}$ is obtained from the open line-segment of slope q/p extending from (0,0) to (p,q) by "reading" its intersection with the vertical lattice lines (see Figure 4). The open line-segment cuts the vertical lattice line x = i at the point P_i with height iq/p. Note that $\lfloor iq/p \rfloor$ is the height of the integer lattice point just beneath P_i . Each time the line passes through another horizontal lattice line, the signs of the ϵ_i 's change. Similarly, the word $u_{q/p}$ can be read from the closed line-segment which is obtained by slightly shifting the closed line-segment of slope q/p joining (0,0) with (2p, 2q) to the upper-left direction (cf. Proof of Lemma 4.7).

(2) For r = 0/1 and r = 1/0, we have $u_{0/1} = ab$ and $u_{1/0} = 1$.

In the remainder of this section, we prove Main Theorem 2.4 by assuming Main Theorem 2.3. To this end we prepare the following lemma.

Lemma 3.2. (1) Let φ be the automorphism of the free group $\pi_1(B^3 - t(\infty)) = \langle a, b \rangle$ which sends the generating pair (a, b) to (a^{-1}, b^{-1}) , (b, a) or (b^{-1}, a^{-1}) . Then $\varphi(u_s)$ is conjugate to u_s or u_s^{-1} for any $s \in \hat{\mathbb{Q}}$.

(2) Let φ be the automorphism of the free group $\pi_1(B^3 - t(\infty)) = \langle a, b \rangle$ which sends the generating pair (a, b) to (a, b^{-1}) , (a^{-1}, b) , (b^{-1}, a) or (b, a^{-1}) . Then $\varphi(u_s)$ is conjugate to u_{s+1} or u_{s+1}^{-1} for any $s \in \hat{\mathbb{Q}}$.



FIGURE 4. The line of slope 4/7 gives $\hat{u}_{4/7} = ba^{-1}b^{-1}aba^{-1}$, so the free homotopy class of $\alpha_{4/7}$ is represented by the cyclic word $(u_{4/7}) = (a\hat{u}_{4/7}b\hat{u}_{4/7}^{-1}) = (aba^{-1}b^{-1}aba^{-1}bab^{-1}a^{-1}bab^{-1})$. Since the inverse image of γ_1 (resp. γ_2) in \mathbb{R}^2 is the union of the single arrowed (resp. double arrowed) vertical edges, a positive intersection with a single arrowed (resp. double arrowed) edge corresponds to a (resp. b).

Proof. (1) Observe that $(B^3, t(\infty))$ admits a natural $(\mathbb{Z}/2\mathbb{Z})^2$ -action, whose generators induce the automorphisms of $\pi_1(B^3-t(\infty))$ sending (a,b) to (a^{-1},b^{-1}) and (b,a), respectively. Moreover, the action preserves the isotopy class of the (unoriented) loop α_s for every $s \in \hat{\mathbb{Q}}$. Since any automorphism φ satisfying the assumption is induced by an element of the $(\mathbb{Z}/2\mathbb{Z})^2$ -action, we obtain the desired result.

(2) Let φ be an automorphism of $\pi_1(B^3 - t(\infty))$ satisfying the assumption. Then it is a composition of an automorphism in (1) and the automorphism, ψ , sending (a, b) to (a, b^{-1}) . Observe that ψ is induced by the half-Dehn twist along the meridian disk of $(B^3, t(\infty))$ and that the half-Denn twist maps α_s to α_{s+1} . Hence we see $\psi(u_s) = u_{s+1}$. This, together with (1), implies the desired result.

Proof of Main Theorem 2.4 assuming Main Theorem 2.3. The if part is essentially equivalent to [25, Theorem 1.1] and is proved as follows. If s belongs to the $\hat{\Gamma}_r$ -orbit of r or ∞ , then Main Theorem 2.3 implies that $u_s = 1$ in $G(K(r)) = \langle a, b | u_r \rangle$. Thus there is an epimorphism from $G(K(s)) = \langle a, b | u_s \rangle$ to $G(K(r)) = \langle a, b | u_r \rangle$ which sends the upper-meridian-pair (a, b) of G(K(s))to the upper-meridian-pair (a, b) of G(K(r)). To prove the remaining case, note that there is a homeomorphism $g: (S^3, K(s)) \to (S^3, K(s+1))$ preserving the upper/lower tangles, such that the restriction of g to $(B^3, t(\infty))$ is a half-Dehn twist. Thus g induces an isomorphism from $G(K(s)) = \langle a, b | u_s \rangle$ to $G(K(s+1)) = \langle a, b | u_{s+1} \rangle$ which sends the upper-meridian-pair (a, b) of G(K(s)) to the upper-meridian-pair (a, b^{-1}) of G(K(s+1)). So, if s+1 belongs to the $\hat{\Gamma}_r$ -orbit of r or ∞ , then we have an epimorphism $G(K(s)) \cong$ $G(K(s+1)) \to G(K(r))$ sending (a, b) to (a, b^{-1}) .

Next, we prove the only if part. Suppose that there is an upper-meridianpair preserving epimorphism f from $G(K(s)) = \langle a, b | u_s \rangle$ to $G(K(r)) = \langle a, b | u_r \rangle$. Then f lifts to an automorphism φ of the free group $\pi_1(B^3 - t(\infty)) = \langle a, b \rangle$ satisfying the condition in Lemma 3.2, modulo post composition of an inner-automorphism. Thus $\varphi(u_s)$ is conjugate to u_s , u_s^{-1} , u_{s+1} or u_{s+1}^{-1} by Lemma 3.2. Since φ is a lift of the homomorphism f, u_s or u_{s+1} represents the trivial element of G(K(r)), accordingly. Hence, by Main Theorem 2.3, we see that s or s + 1 belongs to the $\hat{\Gamma}_r$ -orbit of r or ∞ , accordingly.

4. Sequences associated with 2-bridge links

In this section, we define two sequences S(r) and T(r) of slope r and two cyclic sequences CS(r) and CT(r) of slope r all of which arise from the single relator u_r of the presentation $G(K(r)) = \langle a, b | u_r \rangle$ given in Section 3, and observe several important properties of these sequences, so that we can adopt small cancellation theory in the succeeding sections.

We first fix some definitions and notation. Let X be a set. By a word in X, we mean a finite sequence $x_1^{\epsilon_1} x_2^{\epsilon_2} \cdots x_n^{\epsilon_n}$ where $x_i \in X$ and $\epsilon_i = \pm 1$. Here we call $x_i^{\epsilon_i}$ the *i*-th letter of the word. For two words u, v in X, by $u \equiv v$ we denote the visual equality of u and v, meaning that if $u = x_1^{\epsilon_1} \cdots x_n^{\epsilon_n}$ and $v = y_1^{\delta_1} \cdots y_m^{\delta_m} (x_i, y_j \in X; \epsilon_i, \delta_j = \pm 1)$, then n = m and $x_i = y_i$ and $\epsilon_i = \delta_i$ for each $i = 1, \ldots, n$. For example, two words $x_1 x_2 x_2^{-1} x_3$ and $x_1 x_3 (x_i \in X)$ are not visually equal, though they are equal as elements of the free group with basis X. The length of a word v is denoted by |v|. A word v in X is said to be reduced if v does not contain xx^{-1} or $x^{-1}x$ for any $x \in X$. A word is called cyclically reduced if all its cyclic permutations are reduced. A cyclic word is defined to be the set of all cyclic permutations of a cyclically reduced word v. Also by $(u) \equiv (v)$ we mean the visual equality of two cyclic words (u) and (v). In fact, $(u) \equiv (v)$ if and only if v is visually a cyclic shift of u.

Definition 1. (1) Let v be a reduced word in $\{a, b\}$. Decompose v into

$$v \equiv v_1 v_2 \cdots v_t,$$

where, for each i = 1, ..., t - 1, all letters in v_i have positive (resp. negative) exponents, and all letters in v_{i+1} have negative (resp. positive) exponents. Then the sequence of positive integers $S(v) := (|v_1|, |v_2|, ..., |v_t|)$ is called the *S*-sequence of v.

(2) Let (v) be a cyclic word in $\{a, b\}$. Decompose (v) into

$$(v) \equiv (v_1 v_2 \cdots v_t),$$

where all letters in v_i have positive (resp. negative) exponents, and all letters in v_{i+1} have negative (resp. positive) exponents (taking subindices modulo t). Then the *cyclic* sequence of positive integers $CS(v) := ((|v_1|, |v_2|, \ldots, |v_t|))$ is called the *cyclic S-sequence of* (v). Here the double parentheses denote that the sequence is considered modulo cyclic permutations.

(3) A reduced word v in $\{a, b\}$ is said to be *alternating* if $a^{\pm 1}$ and $b^{\pm 1}$ appear in v alternately, i.e., neither $a^{\pm 2}$ nor $b^{\pm 2}$ appears in v. A cyclic word (v) is said to be *alternating* if all cyclic permutations of v are alternating. In the latter case, we also say that v is cyclically alternating.

The following proposition is obvious from the definition.

Proposition 4.1. (1) An alternating word in $\{a, b\}$ is completely determined by the initial letter and the associated S-sequence.

(2) Let v be a cyclically reduced word in $\{a, b\}$ of length ≥ 2 . Then the S-sequence S(v) represents the cyclic S-sequence CS(v) of (v) if and only if the initial exponent of v is different from the terminal exponent of v.

Definition 2. For a rational number r with $0 < r \leq 1$, let u_r be the word in $\{a, b\}$ defined in Lemma 3.1. Then the symbol S(r) (resp. CS(r)) denotes the S-sequence $S(u_r)$ of u_r (resp. cyclic S-sequence $CS(u_r)$ of (u_r)), which is called the S-sequence of slope r (resp. the cyclic S-sequence of slope r).

We shall first state Propositions 4.2, 4.3, 4.4 and 4.5 below concerning the sequences defined in the above, and then prove the propositions in the remainder of this section. Propositions 4.4 and 4.5 play crucial roles in the proof of Main Theorem 2.3. Though we need those propositions only for the sequences S(r) and CS(r) with $0 < r \le 1$, we need to extend the definitions of S(r) and CS(r) to an arbitrary positive rational number r (Definition 4), in order to prove these propositions. Thus Propositions 4.2, 4.3, 4.4 and 4.5 below should be regarded as propositions for every positive rational number r.

Throughout the remainder of this section, r = q/p denotes a positive rational number, where p and q are relatively prime positive integers. Then r has a continued fraction expansion

$$r = q/p = \frac{1}{m_1 + \frac{1}{m_2 + \dots + \frac{1}{m_k}}} =: [m_1, m_2, \dots, m_k],$$

where $k \ge 1$, $m_1 \in \mathbb{Z}_+ \cup \{0\}$, $(m_2, \ldots, m_k) \in (\mathbb{Z}_+)^{k-1}$ and $m_k \ge 2$ unless k = 1. Note that $m_1 \ge 1$ if $0 < r \le 1$, whereas $m_1 = 0$ if r > 1.

Proposition 4.2. For the positive rational number r = q/p, the sequence S(r) has length 2q, and it represents the cyclic sequence CS(r). Moreover the cyclic sequence CS(r) is invariant by the half-rotation; that is, if $s_j(r)$ denotes the *j*-th term of S(r) $(1 \le j \le 2q)$, then $s_j(r) = s_{q+j}(r)$ for every integer j $(1 \le j \le q)$.

Proposition 4.3. For the positive rational number $r = [m_1, m_2, \ldots, m_k]$, putting $m = m_1$, we have the following.

- (1) Suppose k = 1, i.e., r = 1/m. Then S(r) = (m, m).
- (2) Suppose $k \ge 2$. Then each term of S(r) is either m or m + 1, and S(r) begins with m + 1 and ends with m. Moreover, the following hold.
 - (a) If $m_2 = 1$, then no two consecutive terms of S(r) can be (m, m), so there is a sequence of positive integers (t_1, t_2, \ldots, t_s) such that

$$S(r) = (t_1 \langle m+1 \rangle, m, t_2 \langle m+1 \rangle, m, \dots, t_s \langle m+1 \rangle, m).$$

Here, the symbol " $t_i \langle m+1 \rangle$ " represents t_i successive m+1's.

(b) If $m_2 \ge 2$, then no two consecutive terms of S(r) can be (m+1, m+1), so there is a sequence of positive integers (t_1, t_2, \ldots, t_s) such that

$$S(r) = (m+1, t_1 \langle m \rangle, m+1, t_2 \langle m \rangle, \dots, m+1, t_s \langle m \rangle).$$

Here, the symbol "
$$t_i \langle m \rangle$$
" represents t_i successive m's.

Remark 2. In [9], Hirasawa and Murasugi defined, as one of the key notions of their paper, the sequence of signs for a pair (p, q), which actually gives rise to our S-sequence S(q/p) of slope q/p. They also observed several properties for the sequence of signs for (p, q), which are very similar to the properties of S(q/p) stated in Proposition 4.3.

Definition 3. If $k \ge 2$, the symbol T(r) denotes the sequence (t_1, t_2, \ldots, t_s) in Proposition 4.3, which is called the *T*-sequence of slope *r*. The symbol CT(r)denotes the cyclic sequence represented by T(r), which is called the cyclic *T*-sequence of slope *r*.

Example 1. (1) Let r = 10/37 = [3, 1, 2, 3]. By Lemma 3.1, we see that the *S*-sequence of \hat{u}_r is

$$S(\hat{u}_r) = (3, 4, 4, 3, 4, 4, 3, 4, 4, 3).$$

By the formula for u_r in Lemma 3.1, this implies

$$S(r) = S(u_r) = (\underbrace{4, 4, 4}_{3}, 3, \underbrace{4, 4}_{2}, 3, \underbrace{4, 4}_{2}, 3, \underbrace{4, 4}_{3}, 3, \underbrace{4, 4}_{2}, 3, \underbrace{4, 4}_{2}, 3, \underbrace{4, 4}_{2}, 3).$$

So T(r) = (3, 2, 2, 3, 2, 2) and CT(r) = ((3, 2, 2, 3, 2, 2)).

(2) Let r = 8/35 = [4, 2, 1, 2]. Again by Lemma 3.1, we obtain that the S-sequence of \hat{u}_r is

$$S(\hat{u}_r) = (4, 4, 5, 4, 4, 5, 4, 4).$$

By the formula for u_r in Lemma 3.1, this implies

$$S(r) = S(u_r) = (5, \underbrace{4}_1, 5, \underbrace{4, 4}_2, 5, \underbrace{4, 4}_2, 5, \underbrace{4, 4}_1, 5, \underbrace{4, 4}_2, 5, \underbrace{4, 4}_2, 5, \underbrace{4, 4}_2).$$

$$T(r) = (1, 2, 2, 1, 2, 2) \text{ and } CT(r) = ((1, 2, 2, 1, 2, 2)).$$

Proposition 4.4. For the rational number $r = [m_1, m_2, ..., m_k]$, let r' be the rational number defined as

$$r' = \begin{cases} [m_3, \dots, m_k] & \text{if } m_2 = 1; \\ [m_2 - 1, m_3, \dots, m_k] & \text{if } m_2 \ge 2. \end{cases}$$

Then we have

So

$$T(r) = \begin{cases} S(r') & \text{if } m_2 = 1; \\ \overleftarrow{S}(r') & \text{if } m_2 \geq 2, \end{cases}$$

where $\overleftarrow{S}(r')$ denotes the sequence obtained from S(r') reversing its order.

Proposition 4.5. For the positive rational number $r = [m_1, m_2, \ldots, m_k]$, putting $m = m_1$, the sequence S(r) has a decomposition (S_1, S_2, S_1, S_2) which satisfies the following.

- (1) Each S_i is symmetric, i.e., the sequence obtained from S_i by reversing the order is equal to S_i . (Here, S_1 is empty if k = 1.)
- (2) Each S_i occurs only twice in the cyclic sequence CS(r).
- (3) S_1 begins and ends with m + 1.

(4) S_2 begins and ends with m.

Corollary 4.6. CS(r) is symmetric, i.e., the cyclic sequence obtained from CS(r) by reversing its cyclic order is equivalent to CS(r) (as a cyclic sequence). In particular, in Proposition 4.4, we actually have

$$CT(r) = CS(r')$$

Example 2. (1) Let r = 10/37 = [3, 1, 2, 3]. Recall from Example 1 that

S(r) = (4, 4, 4, 3, 4, 4, 3, 4, 4, 3, 4, 4, 3, 4, 4, 3, 4, 4, 3, 4, 4, 3).

Putting $S_1 = (4, 4, 4)$ and $S_2 = (3, 4, 4, 3, 4, 4, 3)$, we have

$$S(r) = (S_1, S_2, S_1, S_2),$$

where S_1 and S_2 satisfy all the assertions in Proposition 4.5.

(2) Let r = 8/35 = [4, 2, 1, 2]. Recall also from Example 1 that

$$S(r) = (5, 4, 5, 4, 4, 5, 4, 4, 5, 4, 5, 4, 4, 5, 4, 4).$$

Putting $S_1 = (5, 4, 5)$ and $S_2 = (4, 4, 5, 4, 4)$, we also have

$$S(r) = (S_1, S_2, S_1, S_2)$$

where S_1 and S_2 satisfy all the assertions in Proposition 4.5.

The remainder of this section is devoted to the proof of the propositions. We first prepare a few symbols. For a real number t, let $\lfloor t \rfloor$ be the greatest integer not exceeding t, $\lfloor t \rfloor_*$ the greatest integer smaller than t, and $\lceil t \rceil^*$ be the smallest integer greater than t. Then, $\lfloor t \rfloor_* = \lfloor t \rfloor < \lceil t \rceil^*$ for a non-integral real number t, whereas $n - 1 = \lfloor n \rfloor_* < \lfloor n \rfloor < \lceil n \rceil^* = n + 1$ for an integer n. We also note that $\lfloor t + n \rfloor_* = \lfloor t \rfloor_* + n$ and $\lceil t + n \rceil^* = \lceil t \rceil^* + n$ for every $t \in \mathbb{R}$ and $n \in \mathbb{Z}$. By using this symbol, we have the following formula for the relator u_r in the group presentation of G(K(r)) given in Section 3.

Lemma 4.7. For the positive rational number r = q/p, the word u_r is given by the following formula:

$$u_r = a^{\varepsilon_1} b^{\varepsilon_2} \cdots a^{\varepsilon_{2p-1}} b^{\varepsilon_{2p}},$$

where $\varepsilon_i = (-1)^{\lceil (i-1)q/p \rceil^* - 1}$. In particular, u_r is alternating and cyclically reduced.

To prove Lemma 4.7, let L(r) be the line in \mathbb{R}^2 of slope r passing through the origin, and let $L^+(r)$ be the line obtained by translating L(r) by the vector $(0, \eta)$ for sufficiently small positive real number η . Then $L^+(r)$ lies in $\mathbb{R}^2 - \mathbb{Z}^2$ and projects to the simple loop α_r . Pick a base point, z, from the intersection of $L^+(r)$ with the second quadrant, and consider the sub-line-segment of $L^+(r)$ bounded by z and z + (2p, 2q). Then it forms a fundamental domain of the covering $L^+(r) \to \alpha_r$, and the word u_r is obtained by reading the intersection of the line-segment with the vertical lattice lines. To be precise, for each integer $0 \le i \le 2p-1$, let P_i^+ be the intersection of the line-segment with the vertical lattice line x = i. We define the *letter* at P_i^+ to be a or b according as P_i^+ lies on a vertical edge with a single arrow or double arrow in Figure 4, namely according as i is even or odd. We define the sign of P_i^+ to be +1 or -1 according as the corresponding arrow is upward or downward. Then the letter and the sign of P_i^+ , respectively, give the letter and the exponent of the (i+1)th term of the word u_r for each $0 \le i \le 2p-1$. To describe the sign of P_i^+ , note that the y-coordinate of P_i^+ is equal to $iq/p + \eta$, where η is a sufficiently small positive real. Thus it is contained in the open interval $(\lceil iq/p \rceil^* - 1, \lceil iq/p \rceil^*)$. Thus the corresponding arrow is upward or downward according as $\lceil iq/p \rceil^* - 1$ is even or odd. Hence the sign of P_i^+ is equal to $(-1)^{\lceil iq/p \rceil^* - 1}$. This means that the exponent, ε_i , of the *i*-th term of u_r is $(-1)^{\lceil (i-1)q/p \rceil^* - 1}$. Thus we obtain the first assertion of Lemma 4.7. The second assertion is a direct consequence of the first assertion.

Remark 3. For $1 \leq i \leq p-1$, the sign $\epsilon_i = (-1)^{\lfloor iq/p \rfloor}$ in Lemma 3.1 is of course equal to the sign $\varepsilon_{i+1} = (-1)^{\lceil iq/p \rceil^* - 1}$ in Lemma 4.7

Lemma 4.8. If $0 < r \le 1$, then the sequence S(r) has length 2q, and its *j*-th term $s_j(r)$ is given by the following formula $(1 \le j \le 2q)$:

$$s_j(r) = \#\{i \in \{0, 1, \dots, 2q - 1\} \mid P_i^+ \in \mathbb{R} \times (j - 1, j)\}$$

= $\#\{i \in \{0, 1, \dots, 2q - 1\} \mid \lceil iq/p \rceil^* = j\}$
= $|jp/q|_* - |(j - 1)p/q|_*,$

where # denotes the number of elements of the set.

Proof. Suppose $0 < r \leq 1$. Then, for each integer j with $1 \leq j \leq 2q$, the horizontal strip $\mathbb{R} \times (j-1,j)$ contains some P_i^+ , namely, the right hand side of the first identity is a positive integer. By this fact and by the above geometric description of u_r and the definition of $S(r) = S(u_r)$, we see that S(r) has length 2q and that $s_j(r)$ is equal to the number of the points P_i^+ 's which are contained in the horizontal strip $\mathbb{R} \times (j-1,j)$. So we obtain the first identity. As noted in the preceding argument, the condition $P_i^+ \in \mathbb{R} \times (j-1,j)$ is equivalent to the condition $j-1 < iq/p + \eta < j$, where η is a sufficiently small positive real. This condition is equivalent to the condition that $\lceil iq/p \rceil^* = j$.

Thus we obtain the second identity of the lemma. To show the last identity, note that the above condition is equivalent to the condition

 $(j-1)p/q - \eta' < i < jp/q - \eta'$ for a sufficiently small positive real η' . This in turn is equivalent to the following condition:

$$\lfloor (j-1)p/q \rfloor_* < i \le \lfloor jp/q \rfloor_*.$$

Hence we have $s_j(r) = \lfloor jp/q \rfloor_* - \lfloor (j-1)p/q \rfloor_*$, completing the proof of Lemma 4.8.

The above argument also shows that the three numbers on the right hand side of the identity in the above lemma are equal even if r > 1. Thus the following definition makes sense.

Definition 4. We extend the definition of S(r), CS(r), T(r) and CT(r) to an arbitrary positive rational number r by the formula in the above definition. Namely, for a positive rational number r = q/p, the *S*-sequence of slope r, S(r), is defined by

$$S(r) = (s_1(r), s_2(r), \dots, s_{2q}(r)),$$

where

$$s_j(r) = \#\{i \in \{0, 1, \dots, 2q - 1\} \mid P_i^+ \in \mathbb{R} \times (j - 1, j)\} \\ = \#\{i \in \{0, 1, \dots, 2q - 1\} \mid \lceil iq/p \rceil^* = j\} \\ = \lfloor jp/q \rfloor_* - \lfloor (j - 1)p/q \rfloor_*.$$

The cyclic S-sequence, CS(r), the T-sequence, T(r), and the cyclic T-sequence, CT(r), of slope r are defined from the above S(r) as in Definitions 2 and 3.

Remark 4. Though the word u_r for r > 1 is already defined and given by Lemma 4.7, the sequence $S(u_r)$ is different from the sequence S(r). In fact, $S(u_r)$ consists of only positive integers, whereas S(r) may contain 0.

Proof of Proposition 4.2. By Lemma 4.8 and Definition 4, S(r) has length 2q. Since u_r begins with a and ends with b^{-1} (see Lemma 4.7), it follows that the sequence S(r) represents the cyclic sequence CS(r) (cf. Proposition 4.1(2)). The symmetry $s_{q+j}(r) = s_j(r)$ is proved as follows:

$$s_{q+j}(r) = \lfloor (q+j)p/q \rfloor_* - \lfloor (q+j-1)p/q \rfloor_*$$
$$= \lfloor p + (jp/q) \rfloor_* - \lfloor p + (j-1)p/q \rfloor_*$$
$$= \lfloor jp/q \rfloor_* - \lfloor (j-1)p/q \rfloor_*$$
$$= s_j(r).$$

We note that the symmetry also follows from the fact that the translation of \mathbb{R}^2 by the vector (p,q) preserves the line $L^+(r)$ and maps the horizontal strip bounded by lattice lines to another such strip.

For the positive rational number $r = q/p = [m_1, m_2, \ldots, m_k]$, let c be the non-negative integer defined by

 $p = m_1 q + c.$

If k = 1 then c = 0, and if $k \ge 2$ then 0 < c < q.

Lemma 4.9. We have the following continued fraction expansions:

$$q/c = [0, m_2, m_3, \dots, m_k],$$

$$c/q = [m_2, m_3, \dots, m_k],$$

$$(q-c)/c = [m_3, \dots, m_k] \qquad if \ m_2 = 1,$$

$$c/(q-c) = [m_2 - 1, m_3, \dots, m_k],$$

$$q/(q-c) = [0, 1, m_2 - 1, m_3, \dots, m_k].$$

Proof. Since $q/p = [m_1, m_2, \ldots, m_k]$, we have $p/q = m_1 + [m_2, \ldots, m_k]$. So, $c/q = (p - m_1 q)/q = [m_2, \ldots, m_k]$ and hence $q/c = [0, m_2, m_3, \ldots, m_k]$. Since $q/c = m_2 + [m_3, \ldots, m_k]$, we have $(q - c)/c = (m_2 - 1) + [m_3, \ldots, m_k]$. So, if $m_2 = 1$, we have $(q - c)/c = [m_3, \ldots, m_k]$. It also implies that $c/(q - c) = [m_2 - 1, m_3, \ldots, m_k]$. Thus $q/(q - c) = 1 + c/(q - c) = 1 + [m_2 - 1, m_3, \ldots, m_k]$, and hence $(q - c)/q = [1, m_2 - 1, m_3, \ldots, m_k]$ and $q/(q - c) = [0, 1, m_2 - 1, m_3, \ldots, m_k]$.

Lemma 4.10. Assume $k \ge 2$ and put $m = m_1$. Then S(q/p) = S(q/c) + (m, ..., m).

Proof. Note that q/c > 1. By Lemma 4.8 and Definition 4, both S(q/p) and S(q/c) have length 2q. Moreover, their components are related as follows:

$$s_j(r) = \lfloor jp/q \rfloor_* - \lfloor (j-1)p/q \rfloor_*$$

= $\lfloor j(mq+c)/q \rfloor_* - \lfloor (j-1)(mq+c)/q \rfloor_*$
= $(jm + \lfloor jc/q \rfloor_*) - ((j-1)m + \lfloor (j-1)c/q \rfloor_*)$
= $m + \lfloor jc/q \rfloor_* - \lfloor (j-1)c/q \rfloor_*$
= $m + s_j(q/c).$

This completes the proof of Lemma 4.10.

Lemma 4.11. Suppose $k \geq 2$. Then, for the rational number

 $q/c = [0, m_2, m_3, \ldots, m_k],$

the conclusion of Proposition 4.3 holds. Namely, each term of S(q/c) is either 0 or 1, and S(q/c) begins with 1 and ends with 0. Moreover, if $m_2 = 1$, no two consecutive terms of S(q/c) can be (0,0), whereas if $m_2 \ge 2$, no two consecutive terms of S(q/c) can be (1,1).

Proof. By Definition 4,

$$s_j(q/c) = \lfloor jc/q \rfloor_* - \lfloor (j-1)c/q \rfloor_*.$$

Since jc/q - (j-1)c/q = c/q is a positive real number less than 1, $s_j(q/c)$ is 0 or 1. Moreover $s_1(q/c) = \lfloor c/q \rfloor_* - \lfloor 0 \rfloor_* = 0 - (-1) = 1$ and $s_{2q}(q/c) = \lfloor 2c \rfloor_* - \lfloor (2q-1)c/q \rfloor_* = (2c-1) - (2c-1) = 0$. Thus S(q/c) begins with 1 and ends with 0.

Note that if $m_2 = 1$ then 1 < q - c < c and hence q < 2c, whereas if $m_2 \ge 2$ then 0 < c < q - c and hence q > 2c. On the other hand, Definition 4 implies

$$s_{j+1}(q/c) + s_j(q/c) = \lfloor (j+1)c/q \rfloor_* - \lfloor (j-1)c/q \rfloor_*.$$

Since (j+1)c/q - (j-1)c/q = 2c/q is greater than 1 or less than 1 according as $m_2 = 1$ or $m_2 \ge 2$, we see that $\lfloor (j+1)c/q \rfloor_* - \lfloor (j-1)c/q \rfloor_*$ is at least 1 or at most 1, accordingly. In the first case, it is impossible for both $s_{j+1}(q/c)$ and $s_j(q/c)$ to be 0, whereas in the second case, it is impossible for both $s_{j+1}(q/c)$ and $s_j(q/c)$ to be 1. This completes the proof. \Box

Proof of Proposition 4.3. If k = 1, then r = 1/m and the assertion is obvious. If $k \ge 2$, then the assertion is a direct consequence of Lemmas 4.10 and 4.11.

In order to prove Proposition 4.4, we prepare the following lemma.

Lemma 4.12. Suppose $k \ge 2$ and $m_2 = 1$. Assume that $S((q - c)/c) = (t_1, \ldots, t_{2(q-c)})$. Then

$$S(q/c) = (t_1\langle 1 \rangle, 0, t_2\langle 1 \rangle, 0, \dots, t_{2(q-c)}\langle 1 \rangle, 0).$$

In particular, T(q/c) = S((q-c)/c).

Proof. Since the first term $s_1((q-c)/c)$ of S((q-c)/c) is equal to t_1 , we see, by Definition 4 that $\lceil i(q-c)/c \rceil^* = 1$ for every integer *i* such that $0 \le i \le t_1 - 1$. This together with the condition $s_2((q-c)/c) = t_2$ implies that $\lceil i(q-c)/c \rceil^* = 2$ for every integer *i* such that $t_1 \le i \le t_1 + t_2 - 1$. Similarly, for each integer ℓ $(1 \le \ell \le 2(q-c))$, we have $\lceil i(q-c)/c \rceil^* = \ell$ for every integer *i* such that $\sum_{h=1}^{\ell-1} t_h \leq i \leq \sum_{h=1}^{\ell} t_h - 1. \text{ Since } \lceil i(q-c)/c \rceil^* = \lceil iq/c \rceil^* - i, \text{ this implies } \lceil iq/c \rceil^* = \ell + i \text{ for every integer } i \text{ such that } \sum_{h=1}^{\ell-1} t_h \leq i \leq \sum_{h=1}^{\ell} t_h - 1. \text{ By Definition 4, this implies that } s_j(q/c) = 1 \text{ if and only if } j = \ell + i \text{ for some integer } \ell \ (1 \leq \ell \leq 2(q-c)) \text{ and some integer } i \ (\sum_{h=1}^{\ell-1} t_h \leq i \leq \sum_{h=1}^{\ell} t_h - 1). \text{ In other words, } s_j(q/c) = 0 \text{ if and only if } j = \ell + \sum_{h=1}^{\ell} t_h \text{ for some integer } \ell \ (1 \leq \ell \leq 2(q-c)). \text{ Hence,}$

$$S(q/c) = (t_1\langle 1 \rangle, 0, t_2\langle 1 \rangle, 0, \dots, t_{2(q-c)}\langle 1 \rangle, 0).$$

Proof of Proposition 4.4 for the case $m_2 = 1$. Suppose $m_2 = 1$. Then $r' = [m_3, \dots, m_k] = (q-c)/c$ by Lemma 4.9. Thus, by Lemma 4.12, T(q/c) = S((q-c)/c) = S(r'). On the other hand, Lemma 4.10 implies T(q/p) = T(q/c). Hence we have T(r) = T(q/c) = S(r').

In order to prove the remaining case of Proposition 4.4, we need the following lemma.

Lemma 4.13. $S(q/c) = \overleftarrow{S}(q/(q-c))_{0\leftrightarrow 1}$, where $\overleftarrow{S}(q/(q-c))_{0\leftrightarrow 1}$ is obtained from S(q/(q-c)) by reading backwards and by replacing 0 and 1.

Proof. By Lemma 4.9, we have

$$q/c = [0, m_2, m_3, \dots, m_k]$$
$$q/(q-c) = [0, 1, m_2 - 1, m_3, \dots, m_k].$$

Thus S(q/c) and S(q/(q-c)) consists of 0 and 1, by Lemma 4.11. On the other hand, for each $1 \le i \le 2q$, we have the following identities by Definition 4:

$$s_{i}(q/(q-c)) = \lfloor i(q-c)/q \rfloor_{*} - \lfloor (i-1)(q-c)/q \rfloor_{*}$$

= $(i + \lfloor -ic/q \rfloor_{*}) - ((i-1) - \lfloor -(i-1)c/q \rfloor_{*})$
= $1 + \lfloor -ic/q \rfloor_{*} - \lfloor (1-i)c/q \rfloor_{*}$
 $s_{q+1-i}(q/c) = \lfloor (q+1-i)c/q \rfloor_{*} - \lfloor (q-i)c/q \rfloor_{*}$
= $(1 + \lfloor (1-i)c/q \rfloor_{*}) - (1 + \lfloor -ic/q \rfloor_{*})$
= $\lfloor (1-i)c/q \rfloor_{*} - \lfloor -ic/q \rfloor_{*}.$

Hence, for each $1 \leq i \leq 2q$,

$$s_i(q/(q-c)) + s_{q+1-i}(q/c) = 1$$

This implies the desired result.

Corollary 4.14. If $m_2 \ge 2$, then $T(q/c) = \overleftarrow{T}(q/(q-c))$.

Proof. Since $q/c = [0, m_2, m_3, \ldots, m_k]$ and since $m_2 \ge 2$, the sequence T(q/c) records the successive occurrences of 0's in S(q/c). On the other hand, since $q/(q-c) = [0, 1, m_2 - 1, m_3, \ldots, m_k]$, T(q/(q-c)) records the successive occurrences of 1's in S(q/(q-c)). Hence Lemma 4.13 implies the desired result. \Box

Proof of Proposition 4.4 for the case $m_2 \ge 2$. Suppose $m_2 \ge 2$. Then $T(q/c) = \overline{T}(q/(q-c))$ by Corollary 4.14. Note that the m_2 for q/(q-c) is equal to 1, and the r' for q/(q-c) is equal to $[m_2 - 1, m_3, \ldots, m_k]$, which is equal to the r' for the original r = q/p. Hence, it follows from Proposition 4.4 for the case $m_2 = 1$ that T(q/(q-c)) = S(r'). Thus we have $T(q/c) = \overline{T}(q/(q-c)) = \overline{S}(r')$. Since T(q/p) = T(q/c) by Lemma 4.10, we obtain the desired identity, $T(r) = \overline{S}(r')$.

Proof of Proposition 4.5. The proof proceeds by induction on $k \ge 1$. If k = 1, S(r) = (m, m). So putting S_1 to be the empty sequence and $S_2 = (m)$, the assertion clearly holds. Now let $k \ge 2$, and let r' be the rational number defined as in Proposition 4.4. We consider four cases separately.

Case 1. $m_2 = 1$ and k = 3.

In this case, $r' = [m_3]$. Thus $S(r') = (T_1, T_2, T_1, T_2)$, where $T_1 = \emptyset$ and $T_2 = (m_3)$. Put

$$S_1 = (m_3 \langle m+1 \rangle), \text{ and } S_2 = (m).$$

Since $T(r) = S(r') = (m_3, m_3)$ by Proposition 4.4, we see $S(r) = (S_1, S_2, S_1, S_2)$ by the definition of T(r). Obviously, S_1 and S_2 satisfy the desired conditions.

Case 2. $m_2 \ge 2$ and k = 2.

In this case, $r' = [m_2 - 1]$. Thus $S(r') = (T_1, T_2, T_1, T_2)$, where $T_1 = \emptyset$ and $T_2 = (m_2 - 1)$. Put

$$S_1 = (m+1)$$
, and $S_2 = ((m_2 - 1)\langle m \rangle)$.

Since $T(r) = \overleftarrow{S}(r') = (m_2 - 1, m_2 - 1)$ by Proposition 4.4, we see $S(r) = (S_1, S_2, S_1, S_2)$ by the definition of T(r). Obviously, S_1 and S_2 satisfy the desired conditions.

Case 3. $m_2 = 1 \text{ and } k \ge 4.$

In this case, $r' = [m_3, \ldots, m_k]$. By the inductive hypothesis,

$$S(r') = (T_1, T_2, T_1, T_2),$$
²⁰

where T_1 and T_2 are symmetric subsequences of S(r') such that each T_i occurs only twice in CS(r'), T_1 begins and ends with $m_3 + 1$, and such that T_2 begins and ends with m_3 . Write

$$T_1 = (t_1, \dots, t_{s_1})$$
 and $T_2 = (t_{s_1+1}, \dots, t_{s_2}),$

and put

$$S_1 = (t_1 \langle m+1 \rangle, m, t_2 \langle m+1 \rangle, \dots, t_{s_1-1} \langle m+1 \rangle, m, t_{s_1} \langle m+1 \rangle);$$

$$S_2 = (m, t_{s_1+1} \langle m+1 \rangle, m, \dots, m, t_{s_2} \langle m+1 \rangle, m).$$

Since T(r) = S(r') by Proposition 4.4, we see $S(r) = (S_1, S_2, S_1, S_2)$ by the definition of T(r). Since T_1 and T_2 are symmetric by the inductive hypothesis, we see that S_1 and S_2 are symmetric subsequences of S(r) such that S_1 begins and ends with m + 1, and S_2 begins and ends with m.

It remains to show that each S_i occurs only twice in CS(r). Recall that S_1 begins and ends with $m_3 + 1$ consecutive m + 1's, and that the maximum number of consecutive occurrences of m+1 in $((S_1, S_2, S_1, S_2))$ is m_3+1 (apply Proposition 4.3 to T(r) = S(r') and use the definition of T(r)). So, if S_1 occurred more than twice in $((S_1, S_2, S_1, S_2))$, T_1 also would occur more than twice in $((T_1, T_2, T_1, T_2))$, a contradiction. On the other hand, recall that m's are isolated in CS(r), and that S_2 begins and ends with m. So if S_2 occurred more than twice in $((S_1, S_2, S_1, S_2))$, T_2 also would occur more than twice in $((T_1, T_2, T_1, T_2))$, a contradiction.

Case 4. $m_2 \geq 2$ and $k \geq 3$.

In this case, $r' = [m_2 - 1, m_3, \dots, m_k]$. By the inductive hypothesis,

 $S(r') = (T_1, T_2, T_1, T_2),$

where T_1 and T_2 are symmetric subsequences of CS(r') such that each T_i occurs only twice in CS(r'), T_1 begins and ends with m_2 , and such that T_2 begins and ends with $m_2 - 1$. Write

$$T_1 = (t_1, \dots, t_{s_1})$$
 and $T_2 = (t_{s_1+1}, \dots, t_{s_2}),$

and put

$$S_1 = (m+1, t_{s_1+1} \langle m \rangle, m+1, \dots, m+1, t_{s_2} \langle m \rangle, m+1);$$

$$S_2 = (t_1 \langle m \rangle, m+1, t_2 \langle m \rangle, \dots, t_{s_1-1} \langle m \rangle, m+1, t_{s_1} \langle m \rangle).$$

Since $T(r) = \overleftarrow{S}(r')$ by Proposition 4.4, we see $S(r) = (S_1, S_2, S_1, S_2)$ by the definition of T(r) and by using the fact that T_1 and T_2 are symmetric. By using the inductive hypothesis, we see that S_1 and S_2 are symmetric subsequences

of S(r) such that S_1 begins and ends with m + 1, and S_2 begins and ends with m. Furthermore, arguing as in Case 3, we can show that each S_i occurs only twice in CS(r). To show the assertion for S_2 , we use the fact that S_2 begins and ends with m_2 consecutive m's, and that the maximum number of consecutive occurrences of m in $((S_1, S_2, S_1, S_2))$ is m_2 .

5. Small cancellation conditions for 2-bridge link groups

Let F(X) be the free group with basis X. A subset R of F(X) is called *symmetrized*, if all elements of R are cyclically reduced and, for each $w \in R$, all cyclic permutations of w and w^{-1} also belong to R.

Definition 5. Suppose that R is a symmetrized subset of F(X). A nonempty word b is called a *piece* if there exist distinct $w_1, w_2 \in R$ such that $w_1 \equiv bc_1$ and $w_2 \equiv bc_2$. Small cancellation conditions C(p) and T(q), where p and q are integers such that $p \geq 2$ and $q \geq 3$, are defined as follows (see [24]).

- (1) Condition C(p): If $w \in R$ is a product of n pieces, then $n \ge p$.
- (2) Condition T(q): For $w_1, \ldots, w_n \in R$ with no successive elements w_i, w_{i+1} an inverse pair $(i \mod n)$, if n < q, then at least one of the products $w_1w_2, \ldots, w_{n-1}w_n, w_nw_1$ is freely reduced without cancellation.

In this section, we prove the following key theorem.

Theorem 5.1. Let r be a rational number such that 0 < r < 1. Recall the presentation $\langle a, b | u_r \rangle$ of G(K(r)) given in Section 3, and let R be the symmetrized subset of F(a, b) generated by the single relator u_r . Then R satisfies C(4) and T(4).

In the remainder of this section, r denotes a rational number such that 0 < r < 1, and (S_1, S_2, S_1, S_2) denotes the decomposition of $S(r) = S(u_r)$ given by Proposition 4.5. We decompose $u_r \equiv v_1 v_2 v_3 v_4$, where subwords v_1 and v_3 correspond to S_1 , and subwords v_2 and v_4 correspond to S_2 . As in Section 4, we consider the continued fraction expansion $r = [m_1, m_2, \ldots, m_k]$, where $k \ge 1$, $(m_1, m_2, \ldots, m_k) \in (\mathbb{Z}_+)^k$ and $m_k \ge 2$ unless k = 1. It should be noted that if k = 1 then both v_1 and v_3 are empty words.

We begin with the following lemma.

Lemma 5.2. Let w be an arbitrary cyclic permutation of the single relator u_r of the group presentation of G(K(r)). Then the set

{the initial letter of
$$w' \mid (w') \equiv (u_r^{\pm 1})$$
 and $S(w') = S(w)$ }

equals $\{a, a^{-1}, b, b^{-1}\}.$

Proof. We first prove the lemma when $w \equiv u_r (\equiv v_1 v_2 v_3 v_4)$. Consider the cyclic permutation $w_1 := v_3 v_4 v_1 v_2$ of w and the cyclic permutations $w_2 := v_1^{-1} v_4^{-1} v_3^{-1} v_2^{-1}$ and $w_3 := v_3^{-1} v_2^{-1} v_1^{-1} v_4^{-1}$ of u_r^{-1} . Then by Proposition 4.5, w_1 , w_2 and w_3 share the same S-sequence with w. We show that the initial letters of w, w_1 , w_2 and w_3 are all distinct. By Lemma 3.1, $w \equiv u_r$ has the initial letter a, and w_1 has initial letter a^{-1} , b or b^{-1} . Thus w and w_1 have different initial letters. This also implies that w_2 and w_3 have different initial letters, as follows. Suppose w_2 and w_3 share the same initial letter. Then, since $S(w_2) = S(w_3)$, we have $w_2 \equiv w_3$ by Proposition 4.1(1). However, this implies $v_1 \equiv v_3$ and $v_2 \equiv v_4$, and hence $w \equiv w_1$, a contradiction. Next, we show that the initial letters of the contrary that this is not the case. Then, since these four words have the same S-sequences, it follows from Proposition 4.1(1) that w_2 or w_3 is equal to w or w_1 . This implies that u_r^{-1} is a cyclic permutation of u_r . However, this is impossible by the following claim, and this completes the proof of the lemma when $w \equiv u_r$.

Claim. u_r^{-1} cannot be a cyclic permutation of u_r .

Proof of Claim. If u_r^{-1} were a cyclic permutation of u_r , then there would be decompositions such as $u_r \equiv z_1 z_2$ and $u_r^{-1} \equiv z_2 z_1$. Since $u_r^{-1} \equiv (z_1 z_2)^{-1} \equiv z_2^{-1} z_1^{-1}$, we would have $z_i \equiv z_i^{-1}$ yielding that $z_i^2 = 1$ (i = 1, 2) in the free group F(a, b). Since F(a, b) is torsion free, we have $z_i = 1$ (i = 1, 2) and hence $u_r = z_1 z_2 = 1$ in F(a, b), a contradiction.

Now, let w be an arbitrary cyclic permutation of u_r . Let d be an integer such that w is obtained from u_r by cyclical shift of d-digits. For each i = 1, 2, 3, let \hat{w}_i be the word obtained from the word w_i in the previous paragraph by cyclic shift of d-digits. Then, since $S(w_i) = S(u_r)$, we have $S(\hat{w}_i) = S(w)$ (i = 1, 2, 3). This implies that the initial letters of w and \hat{w}_i (i = 1, 2, 3) are all distinct. Because, otherwise, w and \hat{w}_i (i = 1, 2, 3) are not all distinct by Proposition 4.1(1), and hence u_r and w_i (i = 1, 2, 3) are not all distinct, a contradiction. Moreover, \hat{w}_i (i = 1, 2, 3) are cyclic permutations of w^{\pm} , because w_i (i = 1, 2, 3) are cyclic permutations of $u_r^{\pm 1}$ and w is a cyclic permutation of u_r . Hence we obtain the desired result.

Lemma 5.3. For the relator $u_r \equiv v_1 v_2 v_3 v_4$ with $r = [m_1, m_2, \ldots, m_k]$, the following hold.

(1) If k = 1, then the following hold.
(a) No piece can contain v₂ or v₄.

- (b) No piece is of the form $v_{2e}v_{4b}$ or $v_{4e}v_{2b}$, where v_{ib} and v_{ie} are nonempty initial and terminal subwords of v_i , respectively.
- (c) Every subword of the form v_{2b} , v_{2e} , v_{4b} , or v_{4e} is a piece, where v_{ib} and v_{ie} are nonempty initial and terminal subwords of v_i with $|v_{ib}|, |v_{ie}| \leq |v_i| 1$, respectively.
- (2) If $k \ge 2$, then the following hold.
 - (a) No piece can contain v_1 or v_3 .
 - (b) No piece is of the form $v_{1e}v_2v_{3b}$ or $v_{3e}v_4v_{1b}$, where v_{ib} and v_{ie} are nonempty initial and terminal subwords of v_i , respectively.
 - (c) Every subword of the form $v_{1e}v_2$, v_2v_{3b} , $v_{3e}v_4$, or v_4v_{1b} is a piece, where v_{ib} and v_{ie} are nonempty initial and terminal subwords of v_i with $|v_{ib}|, |v_{ie}| \leq |v_i| - 1$, respectively.

Proof. (1a) & (1b) & (1c) The proofs are analogous to the proofs of (2a) & (2b) & (2c) below.

(2a) Suppose to the contrary that there are two distinct cyclic permutations w_1 and w_2 of u_r or u_r^{-1} such that w_1 and w_2 have the same beginning subword y, where either $y \equiv v_1$ or $y \equiv v_3$. Since the cyclic sequence $CS(r) = CS(u_r)$ is symmetric by Corollary 4.6, the cyclic sequence $CS(u_r^{-1})$ is also equal to CS(r). Thus the two cyclic words (w_1) and (w_2) have the same associated cyclic sequence $((S_1, S_2, S_1, S_2))$, regardless of whether w_1 and w_2 are cyclic permutations of u_r or u_r^{-1} . Putting $m = m_1$, note that $((S_1, S_2, S_1, S_2))$ is a cyclic sequence consisting of only m and m+1, S_1 begins and ends with m+1, and the S-sequence of y is S_1 (see Proposition 4.3). This implies that, for each i = 1, 2, the S-sequence $S(w_i)$ begins with S_1 , and that the cyclic S-sequence $CS(w_i)$ is represented by $S(w_i)$. Furthermore, since S_1 appears only twice in $((S_1, S_2, S_1, S_2))$ by Proposition 4.5, we obtain that w_1 and w_2 must have the same associated S-sequence (S_1, S_2, S_1, S_2) , where the first S_1 corresponds to the common beginning subword y. By Proposition 4.1(1), this implies that $w_1 \equiv w_2$, a contradiction.

(2b) Suppose to the contrary that there is a piece, z, which is of the form, say $v_{1e}v_2v_{3b}$, and let w_1 and w_2 be distinct cyclic permutations of u_r or u_r^{-1} such that w_1 and w_2 have the same beginning subword $z \equiv v_{1e}v_2v_{3b}$, namely $w_i \equiv zw'_i \equiv v_{1e}v_2v_{3b}w'_i$ for some subword w'_i of w_i . By the construction of the product $v_1v_2v_3v_4$, the last exponent of v_{1e} , which is equal to the last exponent of v_1 , is different from the first exponent of v_2 . Consider the cyclic permutation $\hat{w}_i := v_2v_{3b}w'_iv_{1e}$. By the observation above, the cyclic sequence $CS(\hat{w}_i)$ is represented by $S(\hat{w}_i)$ (cf. Proposition 4.1(2)). Moreover, since v_{3b} is a nonempty reduced subword of \hat{w}_i whose initial exponent is different from the terminal exponent of v_2 , the sequence $S(\hat{w}_i)$ starts with $S(v_2) = S_2$. Since S_2 appears in $CS(\hat{w}_i) = ((S_1, S_2, S_1, S_2))$ only twice, we see $S(\hat{w}_i) = (S_2, S_1, S_2, S_1)$. This implies $\hat{w}_1 \equiv \hat{w}_2$ by Proposition 4.1(1) and hence $w_1 \equiv w_2$, a contradiction.

(2c) Since every nonempty subword of a piece is also a piece, it is enough to prove the assertion for $v_{1e}v_2$, v_2v_{3b} , $v_{3e}v_4$, or v_4v_{1b} , where v_{ib} and v_{ie} , respectively, are the initial and the terminal subwords of v_i with $|v_{ib}| = |v_{ie}| = |v_i| - 1$.

We show that $v_{1e}v_2$ and v_2v_{3b} are pieces. To this end, we first show that $v_{1e}v_2$ and v_2v_{3b} have the same associated S-sequence. Since the terminal exponent of v_i and the initial exponent of v_{i+1} are different, $S(v_{1e}v_2) = (S(v_{1e}), S(v_2))$ and $S(v_2v_{3b}) = (S(v_2), S(v_{3b}))$. On the other hand, we have $v_{1e}v_2 = \hat{u}_r$, because $u_r \equiv a\hat{u}_r x \hat{u}_r^{-1}$ by Lemma 3.1. Thus we see $S(v_{1e}v_2) = S(\hat{u}_r)$ is symmetric by the following claim.

Claim. The sequence $S(\hat{u}_r)$ is symmetric.

Proof of Claim. Recall that the *i*-th exponent of \hat{u}_r is given by $\epsilon_i = (-1)^{\lfloor iq/p \rfloor}$ (see Lemma 3.1). So we have:

$$\epsilon_{p-i} = (-1)^{\lfloor (p-i)q/p \rfloor} = (-1)^{q+\lfloor -iq/p \rfloor} = \begin{cases} \epsilon_i & \text{if } q \text{ is even,} \\ -\epsilon_i & \text{if } q \text{ is odd.} \end{cases}$$

Hence the sequence $(\epsilon_1, \epsilon_2, \ldots, \epsilon_{p-1})$ is symmetric or skew-symmetric according as q is even or odd. This implies that $S(\hat{u}_r)$ is symmetric.

Hence we have

$$S(v_{1e}v_2) = \overleftarrow{S}(v_{1e}v_2) = (\overleftarrow{S}(v_2), \overleftarrow{S}(v_{1e}))$$
$$= (S(v_2), S(v_{1b})) = (S(v_2), S(v_{3b})) = S(v_2v_{3b}).$$

Here, the third identity follows from the fact that $S(v_1) = S_1$ and $S(v_2) = S_2$ are symmetric and the fourth identity follows from the fact that $S(v_1) = S_1 = S(v_3)$.

Now, let $w_1 := v_{1e}v_2w'_1$ and $w_2 := v_2v_{3b}w'_2$ be cyclic permutations of u_r . Note that the terminal exponent of v_2 and the initial exponent of w'_1 are different, and that the terminal exponent of v_{3b} and the initial exponent of w'_2 are the same. Here, the latter assertion follows from the fact that the last component of $S(v_3) = S_1$ is equal to m_1 or $m_1 + 1$ according as k = 1 or $k \ge 2$ (see Proposition 4.2) and hence it is at least 2. (Recall that $m_1 \ge 2$ or $m_1 \ge 1$ according as k = 1 or $k \ge 2$.) Hence $S(w_1) \ne S(w_2)$. By Lemma 5.2, there is a cyclic permutation \hat{w}_2 of u_r or u_r^{-1} such that \hat{w}_2 has the same initial letter as w_1 and such that $S(\hat{w}_2) = S(w_2)$. Then w_1 and \hat{w}_2 are distinct cyclic permutations of u_r or u_r^{-1} , since $S(\hat{w}_2) \ne S(w_1)$. Note, however, that w_1 and \hat{w}_2 have the same beginning subword $v_{1e}v_2$ (cf. Proposition 4.1(1)). This implies that $v_{1e}v_2$ is a piece. We can also see that v_2v_{3b} is a piece by a similar argument. By using the fact that $v_{3e}v_4 = \hat{u}_r^{-1}$, we can show by a similar argument that $v_{3e}v_4$ and v_4v_{1b} are also pieces.

We now introduce the following definition.

Definition 6. For a positive integer n, a nonempty subword w of the cyclic word (u_r) is called a *maximal n-piece* if w is a product of n pieces and if any subword w' of u_r which properly contains w as an *initial* subword is not a product of n-pieces.

It should be noted that a maximal 1-piece w may not be a maximal piece, because there may exist a piece w' which contains w as a proper terminal subword. (Here a nonempty subword w of the cyclic word (u_r) is called a maximal piece if w is a piece and if any subword w' of u_r which properly contains w is not a piece.) However, every maximal piece is a maximal 1piece.

Corollary 5.4. For the relator $u_r \equiv v_1 v_2 v_3 v_4$ with $r = [m_1, m_2, \ldots, m_k]$, let v_{ib}^* be the maximal proper initial subword of v_i , i.e., the initial subword of v_i such that $|v_{ib}^*| = |v_i| - 1$ (i = 1, 2, 3, 4). Then the following hold, where v_{ib} and v_{ie} are nonempty initial and terminal subwords of v_i with $|v_{ib}|, |v_{ie}| \leq |v_i| - 1$, respectively.

- (1) If k = 1, then the following hold.
 - (a) The following is the list of all maximal 1-pieces of (u_r) , arranged in the order of the position of the initial letter:

 $v_{2b}^*, v_{2e}, v_{4b}^*, v_{4e}.$

(b) The following is the list of all maximal 2-pieces of (u_r) , arranged in the order of the position of the initial letter:

 $v_2, v_{2e}v_{4b}^*, v_4, v_{4e}v_{2b}^*.$

(c) The following is the list of all maximal 3-pieces of (u_r) , arranged in the order of the position of the initial letter:

 $v_2 v_{4b}^*, v_{2e} v_4, v_4 v_{2b}^*, v_{4e} v_2.$

(2) If $k \ge 2$, then the following hold.

(a) The following is the list of all maximal 1-pieces of (u_r) , arranged in the order of the position of the initial letter:

 $v_{1b}^*, v_{1e}v_2, v_2v_{3b}^*, v_{2e}v_{3b}^*, v_{3b}^*, v_{3e}v_4, v_4v_{1b}^*, v_{4e}v_{1b}^*.$

(b) The following is the list of all maximal 2-pieces of (u_r) , arranged in the order of the position of the initial letter:

 $v_1v_2, v_{1e}v_2v_{3b}^*, v_2v_3v_4, v_{2e}v_3v_4, v_3v_4, v_{3e}v_4v_{1b}^*, v_4v_1v_2, v_{4e}v_1v_2.$

(c) The following is the list of all maximal 3-pieces of (u_r) , arranged in the order of the position of the initial letter:

 $v_1v_2v_{3b}^*, v_{1e}v_2v_3v_4, v_2v_3v_4v_{1b}^*, v_{2e}v_3v_4v_{1b}^*, v_3v_4v_{1b}^*, v_{3e}v_4v_1v_2, v_4v_1v_2v_{3b}^*, v_{4e}v_1v_2v_{3b}^*.$

Proof. (1a) & (1b) & (1c) The proofs are analogous to the proofs of (2a) & (2b) & (2c) below.

(2a) This is a direct consequence of Lemma 5.3.

(2b) This is proved by using the fact that if w is a maximal 2-piece, then it has a unique decomposition $w = w_1w_2$ into two maximal 1-pieces w_1 and w_2 . To be precise, if w_1 is equal to v_{1b}^* (resp. $v_{1e}v_2$, $v_2v_{3b}^*$, $v_{2e}v_{3b}^*$), then w_2 is equal to $v_{1e}v_2$ (resp. v_{3b}^* , $v_{3e}v_4$, $v_{3e}v_4$), and hence $w = w_1w_2$ is equal to v_1v_2 (resp. $v_{1e}v_2v_{3b}^*$, $v_2v_3v_4$, $v_{2e}v_3v_4$).

(2c) This is proved by using the fact that if w is a maximal 3-piece, which is a proper subword of the cyclic word (u_r) , then it has a unique decomposition $w = w_1w_2$, where w_1 is a maximal 2-piece and w_2 is a maximal 1-piece. To be precise, if w_1 is equal to v_1v_2 (resp. $v_{1e}v_2v_{3b}^*$, $v_2v_3v_4$, $v_{2e}v_3v_4$), then w_2 is equal to v_{3b}^* (resp. $v_{3e}v_4$, v_{1b}^* , v_{1b}^*), and hence $w = w_1w_2$ is equal to $v_1v_2v_{3b}^*$ (resp. $v_{1e}v_2v_3v_4$, $v_2v_3v_4v_{1b}^*$, $v_{2e}v_3v_4v_{1b}^*$).

Proof of Theorem 5.1. By Corollary 5.4, the cyclic word (u_r) is not a product of 3 pieces. This implies that the cyclic word (u_r^{-1}) as well is not a product of 3 pieces. Hence R satisfies C(4). So we show that R satisfies T(4). To this end, recall that the cyclic word (u_r) is alternating by Lemma 4.7. Now suppose that R does not satisfy T(4). Then there exist $w_1, w_2, w_3 \in R$ such that w_1w_2, w_2w_3 and w_3w_1 are reducible. Let x^{ϵ_1} be the terminal letter of w_1 , where $x \in \{a, b\}$ and $\epsilon_1 = \pm 1$. Then the initial letter of w_2 is $x^{-\epsilon_1}$, because w_1w_2 is reducible. Since w_2 is cyclically alternating, this implies that the terminal letter of w_2 is y^{ϵ_2} for some $\epsilon_2 = \pm 1$, where y is the element of $\{a, b\}$ different from x. Similarly, by using the facts that w_2w_3 is reducible and that w_3 is cyclically alternating, we see that the terminal letter of w_3 is x^{ϵ_3} for some $\epsilon_3 = \pm 1$. Since w_3w_1 is reducible, this implies that the initial letter of w_1 is $x^{-\epsilon_3}$. However, this contradicts the fact that w_1 is cyclically alternating, because the terminal letter of w_1 was x^{ϵ_1} . Hence R satisfies T(4).

6. VAN KAMPEN DIAGRAMS OVER 2-BRIDGE LINK GROUPS

In this section, we investigate the geometric consequences of Theorem 5.1. Let us begin with necessary definitions and notation following [24]. A map Mis a finite 2-dimensional cell complex embedded in \mathbb{R}^2 , namely a finite collection of vertices (0-cells), edges (1-cells), and faces (2-cells) in \mathbb{R}^2 . The boundary (frontier) of M in \mathbb{R}^2 is denoted by ∂M . If D is a face of M, the boundary of D is denoted by ∂D . An edge may be traversed in either of two directions. If v is a vertex of M, $d_M(v)$, the *degree of* v, will denote the number of oriented edges in M having v as initial vertex. A vertex v of M is called an *interior vertex* if $v \notin \partial M$, and an edge e of M is called an *interior edge* if $e \notin \partial M$.

Definition 7. A nonempty map M is called a [p,q]-map if the following conditions hold.

- (1) Every interior vertex of M has degree at least p.
- (2) Every face D of M has at least q edges in ∂D .

A path in M is a sequence of oriented edges e_1, \ldots, e_n such that the initial vertex of e_{i+1} is the terminal vertex of e_i for every $1 \leq i \leq n-1$. A cycle is a closed path, namely a path e_1, \ldots, e_n such that the initial vertex of e_1 is the terminal vertex of e_n . If D is a face of M, any cycle of minimal length which includes all the edges of ∂D is called a *boundary cycle* of D. If M is connected and simply connected, a *boundary cycle* of M is defined to be a cycle of minimal length which contains all the edges of ∂M going around once along the boundary of $\mathbb{R}^2 - M$.

Definition 8. Let R be a symmetrized subset of F(X). An R-diagram is a map M and a function ϕ assigning to each oriented edge e of M, as a label, a reduced word $\phi(e)$ in X such that the following hold.

- (1) If e is an oriented edge of M and e^{-1} is the oppositely oriented edge, then $\phi(e^{-1}) = \phi(e)^{-1}$.
- (2) For any boundary cycle δ of any face of M, $\phi(\delta)$ is a cyclically reduced word representing an element of R. (If $\alpha = e_1, \ldots, e_n$ is a path in M, we define $\phi(\alpha) \equiv \phi(e_1) \cdots \phi(e_n)$.)

In particular, if a group G is presented by $G = \langle X | R \rangle$ with R being symmetrized, then a connected and simply connected R-diagram is called a *van* Kampen diagram over the group presentation $G = \langle X | R \rangle$.

Let D_1 and D_2 be faces (not necessarily distinct) of M with an edge $e \subseteq \partial D_1 \cap \partial D_2$. Let $e\delta_1$ and $\delta_2 e^{-1}$ be boundary cycles of D_1 and D_2 , respectively. Let $\phi(\delta_1) = f_1$ and $\phi(\delta_2) = f_2$. An R-diagram M is called *reduced* if one never has $f_2 = f_1^{-1}$. It should be noted that if M is reduced then $\phi(e)$ is a piece for every interior edge e of M. A boundary label of M is defined to be a word $\phi(\alpha)$ in X for α a boundary cycle of M. It is easy to see that any two boundary labels of M are cyclic permutations of each other.

We recall the following lemma which is a well-known classical result in combinatorial group theory (see [24]).

Lemma 6.1 (van Kampen). Suppose $G = \langle X | R \rangle$ with R being symmetrized. Let v be a word in X. Then v = 1 in G if and only if there exists a reduced van Kampen diagram M over $G = \langle X | R \rangle$ with a boundary label v.

Convention 1. Let R be the symmetrized subset of F(a, b) generated by the single relator u_r of the group presentation of G(K(r)). For any reduced R-diagram M, we assume that M satisfies the following.

- (1) Every interior vertex of M has degree at least three.
- (2) For every edge e of ∂M , the label $\phi(e)$ is a piece.
- (3) For a path e_1, \ldots, e_n in ∂M of length $n \geq 2$ such that the vertex $e_i \cap e_{i+1}$ has degree 2 for $i = 1, 2, \ldots, n-1, \phi(e_1)\phi(e_2)\cdots\phi(e_n)$ cannot be expressed as a product of less than n pieces.

Indeed, we may assume (1), because if there are two interior edges e_1 and e_2 meeting in an interior vertex of degree two, then we can delete the vertex vand unite e_1 and e_2 into a single edge e with label $\phi(e) = \phi(e_1)\phi(e_2)$. To see (2), recall that the assumption that M is reduced implies that $\phi(e)$ is a piece for every interior edge e of M. On the other hand, since the cyclic word (u_r) can be written as a product of pieces, we may also assume that $\phi(e)$ is a piece for every edge e in ∂M . Finally, we may assume (3), because if $\phi(e_1) \cdots \phi(e_n)$ is expressed as a product of less than n pieces, then we can change the cellular structure of the interval $e_1 \cup \cdots \cup e_n$ so that the new cellular structure has fewer vertices compared with the original one.

For the remainder of this section, we assume that r is a rational number such that 0 < r < 1, and let $\langle a, b | u_r \rangle$ be the presentation of G(K(r)) given in Section 3.

The following corollary is immediate from Theorem 5.1 and Convention 1.

Corollary 6.2. Let R be the symmetrized subset of F(a, b) generated by the single relator u_r of the group presentation of G(K(r)). Then every reduced R-diagram is a [4,4]-map.

This corollary enables us to apply the Curvature Formula of Lyndon and Schupp for [p,q]-maps satisfying 1/p + 1/q = 1/2 (see [24]) to obtain the following theorem, the proof of which is deferred to the end of this section. **Theorem 6.3.** Let R be the symmetrized subset of F(a, b) generated by the single relator u_r of the group presentation of G(K(r)). Suppose that M is a reduced van Kampen diagram over $G(K(r)) = \langle a, b | R \rangle$ such that any boundary label of M is cyclically reduced and alternating. Then some boundary label of M contains a subword w of $(u_r^{\pm 1})$ such that the S-sequence of w is (S_1, S_2, ℓ) or (ℓ, S_2, S_1) for some positive integer ℓ , where $S(r) = (S_1, S_2, S_1, S_2)$ is as in Proposition 4.5.

By Lemma 6.1, we obtain the following important corollary which is the main result of this section.

Corollary 6.4. Let s be a rational number such that $0 < s \leq 1$ and that α_s is null-homotopic in $S^3 - K(r)$. Then the cyclic S-sequence CS(s) contains (S_1, S_2) or (S_2, S_1) as a subsequence, where $S(r) = (S_1, S_2, S_1, S_2)$ is as in Proposition 4.5.

In the above corollary (and throughout this paper), we mean by a *sub-sequence* a subsequence without leap. Namely a sequence (a_1, a_2, \ldots, a_p) is called a *subsequence* of a cyclic sequence, if there is a sequence (b_1, b_2, \ldots, b_n) representing the cyclic sequence such that $p \leq n$ and $a_i = b_i$ for $1 \leq i \leq p$.

Proof. Let R be the symmetrized subset of F(a, b) generated by u_r . Since α_s is null-homotopic in $S^3 - K(r)$, the cyclic word (u_s) obtained from α_s (as in Section 3) represents the trivial element of $G(K(r)) = \langle a, b | R \rangle$. Thus, by Lemma 6.1, there is a reduced van Kampen diagram M over $G(K(r)) = \langle a, b | R \rangle$ with a boundary label u_s . Since u_s is cyclically reduced and the cyclic word (u_s) is alternating, Theorem 6.3 implies that the cyclic word (u_s) contains a subword w such that the S-sequence of w is (S_1, S_2, ℓ) or (ℓ, S_2, S_1) for some positive integer ℓ . Recall that S_1 begins and ends with $m_1 + 1$, and S_2 begins and ends with m_1 (see Proposition 4.5). Thus (S_1, S_2, ℓ) is of the form $(m_1 + 1, \ldots, m_1, \ell)$ and (ℓ, S_2, S_1) is of the form $(\ell, m_1, \ldots, m_1 + 1)$. By Proposition 4.3, this yields that $CS(s) = CS(u_s)$ (cf. Definition 2) consists of m_1 and $m_1 + 1$ and that CS(s) contains (S_1, S_2) or (S_2, S_1) as a subsequence.

The remainder of this section is devoted to the proof of Theorem 6.3. An *extremal disk* of a map M is a submap J of M which is topologically a disk and which has a boundary cycle e_1, \ldots, e_n such that the edges e_1, \ldots, e_n occur in order in some boundary cycle of the whole map M. We note that if J is an extremal disk of M, then either J = M or J is connected to the rest of M by a single vertex.

Proof of Theorem 6.3. By Corollary 6.2, a reduced van Kampen diagram M over $G(K(r)) = \langle a, b | R \rangle$ is a [4,4]-map. Since a boundary label of M is cyclically reduced, there is no vertex of degree 1 in ∂M . Moreover, since any boundary label of M is alternating, there is no vertex of degree 3 in ∂M . So every vertex in ∂M must have degree 2 or at least 4.

Choose an extremal disk, say J, of M.

Claim. There are three edges e_1 , e_2 and e_3 in ∂J such that $e_1 \cap e_2 = \{v_1\}$ and $e_2 \cap e_3 = \{v_2\}$, where $d_J(v_i) = 2$ for each i = 1, 2.

Proof of Claim. Clearly J is a connected and simply connected [4, 4]-map having at least one face. By the Curvature Formula of Lyndon and Schupp (see [24, Corollary V.3.4]), we have

(†)
$$\sum_{v \in \partial J} (3 - d_J(v)) \ge 4$$

Putting

$$A = \{ v \in \partial J \mid d_J(v) = 2 \} \text{ and } B = \{ v \in \partial J \mid d_J(v) \ge 4 \},\$$

it is easy to see that A has at least 4 more elements than B does in order to satisfy inequality (†). Since J is an extremal disk of M, either J = M or it is connected to the rest of M by a single vertex. If J = M, then every vertex in $\partial J = \partial M$ belongs to either A or B. On the other hand, if J is connected to the rest of M by a single vertex, say v_0 , then every vertex in ∂J except v_0 belongs to either A or B and $d_J(v_0) = d_M(v_0) - 1 \ge 3$ (note that $d_M(v_0) \ge 4$, since $v_0 \in \partial M$). In either case, we see that there are at least 2 adjacent vertices, say v_1 and v_2 , belonging to A. This proves the claim.

By Claim, there is a face D in M such that $\partial D \cap \partial M$ contains three successive edges e_1 , e_2 and e_3 . By Convention 1(2)–(3), the product $\phi(e_1)\phi(e_2)\phi(e_3)$ which is a subword of the cyclic word $(u_r^{\pm 1})$ cannot be expressed as a product of less than 3 pieces. We may assume without loss of generality that $\phi(e_1)\phi(e_2)\phi(e_3)$ is a subword of the cyclic word (u_r) . We also assume that the length k of the continued fraction $r = [m_1, m_2, \ldots, m_k]$ is greater than 1. (The proof for the case k = 1 is analogous to the proof for the general case $k \geq 2$.) Let w_0 be the maximal 2-piece which forms a proper initial subword of $\phi(e_1)\phi(e_2)\phi(e_3)$. Then w_0 is equal to one of the words in Corollary 5.4(2b) If w_0 is equal to v_1v_2 or $v_{1e}v_2v_{3b}^*$, then $\phi(e_1)\phi(e_2)\phi(e_3)$ contains a subword w such that the S-sequence of w is (S_1, S_2, ℓ) or (ℓ, S_2, S_1) accordingly, for some positive integer ℓ . The remaining possibilities for w_0 can be treated similarly and we obtain Theorem 6.3.

7. Proof of Main Theorem 2.3

In this section, we prove the only if part of Main Theorem 2.3. The if part is Proposition 2.2 ([25, Corollary 4.7]). Though the proof of the main theorem for the trivial knot K(0/1) and the trivial 2-component link K(1/0) is easy, we need to treat them separately. We defer these to the end of this section, and we assume, until the final part of this section, that the slope r of the 2-bridge link K(r) satisfies the condition 0 < r < 1 (cf. Theorem 2.1) and that $r = [m_1, m_2, \ldots, m_k]$, where $k \ge 1, (m_1, \ldots, m_k) \in (\mathbb{Z}_+)^k$, and $m_k \ge 2$.

Recall that the region R bounded by a pair of Farey edges with an endpoint ∞ and a pair of edges with an endpoint r forms a fundamental domain of the action of $\hat{\Gamma}_r$ on \mathbb{H}^2 (see Figure 2). Let I_1 and I_2 be the closed intervals in $\hat{\mathbb{R}}$ obtained as the intersection with $\hat{\mathbb{R}}$ of the closure of R. To be precise, $I_1 = [0, r_1]$ and $I_2 = [r_2, 1]$, where

$$r_{1} = \begin{cases} [m_{1}, m_{2}, \dots, m_{k-1}] & \text{if } k \text{ is odd,} \\ [m_{1}, m_{2}, \dots, m_{k-1}, m_{k} - 1] & \text{if } k \text{ is even,} \end{cases}$$
$$r_{2} = \begin{cases} [m_{1}, m_{2}, \dots, m_{k-1}, m_{k} - 1] & \text{if } k \text{ is odd,} \\ [m_{1}, m_{2}, \dots, m_{k-1}] & \text{if } k \text{ is even.} \end{cases}$$

If r = 1/p (p > 1), then I_1 is degenerate to the singleton $\{0\}$. And if r = (p - 1)/p (p > 1), then I_2 is degenerate to the singleton $\{1\}$. Otherwise, I_1 and I_2 are non-degenerate intervals, and the union $I_1 \cup I_2$ forms a fundamental domain of the action of $\hat{\Gamma}_r$ on the domain of discontinuity of $\hat{\Gamma}_r$, the complement in $\hat{\mathbb{R}}$ of the closure of $\hat{\Gamma}_r\{\infty, r\}$. (In the exceptional case r = 1/p (resp. (p-1)/p), the rational number 0 (resp. 1) lies in the limit set and I_2 (resp. I_1) is a fundamental domain of the action of $\hat{\Gamma}_r$ on the domain of discontinuity.) This fact together with Proposition 2.2 implies the following lemma.

Lemma 7.1. Suppose 0 < r < 1. Then for any $s \in \hat{\mathbb{Q}}$, there is a unique rational number $s_0 \in I_1 \cup I_2 \cup \{\infty, r\}$ such that s is contained in the $\hat{\Gamma}_r$ -orbit of s_0 , and in particular, α_s is homotopic to α_{s_0} in $S^3 - K(r)$.

Proof. Let s be an element of $\hat{\mathbb{Q}}$. Pick a point, z, in the interior of a Farey triangle contained in the fundamental domain R, and consider the geodesic, ℓ , in \mathbb{H}^2 joining z with s. Then ℓ intersects only finitely many Farey edges, and hence it intersects only finitely many $\hat{\Gamma}_r$ -images of the four boundary edges of R. This enables us to find an element $\gamma \in \hat{\Gamma}_r$ such that $\gamma(s) \in \overline{R} \cap \hat{\mathbb{R}} = I_1 \cup I_2 \cup \{\infty, r\}$. Thus s is contained in the $\hat{\Gamma}_r$ -orbit of $s_0 := \gamma(s) \in I_1 \cup I_2 \cup \{\infty, r\}$.

The uniqueness of such an element s_0 can be seen by looking at the quotient space of $\mathbb{H}^2 \cup \Omega(\hat{\Gamma}_r)$ by $\hat{\Gamma}_r$, where $\Omega(\hat{\Gamma}_r)$ is the domain of discontinuity of the action of $\hat{\Gamma}_r$ on $\partial \mathbb{H}^2$. Finally, Proposition 2.2 implies that α_s is homotopic to α_{s_0} in $S^3 - K(r)$.

Thus the only if part of Main Theorem 2.3 is equivalent to the following theorem, except for the the trivial knot K(0) and the trivial 2-component link $K(\infty)$.

Theorem 7.2. Suppose 0 < r < 1. Then, for any $s \in I_1 \cup I_2$, α_s is not null-homotopic in $S^3 - K(r)$.

The following lemma plays an important role in the proof of Theorem 7.2.

Lemma 7.3. Suppose 0 < r < 1, and let $S(r) = (S_1, S_2, S_1, S_2)$ be as in Proposition 4.5. Suppose that a rational number $s \in (0, 1)$ has a continued fraction expansion $s = [l_1, \ldots, l_t]$, where $t \ge 1$, $(l_1, \ldots, l_t) \in (\mathbb{Z}_+)^t$, and $l_t \ge 2$ unless t = 1. If the cyclic S-sequence CS(s) contains (S_1, S_2) or (S_2, S_1) as a subsequence, then the following hold.

- (1) $t \geq k$.
- (2) $l_i = m_i$ for each i = 1, ..., k 1.
- (3) Either $l_k \ge m_k$ or both $l_k = m_k 1$ and t > k.

Proof. From Proposition 4.5, keep in mind that CS(s) consists of l_1 and $l_1 + 1$ (here $l_1 + 1$ appears only if $t \ge 2$). The proof proceeds by induction on $k \ge 1$.

If k = 1, that is, $r = [m_1]$, then $S_1 = \emptyset$ and $S_2 = (m_1)$. So, if CS(s) contains $(S_1, S_2) = (S_2, S_1) = (m_1)$ as a subsequence, then either $l_1 \ge m_1$ or both $l_1 = m_1 - 1$ and $t \ge 2$, proving the base step.

Now let $k \ge 2$. Suppose that CS(s) contains (S_1, S_2) or (S_2, S_1) as a subsequence. By Proposition 4.3, this yields that CS(s) consists of m_1 and $m_1 + 1$. This happens only when $t \ge 2$ and $l_1 = m_1$. We consider three cases separately.

Case 1. $m_2 = 1$.

In this case, $k \ge 3$ and, by Propositions 4.3 and 4.4, (m_1+1, m_1+1) appears in both (S_1, S_2) and (S_2, S_1) as a subsequence, so in CS(s) as a subsequence. Again by Proposition 4.3, $l_2 = 1$ and so $t \ge 3$. Define

$$r' := [m_3, \dots, m_k]$$
 and $s' := [l_3, \dots, l_t].$

Let $S(r') = (T_1, T_2, T_1, T_2)$ be the decomposition of S(r') given by Proposition 4.3, as in the proof of Proposition 4.5. By Corollary 4.6,

$$CT(r) = CS(r')$$
 and $CT(s) = CS(s')$,
33

so it follows that (T_1, T_2) or (T_2, T_1) appears in CS(s') as a subsequence, because (S_1, S_2) or (S_2, S_1) appears in CS(s) as a subsequence, by assumption. Thus the induction completes the case.

Case 2. Both $m_2 = 2$ and k = 2.

In this case, the assertion always holds, because if $l_2 = 1$ then we must have $t \ge 3$, otherwise $l_2 \ge 2 = m_2$.

Case 3. Either $m_2 \ge 3$ or both $m_2 = 2$ and $k \ge 3$.

In this case, by Proposition 4.3, (m_1, m_1) appears in both (S_1, S_2) and (S_2, S_1) as a subsequence, so in CS(s) as a subsequence. Again by Proposition 4.3, $l_2 \geq 2$. Define

$$r' := [m_2 - 1, m_3, \dots, m_k]$$
 and $s' := [l_2 - 1, l_3, \dots, l_t].$

Let $S(r') = (T_1, T_2, T_1, T_2)$ be the decomposition of S(r') given by Proposition 4.3, as in the proof of Proposition 4.5. By Corollary 4.6,

CT(r) = CS(r') and CT(s) = CS(s'),

so it follows that (T_1, T_2) or (T_2, T_1) appears in CS(s') as a subsequence, because (S_1, S_2) or (S_2, S_1) appears in CS(s) as a subsequence, by assumption. As in Case 1, the induction completes the case.

Remark 5. We can easily see that the a rational number $s \in (0, 1]$ satisfies the conclusion of Lemma 7.3 if and only if s lies in the open interval $(r_1, r_2) = (0, 1] - (I_1 \cup I_2)$, where r_1 and r_2 are rational numbers such that $I_1 = [0, r_1]$ and $I_2 = [r_2, 1]$, introduced in the paragraph preceding Lemma 7.1.

We are now in a position to prove Theorem 7.2, i.e., the only if part of Main Theorem 2.3.

Proof of Theorem 7.2. Consider a 2-bridge link K(r) with 0 < r < 1, and pick a rational number s from $I_1 \cup I_2$. Suppose on the contrary that α_s is null-homotopic in $S^3 - K(r)$, namely $u_s = 1$ in G(K(r)). If $s \in (0, 1]$, then we see by Corollary 6.4 that CS(s) contains (S_1, S_2) or (S_2, S_1) as a subsequence. Hence, we see by Lemma 7.3 and Remark 5 that $s \in (r_1, r_2) = (0, 1] - (I_1 \cup I_2)$, a contradiction. So, the only possibility is s = 0. This case can be handled by directly using Theorem 6.3, which implies that u_s must contain a subword w of $(u_r^{\pm 1})$ such that the S-sequence of w is (S_1, S_2, ℓ) or (ℓ, S_2, S_1) for some positive integer ℓ . Note that the length of such a subword w is strictly greater than p, half the length of $(u_r^{\pm 1})$, where r = q/p. Since 0 < r < 1, we have $p \geq 2$. So, the word $u_0 = ab$ cannot contain such a subword, a contradiction. This completes the proof of Theorem 7.2. Thus we have proved Main Theorem 2.3 except for the case r = 0 and $r = \infty$. These exceptional cases are treated as follows. Suppose $r = \infty$, then $K(\infty)$ is the trivial 2-component link, and $G(K(\infty))$ is the free group F(a, b). On the other hand, for every $s \in \mathbb{Q}$, u_s is a non-trivial cyclically reduced word in $\{a, b\}$ and hence it represents a non-trivial element of $G(K(\infty))$. On the other hand, the $\hat{\Gamma}_r$ -orbit of $\{\infty, r\} = \{\infty\}$ is the singleton $\{\infty\}$. Hence Main Theorem 2.3 holds for this case. Next, suppose r = 0. Then $G(K(0)) = \langle a, b | ab \rangle \cong \mathbb{Z}$. Further, $\hat{\Gamma}_r$ is equal to the group generated by the reflections in the edges of any of \mathcal{D} . In particular, any Farey triangle is a fundamental domain for the action of $\hat{\Gamma}_r$ on \mathbb{H}^2 . Hence, any $s \in \hat{\mathbb{Q}}$ belongs to the $\hat{\Gamma}_r$ -orbit of one and only one of $\{0, 1, \infty\}$. On the other hand, $u_1 = ab^{-1} = a^2 \neq 1$ in G(K(0)). Hence, Main Theorem 2.3 holds for this case. This completes the proof of Main Theorem 2.3.

Remark 6. The assertion in [25, Example 4.2] that $\hat{\Gamma}_1$ acts transitively on $\hat{\mathbb{Q}}$ is obviously incorrect. It should be noted that though there is an upper-meridian-pair preserving epimorphism (actually an isomorphism) from $G(K(1)) = \langle a, b | ab^{-1} \rangle$ to $G(K(0)) = \langle a, b | ab \rangle$, it does not send the pair (a, b) to (a, b).

At the end of this section, we describe a geometric intuition behind the proof of the main theorem. Note that a slope s belongs to $I_1 \cup I_2$ if and only if it does not belong to $(-\infty, 0) \cup (1, \infty]$ nor (r_1, r_2) . The condition that $s \notin (-\infty, 0) \cup (1, \infty]$, i.e., $s \in [0, 1]$, implies that the word u_s can be read from a line of slope r in \mathbb{R}^2 "effectively" so that $S(s) = S(u_s)$ (see Remark 4). To describe the geometric meaning of the condition $s \notin (r_1, r_2)$, set p_i and q_i be relatively prime integers such that $r_i = q_i/p_i$ (i = 1, 2). Then $(p,q) = (p_1 + p_2, q_1 + q_2)$, where r = q/p, and the parallelogram in \mathbb{R}^2 spanned by (0,0), (p_1,q_1) , (p_2,q_2) and (p,q) does not contain lattice points in its interior. If $s \in (r_1, r_2)$, then the ray (in the first quadrant) of slope s from the origin passes through the interior of the parallelogram and hence the word u_s shares a long common initial subword with u_r . On the other hand, if $s \notin (r_1, r_2) \cup (-\infty, 0) \cup (1, \infty]$, then the ray (in the first quadrant) of slope s from the origin is disjoint from the interior of the parallelogram, and hence, u_s shares only a short initial subword with u_r . This convinces us that the cyclic word (u_s) , for $s \notin (r_1, r_2) \cup (-\infty, 0) \cup (1, \infty]$, shares only short common subwords with the cyclic word (u_r) . This is the intuition behind the proof of the main theorem.

We realized through discussion with Norbert A'Campo that the decomposition $S(r) = (S_1, S_2, S_1, S_2)$ in Proposition 4.5 has a natural geometric interpretation in terms of the above parallelogram. To describe it, assume $q_1/p_1 < q/p < q_2/p_2$ in the above setting, and consider the infinite broken line, *B*, obtained by joining the lattice points

$$\dots, (0,0), (p_2,q_2), (p,q), (p+p_2,q+q_2), (2p,2q), \dots$$

which is invariant by the translation $(x, y) \mapsto (x + p, y + q)$. By slightly modifying B near the lattice points, we obtain a (topological) line, B^+ , in $\mathbb{R}^2 - \mathbb{Z}^2$, invariant by the translation, which is homotopic to the line $L^+(r)$ in the proof of Lemma 4.7. Pick a point, $z_0 \in B^+$ in the second quadrant, and consider the sub-path of B^+ bounded by z_0 and $z_4 := z_0 + (2p, 2q)$. Then the word u_r is also obtained by reading the intersection of the sub-path with the vertical lattice lines. Pick a point $z_1 \in B^+$ whose x-coordinate is $p_2 +$ (small positive number), and set $z_2 := z_0 + (p, q)$ and $z_3 := z_1 + (p, q)$. Let B_i^+ be the sub-path of B^+ joining z_{i-1} with z_i (i = 1, 2, 3, 4). Then we can see that the subword of u_r corresponding to B_i^+ is equal to the word v_i in Section 5, i.e., $u_r \equiv v_1 v_2 v_3 v_4$, $S_1 = S(v_1) = S(v_3)$ and $S_2 = S(v_2) = S(v_4)$. In particular, $|v_1| = |v_3| = p_2 + 1$ and $|v_2| = |v_4| = p_1 - 1$. We hope to fully describe this on another occasion.

8. Relation with a question by Minsky

In this section, we describe the relation of Main Theorem 2.3 with the question raised by Minsky in [7, Question 5.4]. Let $M = H_+ \cup_S H_-$ be a Heegaard splitting of a 3-manifold M. Let $\Gamma_{\pm} := MCG(H_{\pm})$ be the mapping class group of H_{\pm} , and let Γ_{\pm}^0 be the kernel of the map $MCG(H_{\pm}) \to \text{Out}(\pi_1(H_{\pm}))$. Identify Γ_{\pm}^0 with a subgroup of MCG(S), and consider the subgroup $\langle \Gamma_+^0, \Gamma_-^0 \rangle$ of MCG(S). Now let Δ_{\pm} be the set of (isotopy classes of) simple loops in Swhich bound a disk in H_{\pm} . Let Z be the set of essential simple loops in Swhich are null-homotopic in M. Note that Z contains Δ_{\pm} and invariant under $\langle \Gamma_+^0, \Gamma_-^0 \rangle$. In particular, the orbit $\langle \Gamma_+^0, \Gamma_-^0 \rangle (\Delta_+ \cup \Delta_-)$ is a subset of Z. Then Minsky posed the following question.

Question 2. When is Z equal to the orbit $\langle \Gamma^0_+, \Gamma^0_- \rangle (\Delta_+ \cup \Delta_-)$?

The above question makes sense not only for Heegaard splittings but also bridge decompositions of knots and links. Actually, the groups Γ_{∞} and Γ_r in our setting correspond to the groups Γ_+^0 and Γ_-^0 , and hence the group $\hat{\Gamma}_r$ corresponds to the group $\langle \Gamma_+^0, \Gamma_-^0 \rangle$. To make this precise, recall the bridge decomposition $(S^3, K(r)) = (B^3, t(\infty)) \cup (B^3, t(r))$, and let $\tilde{\Gamma}_+$ (resp. $\tilde{\Gamma}_-$) be the mapping class group of the pair $(B^3, t(\infty))$ (resp. $(B^3, t(r))$), and let $\tilde{\Gamma}_{\pm}^0$ be the kernel of the natural map $\tilde{\Gamma}_+ \to \text{Out}(\pi_1(B^3 - t(\infty)))$ (resp. $\tilde{\Gamma}_{-} \to \operatorname{Out}(\pi_1(B^3 - t(r))))$. Identify $\tilde{\Gamma}^0_{\pm}$ with a subgroup of the mapping class group $MCG(\mathbf{S})$ of the 4-times punctured sphere \mathbf{S} . Recall that the Farey tessellation \mathcal{D} is identified with the curve complex of \mathbf{S} and there is a natural map from $MCG(\mathbf{S})$ to the automorphism group $Aut(\mathcal{D})$ of \mathcal{D} , whose kernel is equal to the image of the $(\mathbb{Z}/2\mathbb{Z})^2$ -action on \mathbf{S} , which appeared in the proof of Lemma 3.2. Then the group Γ_{∞} (resp. Γ_r) introduced in Section 2 is identified with the image of $\tilde{\Gamma}^0_+$ (resp. $\tilde{\Gamma}^0_-$) by this natural map. Moreover, the sets $\{\alpha_{\infty}\}$ and $\{\alpha_r\}$ correspond to the sets Δ_+ and Δ_- , and Main Theorem 2.3 says that the set Z of simple loops in \mathbf{S} which are null-homotopic in $S^3 - K(r)$ is equal to the orbit $\langle \Gamma_{\infty}, \Gamma_r \rangle (\Delta_+ \cup \Delta_-)$. Thus Main Theorem 2.3 may be regarded as an answer to the special variation of Question 2.

Finally, we note that Main Theorem 2.3 is also related to the existence of a possible variation of McShane's identity for 2-bridge knots (see [29]). Related topics are studied in subsequent papers [19, 20, 21, 22]. For an overview of this series of works, please see the research announcement [23].

References

- M. Boileau, S. Boyer, A.W. Reid and S. Wang, Simon's conjecture for 2-bridge knots, Comm. Anal. Geom. 18 (2010), 121–144.
- [2] M. Boileau, J.H. Rubinstein and S. Wang, Finiteness of 3-manifolds associated with non-zero degree mappings, arXiv:math/0511541.
- [3] G. Burde and H. Zieschang, *Knots*, de Gruyter Studies in Math. 5, Walter de Gruyter, 1985.
- [4] R.H. Crowell and R.H. Fox, Introduction to knot theory, Reprint of the 1963 original, Graduate Texts in Math. 57, Springer-Verlag, 1977.
- [5] F. Gonzaléz-Acũna and A. Ramírez, Two-bridge knots with property Q, Quart. J. Math. 52 (2001), 447–454.
- [6] F. Gonzaléz-Acũna and A. Ramírez, Epimorphisms of knot groups onto free products, Topology 42 (2003), 1205–1227.
- [7] C. Gordon, *Problems*, Workshop on Heegaard Splittings, 401–411, Geom. Topol. Monogr. 12, Geom. Topol. Publ., Coventry, 2007.
- [8] C. Gordon and J. Luecke, Knots are determined by their complements, Bull. Amer. Math. Soc. 20 (1989), 83–87.
- [9] M. Hirasawa and K. Murasugi, Fibred double torus knots which are band-sums of torus knots, Osaka J. Math. 44 (2007), 11–70.
- [10] K. Horie, T. Kitano, M. Matsumoto and M. Suzuki, A partial order on the set of prime knots with up to 11 crossings, arXiv:0906.3943.
- [11] J. Hoste and P.D. Shanahan, Epimorphisms and boundary slopes of 2-bridge knots, Algebr. Geom. Topol. 10 (2010), 1221–1244.
- [12] A. Kawauchi, A survey of knot theory, Translated and revised from the 1990 Japanese original, Birkhäuser Verlag, Basel, 1996.

- [13] A. Kawauchi, *Topological imitations*, Lectures at KNOTS '96 (Tokyo), 19–37, Ser. Knots Everything, 15, World Sci. Publishing, River Edge, NJ, 1997.
- [14] R. Kirby, Problems in low-dimensional topology, AMS/IP Stud. Adv. Math., 2.2, Geometric topology (Athens, GA, 1993), 35–473, Amer. Math. Soc., Providence RI, 1997.
- [15] T. Kitano and M. Suzuki, A partial order in the knot table, Experiment. Math. 14 (2005), 385–390.
- [16] T. Kitano and M. Suzuki, A partial order in the knot table (II), Acta Math. Sin. (Engl. Ser.) 24 (2008), 1801–1816.
- [17] T. Kitano, M. Suzuki and M. Wada, Twisted Alexander polynomials and surjectivity of a group homomorphism, Algebr. Geom. Topol. 5 (2005), 1315–1324.
- [18] Y. Komori and C. Series, *The Riley slice revisited*, in the Epstein Birthday Schrift, I. Rivin, C. Rourke and C. Series eds, Geom. Topol. Monogr. 1 (1999), 303–316.
- [19] D. Lee and M. Sakuma, Homotopically equivalent simple loops on 2-bridge spheres in 2-bridge link complements (I), arXiv:1010.2232.
- [20] D. Lee and M. Sakuma, Homotopically equivalent simple loops on 2-bridge spheres in 2-bridge link complements (II), arXiv:1103.0856.
- [21] D. Lee and M. Sakuma, Homotopically equivalent simple loops on 2-bridge spheres in 2-bridge link complements (III), arXiv:1111.3562.
- [22] D. Lee and M. Sakuma, A variation of McShane's identity for 2-bridge links, preliminary notes.
- [23] D. Lee and M. Sakuma, Simple loops on 2-bridge spheres in 2-bridge link complements, Electron. Res. Announc. Math. Sci. 18 (2011), 97–111.
- [24] R.C. Lyndon and P.E. Schupp, Combinatorial Group Theory, Springer-Verlag, Berlin, 1977.
- [25] T. Ohtsuki, R. Riley and M. Sakuma, *Epimorphisms between 2-bridge link groups*, Geom. Topol. Monogr. 14 (2008), 417–450.
- [26] J.G. Ratcliffe, Foundations of Hyperbolic Manifolds, Graduate Texts in Math. 149, Springer-Verlag, 1994.
- [27] A. Reid, S. Wang and Q. Zhou, Generalized Hopfian property, a minimal Haken manifold and epimorphisms between 3-manifold groups, Acta Mathematica Sinica, English Series 18 (2002), 157–172.
- [28] R. Riley, Parabolic representations of knot groups. I, Proc. London Math. Soc. 24 (1972), 217–242.
- [29] M. Sakuma, Variations of McShane's identity for the Riley slice and 2-bridge links, In "Hyperbolic Spaces and Related Topics", R.I.M.S. Kokyuroku 1104 (1999), 103–108.
- [30] M. Sakuma, Epimorphisms between 2-bridge knot groups from the view point of Markoff maps, Intelligence of low dimensional topology 2006, 279–286, Ser. Knots Everything, 40, World Sci. Publ., 2007.
- [31] H. Schubert, Knoten mit zwei Brücken, Math. Z. 65 (1956), 133-170.
- [32] D. Silver and W. Whitten, Knot group epimorphisms, J. Knot Theory Ramifications 15 (2006), 153–166.
- [33] J. Simon, On knots with nontrivial interpolating manifolds, Trans. Amer. Math. Soc. 160 (1971), 467–473.
- [34] T. Soma, Non-zero degree maps to hyperbolic 3-manifolds, J. Differential Geom. 49 (1998), 517–546.

- [35] T. Soma, Epimorphism sequences between hyperbolic 3-manifold groups, Proc. Amer. Math. Soc. 130 (2002), 1221–1223.
- [36] S. Wang, Non-zero degree maps between 3-manifolds. Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 457–468, Higher Ed. Press, Beijing, 2002.

DEPARTMENT OF MATHEMATICS, PUSAN NATIONAL UNIVERSITY, SAN-30 JANGJEON-DONG, GEUMJUNG-GU, PUSAN, 609-735, KOREA

E-mail address: donghi@pusan.ac.kr

Department of Mathematics, Graduate School of Science, Hiroshima University, Higashi-Hiroshima, 739-8526, Japan

 $E\text{-}mail\ address: \texttt{sakuma@math.sci.hiroshima-u.ac.jp}$