A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

Toshimitsu TAKAESU

Faculty of Mathematics, Kyushu University, Fukuoka, 812-8581, Japan

Abstract. In this article we consider linear operators satisfying a generalized commutation relation of a type of the Heisenberg-Lie algebra. It is proven that a generalized inequality of the Hardy's uncertainty principle lemma follows. Its applications to time operators and abstract Dirac operators are also investigated.

Key words : weak commutation relations, Heisenberg-Lie algebra, time operators, Hamiltonians, time-energy uncertainty relation, Dirac operators, essential self-adjointness. **MSC 2010** : 81Q10, 47B25, 46L60.

1 Introduction and Results

In this article we investigate a norm-inequality of the linear operators which obey a generalized weak commutation relation of a type of the Heisenberg-Lie algebra, and consider its application to the theory of the time operator [7, 2], and an abstract Dirac operator. Let $\mathbf{X} = \{X_j\}_{j=1}^N$, $\mathbf{Y} = \{Y_j\}_{j=1}^N$ and $\mathbf{Z} = \{Z_j\}_{j=1}^N$ be symmetric operators on a Hilbert space \mathcal{H} . The weak commutator of operators *A* and *B* is defined for $\psi \in \mathcal{D}(A) \cap \mathcal{D}(B)$ and $\phi \in \mathcal{D}(A^*) \cap \mathcal{D}(B^*)$ by

$$[A,B]^{\mathrm{w}}(\phi,\psi) = (A^*\phi,B\psi) - (B^*\phi,A\psi).$$

Here the inner product has a linearity of $(\eta, \alpha \psi + \beta \phi) = \alpha(\eta, \psi) + \beta(\eta, \phi)$ for $\alpha, \beta \in \mathbb{C}$. We assume that $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ satisfies the following conditions.

(A.1) Z_j , $1 \le j \le N$, is bounded operator.

(A.2) Let
$$\mathcal{D}_{\mathbf{X}} = \bigcap_{j=1}^{N} \mathcal{D}(X_{j})$$
 and $\mathcal{D}_{\mathbf{Y}} = \bigcap_{j=1}^{N} \mathcal{D}(Y_{j})$. It follows that for $\phi, \psi \in \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$,
 $[X_{j}, Y_{l}]^{\mathrm{w}}(\phi, \psi) = \delta_{j,l}(\phi, iZ_{j}\psi),$
 $[X_{j}, Z_{l}]^{\mathrm{w}}(\phi, \psi) = [Y_{j}, Z_{l}]^{\mathrm{w}}(\phi, \psi) = 0$
 $[X_{j}, X_{l}]^{\mathrm{w}}(\phi, \psi) = [Y_{j}, Y_{l}]^{\mathrm{w}}(\phi, \psi) = [Z_{j}, Z_{l}]^{\mathrm{w}}(\phi, \psi) = 0.$

Note that $[Z_j, Z_l] \psi = 0$ follows for $\psi \in \mathcal{H}$, since Z_j , $j = 1, \dots, N$, is bounded. In this article we consider an generalization of the inequality

$$\int_{\mathbf{R}^N} \frac{1}{|\mathbf{r}|^2} |u(\mathbf{r})|^2 d\mathbf{r} \leq \frac{4}{(N-2)^2} \int_{\mathbf{R}^N} |\nabla u(\mathbf{r})|^2 d\mathbf{r}, \qquad N \geq 3$$

This inequality is a basic one of Hardy's uncertainty principle inequalities. For Hardy's uncertainty inequalities, refer to e.g. [5, 6, 13].

Let us introduce the additional conditions.

(A.3) X_j is self-adjoint for all $1 \le j \le N$.

(A.4) X_i and Z_l strongly commutes for all $1 \le j \le N$ and $1 \le l \le N$.

Sicne Z_j , $j = 1, \dots, N$, is bounded self-adjoint operator, we can set $\lambda_{\min}(\mathbf{Z})$ and $\lambda_{\max}(\mathbf{Z})$ by

$$\begin{split} \lambda_{\min}(\mathbf{Z}) &= \min_{1 \leq j \leq N} \inf \sigma(Z_j), \\ \lambda_{\max}(\mathbf{Z}) &= \max_{1 \leq j \leq N} \sup \sigma(Z_j), \end{split}$$

where $\sigma(O)$ denotes the spectrum of the operator O.

Theorem 1 Assume (A.1)-(A.4). Let $\Psi \in \mathcal{D}(|\mathbf{X}|^{-1}) \cap \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$. Then the following (1) and (2) hold (1) If $N\lambda_{min}(\mathbf{Z}) - 2\lambda_{max}(\mathbf{Z}) > 0$, it follows that

$$\left\| \|\mathbf{X}\|^{-1}\Psi \right\|^{2} \leq \frac{4}{\left(N\lambda_{min}(\mathbf{Z}) - 2\lambda_{max}(\mathbf{Z})\right)^{2}} \sum_{j=1}^{N} \left\| Y_{j}\Psi \right\|^{2}.$$
 (1)

(2) If $2\lambda_{min}(\mathbf{Z}) - N\lambda_{max}(\mathbf{Z}) > 0$, it follows that

$$\left\| |\mathbf{X}|^{-1} \Psi \right\| \leq \frac{4}{\left(2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z})\right)^2} \sum_{j=1}^N \left\| Y_j \Psi \right\|^2.$$
⁽²⁾

Before proving Theorem 1, let us consider the replacement of **X** and **Y** in Theorem 1. Let us introduce the following conditions substitute for (**A.3**) and (**A.4**).

(A.5) Y_j is self-adjoint for all $1 \le j \le N$.

(A.6) Y_i and Z_l strongly commutes for all $1 \le j \le N$ and $1 \le l \le N$.

It is seen from (A.2), that

$$[Y_j, X_l]^{\mathsf{w}}(\phi, \psi) = \delta_{j,l}(\phi, i(-Z_j)\psi), \qquad \phi, \psi \in \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}.$$
(3)

Note that $\inf \sigma(-Z_j) = -\sup(Z_j)$ and $\sup(-Z_j) = -\inf \sigma(Z_j)$ follow. Then we obtain a following corollary :

Corollary 2 Assume (A.1)-(A.2) and (A.5)-(A.6). Let $\Psi \in \mathcal{D}(|\mathbf{Y}|^{-1}) \cap \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$. Then the following (1) and (2) hold.

(1) If $2\lambda_{min}(\mathbf{Z}) - N\lambda_{max}(\mathbf{Z}) > 0$, it follows that

$$\left\| |\mathbf{Y}|^{-1} \Psi \right\| \leq \frac{4}{\left(2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z})\right)^2} \sum_{j=1}^N \left\| X_j \Psi \right\|^2.$$
(4)

(2) If $N\lambda_{min}(\mathbf{Z}) - 2\lambda_{max}(\mathbf{Z}) > 0$, it follows that

$$\left\| |\mathbf{Y}|^{-1} \Psi \right\| \leq \frac{4}{\left(N \lambda_{min}(\mathbf{Z}) - 2 \lambda_{max}(\mathbf{Z}) \right)^2} \sum_{j=1}^{N} \left\| X_j \Psi \right\|^2.$$
(5)

(Proof of Theorem 1)

(1)Let $\Psi \in \mathcal{D}(|\mathbf{X}|^{-1}) \cap \mathcal{D}_{\mathbf{X}} \cap \mathcal{D}_{\mathbf{Y}}$. For $\varepsilon > 0$ and t > 0, it is seen that

$$\| (Y_j - itX_j(\mathbf{X}^2 + \varepsilon)^{-1}) \Psi \|^2 = \| Y_j \Psi \|^2 - it [Y_j, X_j(\mathbf{X}^2 + \varepsilon)^{-1}]^w (\Psi, \Psi) + t^2 \| X_j(\mathbf{X}^2 + \varepsilon)^{-1} \Psi \|^2.$$
(6)

We see that

$$[Y_j, X_j(\mathbf{X}^2 + \varepsilon)^{-1}]^{w}(\Psi, \Psi) = [Y_j, X_j]^{w}(\Psi, (\mathbf{X}^2 + \varepsilon)^{-1}\Psi) + [Y_j, (\mathbf{X}^2 + \varepsilon)^{-1}]^{w}(X_j\Psi, \Psi).$$
(7)

From (A.2) and (A.4), we obtain that

$$[Y_j, X_j]^{\mathsf{w}}(\Psi, (\mathbf{X}^2 + \varepsilon)^{-1}\Psi) = -i((\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi, Z_j(\mathbf{X}^2 + \varepsilon)^{-1/2}\Psi).$$
(8)

Note that for a symmetric operator *A* and the non-negative symmetric operator *B*, the resolvent formula $[A, (B+\lambda)^{-1}]^{w}(v, u) = [B, A]^{w}((B+\lambda)^{-1}v, (B+\lambda)^{-1}u)$ for $\lambda > 0$ follows. Then by using this formura, **(A.2)** and **(A.4)** yield that

$$[Y_j, (\mathbf{X}^2 + \varepsilon)^{-1}]^{\mathsf{w}}(X_j \Psi, \Psi) = 2i(X_j(\mathbf{X}^2 + \varepsilon)^{-1}u, Z_j X_j(\mathbf{X}^2 + \varepsilon)^{-1}u)$$
(9)

Since $\left\| \left(Y_j - it X_j (\mathbf{X}^2 + \varepsilon)^{-1} \right) u \right\|^2 \ge 0$ and t > 0, we see from (7), (8) and (9) that

$$||Y_j\Psi||^2$$

$$\geq -t^{2} \left\| X_{j} (\mathbf{X}^{2} + \varepsilon)^{-1} u \right\|^{2} + t \left((\mathbf{X}^{2} + \varepsilon)^{-1/2} \Psi, Z_{j} (\mathbf{X}^{2} + \varepsilon)^{-1/2} u \right) - 2t \left(X_{j} (\mathbf{X}^{2} + \varepsilon)^{-1} u, Z_{j} X_{j} (\mathbf{X}^{2} + \varepsilon)^{-1} \Psi \right) \\ \geq \left(-t^{2} - 2t \lambda_{\max}(\mathbf{Z}) \right) \left\| X_{j} (\mathbf{X}^{2} + \varepsilon)^{-1} u \right\|^{2} + t \lambda_{\min}(\mathbf{Z}) \left\| (\mathbf{X}^{2} + \varepsilon)^{-1/2} \Psi \right\|.$$

$$(10)$$

Then we have that

$$\sum_{j=1}^{N} \|Y_{j}\Psi\|^{2} \geq \left(-t^{2} - 2t\lambda_{\max}(\mathbf{Z})\right) \left\| |\mathbf{X}|(\mathbf{X}^{2} + \varepsilon)^{-1}\Psi \right\|^{2} + tN\lambda_{\min}(\mathbf{Z}) \|(\mathbf{X}^{2} + \varepsilon)^{-1/2}\Psi\|.$$
(11)

Note that $\lim_{\varepsilon \to 0} \left\| |\mathbf{X}| (\mathbf{X}^2 + \varepsilon)^{-1} \Psi \right\|^2 = \| |\mathbf{X}|^{-1} \Psi \|$ and $\lim_{\varepsilon \to 0} \| (\mathbf{X}^2 + \varepsilon)^{-1/2} \Psi \| = \| |\mathbf{X}|^{-1} \Psi \| = 0$ follow from the spectral decomposition theorem. Then we have

$$\sum_{j=1}^{N} \|Y_{j}\Psi\|^{2} \geq \left(-t^{2} + (N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z}))t\right) \| \|\mathbf{X}\|^{-1}\Psi\|.$$
(12)

By taking $t = \frac{N\lambda_{\min}(\mathbf{Z}) - 2\lambda_{\max}(\mathbf{Z})}{2} > 0$ in the right side of (12), we obtain (1). (2) By computing $\left\| \left(Y_j + itX_j(\mathbf{X}^2 + \varepsilon)^{-1} \right) \Psi \right\|^2$ for t > 0 and $\varepsilon > 0$, in a similar way of (1), we see that $\|Y_j\Psi\|^2$

$$\geq -t^{2} \left\| X_{j}(\mathbf{X}^{2} + \varepsilon)^{-1} u \right\|^{2} - t \left((\mathbf{X}^{2} + \varepsilon)^{-1/2} \Psi, Z_{j}(\mathbf{X}^{2} + \varepsilon)^{-1/2} u \right) + 2t \left(X_{j}(\mathbf{X}^{2} + \varepsilon)^{-1} u, Z_{j} X_{j}(\mathbf{X}^{2} + \varepsilon)^{-1} \Psi \right) \\\geq \left(-t^{2} + 2t \lambda_{\min}(\mathbf{Z}) \right) \left\| X_{j}(\mathbf{X}^{2} + \varepsilon)^{-1} u \right\|^{2} - t \lambda_{\max}(\mathbf{Z}) \| (\mathbf{X}^{2} + \varepsilon)^{-1/2} \Psi \|.$$

$$(13)$$

Then by taking $\varepsilon \to 0$ in the right side of (13), it follows that

$$\sum_{j=1}^{N} \|Y_{j}\Psi\|^{2} \geq \left(-t^{2} + (2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}))t\right) \| \|\mathbf{X}\|^{-1}\Psi\|.$$
(14)

By taking $t = \frac{(2\lambda_{\min}(\mathbf{Z}) - N\lambda_{\max}(\mathbf{Z}))}{2} > 0$ in (14), we obtain (2).

2 Applications

2.1 Time-Energy Uncertainty inequality

In this subsection we consider an application to the theory of time operators [2, 7]. Let H, T, and C be linear operators on a Hilbert space H. It is said that H has the weak time operator T with the uncommutative factor C if (H,T,C) satisfy the following conditions.

(**T.1**) H and T are symmetric.

(T.2) C is bounded and self-adjoint.

(T.3) It follows that for $\phi, \psi \in \mathcal{D}(H) \cap \mathcal{D}(T)$,

$$[T,H]^{\mathrm{w}}(\phi,\psi) = (\phi,C\psi).$$

(T.4)

$$\delta_C := \inf_{\psi \in (\ker C)^\perp \setminus \{0\}} rac{|(\Psi, C \Psi)|}{\|\psi\|^2} \, > \, 0.$$

Assume that (H, T, C) satisfies **(T.1)-(T.4)**. Then by using $||Au|| ||Bu|| \ge |\text{Im}(Au, Bu)| \ge \frac{1}{2} |[A, B]^w(u, u)|$, it is seen that (H, T, C) satisfies the time-energy uncertainty inequality ([2], Proposition4.1):

$$\frac{\| (H - \langle H \rangle_{\psi}) \psi \| \| (T - \langle T \rangle_{\psi}) \psi \|}{\| \psi \|^2} \geq \frac{\delta_C}{2}, \qquad \psi \in \mathcal{D}(H) \cap \mathcal{D}(T), \tag{15}$$

where $\langle O \rangle_{\psi} = (\psi, O\psi)$. From (2) in Theorem 1 and (1) in Corollary 2, we obtain another type of the inequality between *T* and *H*:

Corollary 3 (Time-Energy Uncertainty Inequalities)

Assume (T.1)-(T.3). Then the following (i) and (ii) hold.

(i) If T is self-adjoint, C and T strongly commute, and $\sup \sigma(C) < 2 \inf \sigma(C)$, it follows that for $\psi \in \mathcal{D}(|T|^{-1}) \cap \mathcal{D}(T) \cap \mathcal{D}(H)$,

$$\left\| |T|^{-1} \psi \right\| \leq \frac{2}{2 \inf \sigma(C) - \sup \sigma(C)} \left\| H \Psi \right\|.$$
(16)

(ii) If *H* is self-adjoint, *C* and *H* strongly commute, and $\sup \sigma(C) < 2\inf \sigma(C)$, it follows that for $\psi \in \mathcal{D}(|H|^{-1}) \cap \mathcal{D}(H) \cap \mathcal{D}(T)$,

$$\left\| |H|^{-1} \psi \right\| \leq \frac{2}{2 \inf \sigma(C) - \sup \sigma(C)} \left\| T \Psi \right\|.$$
(17)

2.2 Abstract Dirac Operators with Coulomb Potential

Next tlt us consider the application to abstract Dirac operators. We consider the self-adjoint operators $\mathbf{P} = \{P_j\}_{j=1}^N$ and $\mathbf{Q} = \{Q_j\}_{j=1}^N$ on a Hilbert space \mathcal{H} . Let us set a subspace $\mathcal{D} \subset \bigcap_{j,l} (\mathcal{D}(P_j) \cap \mathcal{D}(Q_l))$. It is said that $(\mathcal{H}, \mathcal{D}, \mathbf{P}, \mathbf{Q})_N$ is the weak representation of the CCR with degree *N*, if \mathcal{D} is dense in \mathcal{H} and it follows that for $\phi, \psi \in \mathcal{D}$,

$$\begin{split} & [P_j, Q_l]^{\mathrm{w}}(\phi, \psi) = i\delta_{j,l}(\phi, \psi), \\ & [P_j, P_l]^{\mathrm{w}}(\phi, \psi) = [Q_j, Q_l]^{\mathrm{w}}(\phi, \psi) = 0. \end{split}$$

Let us define an abstract Dirac operator as follows. Let $(\mathcal{H}, \mathcal{D}, \mathbf{P}, \mathbf{Q})_3$ be the weak representation of the CCR with degree three. Let $\mathbf{A} = \{A_j\}_{j=1}^3$ and *B* be the bounded self-adjoint operators on a Hilbert space \mathcal{K} . Here $\mathbf{A} = \{A_j\}_{j=1}^3$ and *B* satisfy the canonical anti-commutation relations $\{A_j, A_l\} = 2\delta_{j,l}, \{A_j, B\} = 0, B^2 = I_{\mathcal{K}}$ where $I_{\mathcal{K}}$ is the identity operator on \mathcal{K} . The state Hilbert space space is defined by $\mathcal{H}_{\text{Dirac}} = \mathcal{K} \otimes \mathcal{H}$. The free abstract Dirac operator is defiend by

$$H_0 = \sum_{j=1}^3 A_j \otimes P_j + B \otimes M.$$

Here we assume the following condition.

(D.1) P_j and P_l strongly commute for $1 \le j \le 3$, $1 \le l \le 3$. P_j , $1 \le j \le 3$, and M strongly commute.

Then it is seen that $H_0^2 \Psi = (\mathbf{P}^2 + M^2) \Psi$ for $\Psi \in \mathcal{D}$. The abstract Dirac Operator with the Coulomb potential is defined by

$$H(\kappa) = H_0 + \kappa I_{\mathcal{K}} \otimes |\mathbf{Q}|^{-1}$$

where $\kappa \in \mathbf{R}$ is a parameter called the coupling constant. We assume that the following condition

(D.2) It follows that $\mathcal{D} \subset \mathcal{D}(\mathbf{Q}|^{-1})$.

Then it follows from (1) in Theorem 1 that for $\psi \in \mathcal{D}$,

$$\|I_{\mathcal{K}} \otimes |\mathbf{Q}|^{-1} \psi\|^2 \le 4 \sum_{j=1}^3 \|P_j \Psi\|^2 \le 4 \|H_0 \Psi\|^2.$$

Hence by the Kato-Rellich theorem, we obtain the following corollary.

Corollary 4 Assume (D.1) and (D.2). Then for $|\kappa| < \frac{1}{2}$, $H(\kappa)$ is essentially self-adjoint on \mathbb{D} .

Acknowledgments

It is pleasure to thank assistant professor Akito Suzuki and associate professor Fumio Hiroshima for their advice and comments.

References

- [1] A.Arai, Mathematical principles of quantum phenamena, Asakura-syoten, 2005. (in japanese)
- [2] A.Arai, Generalized weak Weyl relation and decay of quantum dynamics, *Rev. Math. Phys.* 17 (2005) 1071-1109.
- [3] A.Arai, Heisenberg Operators, invariant domains and Heisenberg equations of motion, *Rev. Math. Phys* 19 (2007) 1045-1069.
- [4] A.Arai, Spectrum of time operators. Lett. Math. Phys. 80 211-221 (2007).
- [5] G.B.Folland and A.Sitaram, The uncertainty principle : A mathematical survey, *J. Fourier Anal. Appl.* **3** (1997) 207-238.
- [6] V.Havin and B.Joricke, The uncertainty principle in harmonic analysis, Springer 1994.
- [7] M.Miyamoto, A generalized Weyl relation approach to the time operator and its connection to the survival probability, *J. Math. Phys* **42** (2001) 1038-1052.
- [8] J.G.Muga, R.S.Mayato, and I.L.Egsquiza (eds.), Time in quantum mechanics. Springer 2002.
- [9] P.Pfeifer and J. Frölich, Generalized time-energy uncertainty relations and bounds on lifetimes of resonances, *Rev. Mod. Phys.* 67 (1995) 759-779.
- [10] M.Reed and B.Simon, Methods of Modern Mathematical Physics Vol.II, Academic Press, 1979.
- [11] K.Schmüdgen, On the Heisenberg commutationn relationhn. I, J. Funct. Anal. 50 (1983) 8-49.
- [12] B. Thaller, *The Dirac equation*, Springer, 1992.
- [13] S.Thangavelu, An introduction to the uncertainty principle : Hardy's theorem on Lie groups, Birkhäuser, 2004.