

**JACOB'S LADDERS AND THE NONLOCAL INTERACTION OF  
THE FUNCTION  $Z^2(t)$  WITH THE FUNCTION  $\tilde{Z}^2(t)$  ON THE  
DISTANCE  $\sim (1 - c)\pi(t)$  FOR THE COLLECTIONS OF  
DISCONNECTED SETS**

JAN MOSER

ABSTRACT. It is shown in this paper that there is a fine correlation of the fourth order between the functions  $Z^2[\varphi_1(t)]$  and  $\tilde{Z}^2(t)$ , respectively. This correlation is with respect to two collections of disconnected sets. Corresponding new asymptotic formulae cannot be obtained within known theories of Balasubramanian, Heath-Brown and Ivic.

1. THE RESULT

In this paper we obtain some new properties of the signal

$$Z(t) = e^{i\vartheta(t)} \zeta\left(\frac{1}{2} + it\right)$$

generated by the Riemann zeta-function, where

$$(1.1) \quad \vartheta(t) = -\frac{t}{2} \ln \pi + \operatorname{Im} \ln \Gamma\left(\frac{1}{4} + i\frac{t}{2}\right) = \frac{t}{2} \ln \frac{t}{2\pi} - \frac{t}{2} - \frac{\pi}{8} + \mathcal{O}\left(\frac{1}{t}\right).$$

Let (see [3])

$$(1.2) \quad \begin{aligned} G_3(x) &= G_3(x; T, U) = \\ &= \bigcup_{T \leq g_{2\nu} \leq T+U} \{t : g_{2\nu}(-x) \leq t \leq g_{2\nu}(x)\}, \quad 0 < x \leq \frac{\pi}{2}, \\ G_4(y) &= G_4(y; T, U) = \\ &= \bigcup_{T \leq g_{2\nu+1} \leq T+U} \{t : g_{2\nu+1}(-y) \leq t \leq g_{2\nu+1}(y)\}, \quad 0 < y \leq \frac{\pi}{2}, \end{aligned}$$

$$(1.3) \quad U = T^{5/12} \ln^3 T,$$

and the collection of sequences  $\{g_\nu(\tau)\}$ ,  $\tau \in [-\pi, \pi]$ ,  $\nu = 1, 2, \dots$  is defined by the equation (see [2], [3], (6))

$$\vartheta_1[g_\nu(\tau)] = \frac{\pi}{2}\nu + \frac{\tau}{2}; \quad g_\nu(0) = g_\nu,$$

where (comp. (1.1))

$$\vartheta_1(t) = \frac{t}{2} \ln \frac{t}{2\pi} - \frac{t}{2} - \frac{\pi}{8}.$$

Let

$$(1.4) \quad G_3(x) = \varphi_1[\hat{G}_3(x)], \quad G_4(y) = \varphi_1[\hat{G}_4(y)],$$

---

*Key words and phrases.* Riemann zeta-function.

where  $y = \varphi_1(T)$ ,  $T \geq T_0[\varphi_1]$  is the Jacob's ladder. The following theorem holds true.

**Theorem.**

$$(1.5) \quad \begin{aligned} & \int_{\mathring{G}_3(x)} Z^2[\varphi_1(t)] \tilde{Z}^2(t) dt = \\ & \frac{x}{\pi} U \ln \frac{T}{2\pi} + \frac{2x}{\pi} \left( c + \frac{\sin x}{x} \right) U + \mathcal{O}(xT^{5/12} \ln^2 T), \\ & \int_{\mathring{G}_4(y)} Z^2[\varphi_1(t)] \tilde{Z}^2(t) dt = \\ & \frac{y}{\pi} U \ln \frac{T}{2\pi} + \frac{2y}{\pi} \left( c - \frac{\sin y}{y} \right) U + \mathcal{O}(yT^{5/12} \ln^2 T), \end{aligned}$$

where

$$(1.6) \quad t - \varphi_1(t) \sim (1 - c)\pi(t), \quad t \rightarrow \infty,$$

and  $c$  is the Euler's constant and  $\pi(t)$  is the prime-counting function.

Let (comp. (1.4))  $T = \varphi_1(\mathring{T})$ ,  $T + U = \varphi_1(\widehat{\mathring{T} + \mathring{U}})$ . Similarly to [14], (1.8) we have

$$(1.7) \quad \rho\{[T, T + U]; [\mathring{T}, \widehat{\mathring{T} + \mathring{U}}]\} \sim (1 - c)\pi(T),$$

where  $\rho$  stands for the distance of the corresponding segments.

*Remark 1.* Some nonlocal interaction of the functions  $Z^2[\varphi_1(t)]$ ,  $\tilde{Z}^2(t)$  is expressed by eq. (1.5). This interaction is connected with two collections of disconnected sets unboundedly receding each from other (see (1.6), (1.7);  $\rho \rightarrow \infty$  as  $T \rightarrow \infty$ ) - like mutually receding galaxies (the Hubble law).

*Remark 2.* The asymptotic formulae (1.5) (comp. (1.3)) cannot be received by methods of Balasubramanian, Heath-Brown and Ivic (comp. [1]).

This paper is a continuation of the series of works [4]-[18].

2. ON BIG ASYMMETRY IN THE BEHAVIOUR OF THE FUNCTION  $Z^2[\varphi_1(t)] \tilde{Z}^2(t)$  RELATIVELY TO THE SETS  $\mathring{G}_3(x)$  AND  $\mathring{G}_4(x)$

We obtain from (1.5)

**Corollary.**

$$(2.1) \quad \begin{aligned} & \int_{\mathring{G}_3(x)} Z^2[\varphi_1(t)] \tilde{Z}^2(t) dt - \int_{\mathring{G}_4(x)} Z^2[\varphi_1(t)] \tilde{Z}^2(t) dt = \\ & \frac{4}{\pi} U \sin x + \mathcal{O}(xT^{5/12} \ln^2 T), \quad x \in (0, \pi/2], \end{aligned}$$

especially, in the case  $x = \pi/2$ , we have

$$(2.2) \quad \int_{\mathring{G}_3(\pi/2)} Z^2[\varphi_1(t)] \tilde{Z}^2(t) dt - \int_{\mathring{G}_4(\pi/2)} Z^2[\varphi_1(t)] \tilde{Z}^2(t) dt \sim \frac{4}{\pi} U,$$

where  $[\mathring{T}, \widehat{\mathring{T} + \mathring{U}}] \subset \mathring{G}_3(\pi/2) \cup \mathring{G}_4(\pi/2)$ .

*Remark 3.* The formulae (2.1), (2.2) represent the big difference of the areas (measures) of the figures which correspond to the functions

$$Z^2[\varphi_1(t)]\tilde{Z}^2(t), t \in \mathring{G}_3(x); \quad Z^2[\varphi_1(t)]\tilde{Z}^2(t), t \in \mathring{G}_4(x).$$

The reason for this big asymmetry given by (2.2) is probably the fact that the zeroes of

$$\zeta\left(\frac{1}{2} + it\right), t \in \mathring{G}_3\left(\frac{\pi}{2}\right) \cup \mathring{G}_4\left(\frac{\pi}{2}\right)$$

lie preferably in the set  $\mathring{G}_4(\pi/2)$ .

### 3. PROOF OF THE THEOREM

3.1. The following lemma holds true (see [6], (2.5); [7], (3.3); [14], (4.1))

**Lemma.** For every integrable function (in the Lebesgue sense)  $f(x)$ ,  $x \in [\varphi_1(T), \varphi_1(T+U)]$  the following is true

$$(3.1) \quad \int_T^{T+U} f[\varphi_1(t)]\tilde{Z}^2(t)dt = \int_{\varphi_1(T)}^{\varphi_1(T+U)} f(x)dx, U \in \left(0, \frac{T}{\ln T}\right],$$

where  $t - \varphi_1(t) \sim (1 - c)\pi(t)$ ,  $c$  is the Euler's constant,  $\pi(t)$  is the prime-counting function and

$$\begin{aligned} \tilde{Z}^2(t) &= \frac{d\varphi_1(t)}{dt}, \quad \varphi_1(t) = \frac{1}{2}\varphi(t), \\ \tilde{Z}^2(t) &= \frac{Z^2(t)}{2\Phi'_\varphi[\varphi(t)]} = \frac{Z^2(t)}{\{1 + \mathcal{O}\left(\frac{\ln \ln t}{\ln t}\right)\} \ln t}, \end{aligned}$$

(see [7], (1.1), (3.1), (3.2)).

*Remark 4.* The formula (3.1) remains true also in the case when the integral on the right-hand side of (3.1) is only relatively convergent integral of the second kind (in the Riemann sense).

In the case (comp. (1.4)  $T = \varphi_1(\overset{\circ}{T})$ ,  $T + U = \varphi_1(\overset{\circ}{T+U})$ ) we obtain from (3.1)

$$(3.2) \quad \int_{\overset{\circ}{T}}^{\overset{\circ}{T+U}} f[\varphi_1(t)]\tilde{Z}^2(t)dt = \int_T^{T+U} f(x)dx.$$

3.2. First of all, we have from (3.2), for example

$$(3.3) \quad \int_{\mathring{g}_{2\nu}(-x)}^{\mathring{g}_{2\nu}(x)} f[\varphi_1(t)]\tilde{Z}^2(t)dt = \int_{g_{2\nu}(-x)}^{g_{2\nu}(x)} f(t)dt$$

(see (1.4)). Next, in the case  $f(t) = Z^2(t)$ , we have the following  $\tilde{Z}^2$ -transformation

$$(3.4) \quad \begin{aligned} \int_{\mathring{G}_3(x)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)dt &= \int_{G_3(x)} Z^2(t)dt, \\ \int_{\mathring{G}_4(y)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)dt &= \int_{G_4(y)} Z^2(t)dt. \end{aligned}$$

Let us remind that we have proved in the paper [3] the following mean-value formulae

$$(3.5) \quad \int_{G_3(x)} Z^2(t)dt = \frac{x}{\pi}U \ln \frac{T}{2\pi} + \frac{2x}{\pi} \left( c + \frac{\sin x}{x} \right) U + \mathcal{O}(xT^{5/12} \ln^2 T),$$

$$\int_{G_4(y)} Z^2(t)dt = \frac{y}{\pi}U \ln \frac{T}{2\pi} + \frac{2y}{\pi} \left( c - \frac{\sin y}{y} \right) U + \mathcal{O}(yT^{5/12} \ln^2 T).$$

Now, the formula (1.5) follows from (3.4), (3.5).

*Remark 5.* The formulae (3.5) are the consequences of their discrete form

$$\sum_{T \leq g_{2\nu} \leq T+U} Z^2[g_{2\nu}(\tau)] =$$

$$\frac{1}{2\pi}U \ln^2 \frac{T}{2\pi} + \frac{1}{\pi}(c + \cos \tau)U \ln \frac{T}{2\pi} + \mathcal{O}(T^{5/12} \ln^3 T),$$

$$\sum_{T \leq g_{2\nu+1} \leq T+U} Z^2[g_{2\nu+1}(\tau)] =$$

$$\frac{1}{2\pi}U \ln^2 \frac{T}{2\pi} + \frac{1}{\pi}(c - \cos \tau)U \ln \frac{T}{2\pi} + \mathcal{O}(T^{5/12} \ln^3 T)$$

(see [3], (10)).

I would like to thank Michal Demetrian for helping me with the electronic version of this work.

REFERENCES

[1] A. Ivic, ‘The Riemann zeta-function’, A Willey-Interscience Pub., New York, 1985.  
 [2] J. Moser, ‘An improvement of the theorem of Hardy-Littlewood on density of zeroes of the function  $\zeta\left(\frac{1}{2} + it\right)$ ’, Acta Arithmetica, 43 (1983), 21-47 (in russian).  
 [3] J. Moser, ‘New mean-value theorems for the function  $|\zeta\left(\frac{1}{2} + it\right)|^2$ ’, Acta Math. Univ. Comen., 46-47 (1985), pp. 21-40, (in russian).  
 [4] J. Moser, ‘Jacob’s ladders and the almost exact asymptotic representation of the Hardy-Littlewood integral’, (2008), arXiv:0901.3973.  
 [5] J. Moser, ‘Jacob’s ladders and the tangent law for short parts of the Hardy-Littlewood integral’, (2009), arXiv:0906.0659.  
 [6] J. Moser, ‘Jacob’s ladders and the multiplicative asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral’, (2009), arXiv:0907.0301.  
 [7] J. Moser, ‘Jacob’s ladders and the quantization of the Hardy-Littlewood integral’, (2009), arXiv:0909.3928.  
 [8] J. Moser, ‘Jacob’s ladders and the first asymptotic formula for the expression of the sixth order  $|\zeta(1/2 + i\varphi(t)/2)|^4|\zeta(1/2 + it)|^2$ ’, (2009), arXiv:0911.1246.  
 [9] J. Moser, ‘Jacob’s ladders and the first asymptotic formula for the expression of the fifth order  $Z[\varphi(t)/2 + \rho_1]Z[\varphi(t)/2 + \rho_2]Z[\varphi(t)/2 + \rho_3]\hat{Z}^2(t)$  for the collection of disconnected sets’, (2009), arXiv:0912.0130.  
 [10] J. Moser, ‘Jacob’s ladders, the iterations of Jacob’s ladder  $\varphi_1^k(t)$  and asymptotic formulae for the integrals of the products  $Z^2[\varphi_1^n(t)]Z^2[\varphi_1^{n-1}(t)] \dots Z^2[\varphi_1^0(t)]$  for arbitrary fixed  $n \in \mathbb{N}$ ’, (2010), arXiv:1001.1632.  
 [11] J. Moser, ‘Jacob’s ladders and the asymptotic formula for the integral of the eight order expression  $|\zeta(1/2 + i\varphi_2(t))|^4|\zeta(1/2 + it)|^4$ ’, (2010), arXiv:1001.2114.  
 [12] J. Moser, ‘Jacob’s ladders and the asymptotically approximate solutions of a nonlinear diophantine equation’, (2010), arXiv: 1001.3019.  
 [13] J. Moser, ‘Jacob’s ladders and the asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral of the function  $|\zeta(1/2 + it)|^4$ ’, (2010), arXiv:1001.4007.

- [14] J. Moser, 'Jacob's ladders and the nonlocal interaction of the function  $|\zeta(1/2 + it)|$  with  $\arg \zeta(1/2 + it)$  on the distance  $\sim (1 - c)\pi(t)$ ', (2010), arXiv: 1004.0169.
- [15] J. Moser, 'Jacob's ladders and the  $\tilde{Z}^2$  - transformation of polynomials in  $\ln \varphi_1(t)$ ', (2010), arXiv: 1005.2052.
- [16] J. Moser, 'Jacob's ladders and the oscillations of the function  $|\zeta(\frac{1}{2} + it)|^2$  around the main part of its mean-value; law of the almost exact equality of the corresponding areas', (2010), arXiv: 1006.4316
- [17] J. Moser, 'Jacob's ladders and the nonlocal interaction of the function  $Z(t)$  with the function  $\tilde{Z}^2(t)$  on the distance  $\sim (1 - c)\pi(t)$  for a collection of disconnected sets', (2010), arXiv: 1006.5158
- [18] J. Moser, 'Jacob's ladders and the  $\tilde{Z}^2$ -transformation of the orthogonal system of trigonometric functions', (2010), arXiv: 1007.0108.

DEPARTMENT OF MATHEMATICAL ANALYSIS AND NUMERICAL MATHEMATICS, COMENIUS UNIVERSITY, MLYNSKA DOLINA M105, 842 48 BRATISLAVA, SLOVAKIA

*E-mail address:* `jan.moser@fmph.uniba.sk`