

HILBERT SERIES AND HILBERT DEPTH OF SQUAREFREE VERONESE IDEALS

Maorong Ge, Jiayuan Lin and Yulan Wang

Abstract In this paper, we obtain explicit formulas for the Hilbert series and Hilbert depth of squarefree Veronese ideals in a standard graded polynomial ring.

1 Introduction

In recent years, *depth*, *Stanley depth*, *Hilbert depth* and their relations of graded modules over polynomial ring $R = K[x_1, \dots, x_n]$ have been extensively studied (e.g. [1]-[12]). One of the motivations behind these researches is the famous Stanley conjecture. It says that $\text{sdepth } M \geq \text{depth } M$ for all finitely generated graded R -module M . Although Stanley conjecture has been confirmed in some special cases, it is still widely open. Analogous to *depth* and *Stanley depth*, W. Bruns et al [3] introduced *Hilbert depth*. For the reader's convenience, let us recall the definition of *Hilbert depth* from [3].

Definition 1.1. Let M be a finitely generated graded S -module. A *Hilbert decomposition* of M is a finite family

$$\mathcal{H} = (S_i, s_i)$$

such that $s_i \in \mathbb{Z}^m$ (where $m = 1$ in the standard graded case and $m = n$ in the multigraded case), S_i is a graded K -algebra retract of R for each i , and

$$M \cong \bigoplus_i S_i(-s_i)$$

as a graded K -vector space.

The number $\text{Hdepth } \mathcal{H} = \min_i \text{depth } S_i(-s_i)$ is called the *Hilbert depth* of \mathcal{H} . The *Hilbert depth* of M is defined to be

$$\text{Hdepth } M = \max\{\text{Hdepth } \mathcal{H} : \mathcal{H} \text{ is a Hilbert decomposition of } M\}.$$

Note that a Stanley decomposition breaks M into a direct sum of submodules over suitable subalgebras, while a Hilbert decomposition only requires an isomorphism to the direct sum of modules over such subalgebras. The latter one depends only on the Hilbert series of M . Since Stanley decompositions are Hilbert decompositions, Hilbert depth provides an upper bound for the Stanley depth.

For a finitely generated standard graded R -module $M = \bigoplus_{k \in \mathbb{Z}} M_k$, the *Hilbert series* of M is given by the Laurent series $H_M(T) = \sum_{k \in \mathbb{Z}} H(M, k)T^k$, where $H(M, k) = \dim_K M_k$ is the *Hilbert function* of M . We say a Laurent series $\sum_{k \in \mathbb{Z}} a_k T^k$ is *positive* if $a_k \geq 0$ for all k . It is easy to see that any Hilbert series is positive. In [12], J. Uliczka proved that the Hilbert depth of M is equal to $\max\{r : (1 - T)^r H_M(T) \text{ positive}\}$. Using this, W. Bruns et

al [4] computed the Hilbert depth of the powers of the irrelevant maximal ideal in R . In this paper we deal with Hilbert series and Hilbert depth of squarefree Veronese ideals $I_{n,d}$. We prove the following theorem.

Theorem 1.2. *Let $R = K[x_1, \dots, x_n]$ be the standard graded polynomial ring in n variables over a field K . Let $I_{n,d}$ be the squarefree Veronese ideal generated by all squarefree monomials of degree d in R . Then*

$$(1) \text{ The Hilbert series of } I_{n,d} \text{ is } H_{I_{n,d}}(T) = \sum_{i=d-1}^{n-1} \binom{i}{d-1} T^d (1-T)^{-n+i-d+1};$$

$$(2) \text{ The Hilbert depth of } I_{n,d} \text{ is equal to } d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor = d + \lfloor \frac{n-d}{d+1} \rfloor = d - 1 + \left\lceil \frac{n-(d-1)}{d+1} \right\rceil.$$

Remark 1.3. The (multigraded) Stanley depth of $I_{n,d}$ has been conjectured (see [6] and [10]) to be equal to $d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor$. M. Keller et al [10] and M. Ge et al [8] partially confirmed this conjecture but it is still open. We wonder whether a suitable modification of a Hilbert decomposition \mathcal{H} with $\text{Hdepth } \mathcal{H} = d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor$ will bring us a solution to that conjecture.

To prove Theorem 1.2(2), it is sufficient to show that $\max\{r : (1-T)^r H_{I_{n,d}}(T) \text{ positive}\} = d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor$. In order to do so, we need a nice formula for $H_{I_{n,d}}(T)$. Surprisingly, we cannot find such a formula in literature. In section 2 we use generating functions to obtain an explicit formula for $H_{I_{n,d}}(T)$. We refer the reader to [13] for the basic theory of generating functions. In section 3, we prove that $(1-T)^r H_{I_{n,d}}(T)$ is *positive* if and only if $r \leq d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor$. This implies Theorem 1.2(2).

Notations: In this paper, we use the generalized binomial coefficient $\binom{m}{r} = \frac{m(m-1)\dots(m-r+1)}{r!}$ when $r \geq 0$. The binomial coefficient $\binom{m}{r} = 0$ when $r < 0$.

2 Hilbert series of squarefree Veronese ideals

Let $H_{I_{n,d}}(T) = \sum_{k=d}^{\infty} a_{n,d,k} T^k$ be the Hilbert series of the squarefree Veronese ideal $I_{n,d}$, where $a_{n,d,k} = \dim_K(I_{n,d})_k$. Then $a_{n,d,k}$ satisfying the following properties.

Proposition 2.1. (1) $a_{n,1,k} = \binom{n+k-1}{k}$;

(2) $a_{n,d,k} = a_{n-1,d,k} + \sum_{s=1}^{k-d+1} a_{n-1,d-1,k-s}$ for $\min\{n, k\} \geq d \geq 2$; and

(3) $a_{n,d,k} = a_{n-1,d,k} + a_{n-1,d-1,k-1} + a_{n,d,k-1} - a_{n-1,d,k-1}$ for $\min\{n, k\} \geq d \geq 2$.

Proof. (1) is easy because $I_{n,1}$ is just the irrelevant maximal ideal \mathfrak{m} in R and its Hilbert series is $H_{I_{n,d}}(T) = \sum_{k=d}^{\infty} \binom{n+k-1}{k} T^k$. To prove (2) and (3), let

$$P_{n,d,k} = \{u : u \in I_{n,d} \text{ monomial, } \deg u = k \text{ and } x_{i_1} x_{i_2} \dots x_{i_d} | u \text{ for some } \{i_1, \dots, i_d\} \subseteq [n]\}$$

Then $a_{n,d,k} = \dim_K(I_{n,d})_k = |P_{n,d,k}|$.

Let $x_1^s || u$ be the highest power of x_1 in $u \in P_{n,d,k}$. Then s can take values $0, \dots, k-d+1$. We can divide $P_{n,d,k}$ into $k-d+2$ classes $P_{n,d,k,s}$ according to these s values. It is easy to see that $P_{n,d,k,s}$ are pairwise disjoint, $|P_{n,d,k,0}| = a_{n-1,d,k}$ and $|P_{n,d,k,s}| = a_{n-1,d-1,k-s}$ for $s > 0$. So (2) follows easily.

(3) is an immediate consequence of (2) because $a_{n,d,k-1} = a_{n-1,d,k-1} + \sum_{s=1}^{k-d} a_{n-1,d-1,k-s-1}$. \square

Let $F(x, y, z) = \sum_{\min\{n,k\} \geq d \geq 1} a_{n,d,k} x^n y^d z^k$ be the generating function of $a_{n,d,k}$. Then the following is true.

Proposition 2.2. $F(x, y, z) = \frac{xyz(1-z)}{(1-x-xyz-z+xz)(1-z-x)}$

Proof. By the definition of $F(x, y, z)$ and Proposition 2.1(3), we have that $F(x, y, z) = \sum_{\min\{n,k\} \geq d \geq 1} a_{n,d,k} x^n y^d z^k = \sum_{\min\{n,k\} \geq 1} a_{n,1,k} x^n y z^k + \sum_{\min\{n,k\} \geq d \geq 2} a_{n,d,k} x^n y^d z^k$
 $= y \sum_{\min\{n,k\} \geq 1} \binom{n+k-1}{k} x^n z^k + \sum_{\min\{n,k\} \geq d \geq 2} (a_{n-1,d,k} + a_{n-1,d-1,k-1} + a_{n,d,k-1} - a_{n-1,d,k-1}) x^n y^d z^k$
 $= y \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \binom{n+k-1}{k} z^k x^n + \sum_{\min\{n,k\} \geq d \geq 2} a_{n-1,d,k} x^n y^d z^k + \sum_{\min\{n,k\} \geq d \geq 2} a_{n-1,d-1,k-1} x^n y^d z^k +$
 $\sum_{\min\{n,k\} \geq d \geq 2} a_{n,d,k-1} x^n y^d z^k - \sum_{\min\{n,k\} \geq d \geq 2} a_{n-1,d,k-1} x^n y^d z^k$
 $= \frac{xyz}{(1-z-x)(1-x)} + x[F(x, y, z) - \frac{xyz}{(1-z-x)(1-x)}] + xyzF(x, y, z) + z[F(x, y, z) - \frac{xyz}{(1-z-x)(1-x)}] -$
 $xz[F(x, y, z) - \frac{xyz}{(1-z-x)(1-x)}].$

Solving for $F(x, y, z)$ from the previous equation gives

$$F(x, y, z) = \frac{xyz(1-z)}{(1-x-xyz-z+xz)(1-z-x)}$$

\square

Now we deduce a nice formula for $H_{I_{n,d}}(T)$ from Proposition 2.2.

Theorem 2.3. *The Hilbert series of the squarefree Veronese ideal $I_{n,d}$ is given by*

$$H_{I_{n,d}}(T) = \sum_{i=d-1}^{n-1} \binom{i}{d-1} T^d (1-T)^{-n+i-d+1}$$

Proof. By Proposition 2.2, $F(x, y, z) = \sum_{\min\{n,k\} \geq d \geq 1} a_{n,d,k} x^n y^d z^k = \frac{xyz(1-z)}{(1-x-xyz-z+xz)(1-z-x)}$.

$$\text{Now } \frac{xyz(1-z)}{(1-x-xyz-z+xz)(1-z-x)} = \frac{xyz}{1-z} \cdot \frac{1}{1-x-\frac{xyz}{1-z}} \cdot \frac{1}{1-\frac{x}{1-z}} = \frac{xyz}{1-z} \sum_{i=0}^{\infty} x^i \left(1 + \frac{yz}{1-z}\right)^i \sum_{j=0}^{\infty} \frac{x^j}{(1-z)^j}$$

The coefficient of x^n in $F(x, y, z)$ is $\frac{yz}{1-z} \sum_{i+j=n-1} \left(1 + \frac{yz}{1-z}\right)^i \frac{1}{(1-z)^j} = yz \sum_{i=0}^{n-1} \left(1 + \frac{yz}{1-z}\right)^i$

$\frac{1}{(1-z)^{n-i}}$. The coefficient of y^d in $yz \sum_{i=0}^{n-1} \left(1 + \frac{yz}{1-z}\right)^i \frac{1}{(1-z)^{n-i}}$ is $\sum_{i=0}^{n-1} \binom{i}{d-1} z \left(\frac{z}{1-z}\right)^{d-1}$

$$\frac{1}{(1-z)^{n-i}} = \sum_{i=d-1}^{n-1} \binom{i}{d-1} z^d (1-z)^{-n+i-d+1}. \text{ Replacing } z \text{ with } T \text{ gives the formula } H_{I_{n,d}}(T) = \sum_{i=d-1}^{n-1} \binom{i}{d-1} T^d (1-T)^{-n+i-d+1}. \quad \square$$

As a corollary of Theorem 2.3, we have

$$\text{Corollary 2.4. } a_{n,d,k} = \sum_{i=d-1}^{n-1} \binom{i}{d-1} \binom{n-i+k-2}{k-d}.$$

Proof. It follows directly from the expansion of $H_{I_{n,d}}(T) = \sum_{i=d-1}^{n-1} \binom{i}{d-1} T^d (1-T)^{-n+i-d+1}$. □

In the next section, we will prove that the Hilbert depth of $I_{n,d}$ is equal to $d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor$.

3 Hilbert depth of squarefree Veronese ideals

By J. Uliczka [12], the Hilbert depth of $I_{n,d}$ is equal to $\max\{r : (1-T)^r H_{I_{n,d}}(T) \text{ positive}\}$. The coefficient of T^{d+1} in $(1-T)^r H_{I_{n,d}}(T)$ is $a_{n,d,d+1} - r a_{n,d,d}$. By Corollary 2.4, $a_{n,d,d} = \sum_{i=d-1}^{n-1} \binom{i}{d-1} = \binom{n}{d}$ and $a_{n,d,d+1} = \sum_{i=d-1}^{n-1} \binom{i}{d-1} (n-i+d-1) = n \sum_{i=d-1}^{n-1} \binom{i}{d-1} - \sum_{i=d-1}^{n-1} \binom{i}{d-1} (i-d+1) = n \binom{n}{d} - d \sum_{i=d-1}^{n-1} \binom{i}{d} = n \binom{n}{d} - d \binom{n}{d+1} = d \binom{n}{d} + (n-d) \binom{n}{d} - d \binom{n}{d+1} = d \binom{n}{d} + (d+1) \binom{n}{d+1} - d \binom{n}{d+1} = d \binom{n}{d} + \binom{n}{d+1}$. So $a_{n,d,d+1} - r a_{n,d,d} \geq 0$ implies that $r \leq \frac{a_{n,d,d+1}}{a_{n,d,d}} = \frac{d \binom{n}{d} + \binom{n}{d+1}}{\binom{n}{d}} = d + \frac{\binom{n}{d+1}}{\binom{n}{d}}$. Therefore, $r \leq d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor$ is a necessary condition for the positivity of $(1-T)^r H_{I_{n,d}}(T)$. We will show that it is also a sufficient condition.

Denote $(1-T)^r H_{I_{n,d}}(T) = \sum_{k=d}^{\infty} b_{n,d,k,r} T^k$. The following lemma is true.

$$\text{Lemma 3.1. } b_{n,d,k,r} = \sum_{i=d-1}^{n-1} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d}.$$

Proof. It follows directly from the expansion of $(1-T)^r H_{I_{n,d}}(T) = (1-T)^r \sum_{i=d-1}^{n-1} \binom{i}{d-1} T^d (1-T)^{-n+i-d+1} = \sum_{i=d-1}^{n-1} \binom{i}{d-1} T^d (1-T)^{-n+i+r-d+1}$. □

Lemma 3.2. For all positive integers $\min\{n, k\} \geq d$, we have

- (1) $b_{n-r+p-1,p,k,p-1} = \binom{n+k-r-1}{k}$ for any integer $1 \leq p \leq d$;
- (2) $b_{n,d,k,r} = (-1)^{k-d} \sum_{t=1}^{r-k+1} \binom{n-(d-1)-t+d-1}{d-1} \binom{r-(d-1)-t}{k-d} + \binom{n+k-r-1}{k}$.

Proof. We use induction on p to prove (1).

When $p = 1$, we have that $b_{n-r+p-1,p,k,p-1} = b_{n-r,1,k,0} = a_{n-r,1,k}$, which is equal to $\sum_{i=0}^{n-r-1} \binom{n-i+k-2}{k-1} = \binom{n+k-r-1}{k}$ by Corollary 2.4. So (1) holds true in this case.

Suppose that $b_{n-r+q-1,q,k,q-1} = \binom{n+k-r-1}{k}$ for any $1 \leq q \leq p$. If $p = d$, we are done. Otherwise $p \leq d-1$. By Lemma 3.1, we have that

$$\begin{aligned}
b_{n-r+p,p+1,k,p} &= \sum_{i=p}^{n-r+p-1} \binom{i}{p} \binom{n-i+k-r-2}{k-p-1} = \sum_{i=p}^{n-r+p-1} \binom{i}{p} \left[\binom{n-i+k-r-1}{k-p} - \binom{n-i+k-r-2}{k-p} \right] \\
&= \sum_{i=p}^{n-r+p-1} \binom{i}{p} \binom{n-i+k-r-1}{k-p} - \sum_{i=p}^{n-r+p-1} \binom{i}{p} \binom{n-i+k-r-2}{k-p} \\
&= \sum_{i=p}^{n-r+p-1} \binom{i}{p} \binom{n-i+k-r-1}{k-p} - \sum_{i=p+1}^{n-r+p-1} \binom{i-1}{p} \binom{n-i+k-r-1}{k-p} \\
&= \binom{n-p+k-r-1}{k-p} + \sum_{i=p+1}^{n-r+p-1} \binom{i-1}{p-1} \binom{n-i+k-r-1}{k-p} \\
&= \binom{n-p+k-r-1}{k-p} + \sum_{i=p}^{n-r+p-2} \binom{i}{p-1} \binom{n-i+k-r-2}{k-p} \\
&= \sum_{i=p-1}^{n-r+p-2} \binom{i}{p-1} \binom{n-i+k-r-2}{k-p} = b_{n-r+p-1,p,k,p-1}, \text{ which is equal to } \binom{n+k-r-1}{k} \text{ by the induc-} \\
\end{aligned}$$

tive assumption.

$$\begin{aligned}
\text{To prove (2), note that } b_{n,d,k,r} &= \sum_{i=d-1}^{n-1} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d} \\
&= \sum_{i=n+k-r-1}^{n-1} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d} + \sum_{i=d-1}^{n+k-r-2} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d} \\
&= (-1)^{k-d} \sum_{i=n+k-r-1}^{n-1} \binom{i}{d-1} \binom{r+i+1-n-d}{k-d} + \sum_{i=d-1}^{n+k-r-2} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d} \\
&\stackrel{i=n-t+d-1}{=} (-1)^{k-d} \sum_{t=d}^{d+r-k} \binom{n-t+d-1}{d-1} \binom{r-t}{k-d} + \sum_{i=d-1}^{n+k-r-2} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d} \\
&= (-1)^{k-d} \sum_{t=1}^{r-k+1} \binom{n-(d-1)-t+d-1}{d-1} \binom{r-(d-1)-t}{k-d} + \sum_{i=d-1}^{n+k-r-2} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d} \\
&= (-1)^{k-d} \sum_{t=1}^{r-k+1} \binom{n-(d-1)-t+d-1}{d-1} \binom{r-(d-1)-t}{k-d} + \sum_{i=d-1}^{n+d-r-2} \binom{i}{d-1} \binom{n-i+k-r-2}{k-d} \\
&= (-1)^{k-d} \sum_{t=1}^{r-k+1} \binom{n-(d-1)-t+d-1}{d-1} \binom{r-(d-1)-t}{k-d} + \sum_{i=d-1}^{n+d-r-2} \binom{i}{d-1} \binom{(n+d-r-1)-i+k-(d-1)-2}{k-d} \\
&= (-1)^{k-d} \sum_{t=1}^{r-k+1} \binom{n-(d-1)-t+d-1}{d-1} \binom{r-(d-1)-t}{k-d} + b_{n+d-r-1,d,k,d-1}. \text{ By (1), } b_{n+d-r-1,d,k,d-1} = \\
&\binom{n+k-r-1}{k}. \text{ So (2) follows.} \quad \square
\end{aligned}$$

To prove the positivity of $(1-T)^r H_{I_{n,d}}(T)$ for $r \leq d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor = d-1 + \left\lceil \frac{n-(d-1)}{d+1} \right\rceil$, it is sufficient to show the following.

Proposition 3.3. *Let n and d be positive integers with $n \geq d$, and let $r = d - 1 + \left\lceil \frac{n - (d - 1)}{d + 1} \right\rceil$. Then for all $k = d + 1, \dots, r$ we have*

$$\binom{n + k - r - 1}{k} \geq \sum_{t=1}^{r-k+1} \binom{n - (d - 1) - t + d - 1}{d - 1} \binom{r - (d - 1) - t}{k - d} \quad (3.1)$$

Remark 3.4. The inequality in Proposition is trivially true if $r + 1 \leq k$.

Proof. Replacing r , n and s in Proposition 3.1 in [4] with $r - (d - 1)$, $n - (d - 1)$ and d respectively, we have

$$\binom{[n - (d - 1)] + k - [r - (d - 1)] - 1}{k} \geq \sum_{t=1}^{r - (d - 1)} \binom{n - (d - 1) - t + d - 1}{d - 1} \binom{r - (d - 1) - t}{k - d}$$

Because $\binom{r - (d - 1) - t}{k - d} = 0$ for $r - k + 1 < t \leq r - (d - 1)$, after simplification, the above inequality becomes

$$\binom{n + k - r - 1}{k} \geq \sum_{t=1}^{r-k+1} \binom{n - (d - 1) - t + d - 1}{d - 1} \binom{r - (d - 1) - t}{k - d}.$$

This completes the proof of Proposition 3.3. \square

By Lemma 3.2(2), Proposition 3.3, Remark 3.4 and the necessary condition $r \leq d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor$ for the positivity of $(1 - T)^r H_{I_{n,d}}(T)$, we conclude that the Hilbert depth of $I_{n,d}$ is equal to $d + \lfloor \binom{n}{d+1} / \binom{n}{d} \rfloor = d + \lfloor \frac{n-d}{d+1} \rfloor = d - 1 + \left\lceil \frac{n - (d - 1)}{d + 1} \right\rceil$. This completes the proof of Theorem 1.2(2).

References

- [1] I. Anwar and D. Popescu, *Stanley Conjecture in small embedding dimension*, J. Algebra **318**, 1027-1031, 2007.
- [2] C. Biró, D. Howard, M. Keller, W. Trotter and S. Young, *Interval partition and Stanley depth*, J. Combin. Theory Ser. A. **117** (4), 475-482, 2010.
- [3] W. Bruns, C. Krattenthaler and J. Uliczka, *Stanley decompositions and Hilbert depth in the Koszul complex*, arXiv:0909.0686.
- [4] W. Bruns, C. Krattenthaler and J. Uliczka, *Hilbert depth of powers of the maximal ideal*, arXiv:1002.1400.
- [5] M. Cimpoeaş, *Stanley depth of complete intersection monomial ideals*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **51** (99), no. 3, 205-211, 2008.

- [6] M. Cimpoeaş, *Stanley depth of square free Veronese ideals*, arXiv:0907.1232.
- [7] G. Fløystad and J. Herzog, *Gröbner bases of syzygies and Stanley depth*, arXiv:1003.4495.
- [8] M. Ge, J. Lin, and Y. Shen, *On a conjecture of Stanley depth of squarefree Veronese ideals*, arXiv:0911.5458.
- [9] J. Herzog, M. Vladioiu and X. Zheng, *How to compute the Stanley depth of a monomial ideal*, J. Algebra **322** (9), 3151-3169, 2009.
- [10] M. Keller, Y. Shen, N. Streib and S. Young, *On the Stanley depth of squarefree Veronese ideals*, to appear in J. Algebr. Comb.
- [11] D. Popescu, *Stanley depth of multigraded modules*, J. Algebra **321** (10), 2782-2797, 2009.
- [12] J. Uliczka, *Remarks on Hilbert series of graded modules over polynomial rings*, Manuscr. Math. **132**, no. 1-2, 159-168, 2010.
- [13] H. Wilf, *generatingfunctionology*, Third edition. A K Peters, Ltd., Wellesley, MA, 2006.

DEPARTMENT OF MATHEMATICS, ANHUI UNIVERSITY, HEFEI, ANHUI, 230039, CHINA
E-mail address: ge1968@126.com

DEPARTMENT OF MATHEMATICS, SUNY CANTON, 34 CORNELL DRIVE, CANTON,
 NY 13617, USA

E-mail address: linj@canton.edu

DEPARTMENT OF MATHEMATICS, SUNY CANTON, 34 CORNELL DRIVE, CANTON,
 NY 13617, USA

DEPARTMENT OF MATHEMATICS, ANHUI ECONOMIC MANAGEMENT INSTITUTE, HEFEI,
 ANHUI, 230059, CHINA

E-mail address: wangy@canton.edu