

A CHARACTERIZATION OF INNER PRODUCT SPACES

MOHAMMAD SAL MOSLEHIAN¹ AND JOHN M. RASSIAS²

ABSTRACT. In this paper we present a new criterion on characterization of real inner product spaces. We conclude that a real normed space $(X, \|\cdot\|)$ is an inner product space if

$$\sum_{\varepsilon_i \in \{-1,1\}} \left\| x_1 + \sum_{i=2}^k \varepsilon_i x_i \right\|^2 = \sum_{\varepsilon_i \in \{-1,1\}} \left(\|x_1\| + \sum_{i=2}^k \varepsilon_i \|x_i\| \right)^2,$$

for some positive integer $k \geq 2$ and all $x_1, \dots, x_k \in X$. Conversely, if $(X, \|\cdot\|)$ is an inner product space, then the equality above holds for all $k \geq 2$ and all $x_1, \dots, x_k \in X$

1. INTRODUCTION

There are a lot of significant natural geometric properties, which fail in general normed spaces as non Euclidean spaces. Some of these interesting properties hold just when the space is an inner product space. This is the most important motivation for study of characterizations of inner product spaces.

The first norm characterization of inner product spaces was given by Fréchet [9] in 1935. He proved that a normed space $(X, \|\cdot\|)$ is an inner product space if and only if

$$\|x + y + z\|^2 + \|x\|^2 + \|y\|^2 + \|z\|^2 - \|x + y\|^2 - \|y + z\|^2 - \|x + z\|^2 = 0$$

for all $x, y, z \in X$. In 1936 Jordan and von Neumann [10] showed that a normed space X is an inner product space if and only if the parallelogram law $\|x - y\|^2 + \|x + y\|^2 = 2\|x\|^2 + 2\|y\|^2$ holds for all $x, y \in X$. Later, Day [6] showed that a normed linear space X is an inner product space if one requires only that the parallelogram equality holds for x and y on the unit sphere. In other words, he showed that the parallelogram equality may be replaced by the condition $R = 4$ ($\|x\| = 1, \|y\| = 1$), where $R = \|x - y\|^2 + \|x + y\|^2$. There are several

2010 *Mathematics Subject Classification.* Primary 46C15; Secondary 46B20, 46C05.

Key words and phrases. inner product space; Euler–Lagrange identity; Day’s condition; normed space; characterization; operator.

characterizations of inner product spaces introduced by many mathematicians some of which are [1–16].

In this paper we present a new criterion on characterization of inner product spaces and give an operator version of it. The notion of inner product space plays an essential role in quantum mechanics, since every physical system is associated with a Hilbert space and self-adjoint operators associated to a system represent physical quantities; see [17].

2. MAIN RESULT

Theorem 2.1. *Let $(X, \|\cdot\|)$ be a real normed space, n be a positive real number and $k \geq 2$ be a positive integer. If*

$$R_{k,n} = \sum_{\varepsilon_i \in \{-1,1\}} \left\| x_1 + \sum_{i=2}^k \varepsilon_i x_i \right\|^n$$

and

$$A_{k,n} = \sum_{\varepsilon_i \in \{-1,1\}} \left(\|x_1\| + \sum_{i=2}^k \varepsilon_i \|x_i\| \right)^n,$$

then a necessary and sufficient condition for that the norm $\|\cdot\|$ over X is induced by an inner product is that

(I) $R_{k,n} \leq A_{k,n}$ if $n \geq 2$

and

(II) $R_{k,n} \geq A_{k,n}$ if $0 < n \leq 2$

for any $x_1, \dots, x_k \in X$.

Proof. Necessity.

Assume that the norm $\|\cdot\|$ on X is induced by an inner product $\langle \cdot, \cdot \rangle$. Hence

$\|x\|^2 = \langle x, x \rangle$ ($x \in X$). We have

$$\begin{aligned}
R_{k,n} &= \sum_{\varepsilon_i \in \{-1,1\}} \left\| x_1 + \sum_{i=2}^k \varepsilon_i x_i \right\|^n \\
&= \frac{1}{2} \sum_{\varepsilon_i \in \{-1,1\}} \left(\left\| \sum_{i=1}^k \varepsilon_i x_i \right\|^2 \right)^{n/2} \\
&= \frac{1}{2} \sum_{\varepsilon_i, \varepsilon_j \in \{-1,1\}} \left(\sum_{i=1}^k \|x_i\|^2 + 2 \sum_{1 \leq i < j \leq k} \varepsilon_i \varepsilon_j \langle x_i, x_j \rangle \right)^{n/2} \\
&= \frac{1}{2} \sum_{\varepsilon_i, \varepsilon_j \in \{-1,1\}} \left(a + \sum_{1 \leq i < j \leq k} \varepsilon_i \varepsilon_j a_{i,j} \cos(p_{i,j}) \right)^{n/2} \\
&= R_{k,n}(P),
\end{aligned}$$

where $a := \sum_{i=1}^k \|x_i\|^2$, $a_{i,j} := 2\|x_i\| \|x_j\|$ and $p_{i,j}$'s are defined in such a way that $\langle x_i, x_j \rangle = \|x_i\| \|x_j\| \cos(p_{i,j})$. Let P denote the $\frac{k^2-k}{2}$ -tuple consisting of $p_{i,j}$ ($1 \leq i < j \leq k$) by going row-by-row throughout the matrix

$$\begin{bmatrix}
\star & \langle x_1, x_2 \rangle & \langle x_1, x_3 \rangle & \cdots & \langle x_1, x_k \rangle \\
\star & \star & \langle x_2, x_3 \rangle & \cdots & \langle x_2, x_k \rangle \\
\vdots & \vdots & \vdots & \vdots & \vdots \\
\star & \star & \cdots & \star & \langle x_{k-1}, x_k \rangle \\
\star & \star & \cdots & \star & \star
\end{bmatrix}.$$

For each fixed $1 \leq t < s \leq k$, we have

$$\begin{aligned}
\frac{\partial R_{k,n}(P)}{\partial p_{t,s}} &= \frac{1}{2} \sum_{\varepsilon_i, \varepsilon_j \in \{-1,1\}} \left[-\frac{n}{2} \left(a + \sum_{1 \leq i < j \leq k} \varepsilon_i \varepsilon_j a_{i,j} \cos(p_{i,j}) \right)^{\frac{n-2}{2}} \varepsilon_t \varepsilon_s a_{t,s} \sin(p_{t,s}) \right] \\
&= \frac{n}{4} \varphi(P) a_{t,s} \sin(p_{t,s}),
\end{aligned}$$

in which

$$\varphi(P) := \sum_{\varepsilon_i, \varepsilon_j \in \{-1,1\}} -\varepsilon_t \varepsilon_s \left(a + \sum_{1 \leq i < j \leq k} \varepsilon_i \varepsilon_j a_{i,j} \cos(p_{i,j}) \right)^{\frac{n-2}{2}}.$$

The solution of the system of equations $\frac{\partial R_{k,n}(P)}{\partial p_{t,s}}$ where t, s run throughout $1 \leq t < s \leq k$ is $P_0 = (K_1 \pi, \dots, K_{\frac{k^2-k}{2}} \pi)$, where $K_1, \dots, K_{\frac{k^2-k}{2}} \in \{0, \pm 1, \pm 2, \dots\}$.

We use the second partial test to show that P_0 is an extremum point of $R_{k,n}(P)$. For $(u, v) \neq (t, s)$, we have

$$\begin{aligned} \frac{\partial^2 R_{k,n}(P)}{\partial p_{u,v} \partial p_{t,s}} &= \frac{\partial}{\partial p_{u,v}} \left(\frac{n}{4} \varphi(P) a_{t,s} \sin(p_{t,s}) \right) \\ &= \frac{n}{4} a_{t,s} \sin(p_{t,s}) \frac{\partial}{\partial p_{u,v}} \varphi(P) \end{aligned} \quad (2.1)$$

and

$$\begin{aligned} \frac{\partial^2 R_{k,n}(P)}{\partial p_{t,s}^2} &= \frac{\partial}{\partial p_{t,s}} \left(\frac{n}{4} \varphi(P) a_{t,s} \sin(p_{t,s}) \right) \\ &= \frac{n}{4} a_{t,s} \sin(p_{t,s}) \frac{\partial}{\partial p_{t,s}} \varphi(P) + \frac{n}{4} \varphi(P) a_{t,s} \cos(p_{t,s}). \end{aligned} \quad (2.2)$$

It follows from (2.1) that

$$\frac{\partial^2 R_{k,n}}{\partial p_{u,v} \partial p_{t,s}}(P_0) = 0$$

and from (2.2) that

$$\begin{aligned} \frac{\partial^2 R_{k,n}}{\partial p_{t,s}^2}(P_0) &= \frac{n}{4} a_{t,s} \varphi(P_0) \\ &= \frac{n}{4} a_{t,s} \sum_{\varepsilon_i, \varepsilon_j \in \{-1, 1\}} -\varepsilon_t \varepsilon_s \left(a + \sum_{1 \leq i < j \leq k} \varepsilon_i \varepsilon_j a_{i,j} \right)^{\frac{n-2}{2}} \\ &:= \gamma_{t,s}. \end{aligned}$$

We also consider the determinants

$$\begin{aligned}
D_1(P_0) &:= \frac{\partial^2 R_{k,n}}{\partial p_{1,2}^2}(P_0) = \gamma_{1,2} \\
D_2(P_0) &:= \begin{vmatrix} \frac{\partial^2 R_{k,n}}{\partial p_{1,2}^2}(P_0) & \frac{\partial^2 R_{k,n}}{\partial p_{1,2} \partial p_{1,3}}(P_0) \\ \frac{\partial^2 R_{k,n}}{\partial p_{1,3} \partial p_{1,2}}(P_0) & \frac{\partial^2 R_{k,n}}{\partial p_{1,3}^2}(P_0) \end{vmatrix} = \begin{vmatrix} \gamma_{1,2} & 0 \\ 0 & \gamma_{1,3} \end{vmatrix} = \gamma_{1,2} \gamma_{1,3} \\
&\vdots \\
D_{\frac{k^2-k}{2}}(P_0) &:= \begin{vmatrix} \frac{\partial^2 R_{k,n}}{\partial p_{1,2}^2}(P_0) & \frac{\partial^2 R_{k,n}}{\partial p_{1,2} \partial p_{1,3}}(P_0) & \cdots & \frac{\partial^2 R_{k,n}}{\partial p_{1,2} \partial p_{k-1,k}}(P_0) \\ \frac{\partial^2 R_{k,n}}{\partial p_{1,3} \partial p_{1,2}}(P_0) & \frac{\partial^2 R_{k,n}}{\partial p_{1,3}^2}(P_0) & \cdots & \frac{\partial^2 R_{k,n}}{\partial p_{1,3} \partial p_{k-1,k}}(P_0) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 R_{k,n}}{\partial p_{k-1,k} \partial p_{1,2}}(P_0) & \frac{\partial^2 R_{k,n}}{\partial p_{k-1,k} \partial p_{1,3}}(P_0) & \cdots & \frac{\partial^2 R_{k,n}}{\partial p_{k-1,k}^2}(P_0) \end{vmatrix} \\
&= \begin{vmatrix} \gamma_{1,2} & 0 & 0 & 0 \\ 0 & \gamma_{1,3} & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \gamma_{k-1,k} \end{vmatrix} = \gamma_{1,2} \gamma_{1,3} \cdots \gamma_{k-1,k}.
\end{aligned}$$

It is not hard to see that for each t, s , $\gamma_{t,s} < 0$ if $n > 2$ and $\gamma_{t,s} > 0$ if $0 < n < 2$. Hence $(-1)^i D_i(P_0) > 0$ ($i = 1, 2, \dots, (k^2 - k)/2$) for $n > 2$, whence, by utilizing the second partial test, we infer that

$$\begin{aligned}
\max_P R_{k,n}(P) &= \max_{P_0} R_{k,n}(P_0) \\
&= \frac{1}{2} \sum_{\varepsilon_i, \varepsilon_j \in \{-1, 1\}} \left(2 \sum_{1 \leq i, j \leq k} \varepsilon_i \varepsilon_j \|x_i\| \|x_j\| \right)^{n/2} \\
&= \frac{1}{2} \sum_{\varepsilon_i \in \{-1, 1\}} \left[\left(\sum_{i=1}^k \varepsilon_i \|x_i\| \right)^2 \right]^{n/2} \\
&= \frac{1}{2} \sum_{\varepsilon_i \in \{-1, 1\}} \left(\sum_{i=1}^k \varepsilon_i \|x_i\| \right)^n \\
&= \sum_{\varepsilon_i \in \{-1, 1\}} \left(\|x_1\| + \sum_{i=2}^k \varepsilon_i \|x_i\| \right)^n \\
&= A_{k,n},
\end{aligned}$$

which yields (I). Similarly, $D_i(P_0) > 0$ ($i = 1, 2, \dots, (k^2 - k)/2$) for $0 < n < 2$, whence, by utilizing the second partial test, we deduce that

$$\min_P R_{k,n}(P) = \min_{P_0} R_{k,n}(P_0) = A_{k,n},$$

which gives us (II).

Sufficiency.

Assume that condition (I) to be held. The continuity of the function $n \mapsto \|\cdot\|^n$ implies that

$$R_{k,2} \leq A_{k,2} = k2^{k-1}$$

for $\|x_1\| = \dots = \|x_k\| = 1$. From the pertinent sufficient condition of M.M. Day, it can be proved the following criterion [6]:

“The necessary and sufficient condition for a norm defined over a vector space X to spring from an inner product is that $R_{k,2} \leq k2^{k-1}$ where $k \geq 2$ is a positive integer and $\|x_1\| = \dots = \|x_k\| = 1$ ”. Due to this condition holds, we conclude that the norm $\|\cdot\|$ on X can be deduced from an inner product.

Similarly, if condition (II) holds, then we get

$$R_{k,2} \geq A_{k,2} = k2^{k-1}$$

for $\|x_1\| = \dots = \|x_k\| = 1$. Applying the same statement as the above criterion except that $R_{k,2} \geq k2^{k-1}$, we conclude that the norm $\|\cdot\|$ on X can be deduced from an inner product. \square

Corollary 2.2. *A normed space $(X, \|\cdot\|)$ is an inner product space if*

$$\sum_{\varepsilon_i \in \{-1,1\}} \left\| x_1 + \sum_{i=2}^k \varepsilon_i x_i \right\|^2 = \sum_{\varepsilon_i \in \{-1,1\}} \left(\|x_1\| + \sum_{i=2}^k \varepsilon_i \|x_i\| \right)^2 \quad (2.3)$$

for some $k \geq 2$ and all $x_1, \dots, x_k \in X$. The converse is true if (2.3) holds for all $k \geq 2$ and all $x_1, \dots, x_k \in X$.

We can have an operator version of Corollary above. In fact a straightforward computation shows that

Corollary 2.3. *Let $k \geq 2$ and T_1, T_2, \dots, T_k be bounded linear operators acting on a Hilbert space. Then*

$$\sum_{\varepsilon_i \in \{-1,1\}} \left| T_1 + \sum_{i=2}^k \varepsilon_i T_i \right|^2 = 2^{k-1} \sum_{i=1}^k |T_i|^2 = \sum_{\varepsilon_i \in \{-1,1\}} \left(|T_1| + \sum_{i=2}^k \varepsilon_i |T_i| \right)^2,$$

where $|T| = (T^*T)^{1/2}$ denotes the absolute value of T .

REFERENCES

- [1] J. Alonso and A. Ullán, *Moduli in normed linear spaces and characterization of inner product spaces*, Arch. Math. (Basel) **59** (1992), no. 5, 487–495.
- [2] C. Alsina, J. Sikorska and M.S. Justyna, *Norm derivatives and characterizations of inner product spaces*, Hackensack, NJ: World Scientific, 2010.
- [3] D. Amir, *Characterizations of inner product spaces*, Operator Theory: Advances and Applications, 20 Birkhäuser Verlag, Basel, 1986.
- [4] M. Baronti and E. Casini, *Characterizations of inner product spaces by orthogonal vectors*, J. Funct. Spaces Appl. **4** (2006), no. 1, 1–6.
- [5] F. Dadipour and M.S. Moslehian, *An approach to operator Dunkl–Williams inequality*, J. Math. Anal. Appl. (to appear).
- [6] M.M. Day, *Some characterizations of inner-product spaces*, Trans. Amer. Math. Soc. **62** (1947), 320–337.
- [7] C.R. Diminnie, E.Z. Andalafte and R. Freese, *Triangle congruence characterizations of inner product spaces*, Math. Nachr. **144** (1989), 81–86.
- [8] S.S. Dragomir, *Some characterizations of inner product spaces and applications*, Studia Univ. Babeş-Bolyai Math. **34** (1989), no. 1, 50–55.
- [9] M. Fréchet, *Sur la définition axiomatique d’une classe d’espaces vectoriels distanciés applicables vectoriellement sur l’espace de Hilbert*, Ann. of Math. (2) **36** (1935), no. 3, 705–718.
- [10] P. Jordan and J. von Neumann, *On inner products in linear, metric spaces*, Ann. of Math. (2) **36** (1935), no. 3, 719–723.
- [11] M.S. Moslehian and J.M. Rassias, *A characterization of inner product spaces concerning an Euler–Lagrange identity*, Commun. Math. Anal. **8** (2010), no. 2, 16–21.
- [12] K. Nikodem and Z. Pales, *Characterizations of inner product spaces by strongly convex functions*, Banach J. Math. Anal. (to appear).
- [13] P.L. Papini, *Inner products and norm derivatives*, J. Math. Anal. Appl. **91** (1983), no. 2, 592–598.
- [14] J.M. Rassias, *Two new criteria on characterizations of inner products* Discuss. Math. **9** (1988), 255–267 (1989).
- [15] J.M. Rassias, *Four new criteria on characterizations of inner products*, Discuss. Math. **10** (1990), 139–146 (1991).
- [16] Th.M. Rassias, *New characterizations of inner product spaces*, Bull. Sci. Math. (2) **108** (1984), no. 1, 95–99.
- [17] N. Zettili, *Quantum Mechanics: Concepts and Applications*, Chichester: John Wiley & Sons, Ltd, 2009.

¹ DEPARTMENT OF PURE MATHEMATICS, FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASHHAD 91775, IRAN.

E-mail address: moslehian@ferdowsi.um.ac.ir and moslehian@ams.org

URL: <http://profsite.um.ac.ir/~moslehian/>

² PEDAGOGICAL DEPARTMENT, NATIONAL AND CAPODISTRIAN UNIVERSITY OF ATHENS, SECTION OF MATHEMATICS AND INFORMATICS, 4, AGAMEMNONOS STR., AGHIA PARASKEVI, ATTIKIS 15342, ATHENS, GREECE.

E-mail address: jrassias@primedu.uoa.gr and Ioannis.Rassias@primedu.uoa.gr

URL: <http://www.primedu.uoa.gr/~jrassias/>