

BEURLING-FOURIER ALGEBRAS, OPERATOR AMENABILITY AND ARENS REGULARITY

HUN HEE LEE AND EBRAHIM SAMEI

ABSTRACT. We introduce the class of Beurling-Fourier algebras on locally compact groups and show that they are non-commutative analogs of classical Beurling algebras. We obtain various results with regard to the operator amenability, operator weak amenability and Arens regularity of Beurling-Fourier algebras on compact groups and show that they behave very similarly to the classical Beurling algebras of discrete groups. We then apply our results to study explicitly the Beurling-Fourier algebras on $SU(2)$, the 2×2 unitary group. We demonstrate that how Beurling-Fourier algebras are closely connected to the amenability of the Fourier algebra of $SU(2)$. Another major consequence of our results is that our investigation allows us to construct families of unital infinite-dimensional closed Arens regular subalgebras of the Fourier algebra of certain products of $SU(2)$.

Beurling algebras play an important role in different areas of harmonic analysis. These are L^1 -algebras associated to locally compact groups when we put extra “weight” on the groups (see Section 1.2). The basic properties of these algebras are well-known since the works of Beurling [1], [2], and Domar [8], for abelian groups, and Reiter [35] for the general case (see also [7], [12], [22], [23], [24], and [37]). For example, it is shown in [8] that the Beurling algebra $L^1(G, \omega)$ is $*$ -regular for G abelian if and only if the weight ω is symmetric and non-quasianalytic. Also various aspects of cohomologies and Arens regularities of Beurling algebras have been studied by several authors, most notably Grønbaek [22], [23], and Dales and Lau [7]. It is shown that $L^1(G, \omega)$ is amenable as a Banach algebra if and only if G is amenable as a locally compact group and $\{\omega(x)\omega(x^{-1}) : x \in G\}$ is bounded [23]. This demonstrates that in most cases, the amenability of Beurling algebras forces the weight to be trivial. On the other hand, even though the group algebra $L^1(G)$ is not Arens regular when G is infinite, for a large classes of weights, it can happen that $\ell^1(G, \omega)$ will be Arens regular [7].

The aim of the present paper is to develop the corresponding “dual theory” for the classical Beurling algebras. That is, we consider the Fourier algebra $A(G)$ of a locally compact group G , and the question of how could

1991 *Mathematics Subject Classification.* Primary 43A30, 46J10; Secondary 22E15, 43A75.

Key words and phrases. locally compact groups, Beurling algebras, Beurling-Fourier algebras, operator amenability, operator weak amenability, Arens regularity, 2×2 special unitary group, Heisenberg groups.

we interpret Beurling algebras in this context and what would be their basic properties? In the language of Kac algebras [10] (or more generally locally compact quantum groups - see [30]), $A(G)$ is interpreted as the dual object of $L^1(G)$ in the sense of generalized Pontryagin duality. In particular, when G is abelian, with dual group \widehat{G} , then $A(G) \cong L^1(\widehat{G})$ via the Fourier transform. Thus for an abelian group G and a weight ω on \widehat{G} , we define the Beurling-Fourier algebra $A(G, \omega)$ to be the Fourier transform of Beurling algebra $L^1(\widehat{G}, \omega)$ [35, Section 6.3]. In the general non-abelian setting though, \widehat{G} is not a group and so the extension of this idea is more delicate!

In order to achieve our goal, we need to focus on the somewhat non-standard interpretation of the weight ω . Consider the co-multiplication

$$\Gamma : L^\infty(G) \rightarrow L^\infty(G \times G), f \mapsto \Gamma f,$$

where

$$\Gamma f(s, t) = f(st).$$

This Γ can be easily extended to unbounded Borel measurable functions on G using the same formula. Now let $\omega : G \rightarrow (0, \infty)$ be a continuous function. Then the submultiplicativity of ω (i.e. ω being a weight) is clearly equivalent to the condition

$$(0.1) \quad \Gamma(\omega)(\omega^{-1} \otimes \omega^{-1}) \leq 1.$$

Now let $VN(G)$ be the group von Neumann algebra of G , and let Γ be the usual co-multiplication on $VN(G)$ defined by

$$(0.2) \quad \Gamma : VN(G) \rightarrow VN(G \times G), \lambda(s) \mapsto \lambda(s) \otimes \lambda(s),$$

where λ is the left regular representation of G . In Section 2.1, we consider a dual version of weight functions satisfying a dual version of (0.1), which requires an extension of the $*$ -isomorphism Γ in (0.2) for certain unbounded operators. For a fixed representation $VN(G) \subset B(H)$, we define a *weight on the dual of G* to be a “suitable” densely defined (possibly unbounded) operator W acting on H which is affiliated to $VN(G)$ (Definition 2.4). To simplify our computation, we make a further assumption that W has a bounded inverse $W^{-1} \in VN(G)$. One major condition that W has to satisfy is the corresponding dual version of (0.1):

$$\Gamma(W)(W^{-1} \otimes W^{-1}) \leq 1_{VN(G \times G)}.$$

We show that this is the natural extension of a weight on duals of non-abelian groups. Furthermore, (see Definition 2.6), we define

$$VN(G, W^{-1}) := \{AW : A \in VN(G)\}$$

and equip $VN(G, W^{-1})$ with an operator space structure induced by the natural linear isomorphism

$$\Phi : VN(G) \rightarrow VN(G, W^{-1}), A \mapsto AW.$$

We will denote the predual of $VN(G, W^{-1})$ by $A(G, W)$ and show that it is a completely contractive Banach algebra. We call $A(G, W)$ the **Beurling-Fourier algebras** on G .

In the reminder of Section 2, we show that our approach allows us to construct various classes of weights on duals of not necessary abelian groups, namely compact groups and Heisenberg groups. In Sections 2.2, we compute certain *central* weights on duals of compact groups. By central weights, we mean those weights that roughly speaking commute with elements of $VN(G)$ (Definition 2.4). We show that these central weight on \widehat{G} , the dual of a compact group G , are of the form

$$(0.3) \quad W = \bigoplus_{\pi \in \widehat{G}} \omega(\pi) 1_{M_{d_\pi}},$$

where $\omega : \widehat{G} \rightarrow (\delta, \infty)$, for some $\delta > 0$, is a function satisfying (2.12). In this case, we write $A(G, \omega)$ instead of $A(G, W)$. When G is abelian, the relation (2.12) is exactly the submultiplicity of ω . However we also construct central weights on duals of non-abelian compact groups using (2.12)(see Example 2.15). One family of weights which are of particular interest to us is ($a \geq 0$),

$$(0.4) \quad \omega_a(\pi) = d_\pi^a \quad (\pi \in \widehat{G}).$$

We also characterize certain forms of central weights on duals of Heisenberg groups in terms of weights on their center (Section 2.3 and Definition 2.18).

Section 3 is devoted to study operator amenability, operator weak amenability, and Arens regularity of the Beurling-Fourier algebra $A(G, \omega)$ when G is compact and W is the central weight (0.3). In Section 3.2, we first compute the operator amenability constant of $A(G, \omega)$ for G finite and use it to characterize the operator amenability of $A(G, \omega)$ when G is an arbitrary product of finite groups and ω is the corresponding weight associated to this product. By applying this result to products of S_3 , the permutation group on $\{1, 2, 3\}$, we construct Beurling-Fourier algebra with arbitrary operator amenability constant. This is in contrast to the Fourier algebra of compact group since the operator amenability constant is always 1 [36]. We then change our focus and show that for a compact group G , $A(G, \omega)$ fails to be operator amenable if $\Omega(\pi) = \omega(\pi)\omega(\bar{\pi}) \rightarrow \infty$ whenever $\pi \rightarrow \infty$ in the discrete topology. This provides, for instance, central weights (such as the one defined in (0.4)) on compact connected semisimple Lie groups whose Beurling-Fourier algebras are not operator amenable. On some other direction, we show that $A(G, \omega)$ is always operator weakly amenable if G is totally disconnected (Section 3.3). Finally in Section 3.4, we present various classes of central weights whose Beurling-Fourier algebras are Arens regular or fails to be Arens regular. For instance, we show that $A(G, \omega_a)$ is Arens regular if G is a compact connected semi-simple Lie group and ω_a is a weight satisfying (0.4). All of these results go parallel to the analogous results in [7], [22], and [23] for classical Beurling algebras.

In Section 4, we apply the results of the preceding section to study explicitly Beurling-Fourier algebras on $SU(2)$. We present various classes weights on $\widehat{SU(2)}$ and show the interesting fact that their Beurling-Fourier algebras behave vary similarly to the corresponding Beurling algebras on the $\mathbb{Z} = \widehat{\mathbb{T}}$, where we regard \mathbb{T} as the maximal torus of $SU(2)$. In Section 4.3, we explain in details the intriguing connection between Beurling-Fourier algebras on $SU(2)$ and the fundamental work of B. E. Johnson in [28] on non-amenability of the Fourier algebra $A(G)$ for a compact connected non-abelian Lie group G . We should say that this was one of the major motivations for us to do this project.

The final Section 4.5 is perhaps the most surprising to us because there are no corresponding results in the classical Beurling algebras! We construct unital infinite-dimensional closed subalgebras of the Fourier algebra of certain products of $SU(2)$ which are *Arens regular*. We actually show that they are of the form $A(SU(2), \omega_{2^n})$, where $n \in \mathbb{N}$ and ω_{2^n} is the weight defined in (0.4). This is remarkable because this can not happen for the classical Beurling algebras! There are unital infinite-dimensional Arens regular Beurling algebras but they can never be closed subalgebras of some group algebra. These connections are certainly worthwhile further investigations.

In collaboration with M. Ghandehari, we have obtained further results concerning Beurling-Fourier algebras of Heisenberg groups $H_d = \mathbb{C}^d \times \mathbb{R}$ ($d \in \mathbb{N}$) and $n \times n$ special unitary groups $SU(n)$ which will appear in the subsequent article [20]. We would like to point out that J. Ludwig, N. Spronk, and L. Turowska in [31] have also considered and studied the properties of Beurling-Fourier algebras on compact groups. However they have mainly focused on the question of determining the spectrum of Beurling-Fourier algebras. Their investigation is parallel to ours and provides a very good complement to our paper.

1. PRELIMINARIES

1.1. Fourier algebras. Let G be a locally compact group with a fixed left Haar measure. We denote the group algebra of G with $L^1(G)$. Given a function f on G the left and right translation of f by $x \in G$ is denoted by $(L_x f)(y) = f(xy)$ and $(R_x f)(y) = f(yx)$, respectively. Let $P(G)$ be the set of all continuous positive definite functions on G and let $B(G)$ be its linear span. The space $B(G)$ can be identified with the dual of the group C^* -algebra $C^*(G)$, this latter being the completion of $L^1(G)$ under its largest C^* -norm. With the pointwise multiplication and the dual norm, $B(G)$ is a commutative regular semisimple Banach algebra. The Fourier algebra $A(G)$ is the closure of $B(G) \cap C_c(G)$ in $B(G)$. It was shown in [11] that $A(G)$ is a commutative regular semisimple Banach algebra whose carrier space is G . Also, if λ is the left regular representation of G on $L^2(G)$ then, up to isomorphism, $A(G)$ is the unique predual of $VN(G)$, the von Neumann algebra generated by the representation λ .

Let \widehat{G} be the collection of all equivalence classes of weakly continuous irreducible unitary representations of G into $B(H_\pi)$ for some Hilbert space H_π . \widehat{G} can be regarded as the *dual* of G . If G is abelian, the \widehat{G} is the set of continuous characters from G into $\mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$ which forms a locally compact abelian group with compact-open topology. The well-known Fourier transform gives us the identification $L^1(G) \cong A(\widehat{G})$ isometrically as Banach algebras.

If G is a compact group, then for all $\pi \in \widehat{G}$, H_π is finite-dimensional. We denote $d_\pi = \dim H_\pi$, M_{d_π} to be the matrix representation of $B(H_\pi)$, and use the convention that d_π is the dimension of π . If $\pi \in \widehat{G}$, we fix an orthonormal basis $\{\xi_1^\pi, \dots, \xi_{d_\pi}^\pi\}$ for H_π and define

$$(1.1) \quad \pi_{ij} : G \rightarrow \mathbb{C} \quad , \quad \pi_{ij}(s) = \langle \pi(s)\xi_j^\pi \mid \xi_i^\pi \rangle$$

for $i, j = 1 \dots d_\pi$. We recall the well-known fact that

$$(1.2) \quad \mathcal{T}(G) = \text{span}\{\pi_{ij} : \pi \in \widehat{G}, i, j = 1, \dots, d_\pi\}$$

is uniformly dense in $C(G)$, the space of continuous functions on G . The Fourier transform on $L^1(G)$ is the one-to-one $*$ -linear mapping \mathcal{F} defined by

$$(1.3) \quad \mathcal{F} : L^1(G) \rightarrow \bigoplus_{\pi \in \widehat{G}}^{\infty} M_{d_\pi} \quad , \quad f \mapsto (\widehat{f}(\pi))_{\pi \in \widehat{G}} \quad ,$$

where $\widehat{f}(\pi) = \int_G f(t)\overline{\pi}(t)dt \in M_{d_\pi}$ and $\overline{\pi}$ is the conjugate representation of π . Moreover,

$$(1.4) \quad \mathcal{F}(\mathcal{T}(G)) = \left\{ \bigoplus_{\pi \in F} A_\pi : A_\pi \in M_{d_\pi}, F \subset \widehat{G} \text{ is finite} \right\}.$$

Note that if $A = \bigoplus_{\pi \in F} A_\pi$ with $F \subset \widehat{G}$ finite, then

$$(1.5) \quad f(x) = \mathcal{F}^{-1}(A)(x) = \sum_{\pi \in F} d_\pi \text{tr}(A_\pi \pi(x)), \quad x \in G.$$

Also if we regard $L^1(G)$ as convolution operators on $L^2(G)$, then $L^1(G)$ is a subalgebra of $VN(G)$ and \mathcal{F} induces an $*$ -isomorphism

$$(1.6) \quad \mathcal{F} : VN(G) \cong \bigoplus_{\pi \in \widehat{G}}^{\infty} M_{d_\pi}.$$

Note that the above direct sums over \widehat{G} assume the repetition of the same component d_π -times for $\pi \in \widehat{G}$. It follows from the preceding identification that

$$A(G) = \{f \in C(G) : \|f\|_{A(G)} = \sum_{\pi \in \widehat{G}} d_\pi \|\widehat{f}(\pi)\|_1 < \infty\},$$

where $\|\cdot\|_1$ is the trace-class norm on M_{d_π} . See [25, sections 27 and 34] for complete details.

1.2. Beurling algebras. Let G be a locally compact group. A weight on G is a continuous function $\omega : G \rightarrow (0, \infty)$ such that

$$\omega(st) \leq \omega(s)\omega(t) \quad (s, t \in G).$$

Sometimes we allow a weight ω just to be measurable and locally finite (i.e. bounded on every compact subset of G), but it is known that ([35, Theorem 3.7.5]) for every measurable weight ω there is a continuous weight ω' equivalent to ω .

For a (continuous) weight w we define weighted spaces

$$L^1(G, \omega) := \{f \text{ Borel measurable} : \|f\|_{L^1(G, \omega)} = \|\omega f\|_{L^1(G)} < \infty\}$$

and

$$L^\infty(G, \frac{1}{\omega}) := \{f \text{ Borel measurable} : \|f\|_{L^\infty(G, \frac{1}{\omega})} = \left\| \frac{f}{\omega} \right\|_{L^\infty(G)} < \infty\},$$

which are isometric to $L^1(G)$ and $L^\infty(G)$, respectively. Moreover, $L^\infty(G, \frac{1}{\omega})$ is the dual of $L^1(G, \omega)$ with the duality bracket

$$\langle f, g \rangle = \int_G f(x)g(x)d\mu(x), \quad f \in L^1(G, \omega), \quad g \in L^\infty(G, \frac{1}{\omega}),$$

where μ is the left Haar measure on G .

For discrete G we denote $L^1(G, \omega)$ by $\ell^1(G, \omega)$. With the convolution multiplication $L^1(G, \omega)$ becomes a Banach algebra (due to the multiplicativity of the weight), and the algebras $L^1(G, \omega)$ are called the *Beurling algebras* on G . For more details see [7, Chapter 7].

1.3. Operator spaces. We will now briefly remind the reader about the basic properties of operator spaces. We refer the reader to [9] for further details concerning the notions presented below.

Let \mathcal{H} be a Hilbert space. Then there is a natural identification between the space $M_n(B(\mathcal{H}))$ of $n \times n$ matrices with entries in $B(\mathcal{H})$ and the space $B(\mathcal{H}^n)$. This allows us to define a sequence of norms $\{\|\cdot\|_n\}$ on the spaces $\{M_n(B(\mathcal{H}))\}$. If V is any subspace of $B(\mathcal{H})$, then the spaces $M_n(V)$ also inherit the above norm. A subspace $V \subseteq B(\mathcal{H})$ together with the family $\{\|\cdot\|_n\}$ of norms on $\{M_n(V)\}$ is called a *concrete operator space*. This leads us to the following abstract definition of an operator space:

Definition 1.1. An operator space is a vector space V together with a family $\{\|\cdot\|_n\}$ of Banach space norms on $M_n(V)$ such that for each $A \in M_n(V)$, $B \in M_m(V)$ and $[a_{ij}], [b_{ij}] \in M_n(\mathbb{C})$

$$i) \quad \left\| \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \right\|_{n+m} = \max\{\|A\|_n, \|B\|_m\}$$

$$ii) \quad \|[a_{ij}]A[b_{ij}]\|_n \leq \|[a_{ij}]\| \|A\|_n \|[b_{ij}]\|$$

Let V, W be operator space, $\varphi : V \rightarrow W$ be linear. Then

$$\|\varphi\|_{cb} = \sup_n \{\|\varphi_n\|\}$$

where $\varphi_n : M_n(V) \rightarrow M_n(W)$ is given by

$$\varphi_n([v_{ij}]) = [\varphi(v_{ij})].$$

We say that φ is completely bounded if $\|\varphi\|_{cb} < \infty$; is completely contractive if $\|\varphi\|_{cb} \leq 1$ and is a complete isometry if each φ_n is an isometry.

Given two operator spaces V and W , we let $CB(V, W)$ denote the space of all completely bounded maps from V to W . Then $CB(V, W)$ becomes a Banach space with respect to the norm $\|\cdot\|_{cb}$ and is in fact an operator space via the identification $M_n(CB(V, W)) \cong CB(V, M_n(W))$.

It is well-known that every Banach space can be given an operator space structure, though not necessarily in a unique way. It is also clear that any subspace of an operator space is also an operator space with respect to the inherited norms. Moreover, for duals and preduals of operator spaces, there are canonical operator space structures. As such the predual of a von Neumann algebra and the dual of a C^* -algebras respectively, the Fourier and Fourier-Stieltjes algebras inherit natural operator space structures.

Given two Banach spaces V and W , there are many ways to define a norm on the algebraic tensor product $V \otimes W$. Distinguished amongst such norms is the *Banach space projective tensor product norm* which we denote by $V \otimes^\gamma W$. A fundamental property of the projective tensor product is that there is a natural isometry between $(V \otimes^\gamma W)^*$ and $B(V, W^*)$. Given two operator spaces V and W , there is an operator space analog of the projective tensor product norm which we denote by $V \widehat{\otimes} W$. In this case, we have a natural complete isometry between $(V \widehat{\otimes} W)^*$ and $CB(V, W^*)$.

Definition 1.2. A Banach algebra A that is also an operator space is called a *completely contractive Banach algebra* if the multiplication map

$$m : A \widehat{\otimes} A \rightarrow A, \quad u \otimes v \mapsto uv$$

is completely contractive. In particular, both $B(G)$ and $A(G)$ are completely contractive Banach algebras (see [11]).

Let A be a completely contractive Banach algebra. An operator space X is called a *completely bounded A -bimodule*, if X is a Banach A -bimodule and if the maps

$$A \widehat{\otimes} X \rightarrow X, \quad u \otimes x \mapsto ux$$

and

$$X \widehat{\otimes} A \rightarrow X, \quad x \otimes u \mapsto xu$$

are completely bounded. In general, if X is a completely bounded A -bimodule, then its dual space X^* is a completely bounded A -bimodule via the actions

$$(u \cdot T)(x) = T(xu) \quad , \quad (T \cdot u)(x) = T(ux)$$

for every $u \in A$, $x \in X$, and $T \in X^*$.

A is *operator amenable* if, for every completely contractive Banach X -bimodules, every completely bounded derivation from A into X^* is inner. One characterization of operator amenability is that A is operator amenable if and only if it has a *virtual diagonal* [27] i.e. there is $M \in (A \widehat{\otimes} A)^{**}$ such that

$$a \cdot M = M \cdot a, \quad am^{**}(M) = m^{**}(M)a = a \quad (a \in A),$$

where $a \cdot (b \otimes c) = ab \otimes c$, $(b \otimes c) \cdot a = b \otimes ca$, and $\pi : A \otimes A \rightarrow A$ is the multiplication operator. A is *operator weakly amenable* if every completely bounded derivation from A into A^* is inner [19].

1.4. Arens regular Banach algebras. Let A be a (completely contractive) Banach algebra. We can define two products on A^{**} , the second dual of A , known as the *first and second Arens products* as follows: For every $F, E \in A^{**}$ with $F = w^* - \lim_{\alpha} f_{\alpha}$, and $E = w^* - \lim_{\beta} g_{\beta}$, $\{f_{\alpha}\}, \{g_{\beta}\} \subset A$, we let the first (second) Arens product be

$$F \square E = w^* - \lim_{\alpha} \lim_{\beta} f_{\alpha} g_{\beta} \quad \text{and} \quad F \diamond E = w^* - \lim_{\beta} \lim_{\alpha} g_{\beta} f_{\alpha}.$$

We say that A is **Arens regular** if the first and second Arens products always coincide i.e.

$$F \square E = F \diamond E, \quad \forall F, E \in A^{**}.$$

If A is Arens regular, then every closed subalgebra of A or a quotient of A is also Arens regular. It is well-known that C^* -algebras (or more generally, operator algebras) are Arens regular. However the group algebra $L^1(G)$ is Arens regular if and only if G is finite [7].

Also the Arens regularity of the Fourier algebra $A(G)$ implies that G is discrete, non-amenable, and does not contain a copy of \mathbb{F}_2 , the free group on two generators [14], [15]. It is still an open question whether the Arens regularity of $A(G)$ implies that G is finite.

2. BEURLING-FOURIER ALGEBRA ON A LOCALLY COMPACT GROUP

2.1. General construction. We begin the construction of a dual object of classical Beurling algebras by the following reformulation of the multiplicativity of weight functions.

Let G be a locally compact group and recall the co-multiplication

$$\Gamma : L^{\infty}(G) \rightarrow L^{\infty}(G \times G), \quad f \mapsto \Gamma f,$$

where $\Gamma f(s, t) = f(st)$. This Γ can be easily extended for unbounded Borel measurable functions on G using the same formula.

Now let $\omega : G \rightarrow (0, \infty)$ be a continuous function. Then the submultiplicativity of ω is clearly equivalent to the condition

$$(2.1) \quad \Gamma(\omega)(\omega^{-1} \otimes \omega^{-1}) \leq 1.$$

Our aim is first to define a dual version of weight functions satisfying a dual version of (2.1), which requires an extension of a $*$ -isomorphism for certain unbounded operators. We will describe the process in the following

lemma. We refer the reader to [4, Chapters X.1 and X.2] and [29, Chapter 5.5.6] for the definition and basic properties of unbounded operators.

Lemma 2.1. *Let $\mathcal{M} \subseteq B(H)$ and $\mathcal{N} \subseteq B(K)$ be von Neumann algebras and $\Phi : \mathcal{M} \rightarrow \mathcal{N}$ be a $*$ -isomorphism. We suppose that*

- (1) *there is an increasing net of projections $(E_i)_{i \in \mathcal{I}} \subseteq \mathcal{M}$ such that $\mathcal{D} := \bigcup_{i \in \mathcal{I}} E_i(H)$ is dense in H ,*
- (2) *there is a closed operator W on H with the domain containing \mathcal{D} such that WE_i 's are bounded self-adjoint operators in \mathcal{M} , and*
- (3) *$\mathcal{D}' := \bigcup_{i \in \mathcal{I}} \Phi(E_i)(K)$ is dense in K .*

Then, the linear operator B defined on \mathcal{D}' by

$$B(k) := \Phi(WE_i)(k) \text{ for } k \in \Phi(E_i)(K)$$

is a closable operator on K , whose closure is self-adjoint.

Proof. Since $(E_i)_{i \in \mathcal{I}}$ is increasing, $(\Phi(E_i))_{i \in \mathcal{I}}$ is also an increasing net of projections in \mathcal{N} , so that B is well-defined. Now we can apply the same argument as in [29, Lemma 5.6.1] to show that B is closable with the self-adjoint closure acting on K . \square

Definition 2.2. Suppose that we are in the same situation as in Lemma 2.1. We define $\Phi(W)$ acting on K by $\Phi(W) := \overline{B}$, where \overline{B} is the closure of B .

Remark 2.3. (1) The above definition of $\Phi(W)$ is an extension of Φ in the following sense. If W is bounded with the domain H , then $\Phi(W)$ defined in Definition 2.2 (denoted by T) and the original $\Phi(W)$ (denoted by S) coincide on a dense subspace of K . Indeed, if we put $W_i = WE_i$, $i \in \mathcal{I}$, where E_i 's are the projection in Lemma 2.1, then we have $W_i \rightarrow W$ strongly, and so, $W \in \mathcal{M}$. Moreover since W_i 's and W are uniformly bounded, we have actually $W_i \rightarrow W$ σ -strongly. Thus, $\Phi(W_i) \rightarrow S$ σ -strongly. From the definition it is clear that $\Phi(W_i)x \rightarrow Tx$ for all $x \in \mathcal{D}'$, so that $Tx = Sx$ for all $x \in \mathcal{D}'$, and \mathcal{D}' is dense in K .

(2) We will use the convention that if two bounded operators S, T , acting on a Hilbert space, coincide on a dense subspace, then we identify S and T , and we use the notation $S = T$.

Now we go back to the definition of a dual version of weight functions. Let $VN(G)$ be the group von Neumann algebra, and let Γ be the usual co-multiplication on $VN(G)$ defined by

$$\Gamma : VN(G) \rightarrow VN(G \times G), \lambda(s) \mapsto \lambda(s) \otimes \lambda(s),$$

where λ is the left regular representation of G . Recall that a densely defined (possibly unbounded) operator T acting on H is said to be *affiliated to \mathcal{M}* , a von Neumann algebra in $B(H)$, if $UTU^* = T$ for any unitary $U \in \mathcal{M}'$ [29, Chapter 5.5.6], and that T is called *boundedly invertible* if there is a bounded operator $S : H \rightarrow H$ such that $TS = id_H$ and $ST \subseteq id_H$ [4, 1.14

Definition]. In the latter case, the choice of S is unique so we denote S by T^{-1} and call it the *bounded inverse of T* .

Definition 2.4. Let G be a locally compact group, and let $VN(G) \subseteq B(H)$ be a fixed representation of $VN(G)$. A closed densely defined positive operator W on H affiliated to $VN(G)$ with the bounded inverse $W^{-1} \in VN(G)$ is called a **weight on the dual of G** if

- (1) W satisfies the conditions in Lemma 2.1 with $\mathcal{M} = VN(G)$, $\mathcal{N} = VN(G \times G) \subseteq B(H \otimes_2 H)$, and $\Phi = \Gamma$,
- (2) $\mathcal{D}_0 := \{x \in H \otimes_2 H : (W^{-1} \otimes W^{-1})x \in \mathcal{D}'\}$ is dense in $H \otimes_2 H$,
- (3) $\Gamma(W)(W^{-1} \otimes W^{-1})$ is bounded on \mathcal{D}_0 (we still denote its unique extension to $H \otimes_2 H$ by $\Gamma(W)(W^{-1} \otimes W^{-1})$),
- (4) $\Gamma(W)(W^{-1} \otimes W^{-1}) \leq 1_{VN(G \times G)}$, and
- (5) $VN(G)W^{-1} := \{AW^{-1} : A \in VN(G)\}$ is w^* -dense in $VN(G)$.

We say that a weight W on the dual of G is **central** if $WE_i \in VN(G)'$ for any $i \in \mathcal{I}$, where $(E_i)_{i \in \mathcal{I}}$ is the net of projections in Lemma 2.1.

Remark 2.5. (1) In this paper, we will usually exploit the representation of $VN(G)$ coming from the representation theory of the group G in the concrete examples, namely the case of compact groups and the case of Heisenberg groups.

(2) We require our weight W to be boundedly invertible in order to avoid unnecessary difficulties of unbounded inverses. Of course, we sacrifice some generality here, but all of our examples show that this is a reasonable restriction.

Definition 2.6. For a weight W on the dual of G we define

$$(2.2) \quad VN(G, W^{-1}) := \{AW : A \in VN(G)\}.$$

Hence each element of $VN(G, W^{-1})$ is a densely defined operator on H . We put the canonical linear structure on $VN(G, W^{-1})$. Since $W^{-1} \in VN(G)$, it follows that the mapping

$$(2.3) \quad \Phi : VN(G) \rightarrow VN(G, W^{-1}), \quad A \mapsto AW$$

is a linear isomorphism. We endow an operator space structure on $VN(G, W^{-1})$ so that Φ induces a complete isometry. In particular,

$$\|AW\|_{VN(G, W^{-1})} = \|A\|_{VN(G)}.$$

We will denote the predual of $VN(G, W^{-1})$ by $A(G, W)$.

Finally we define $C_r^*(G, W^{-1})$ by

$$C_r^*(G, W^{-1}) := \{AW : A \in C_r^*(G)\}.$$

Clearly $\Phi|_{C_r^*(G)}$ is a complete isometry between $C_r^*(G)$ and $C_r^*(G, W^{-1})$.

Remark 2.7. (1) The above definition of $A(G, W)$ is an abstract one, but we have a natural realization of $A(G, W)$ as follows. For any $\phi \in A(G)$, $W^{-1}\phi$ is an element in $A(G)$ satisfying

$$(W^{-1}\phi)(A) = \phi(AW^{-1}), \quad A \in VN(G).$$

Hence we have

$$(2.4) \quad A(G, W) = \{W^{-1}\phi : \phi \in A(G)\}$$

with the duality bracket

$$(2.5) \quad \langle W^{-1}\phi, AW \rangle = \phi(A)$$

for $\phi \in A(G)$ and $A \in VN(G)$. Moreover, Φ is w^* - w^* continuous and its preadjoint $\Phi_* : A(G, W) \rightarrow A(G)$ is given by

$$\Phi_*(W^{-1}\phi) = \phi.$$

(2) The condition (5) of Definition 2.4 is redundant if the weight W is central. Indeed, $VN(G)W^{-1}$ is w^* -dense in $VN(G)$ if and only if the map $A(G) \rightarrow A(G)$, $\varphi \mapsto W^{-1}\varphi$ is one-to-one. Now suppose that $W^{-1}\varphi = 0$. Then $WE_iW^{-1}\varphi = 0$, $i \in \mathcal{I}$, where E_i 's are the projection in Lemma 2.1. However $WE_i \in VN(G)'$, and so, $WE_iW^{-1}\varphi = W^{-1}\varphi WE_i = E_i \rightarrow 1_{VN(G)}$ strongly. Hence $\varphi = 0$.

(3) Since $W^{-1} \in VN(G)$, the inclusion map (or the formal identity) $j : VN(G) \rightarrow VN(G, W^{-1})$, $A \mapsto (AW^{-1})W$ is a completely bounded w^* - w^* continuous map with $\|j\|_{cb} \leq \|W^{-1}\|$. Moreover, j has a dense range since $\Phi^{-1} \circ j : VN(G) \rightarrow VN(G)$, $A \mapsto AW^{-1}$ has a dense range by Definition 2.4 (5). This implies that the preadjoint of j , $j_* : A(G, W) \rightarrow A(G)$ is completely bounded and one-to-one. Note that j_* is clearly the formal identity. Thus we can (and will) assume that $A(G, W) \subseteq A(G)$ and view any element $\phi \in A(G, W)$ as a continuous function on G vanishing at infinity.

(4) We do not know whether $W \otimes W$ always defines a weight on the dual of $G \times G$. Nevertheless we can formally define $VN(G \times G, W^{-1} \otimes W^{-1})$ and $A(G \times G, W \otimes W)$ similar to (2.2) and (2.4), respectively. This induces the natural complete isometry

$$(2.6) \quad \Psi : VN(G \times G) \rightarrow VN(G \times G, W^{-1} \otimes W^{-1}), \quad A \otimes B \mapsto (A \otimes B)(W \otimes W).$$

In fact, we can identify

$$(A(G, W) \widehat{\otimes} A(G, W))^* = VN(G \times G, W^{-1} \otimes W^{-1}).$$

Indeed, from (2.3) and (2.6) we have the following composition of complete isometries

$$A(G, W) \widehat{\otimes} A(G, W) \xrightarrow{\Phi_* \otimes \Phi_*} A(G) \widehat{\otimes} A(G) \cong A(G \times G) \xrightarrow{\Psi_*^{-1}} A(G \times G, W \otimes W),$$

which can be easily checked to be the formal identity.

Now we would like to endow a completely contractive Banach algebra structure on $A(G, W)$. Recall that the Banach algebra structure of $A(G)$ comes from the co-multiplication Γ , so that we will consider an appropriate map $VN(G, W^{-1}) \rightarrow VN(G \times G, W^{-1} \otimes W^{-1})$, which is essentially the extension of Γ . By (3) in Definition 2.6 we have a normal complete contraction

$$\tilde{\Gamma} : VN(G) \rightarrow VN(G \times G),$$

defined by

$$A \mapsto \Gamma(A)\Gamma(W)(W^{-1} \otimes W^{-1}).$$

We define the w^* - w^* continuous complete contraction

$$\Gamma^W : VN(G, W^{-1}) \rightarrow VN(G \times G, W^{-1} \otimes W^{-1})$$

by

$$(2.7) \quad \Gamma^W := \Psi \circ \tilde{\Gamma} \circ \Phi^{-1}.$$

We can say that Γ^W is essentially an extension of Γ in the following sense.

Theorem 2.8. *Let G be a locally compact group, and let W be a weight on the dual of G . Then the following diagram is commutative:*

$$\begin{array}{ccc} VN(G) & \xrightarrow{\Gamma} & VN(G \times G) \\ \downarrow j & & \downarrow j \otimes j \\ VN(G, W^{-1}) & \xrightarrow{\Gamma^W} & VN(G \times G, W^{-1} \otimes W^{-1}). \end{array}$$

Proof. It suffices to show that for every $A \in VN(G)$,

$$(2.8) \quad \Gamma^W(A)x = \Gamma(A)x$$

for all $x \in \mathcal{D}(W) \otimes \mathcal{D}(W)$, where $\mathcal{D}(W)$ is the domain of W . Let $W_i = WE_i$, $i \in \mathcal{I}$, where E_i 's are the projection in Lemma 2.1. Then we have $W^{-1}W_i x \rightarrow x$ for all $x \in \mathcal{D} = \bigcup_{i \in \mathcal{I}} E_i(H)$. Since $W^{-1}W_i$'s are uniformly bounded and \mathcal{D} is dense in H , we have $W^{-1}W_i \rightarrow 1_{VN(G)}$ σ -strongly. Thus $\Gamma(W^{-1})\Gamma(W_i) \rightarrow 1_{VN(G \times G)}$ σ -strongly. Since

$$\Gamma(W_i)x \rightarrow \Gamma(W)x$$

for all $x \in \mathcal{D}' = \bigcup_{i \in \mathcal{I}} \Gamma(E_i)(H \otimes_2 H)$, we have $\Gamma(W^{-1})\Gamma(W)x = x$ for all $x \in \mathcal{D}'$, so that for every $A \in VN(G)$

$$\begin{aligned} \Gamma(AW^{-1})\Gamma(W)(W^{-1} \otimes W^{-1})x &= \Gamma(A)\Gamma(W^{-1})\Gamma(W)(W^{-1} \otimes W^{-1})x \\ &= \Gamma(A)(W^{-1} \otimes W^{-1})x \end{aligned}$$

for all $x \in \mathcal{D}_0 = \{x \in H \otimes_2 H : (W^{-1} \otimes W^{-1})x \in \mathcal{D}'\}$. Since \mathcal{D}_0 is dense in $H \otimes_2 H$ ((2) of Definition 2.4) and both operators are bounded ((3) of Definition 2.4), we have

$$\Gamma(AW^{-1})\Gamma(W)(W^{-1} \otimes W^{-1}) = \Gamma(A)(W^{-1} \otimes W^{-1}).$$

Thus

$$\begin{aligned}
\Gamma^W(A) &= \Psi(\tilde{\Gamma}(\Phi^{-1}(A))) \\
&= \Psi(\tilde{\Gamma}(AW^{-1})) \\
&= \Psi(\Gamma(AW^{-1})\Gamma(W)(W^{-1} \otimes W^{-1})) \\
&= \Psi(\Gamma(A)(W^{-1} \otimes W^{-1})) \\
&= \Gamma(A)(W^{-1} \otimes W^{-1})(W \otimes W).
\end{aligned}$$

Hence (2.8) follows. \square

We are now ready to define a suitable completely contractive Banach algebra structure on $A(G, W)$. Indeed, since Γ^W is a complete contraction and also a w^* - w^* continuous mapping, the preadjoint Γ_*^W of Γ^W defines a completely contractive Banach algebra structure on $A(G, W)$. This will allow us to present the following definition.

Definition 2.9. Let G be a locally compact group, and let W be a weight on the dual of G . The completely contractive Banach algebra $A(G, W)$ defined in Definition 2.6 with the multiplication

$$\Gamma_*^W : A(G, W) \widehat{\otimes} A(G, W) \rightarrow A(G, W)$$

is called the **Beurling-Fourier algebra** on G .

We will use the notation

$$\phi \cdot_{A(G, W)} \psi = \Gamma_*^W(\phi \otimes \psi), \quad \phi, \psi \in A(G, W),$$

while

$$\phi \cdot_{A(G)} \psi = \Gamma_*(\phi \otimes \psi), \quad \phi, \psi \in A(G).$$

Remark 2.10. (1) It follows from the commuting diagram in Theorem 2.8 that the following diagram is also commutative:

$$\begin{array}{ccc}
A(G \times G, W \otimes W) & \xrightarrow{\Gamma_*^W} & A(G, W) \\
\downarrow \iota_* \otimes \iota_* & & \downarrow \iota_* \\
A(G \times G) & \xrightarrow{\Gamma_*} & A(G).
\end{array}$$

This implies that for every $\phi, \psi \in A(G, W)$,

$$\phi \cdot_{A(G, W)} \psi = \Gamma_*^W(\phi \otimes \psi) = \Gamma_*(\phi \otimes \psi) = \phi \cdot_{A(G)} \psi,$$

or equivalently, the multiplication on $A(G, W)$ can be understood as the pointwise multiplication of continuous functions so that $A(G, W)$ can be viewed as a subalgebra of $A(G)$.

(2) The definition of the Banach algebra structure on $A(G, W)$ for a weight W on the dual of G is somewhat technical since we are working with general unbounded operators. If W is bounded or at least $VN(G)$ is semifinite with a trace τ and W is τ -measurable, then the above construction becomes much easier, since the extension of $*$ -isomorphism can be easily understood ([33,

Lemma 2.4]). However, the weight W we are interested in is usually pretty much unbounded, so that W is not even τ -measurable.

2.2. Central weights on the dual of compact groups. We will show in this section how we can construct central weights on the duals of compact groups. We will see, eventually, that they are a generalization of classical weights on discrete groups.

Let G be a compact group. Then from (1.6),

$$VN(G) \cong \bigoplus_{\pi \in \widehat{G}} M_{d_\pi} \subseteq B(H),$$

where $H = \bigoplus_{\pi \in \widehat{G}} \ell_{d_\pi}^2$. Note that the above direct sums over \widehat{G} assume the repetition of the same component d_π -times for $\pi \in \widehat{G}$. For the rest of this article, we always consider the above representation of $VN(G)$.

Before proceeding further we need to know how the co-multiplication on $VN(G)$ is translated in the above representation of $VN(G)$. Note that the left regular representation λ has the decomposition $\lambda \cong \bigoplus_{\pi \in \widehat{G}} \overline{\pi}$. Consider a central element $W \in VN(G)$ defined by

$$W = \bigoplus_{\pi \in \widehat{G}} \omega(\pi) 1_{M_{d_\pi}},$$

where $\omega(\pi)$'s are positive numbers and $F = \{\pi \in \widehat{G} : \omega(\pi) > 0\}$ is a finite set. Then from (1.3), (1.4) and the Fourier inversion formula (1.5) we have that

$$\mathcal{F}(f) = (\widehat{f}(\pi))_{\pi \in \widehat{G}} = W \text{ or } W = \mathcal{F}\left(\int_G f(x)\lambda(x)dx\right),$$

where

$$f(x) = \sum_{\sigma \in \widehat{G}} d_\sigma \omega(\sigma) \text{tr}(\sigma(x)), \quad (x \in G).$$

Thus

$$\begin{aligned} \Gamma(W) &= \mathcal{F}\left(\int_G f(x)\lambda(x) \otimes \lambda(x)dx\right) \\ &= \bigoplus_{\pi, \pi' \in \widehat{G}} \int_G f(x) \overline{\pi(x)} \otimes \overline{\pi'(x)} dx \\ &= \bigoplus_{\pi, \pi' \in \widehat{G}} \bigoplus_{k=1}^N \int_G f(x) \overline{\tau^k(x)} dx, \end{aligned}$$

where

$$(2.9) \quad \pi \otimes \pi' \cong \bigoplus_{k=1}^N \tau^k$$

for some $(\tau^k)_{k=1}^N \subseteq \widehat{G}$. Note that we are allowing the repetition of τ^k 's, so that it is possible that $\tau^k \cong \tau^l$ for some $k \neq l$. By the Schur orthogonality relation,

$$\begin{aligned} \Gamma(W) &= \bigoplus_{\pi, \pi' \in \widehat{G}} \bigoplus_{k=1}^N \int_G \sum_{\sigma} d_{\sigma} w(\sigma) \sum_{i=1}^{d_{\sigma}} \sigma_{ii}(x) \overline{\tau^k(x)} dx \\ &= \bigoplus_{\pi, \pi' \in \widehat{G}} \bigoplus_{k=1}^N w(\tau^k) 1_{M_{d_{\tau^k}}}. \end{aligned}$$

We can change the order of the direct sum using the following notation.

Definition 2.11. Let ρ be a continuous finite-dimensional (unitary) representation of G . We recall that the **support** of ρ in \widehat{G} is the (finite) set of continuous finite-dimensional irreducible unitary representation of G that appear in the decomposition of ρ , i.e.

$$\text{supp } \rho = \{\tau_i \in \widehat{G} \mid \rho \cong \bigoplus_{i=1}^n \tau_i\}.$$

Using the preceding definition and the fact that $F = \{\pi \in \widehat{G} : \omega(\pi) > 0\}$, we can write

$$(2.10) \quad \Gamma(W) = \bigoplus_{\sigma \in F} \bigoplus_{\substack{\pi, \pi' \in \widehat{G} \\ \sigma \in \text{supp } \pi \otimes \pi'}} w(\sigma) 1_{M_{d_{\sigma}}}.$$

Now we consider a function $\omega : \widehat{G} \rightarrow (\delta, \infty)$ for some $\delta > 0$. We would like to construct a central weight associated to ω . Let \mathcal{F} be the set of all finite subset of \widehat{G} directed by the inclusion. For every $F \in \mathcal{F}$, let E_F be the projection in $VN(G)$ defined by

$$E_F = \bigoplus_{\pi \in F} 1_{M_{d_{\pi}}}.$$

It is clear that $(E_F)_{F \in \mathcal{F}}$ is an increasing net of projections in $VN(G)$ and $\mathcal{D} = \bigcup_{F \in \mathcal{F}} E_F(H)$ is dense in H . Let W_F be the operator in $VN(G)$ given by

$$W_F := \bigoplus_{\pi \in F} \omega(\pi) 1_{M_{d_{\pi}}}.$$

Consider the linear operator W_0 with the domain \mathcal{D} defined by

$$W_0(h) := W_F(h), \quad h \in E_F(H).$$

If we apply the same argument as in [29, Lemma 5.6.1], then we can show that W_0 is closable with the self-adjoint closure. We will denote this closure by

$$(2.11) \quad W = \bigoplus_{\pi \in \widehat{G}} \omega(\pi) 1_{M_{d_{\pi}}}.$$

We can exactly determine when W is a weight on the dual of G .

Theorem 2.12. *Let G be a compact group, and let $\omega : \widehat{G} \rightarrow (\delta, \infty)$ be a function, where $\delta > 0$. The operator W constructed in (2.11) defines a central weight on the dual of G if and only if*

$$(2.12) \quad \omega(\sigma) \leq \omega(\pi)\omega(\pi')$$

for all $\sigma, \pi, \pi' \in \widehat{G}$ with $\sigma \in \text{supp } \pi \otimes \pi'$.

Proof. Following the construction of W , it is routine to verify that W is a closed densely defined positive operator on $H = \bigoplus_{\pi \in \widehat{G}} \ell^2_{d_\pi}$ affiliated to $VN(G)$. Also W has the inverse

$$W^{-1} = \bigoplus_{\pi \in \widehat{G}} \omega(\pi)^{-1} 1_{M_{d_\pi}} \in VN(G),$$

since ω is bounded away from zero. Moreover, (2.10) implies that

$$\Gamma(E_F) = \bigoplus_{\sigma \in F} \bigoplus_{\substack{\pi, \pi' \in \widehat{G} \\ \sigma \in \text{supp } \pi \otimes \pi'}} 1_{M_{d_\sigma}}.$$

Thus it is clear that $\mathcal{D}' = \bigcup_{F \in \mathcal{F}} \Gamma(E_F)(H \otimes_2 H)$ is dense in $H \otimes_2 H$, so that we can apply Lemma 2.1 to define $\Gamma(W)$. Note that we have

$$\Gamma(W)\Gamma(E_F) = \bigoplus_{\sigma \in F} \bigoplus_{\substack{\pi, \pi' \in \widehat{G} \\ \sigma \in \text{supp } \pi \otimes \pi'}} w(\sigma) 1_{M_{d_\sigma}}.$$

On the other hand,

$$\begin{aligned} W^{-1} \otimes W^{-1} &= \bigoplus_{\pi, \pi' \in \widehat{G}} \omega(\pi)^{-1} \omega(\pi')^{-1} 1_{M_{d_\pi}} \otimes 1_{M_{d_{\pi'}}} \\ &= \bigoplus_{\sigma \in \widehat{G}} \bigoplus_{\substack{\pi, \pi' \in \widehat{G} \\ \sigma \in \text{supp } \pi \otimes \pi'}} \omega(\pi)^{-1} \omega(\pi')^{-1} 1_{M_{d_\sigma}}, \end{aligned}$$

and so the condition (2) of Definition 2.4 is clearly satisfied. Moreover,

$$(2.13) \quad \Gamma(W)(W^{-1} \otimes W^{-1}) = \bigoplus_{\sigma \in \widehat{G}} \bigoplus_{\substack{\pi, \pi' \in \widehat{G} \\ \sigma \in \text{supp } \pi \otimes \pi'}} \omega(\sigma) \omega(\pi)^{-1} \omega(\pi')^{-1} 1_{M_{d_\sigma}}.$$

Hence the condition (4) of Definition 2.4 is equivalent to the relation (2.12). Finally, it is clear that $W_F \in VN(G)'$ for every finite subset F of \widehat{G} , and so, by Remark 2.7(2), the condition (5) of Definition 2.4 is satisfied. Consequently, W is a central weight on the dual of G if and only if (2.12) is satisfied. \square

The preceding theorem leads us to the following definition. This idea was also considered by J. Ludwig, N. Spronk, and L. Turowska [31].

Definition 2.13. Let G be a compact group, and $W = \bigoplus_{\pi \in \widehat{G}} \omega(\pi) 1_{M_{d_\pi}}$ be a central weight on the dual of G for a function $\omega : \widehat{G} \rightarrow (\delta, \infty)$ ($\delta > 0$) satisfying (2.12). For convenience, we use ω to represent W , $A(G, \omega)$ to represent $A(G, W)$, $VN(G, \omega^{-1})$ to represent $VN(G, W^{-1})$, $C_r^*(G, \omega^{-1})$ to represent $C_r^*(G, W^{-1})$, and use the terminology that ω is a **central weight** on \widehat{G} . Finally, we define the **symmetrization** of ω , denoted by Ω , to be

$$\Omega(\pi) = \omega(\pi)\omega(\overline{\pi}) \quad (\pi \in \widehat{G}).$$

In particular, we have the completely isometric identification

$$A(G, W) = C_r^*(G, W^{-1})^*.$$

Remark 2.14. (1) Since $A(G, \omega) \subseteq A(G)$ boundedly, we can understand each element in $A(G, \omega)$ as a continuous function on G . More precisely, we have

$$A(G, \omega) \cong \{f \in C(G) : \|f\|_{A(G, \omega)} = \sum_{\pi \in \widehat{G}} d_\pi \omega(\pi) \|\widehat{f}(\pi)\|_1 < \infty\}.$$

(2) It is easy to verify that the symmetrization Ω of ω is also a central weight on \widehat{G} . It is also easy to check that for any two central weights ω_1 and ω_2 on \widehat{G} , the function $\omega_1\omega_2$ defined by

$$(\omega_1\omega_2)(\pi) = \omega_1(\pi)\omega_2(\pi), \quad \pi \in \widehat{G}$$

is again a central weight on \widehat{G} .

(3) Let $\{G_i\}_{i \in I}$ be a family of compact groups, and $F(I)$ be the set of finite subsets of I . It follows from [25, Theorem 27.43] that the dual of $\prod_{i \in I} G_i$ consist of all the representations

$$(\pi_i)_{i \in F} : \prod_{i \in F} G_i \rightarrow B(\otimes_{i \in F} H_{\pi_i}), \quad (x_i)_{i \in F} \mapsto \otimes_{i \in F} \pi_i(x_i) \quad (F \in F(I)).$$

Now suppose that, for every $i \in I$, ω_i is a central weight on \widehat{G}_i . Then it is straightforward to see that the product function $\prod_{i \in I} \omega_i$ given by

$$\left(\prod_{i \in I} \omega_i\right)((\pi_i)_{i \in F}) = \prod_{i \in F} \omega_i(\pi_i)$$

is again a central weight on the dual of $\prod_{i \in I} G_i$ provided that $\prod_{i \in I} \omega_i$ is bounded away from zero as well.

Let $m \in \mathbb{N}$, and $\mathbb{T}^{(m)}$ denotes the m -times Cartesian product of \mathbb{T} . There are various classical weight associated to $\widehat{\mathbb{T}^{(m)}} = \mathbb{Z}^{(m)}$ such as

$$n \mapsto (1 + \ln(1 + \|n\|))^a, \quad n \mapsto (1 + \|n\|)^a \quad (a > 0),$$

where $\|n\|$ is the natural norm on $\mathbb{Z}^{(m)}$. Since the dual of a non-abelian compact group G is not a group anymore, we can not use this idea to define weights on \widehat{G} . However, as we see in the following example, our generalization allows us to define very natural weights on \widehat{G} .

Example 2.15. Let G be a compact group, and let $a \geq 0$. We define the functions σ_a and ω_a from \widehat{G} into $[1, \infty)$ by

$$(2.14) \quad \sigma_a(\pi) = (1 + \ln d_\pi)^a \quad (\pi \in \widehat{G}),$$

$$(2.15) \quad \omega_a(\pi) = d_\pi^a \quad (\pi \in \widehat{G}).$$

It follows from the tensor formula (2.9) that both σ_a and ω_a satisfy (2.12), and so, they are central weights on \widehat{G} . Since the irreducible representations of abelian groups are 1-dimensional, the preceding weights are trivial if G is abelian. Thus they are interesting for compact non-abelian groups.

2.3. Central weights on the dual of the Heisenberg groups. Let H_d ($d \geq 1$) be the Heisenberg group on $\mathbb{C}^d \times \mathbb{R}$. Our references for the Heisenberg groups are [39, Chapter 1] and [13, Examples 6.7 and 7.6]. For $h \in \mathbb{R}^* (= \mathbb{R} \setminus \{0\})$ we consider the Schrödinger representations of H_d acting on $\mathcal{H} = L^2(\mathbb{R}^d)$ defined by

$$\pi_h(z, t)\varphi(\xi) = e^{iht} e^{ih(x \cdot \xi + \frac{1}{2}x \cdot y)} \varphi(\xi + y),$$

where \cdot is the usual inner product in \mathbb{R}^d , $z = x + iy$, $x, y \in \mathbb{R}^d$ and $\varphi \in \mathcal{H}$. The Haar measure on H_d is just the Lebesgue measure on $\mathbb{C}^d \times \mathbb{R}$, which will be denoted by $dzdt$. The Fourier transform on H_d is defined as follows:

$$\widehat{f}^{H_d}(h) = \int_{H_d} f(z, t) \overline{\pi_h(z, t)} dzdt, \quad h \in \mathbb{R}^*$$

for $f \in L^1(H_d)$, and the Plancherel theorem says

$$\int_{\mathbb{R}^*} \left\| \widehat{f}^{H_d}(h) \right\|_{S_2(\mathcal{H})}^2 d\mu(h) = \int_{H_d} |f(z, t)|^2 dzdt,$$

where $S_2(\mathcal{H})$ is the Hilbert-Schmidt class on \mathcal{H} , $d\mu(h) = \frac{|h|^d}{(2\pi)^{d+1}} dh$ on \mathbb{R}^* and $f \in L^1(H_d) \cap L^2(H_d)$. Moreover, it is well known that

$$(2.16) \quad \lambda \stackrel{\text{unitarily}}{\cong} \int^{\oplus} \overline{\pi_h} d\mu(h),$$

where λ is the left regular representation of H_d , and

$$VN(H_d) \cong L^\infty(\mu; B(\mathcal{H})) \subseteq B(H),$$

where $H = L^2(\mu; S_2(\mathcal{H}))$. Note that the above vector-valued L^∞ space $L^\infty(\mu; B(\mathcal{H}))$ can be naturally identified with the von Neumann algebra tensor product $L^\infty(\mu) \overline{\otimes} B(\mathcal{H})$.

Lastly, we recall the Fourier inverse transform

$$\mathcal{F}^{-1} : L^1(\mu; S_1(\mathcal{H})) \rightarrow L^\infty(H_d), \quad X = (X(h)) \mapsto \mathcal{F}^{-1}(X),$$

where

$$\mathcal{F}^{-1}(X)(z, t) = \int_{\mathbb{R}^*} \text{tr}(\pi_h(z, t)X(h)) d\mu(h),$$

$S_1(\mathcal{H})$ is the trace class on \mathcal{H} and $L^1(\mu; S_1(\mathcal{H}))$ refers to a vector-valued L^1 space.

As in the compact group case, we need to know how the co-multiplication is translated in this setting. In order to achieve this, we first need the following lemma.

We fix an orthonormal basis $(\xi_i)_{i \geq 1}$ for \mathcal{H} , and let $P_{ji} \in B(\mathcal{H})$ is the operator defined by

$$(2.17) \quad P_{ji}(\eta) = \langle \eta, \xi_i \rangle \xi_j, \quad i, j \geq 1.$$

Then we get the following substitute for Schur orthogonality.

Lemma 2.16. *Let g be a function in $\mathcal{S}(\mathbb{R})$, the Schwarz class on \mathbb{R} .*

$$(2.18) \quad \begin{aligned} & g(h') \delta_{ik} \delta_{jl} \\ &= \int_{\mathbb{R}^*} \int_{H_d} g(h) \langle \pi_h(z, t) \xi_j, \xi_i \rangle \overline{\langle \pi_{h'}(z, t) \xi_k, \xi_l \rangle} dz dt d\mu(h), \quad i, j, k, l \geq 1 \end{aligned}$$

for every $h' \in \mathbb{R}^*$.

Proof. Let $X = g \otimes P_{ji}$, $i, j \geq 1$. If we take Fourier inverse transform, then we get

$$\begin{aligned} f(z, t) &= \mathcal{F}^{-1}(X)(z, t) = \int_{\mathbb{R}^*} g(h) \text{tr}(\pi_h(z, t) P_{ji}) d\mu(h) \\ &= \int_{\mathbb{R}^*} g(h) \langle \pi_h(z, t) \xi_j, \xi_i \rangle d\mu(h), \quad (z, t) \in H_d. \end{aligned}$$

From the inversion theorem ([39, theorem 1.3.2]) we recover X as the Fourier transform of f , so that we have

$$\begin{aligned} X(h') &= g(h') P_{ji} \\ &= \int_{\mathbb{R}^*} \int_{H_d} g(h) \langle \pi_h(z, t) \xi_j, \xi_i \rangle \overline{\pi_{h'}(z, t)} dz dt d\mu(h) \end{aligned}$$

for every $h' \in \mathbb{R}^*$. Since $\langle P_{ji} \xi_k, \xi_l \rangle = \langle \xi_k, \xi_i \rangle \langle \xi_j, \xi_l \rangle$, we get the conclusion. \square

Now let $W = w \otimes id_n$ for some strictly positive $w \in \mathcal{S}(\mathbb{R})$ and $n \geq 1$, where $id_n = \sum_{i=1}^n P_{ii}$, the $n \times n$ upper-left corner of $1_{B(\mathcal{H})}$. We set

$$f(z, t) = \mathcal{F}^{-1}(W)(z, t) = \int_{\mathbb{R}^*} w(h) \chi_{\pi_h}^n(z, t) d\mu(h), \quad (z, t) \in H_d,$$

where

$$(2.19) \quad \chi_{\pi_h}^n(z, t) = \sum_{i=1}^n \langle \pi_h(z, t) \xi_i, \xi_i \rangle.$$

Then we have $\Gamma(W) = \int_{H_d} f(z, t) \lambda(z, t) \otimes \lambda(z, t) dz dt$, and if we focus on a particular point $(h', h'') \in \mathbb{R}^* \times \mathbb{R}^*$, then by (2.16) we have

$$\begin{aligned}\Gamma(W)(h', h'') &= \int_{H_d} f(z, t) \overline{\pi_{h'}(z, t)} \otimes \overline{\pi_{h''}(z, t)} dz dt \\ &= \int_{\mathbb{R}^*} \int_{H_d} w(h) \chi_{\pi_h^n}^n(z, t) \overline{\pi_{h'}(z, t)} \otimes \overline{\pi_{h''}(z, t)} dz dt\end{aligned}$$

for almost every $(h', h'') \in \mathbb{R}^* \times \mathbb{R}^*$. By the Stone-von Neumann theorem ([13, Theorem 6.49]) for $h' + h'' \neq 0$ (note that the cases $h' + h'' = 0$ are measure zero with respect to $\mu \times \mu$) we have

$$\pi_{h'} \otimes \pi_{h''} \cong \bigoplus_{\alpha} \pi_{h'+h''}^{\alpha},$$

where $\pi_{h'+h''}^{\alpha}$ are copies of $\pi_{h'+h''}$. Thus, by (2.18) and (2.19) we have

$$\begin{aligned}\Gamma(W)(h', h'') &= \bigoplus_{\alpha} \int_{\mathbb{R}^*} \int_{H_d} w(h) \chi_{\pi_h^n}^n(z, t) \overline{\pi_{h'+h''}^{\alpha}(z, t)} dz dt d\mu(h) \\ &= \bigoplus_{\alpha} w(h' + h'') id_n\end{aligned}$$

for almost every $(h', h'') \in \mathbb{R}^* \times \mathbb{R}^*$. Note that the above equality can be extended to any $\omega \in L^{\infty}(\mathbb{R})$ since $\mathcal{S}(\mathbb{R})$ is w^* -dense in $L^{\infty}(\mathbb{R})$, and we can replace id_n by $1_{B(\mathcal{H})}$ by w^* - w^* continuity. Thus, for any $\omega \in L^{\infty}(\mathbb{R})$ and $W = \omega \otimes 1_{B(\mathcal{H})}$ we have

$$(2.20) \quad \Gamma(W)(h', h'') = w(h' + h'') 1_{B(\mathcal{H})} \otimes 1_{B(\mathcal{H})}$$

for almost every $(h', h'') \in \mathbb{R}^* \times \mathbb{R}^*$. Note that we used the fact that $\bigoplus_{\alpha} 1_{B(\mathcal{H})}$ is identified with $1_{B(\mathcal{H})} \otimes 1_{B(\mathcal{H})}$.

Now we consider a continuous positive function ω on \mathbb{R} which is bounded away from zero. We would like to construct a central weight associated to ω . For $m \in \mathbb{N}$, we consider the projection E_m in $VN(H_d)$ given by

$$E_m = 1_{[-m, m]} \otimes 1_{B(\mathcal{H})}.$$

It is clear that $(E_m)_{m \geq 1}$ is an increasing net of projections in $VN(H_d)$ and $\mathcal{D} = \bigcup_{m \geq 1} E_m(H)$ is dense in $H = L^2(\mu; S_2(\mathcal{H}))$. Let W_m be the operator in $VN(H_d)$ given by

$$W_m := (\omega 1_{[-m, m]}) \otimes 1_{B(\mathcal{H})}.$$

Consider a linear operator W_0 with the domain \mathcal{D} defined by

$$W_0(h) := W_m(h), \quad h \in E_m(H).$$

If we apply the same argument as in [29, Lemma 5.6.1], then we can show that W_0 is closable with the self-adjoint closure. We will denote this closure by

$$(2.21) \quad W = \omega \otimes 1_{B(\mathcal{H})}.$$

Similar to the compact groups, we can exactly determine when W defines a weight on the dual of G .

Theorem 2.17. *Let $w : \mathbb{R} \rightarrow (\delta, \infty)$ be a continuous function, where $\delta > 0$. The operator W constructed in (2.21) defines a central weight on the dual of the Heisenberg group H_d if and only if*

$$(2.22) \quad w(h' + h'') \leq w(h')w(h'')$$

for every h' and h'' in \mathbb{R} .

Proof. Following the construction of W , it is routine to verify that W is a closed densely defined positive operator on H_d affiliated to $VN(H_d)$. Also W has the bounded inverse

$$W^{-1} = \omega^{-1} \otimes 1_{B(\mathcal{H})},$$

since ω is bounded away from zero. Moreover, (2.20) implies that

$$\Gamma(E_m)(h', h'') = 1_{[-m, m]}(h' + h'')1_{B(\mathcal{H})} \otimes 1_{B(\mathcal{H})}$$

for almost every $(h', h'') \in \mathbb{R}^* \times \mathbb{R}^*$. Thus it is clear that

$$\mathcal{D}' = \bigcup_{m \geq 1} \Gamma(E_m)(H \otimes_2 H)$$

is dense in $H \otimes_2 H$, so that we can apply Lemma 2.1 to define $\Gamma(W)$. Note that

$$\Gamma(W)(h', h'') = 1_{[-m, m]}(h' + h'')w(h' + h'')1_{B(\mathcal{H})} \otimes 1_{B(\mathcal{H})}$$

on $\Gamma(E_m)(H \otimes_2 H)$, for every $m \geq 1$ and almost every $(h', h'') \in \mathbb{R}^* \times \mathbb{R}^*$. On the other hand,

$$(W^{-1} \otimes W^{-1})(h', h'') = w^{-1}(h')w^{-1}(h'')1_{B(\mathcal{H})} \otimes 1_{B(\mathcal{H})}$$

for every $(h', h'') \in \mathbb{R}^* \times \mathbb{R}^*$. Then clearly \mathcal{D}' and $\mathcal{D}_0 = \{x \in H \otimes_2 H : (W^{-1} \otimes W^{-1})x \in \mathcal{D}'\}$ both contain $C_{00}(\mathbb{R}^* \times \mathbb{R}^*) \otimes S_2(\mathcal{H} \otimes \mathcal{H})$, where $C_{00}(\mathbb{R}^* \times \mathbb{R}^*)$ refers to the space of continuous functions on $\mathbb{R}^* \times \mathbb{R}^*$ with compact support. Moreover, $W^{-1} \otimes W^{-1}$ preserves $C_{00}(\mathbb{R}^* \times \mathbb{R}^*) \otimes S_2(\mathcal{H} \otimes \mathcal{H})$, and since $C_{00}(\mathbb{R}^* \times \mathbb{R}^*) \otimes S_2(\mathcal{H} \otimes \mathcal{H})$ is dense in $H \otimes_2 H$, the condition (2) of Definition 2.4 is satisfied. Moreover, the condition (4) of Definition 2.4 is equal to

$$w(h' + h'') \leq w(h')w(h'')$$

for almost every h' and h'' in \mathbb{R}^* which is equivalent to the relation (2.22) since w is continuous. Finally, it is clear that $W_m \in VN(G)'$ for every $m \in \mathbb{N}$, and so, by Remark 2.7(2), the condition (5) of Definition 2.4 is satisfied. Consequently, $W = \omega \otimes 1_{B(\mathcal{H})}$ defines a weight on the dual of H_d if and only if (2.22) holds. \square

The preceding theorem is the motivation behind the following definition.

Definition 2.18. Let $W = \omega \otimes 1_{B(\mathcal{H})}$ be a central weight on the dual of H_d for a continuous function $\omega : \mathbb{R} \rightarrow (\delta, \infty)$ ($\delta > 0$) satisfying (2.22). For convenience, we use ω to represent W , $A(H_d, \omega)$ to represent $A(H_d, W)$, and use the terminology that ω is a **central weight** on \widehat{H}_d .

Example 2.19. Let $a \geq 0$. We define the function $\tau_a : \mathbb{R} \rightarrow [1, \infty)$ by

$$(2.23) \quad \tau_a(x) = (1 + |x|)^a \quad (x \in \mathbb{R}).$$

By Theorem 2.17 and Definition 2.18, τ_a is a central weight on \widehat{H}_d .

3. COMPACT GROUPS

Throughout this section, G is always assumed to be a compact group. We start with showing certain functorial property that holds for the Beurling-Fourier algebras.

3.1. Functorial Property. Consider the map $Q : A(G \times G) \rightarrow C(G)$ defined by

$$Qw(s) = \int_G w(sr, r) dr.$$

We denote the image of Q by $A_\Delta(G)$. We endow $A_\Delta(G)$ with the operator space structure which makes Q a complete quotient map. We also note that

$$N : A_\Delta(G) \rightarrow A(G \times G), \quad Nu(s, t) = u(st^{-1})$$

is a complete isometry. As in [17, Theorem 2.6], if we repeat the procedure above we obtain

$$(3.1) \quad A_{\Delta^{n+1}}(G) = Q(A_{\Delta^n}(G \times G)) \quad (n \in \mathbb{N}).$$

We can do a similar construction with

$$\check{Q} : A(G \times G) \rightarrow C(G), \quad \check{Q}w(s) = \int_G w(st, t^{-1}) dt.$$

We denote the image of \check{Q} by $A_\gamma(G)$. We endow $A_\gamma(G)$ with the operator space structure which makes \check{Q} a complete quotient map. We also note that

$$\check{N} : A_\gamma(G) \rightarrow A(G \times G : \check{\Delta}), \quad \check{N}u(s, t) = u(st)$$

is a complete isometry. If we repeat the procedure above we obtain

$$(3.2) \quad A_{\gamma^{n+1}}(G) = \check{Q}(A_{\gamma^n}(G \times G)) \quad (n \in \mathbb{N} \cup \{0\}).$$

It follows immediately that, for each $n \in \mathbb{N}$, $A_{\gamma^n}(G)$ is a closed unital subalgebra of the Fourier algebra $A(G^{(2n)})$. Moreover, by [17, Theorem 4.1],

$$(3.3) \quad \|f\|_{A_{\gamma^n}(G)} = \sum_{\pi \in \widehat{G}} d_\pi^{2^n+1} \left\| \widehat{f}(\pi) \right\|_1.$$

Let ω be a central weight on \widehat{G} , and let Ω be the symmetrization of ω (Definition 2.13). Since $A(G \times G, \omega \times \omega)$ is a subalgebra of $A(G \times G)$, we can restrict the map Q to $A(G \times G, \omega \times \omega)$. We denote

$$A_\Delta(G, \Omega) = Q(A(G \times G, \omega \times \omega))$$

and endow $A_\Delta(G, \Omega)$ with the operator space structure so that it became a complete quotient of $A(G \times G, \omega \times \omega)$. It is clear that $A_\Delta(G, \Omega)$ is a completely contractive Banach algebra. Moreover

$$N : A_\Delta(G, \Omega) \rightarrow A(G \times G, \omega \times \omega), \quad Nu(s, t) = u(st^{-1})$$

induces a completely isometric algebraic monomorphism from $A_\Delta(G, \Omega)$ into $A(G \times G, \omega \times \omega)$.

The following theorem explains the motivation behind using Ω in the preceding definition.

Theorem 3.1. *Let G be a compact group, and let ω be a central weight on \widehat{G} . Then*

$$A_\Delta(G, \Omega) = \{f \in C(G) : \sum_{\pi \in \widehat{G}} d_\pi^{3/2} \Omega(\pi) \|\widehat{f}(\pi)\|_2 < \infty\}.$$

Moreover, for every $f \in A_\Delta(G, \Omega)$, we have:

$$\begin{aligned} \|f\|_{A_\Delta(G, \Omega)} &= \inf\{\|u\|_{\omega \times \omega} : u \in A(G \times G, \omega \times \omega), Qu = f\} \\ &= \sum_{\pi \in \widehat{G}} d_\pi^{3/2} \Omega(\pi) \|\widehat{f}(\pi)\|_2. \end{aligned}$$

Proof. It suffices to show that, for $f \in C(G)$,

$$f \in A_\Delta(G, \Omega) \text{ iff } \sum_{\pi \in \widehat{G}} d_\pi^{3/2} \Omega(\pi) \|\widehat{f}(\pi)\|_2 < \infty.$$

We note that $f \in A_\Delta(G, \Omega)$ if and only if $Nf \in A(G \times G, \omega \times \omega)$, in which case $\|f\|_{A_\Delta(G, \Omega)} = \|Nf\|_{\omega \times \omega}$. On the other hand, following a similar argument as in the proof of Theorem 2.2 in [17], we can compute $\|Nf\|_{\omega \times \omega}$, which is $\sum_{\pi \in \widehat{G}} d_\pi^{3/2} \Omega(\pi) \|\widehat{f}(\pi)\|_2$. This completes the proof. \square

We collect some notations for ideals which we will need in this section.

Definition 3.2. Let E be a closed subset of G . We define

$$E^* := \{(s, t) \in G \times G : st^{-1} \in E\}.$$

For any central weight ω on \widehat{G} we define the ideal $I_\omega(E)$ to be the $\|\cdot\|_{A(G, \omega)}$ -closure of $\{f \in \mathcal{T}(G) : f = 0 \text{ on } E\}$. Similarly,

$$I_{\Delta, \Omega}(E) = \{f \in A_\Delta(G, \Omega) : f = 0 \text{ on } E\}$$

and

$$I_{\omega \times \omega}(E^*) = \{g \in A_\Delta(G \times G, \omega \times \omega) : g = 0 \text{ on } E^*\}.$$

Theorem 3.3. *Let G be a compact group, and let ω be a central weight on \widehat{G} . If E is a closed subset of G , then we have*

- (i) $QI_{\omega \times \omega}(E^*) = I_{\Delta, \Omega}(E)$.
- (ii) $I_{\omega \times \omega}(E^*)$ is the closed ideal generated by $NI_{\Delta, \Omega}(E)$.

Proof. Since ω is bounded away from zero, $A(G \times G, \omega \times \omega)$ satisfies the assumption of [17, Theorem 1.4]. Thus it is a special case of [17, Theorem 1.4]. \square

The following proposition is shown to hold in [35, Theorem 3.7.13] when G is abelian. We prove it for the general case with a different method and later use it to relate the properties of different Beurling-Fourier algebras together. But first we need the following definition.

Definition 3.4. For a closed subgroup H of G , we say that $\pi \in \widehat{G}$ is an **extension** of $\tau \in \widehat{H}$ if $\tau \in \text{supp } \pi|_H$, where $\pi|_H$ is the representation on H obtained by restricting on H .

Proposition 3.5. Let ω be a central weight on \widehat{G} , and let H be a closed subgroup of G . Define the function $\omega_H : \widehat{H} \rightarrow (0, \infty)$ by

$$\omega_H(\pi) = \inf\{\omega(\tilde{\pi}) \mid \tilde{\pi} \text{ is an extension of } \pi\}.$$

Then:

(i) ω_H is a central weight on \widehat{H} .

(ii) The restriction map $R_H : \mathcal{T}(G) \rightarrow \mathcal{T}(H)$ extends to a complete quotient map from $A(G, \omega)$ onto $A(H, \omega_H)$.

Proof. (i) By [25, 27.46], ω_H is well-defined. We will show that ω_H is a weight on \widehat{H} . Let $\pi, \rho \in \widehat{H}$ and $\tau \in \widehat{H}$ such that $\tau \in \text{supp } (\pi \otimes \rho)$. We want to prove that

$$\omega_H(\tau) \leq \omega_H(\pi)\omega_H(\rho),$$

or equivalently,

$$(3.4) \quad \omega_H(\tau) \leq \omega(\tilde{\pi})\omega(\tilde{\rho}),$$

for every extension $\tilde{\pi}$ and $\tilde{\rho}$ of π and ρ , respectively. Let

$$(3.5) \quad \tilde{\pi} \otimes \tilde{\rho} = \bigoplus_{i=1}^n \sigma_i,$$

where $\sigma_i \in \widehat{G}$ (note that σ_i 's may not be all distinct). We claim that, for some i_0 , σ_{i_0} is an extension of τ . To see this, first note that

$$\tilde{\pi}|_H = \bigoplus_{j=1}^m \pi_j, \quad \tilde{\rho}|_H = \bigoplus_{k=1}^p \rho_k,$$

with $\pi_i, \rho_i \in \widehat{H}$ with $\pi_1 = \pi$, and $\rho_1 = \rho$. Thus

$$(\tilde{\pi} \otimes \tilde{\rho})|_H = (\pi \otimes \rho) \oplus \left(\bigoplus_{j+k>2}^{m+p} \pi_j \otimes \rho_k \right).$$

Since, by our assumption, $\tau \in \text{supp } (\pi \otimes \rho)$, τ appears in the decomposition of $(\tilde{\pi} \otimes \tilde{\rho})|_H$ into the irreducible elements of \widehat{H} . Hence, by (3.5), τ appears in

the decomposition of $\bigoplus_{i=1}^n \sigma_i|_H$. Therefore, by Schur orthogonality relation, for some i_0 , σ_{i_0} is an extension of τ . Thus

$$\omega_H(\tau) \leq \omega(\sigma_{i_0}) \leq \omega(\tilde{\pi})\omega(\tilde{\rho}),$$

which proves (3.4). This completes the proof.

(ii) Let $\iota : C_r^*(H) \rightarrow C_r^*(G)$ be the $*$ -isomorphism defined by

$$\iota(L_f) = \int_H f(h)\lambda_G(h)dh, \quad f \in L^1(H),$$

where L_f is the convolution operator by f on $L^2(H)$. It is easy to check that $R_H|_{\mathcal{T}(H)} = \iota^*|_{\mathcal{T}(H)}$, so that it suffices to show that ι extends to a complete isometry from $C_r^*(H, \omega_H^{-1})$ into $C_r^*(G, \omega^{-1})$.

Now we fix $n \geq 1$ and consider a finite sequence of distinct representations $(\sigma_k)_{k=1}^N \subseteq \widehat{H}$. Let A be an element in $M_n(VN(H, \omega_H^{-1}))$ supported on $(\sigma_k)_{k=1}^N$, i.e.

$$A = \sum_{i,j=1}^n e_{ij} \otimes \left[\bigoplus_{k=1}^N A_{ij}^k \right],$$

where $A_{ij}^k \in M_{d_{\sigma_k}}$. From (1.4), (1.5), and (1.6) it follows that for every $1 \leq i, j \leq n$, there is $f_{ij} \in L^1(H)$ such that

$$\mathcal{F}(L_{f_{ij}}) = \bigoplus_{k=1}^N A_{ij}^k.$$

Moreover, for every $h \in H$,

$$f_{ij}(h) = \sum_{k=1}^N d_{\sigma_k} \operatorname{tr}(A_{ij}^k \sigma_k(h)).$$

Hence we have

$$\begin{aligned} \iota_n([L_{f_{ij}}]) &= \left[\int_H f_{ij}(h)\lambda_G(h)dh \right] \\ &= \sum_{i,j=1}^n e_{ij} \otimes \left[\bigoplus_{\pi \in \widehat{G}} \int_H \sum_{k=1}^N d_{\sigma_k} \operatorname{tr}(A_{ij}^k \sigma_k(h)) \overline{\pi(h)} dh \right]. \end{aligned}$$

The integrals in the above formula are zero unless there is some $\sigma_k \in \text{supp } \pi$, equivalently, π extends some σ_k . Thus

$$\begin{aligned}
& \|\iota_n([L_{f_{ij}}])\| \\
&= \sup_{\pi \in \widehat{G}} \left\{ \frac{1}{\omega(\pi)} \left\| \sum_{i,j=1}^n e_{ij} \otimes \int_H \sum_{k=1}^N d_{\sigma_k} \text{tr}(A_{ij}^k \sigma_k(h)) \overline{\pi(h)} dh \right\|_{M_n(M_{d\pi})} \right\} \\
&= \sup_{\sigma_k \in \text{supp } \pi} \left\{ \frac{1}{\omega(\pi)} \left\| \sum_{i,j=1}^n e_{ij} \otimes \int_H \sum_{k=1}^N d_{\sigma_k} \text{tr}(A_{ij}^k \sigma_k(h)) \overline{\sigma_k(h)} dh \right\|_{M_n(M_{d\pi})} \right\} \\
&= \max_{1 \leq k \leq N} \frac{1}{\inf\{\omega(\pi) : \sigma_k \in \text{supp } \pi\}} \left\| \sum_{i,j=1}^n e_{ij} \otimes B_{ij}^k \right\|_{M_n(M_{d\pi})} \\
&= \max_{1 \leq k \leq N} \frac{\left\| \sum_{i,j=1}^n e_{ij} \otimes B_{ij}^k \right\|_{M_n(M_{d\pi})}}{\omega_H^{-1}(\sigma_k)} \\
&= \|L_f\|_{M_n(C_r^*(H, \omega_H^{-1}))},
\end{aligned}$$

where $B_{ij}^k = \int_H \sum_{k=1}^N d_{\sigma_k} \text{tr}(A_{ij}^k \sigma_k(h)) \overline{\sigma_k(h)} dh$. By a standard density argument ι extends to a complete isometry from $C_r^*(H, \omega_H^{-1})$ into $C_r^*(G, \omega^{-1})$. \square

We note that every (infinite) compact group contains (infinite) abelian subgroups [41]. Thus by the preceding proposition, every Beurling-Fourier algebra has certain classical Beurling algebras as complete quotients. This can be very useful particularly when the abelian subgroups can be chosen so that they contain various information about the original compact group. This happens, for example, in the case where G is a compact Lie group and H is any maximal torus of G . We will show in details in Section 4 how this idea can be applied to relate properties of Beurling-Fourier algebra on $SU(2)$ and Beurling algebras on \mathbb{T} .

3.2. Operator Amenability. In this section, we present certain criteria for investigating the operator amenability of $A(G, \omega)$. We will later show that this criteria can be applied to large classes of weights. But first, we need to recall the following terminologies:

We recall that a completely contractive Banach algebra A is K -operator amenable if there is a virtual diagonal $M \in (A \widehat{\otimes} A)^{**}$ such that $\|M\| = K$. The *operator amenability constant* of A is the smallest K such that A is K -operator amenable. We also recall that if A is a Banach algebra of continuous functions on a locally compact space X , then for every $x \in X$, a functional $d \in A^*$ is called a *point derivation* at x if

$$d(ab) = a(x)d(b) + b(x)d(a) \quad (a, b \in A).$$

Theorem 3.6. *Let G be a compact group with the identity e , and let ω be a central weight on \widehat{G} . Then:*

(i) *$A(G, \omega)$ is operator amenable if and only if $I_{\Delta, \Omega}(\{e\})$ has a bounded approximate identity.*

(ii) *$A(G, \omega)$ is K -operator amenable if and only if there is $F \in A_{\Delta}(G, \Omega)^{**}$ such that $\|F\| = K$, $\langle F, \delta_e \rangle = 1$, and*

$$f \cdot F = f(e)F \quad (f \in A_{\Delta}(G, \Omega)).$$

(iii) *$A(G, \omega)$ is operator weakly amenable if and only if $\overline{I_{\Delta, \Omega}(\{e\})^2} = I_{\Delta, \Omega}(\{e\})$ is essential, or equivalently, there is no non-zero continuous point derivation on $A_{\Delta}(G, \Omega)$ at e .*

Proof. (i) and (iii). If we let $m : A(G, \omega) \widehat{\otimes} A(G, \omega) \rightarrow A(G, \omega)$ be the multiplication map, then it is easy to verify that

$$I_{\omega \times \omega}(\Delta) = \ker m.$$

Thus following the arguments in [36] and [38], we see that $A(G, \omega)$ is operator amenable (respectively, operator weakly amenable) if and only if $I_{\omega \times \omega}(\Delta)$ has a bounded approximate identity (respectively, $\overline{I_{\omega \times \omega}(\Delta)^2} = I_{\omega \times \omega}(\Delta)$). Thus the results follows from Theorem 3.3 and the fact that $\{e\}^* = \Delta$.

(ii) Let $A(G, \omega)$ be K -operator amenable, and let M be a virtual diagonal for $A(G, \omega)$ with $\|M\| = K$. Let Q^{**} be the second adjoint of $Q : A(G \times G, \omega \times \omega) \rightarrow A_{\Delta}(G, \Omega)$ defined in Section 3.1, and let $F = Q^{**}(M)$. Then it is routine to verify that F holds the required properties of (ii). Conversely, if such an $F \in A_{\Delta}(G, \Omega)^{**}$ exists, then $M = N^{**}(F)$ is a virtual diagonal for $A(G, \omega)$ with $\|M\| = \|F\| = K$. \square

In [28, Theorem 4.1], B. E. Johnson computed the amenability constant of the Fourier algebra of a finite group. The following theorem is the quantization of Johnson's result to Beurling-Fourier algebras on a finite group and its proof is inspired by that of Johnson's.

Theorem 3.7. *Let G be a finite group, and let ω be a central weight on \widehat{G} . Then $A(G, \omega)$ is operator amenable with the operator amenability constant*

$$\frac{\sum_{\pi \in \widehat{G}} d_{\pi}^2 \Omega(\pi)}{\sum_{\pi \in \widehat{G}} d_{\pi}^2}.$$

Proof. It is straightforward to verify that δ_e , the dirac function at $\{e\}$, is the unique element in $A_{\Delta}(G, \Omega)^{**} = A_{\Delta}(G, \Omega)$ satisfying the assumption of Theorem 3.6(ii). Thus it follows from Theorem 3.6(ii) that $A(G, \omega)$ is operator amenable and the operator amenability constant is the $\|\cdot\|_{A_{\Delta}(G, \Omega)}$ -norm of δ_e . However $|G| \widehat{\delta}_e(\pi) = 1_{B(H_{\pi})}$ for every $\pi \in \widehat{G}$. Therefore, considering the well-known fact that $|G| = \sum_{\pi \in \widehat{G}} d_{\pi}^2$, we have

$$\|\delta_e\|_{A_{\Delta}(G, \Omega)} = \sum_{\pi \in \widehat{G}} d_{\pi}^{\frac{3}{2}} \Omega(\pi) \|\widehat{\delta}_e(\pi)\|_2 = \frac{\sum_{\pi \in \widehat{G}} d_{\pi}^2 \Omega(\pi)}{\sum_{\pi \in \widehat{G}} d_{\pi}^2}.$$

□

Corollary 3.8. *Let $\{G_i\}_{i \in \mathbb{N}}$ be a family of finite groups, and let, for each $i \in \mathbb{N}$, ω_i be the central weight on \widehat{G}_i and Ω_i its symmetrization. Let $G = \prod_{i \in I} G_i$ and $\omega = \prod_{i \in I} \omega_i$. Suppose further that ω is bounded away from zero. Then $A(G, \omega)$ is operator amenable if and only if*

$$M_{G, \omega} := \prod_{i \in \mathbb{N}} \frac{\sum_{\pi \in \widehat{G}_i} d_\pi^2 \Omega_i(\pi)}{\sum_{\pi \in \widehat{G}_i} d_\pi^2}$$

is convergent. In this case, $M_{G, \omega}$ is the operator amenability constant of $A(G, \omega)$.

Proof. It is clear that, for each $n \in \mathbb{N}$, there is a complete quotient map from $A(G, \omega)$ onto $A(\prod_{i=1}^n G_i, \prod_{i=1}^n \omega_i)$. Also it follows from Theorem 3.7 that the amenability constant of $A(\prod_{i=1}^n G_i, \prod_{i=1}^n \omega_i)$ is

$$\prod_{i=1}^n \frac{\sum_{\pi \in \widehat{G}_i} d_\pi^2 \Omega_i(\pi)}{\sum_{\pi \in \widehat{G}_i} d_\pi^2}.$$

Therefore the operator amenability constant of $A(G, \omega)$ is at least $M_{G, \omega}$. In particular, if $A(G, \omega)$ is operator amenable, then $M_{G, \omega}$ is convergent. Conversely, suppose that $M_{G, \omega}$ is convergent. Consider the sequence of continuous functions $\{f_n\}$ on G defined by

$$f_n(\{x_i\}) = \begin{cases} 1 & x_1 = \cdots = x_n = e \\ 0 & \text{otherwise.} \end{cases}$$

For each n , we see that

$$\|f_n\|_{A_\Delta(G, \Omega)} = \prod_{i=1}^n \frac{\sum_{\pi \in \widehat{G}_i} d_\pi^2 \Omega_i(\pi)}{\sum_{\pi \in \widehat{G}_i} d_\pi^2}.$$

In particular, $\{f_n\}$ is bounded in $A_\Delta(G, \Omega)$. Let F be a weak*-cluster point of $\{f_n\}$ in $A_\Delta(G, \Omega)^{**}$. Then it is straightforward to verify that F satisfies the hypothesis of Theorem 3.6(ii). Moreover $\|F\| = M_{G, \omega}$. Thus $A(G, \omega)$ is operator amenable with the operator amenability constant $M_{G, \omega}$. □

The preceding corollary has an interesting application when each G_i is S_3 ; the permutation group on $\{1, 2, 3\}$. It is well-known (e.g. [25, 27.61(a)]) that \widehat{S}_3 have two 1-dimensional elements and one 2-dimensional element. Using this fact, we can construct Beurling-Fourier algebras on countably infinite products of S_3 so that they are operator amenable. Moreover, we can let amenability constant be as large as we would like! This is something that does not happen in the Fourier algebra case since the amenability constant is always 1 [36].

Theorem 3.9. *Let $G_i = S_3$ for every $i \in \mathbb{N}$, and let, ω_{a_i} be the central weight (2.14) on \widehat{G}_i defined in Example 2.15. Let $G = \prod_{i \in \mathbb{N}} G_i$ and $\omega = \prod_{i \in \mathbb{N}} \omega_i$. Then:*

- (i) $A(G, \omega)$ is operator amenable if and only if $\sum_{i=1}^{\infty} (2^{2a_i} - 1)$ is convergent;
(ii) For every $1 \leq K < \infty$, we can choose $\{a_i\}$ so that the amenability constant of $A(G, \omega)$ is K .

Proof. By Corollary 3.8, $A(G, \omega)$ is operator amenable if and only if its amenability constant which is

$$\prod_{i \in \mathbb{N}} \frac{\sum_{\pi \in \widehat{G}_i} d_{\pi}^2 \Omega_i(\pi)}{\sum_{\pi \in \widehat{G}_i} d_{\pi}^2} = \prod_{i \in \mathbb{N}} \frac{1 + 2^{2a_i+1}}{3} = \prod_{i \in \mathbb{N}} \left(1 + \frac{2^{2a_i+1} - 2}{3}\right)$$

is finite. However this happens if and only if $\sum_{i=1}^{\infty} \frac{2^{2a_i+1} - 2}{3}$ is convergent.

Thus (i) holds since

$$\sum_{i=1}^{\infty} \frac{2^{2a_i+1} - 2}{3} = 2/3 \sum_{i=1}^{\infty} (2^{2a_i} - 1).$$

The proof of (ii) is easy. In fact, there are various way to chose the required $\{a_i\}$. For example, we can pick a_1 so that $K = \frac{1 + 2^{2a_1+1}}{3}$ and take $a_i = 0$ for $i = 2, 3, \dots$. \square

In [23], N. Grønbaek has shown that the Beurling algebra $L^1(H, \omega)$ on a locally compact group H is amenable if and only if H is amenable and Ω is bounded, where Ω is the symmetrization of ω given by $\Omega(x) = \omega(x)\omega(x^{-1})$, $x \in H$. In the below we prove a weaker version of Grønbaek's result for Beurling-Foureir algebras on compact groups. This presents a nice duality to Grønbaek's criteria.

Theorem 3.10. *Let G be a compact group, and let ω be a central weight on \widehat{G} . Then the followings holds:*

- (i) *If Ω is bounded, then $A(G, \omega)$ is operator amenable;*
(ii) *If $\lim_{\pi \rightarrow \infty} \Omega(\pi) = \infty$, then $A(G, \omega)$ is not operator amenable.*

Proof. (i) If Ω is bounded, then $A_{\Delta}(G, \Omega) = A_{\Delta}(G)$. However, by [17, Theorem 3.9(iii)], $I_{\Delta}(\{e\})$ has a bounded approximate identity. Hence the result follows from Theorem 3.6(i).

(ii) Suppose that $A(G, \omega)$ is operator amenable. Then by Theorem 3.6(ii) and going to an appropriate subnet, there is a bounded net $\{f_{\alpha}\}_{\alpha} \subset A_{\Delta, \Omega}(G)$ such that $f_{\alpha}(e) = 1$ for all α , and

$$(3.6) \quad f f_{\alpha} = f(e) f_{\alpha} \text{ for all } f \in A_{\Delta}(G, \Omega).$$

Since for every $n \in \mathbb{N}$ and $T \in M_n$, $\|T\|_1 \leq n^{1/2} \|T\|_2$, and so, we have $A_{\Delta}(G, \Omega) \subseteq A(G, \Omega)$. Therefore we can assume that $\{f_{\alpha}\}_{\alpha} \subset A(G, \Omega)$. Now let g be a weak*-cluster point of $\{f_{\alpha}\}$ in $A(G, \Omega) = C_r^*(G, \Omega^{-1})^*$. Let $I : \widehat{G} \rightarrow \oplus_{\pi \in \widehat{G}} M_{d_{\pi}}$ be defined by $I(\pi) = 1_{B(H_{\pi})}$ for all $\pi \in \widehat{G}$. It is clear that I is the inverse Fourier transform of δ_e , the evaluation functional on $A(G, \Omega)$

at $\{e\}$. Now since $\lim_{\pi \rightarrow \infty} \Omega(\pi) = \infty$, it follows that $\delta_e = \check{I} \in C_r^*(G, \Omega^{-1})$, and so,

$$g(e) = \langle g, \delta_e \rangle = \lim_{\alpha} \langle f_{\alpha}, \delta_e \rangle = 1.$$

On the other hand, it follows routinely from (3.6) that $g(t) = 0$ if $t \neq e$. Since g is continuous, it follows that G is finite which contradict the fact that $\lim_{\pi \rightarrow \infty} \Omega(\pi) = \infty$. This completes the proof. \square

Corollary 3.11. *Let G be a compact connected, simple Lie group and let σ_a and ω_a be the central weights on \widehat{G} defined in Example 2.15. Then $A(G, \sigma_a)$ or $A(G, \omega_a)$ is operator amenable if and only if $a = 0$.*

Proof. When $a = 0$, $A(G, \sigma_a) = A(G, \omega_a) = A(G)$. Hence the result follows from [36]. For the converse, it is shown in the proof of [34, Lemma 9.1] that, for any positive integer n , there are only finitely many elements in \widehat{G} whose dimension is n . Thus $d_{\pi} \rightarrow \infty$ as $\pi \rightarrow \infty$. Therefore if $a > 0$, by Theorem 3.10, neither of $A(G, \sigma_a)$ or $A(G, \omega_a)$ is operator amenable. \square

3.3. Operator weak amenability. In this section we consider the question of whether a Beurling-Fourier algebra on a compact group can be operator weakly amenable. Our main tool is to use the criterion presented in Theorem 3.6(iii). We first need the following definition.

Definition 3.12. Let π be a continuous finite-dimensional (unitary) representation of G , and let ω be a central weight on \widehat{G} . For each $n \in \mathbb{N}$, we define

$$n(\omega, \pi) = \max\{\omega(\tau) \mid \tau \in \text{supp } \pi\}.$$

Proposition 3.13. *Let ω be a central weight on \widehat{G} . Suppose that, for every $\pi \in \widehat{G}$,*

$$\inf\left\{\frac{n(\omega, \pi^{\otimes n})}{n} \mid n \in \mathbb{N}\right\} = 0,$$

where $\pi^{\otimes n} = \pi \otimes \cdots \otimes \pi$, n -times. Then $A(G, \omega)$ has no non-zero continuous point derivation at e .

Proof. Let $P^1(G)$ denote the set of continuous positive-definite functions on G such that $\|f\|_{A(G)} = f(e) = 1$. Each function in $P^1(G)$ is uniquely determined by a representation $\pi \in \widehat{G}$ and an orthonormal vector $\xi \in H_{\pi}$ so that $f(x) = \langle \pi(x)\xi \mid \xi \rangle$. Hence, in particular, $A(G, \omega)$ is the $\|\cdot\|_{A(G, \omega)}$ -closure of $\text{span}\{P^1(G)\}$ which is $\mathcal{T}(G)$.

Now let $d : A(G, \omega) \rightarrow \mathbb{C}$ be a continuous point derivation at e . We will show that $d = 0$ by showing that d vanishes on $P^1(G)$. Let $f \in P^1(G)$, and $\pi \in \widehat{G}$ and an orthonormal vector $\xi \in H_{\pi}$ so that $f(x) = \langle \pi(x)\xi \mid \xi \rangle$. It is routine to verify that

$$(3.7) \quad d(f^n) = n[f(e)]^{n-1}d(f) = nd(f) \quad (n \in \mathbb{N}).$$

On the other hand, for each $n \in \mathbb{N}$, by Schur orthogonality relation, $\widehat{f^n}(\tau) = 0$ if $\tau \notin \text{supp } \pi^{\otimes n}$. Thus

$$\begin{aligned} \|\widehat{f^n}\|_{A(G,\omega)} &= \sum_{\tau \in \text{supp } \pi^{\otimes n}} \omega(\tau) d_\tau \|\widehat{f^n}(\tau)\|_1 \\ &\leq n(\omega, \pi^{\otimes n}) \sum_{\tau \in \text{supp } \pi^{\otimes n}} d_\tau \|\widehat{f^n}(\tau)\|_1 \\ &= n(\omega, \pi^{\otimes n}) \|f^n\|_{A(G)} \\ &= n(\omega, \pi^{\otimes n}), \end{aligned}$$

where the last equality follows since f^n is a positive-definite function, and so, $\|f^n\|_{A(G)} = f^n(e) = 1$. Therefore, by (3.7),

$$|d(f)| \leq \frac{\|d\| \|f^n\|_{A(G,\omega)}}{n} \leq \|d\| \frac{n(\omega, \pi^{\otimes n})}{n} \quad (n \in \mathbb{N}).$$

So by hypothesis, $d(f) = 0$. \square

The preceding proposition was proven in [37, Proposition 5.1] for G abelian. Our extension allow us to study operator weak amenability for the case when G is non-abelian. The following theorem is one application of Proposition 3.13. Other applications will be given in Section 4.

Theorem 3.14. *Let G be a compact, totally disconnected group, and let ω be a central weight on G . Then $A(G, \omega)$ is operator weakly amenable.*

Proof. It is well-known that G has a base of the identity consisting of open, normal compact subgroups of G , and so, G is a projective limit of finite groups in the sense of [35, Definition 4.1.4] (see also [35, Theorem 4.1.14]). Hence a similar argument to [25, Theorem 27.43] shows that, for every $\pi \in \widehat{G}$, there is a finite group H (which is the quotient of G by some open normal compact subgroup) so that $\pi \in \widehat{H}$. Hence

$$\{\tau \mid \tau \in \text{supp } \pi^{\otimes n}, n \in \mathbb{N}\} \subset \widehat{H}$$

is finite. Thus for the weight $\Omega_1(\sigma) = \Omega(\sigma) d_\sigma$, $\sigma \in \widehat{G}$ (Remark 2.14 (2)), we have

$$\inf\left\{\frac{n(\Omega_1, \pi^{\otimes n})}{n} \mid n \in \mathbb{N}\right\} = 0,$$

for every $\pi \in \widehat{G}$. Therefore, by Proposition 3.13, $A(G, \Omega_1)$ has no continuous point derivation at e . Since $A(G, \Omega_1) \subseteq A_\Delta(G, \Omega)$, the same holds for $A_\Delta(G, \Omega)$. It follows from Theorem 3.6(iii), $A(G, \omega)$ is operator weakly amenable. \square

3.4. Arens regularity. In this section, we study the Arens regularity of Beurling-Fourier algebras. We provide classes of Beurling-Fourier algebras on compact groups that either satisfy or fail the Arens regularity.

Definition 3.15. Let W be a weight on $A(G)$. We denote the bounded operator $\Gamma(W)(W^{-1} \otimes W^{-1})$ by Θ , and we write

$$\Theta = \bigoplus_{\pi, \sigma \in \widehat{G}} \Theta(\pi, \sigma),$$

where $\Theta(\pi, \sigma) \in M_{d_\pi} \otimes M_{d_\sigma}$.

The following theorem is proven in [7, Theorem 8.11] in the case where G is abelian. We extend it to the general case and apply it to construct Arens regular Beurling-Fourier algebras on non-abelian compact groups.

Theorem 3.16. *Let ω be a central weight on \widehat{G} . Suppose that*

$$\lim_{\pi \rightarrow \infty} \limsup_{\sigma \rightarrow \infty} \|\Theta(\pi, \sigma)\|_{M_{d_\pi} \otimes M_{d_\sigma}} = \lim_{\sigma \rightarrow \infty} \limsup_{\pi \rightarrow \infty} \|\Theta(\pi, \sigma)\|_{M_{d_\pi} \otimes M_{d_\sigma}} = 0.$$

Then $A(G, \omega)$ and all its even duals are Arens regular.

Proof. Since $A(G, \omega)^{**} = A(G, \omega) \oplus C_r^*(G, \omega^{-1})^\perp$, it suffices to show that

$$(3.8) \quad \Phi \square \Psi = \Phi \diamond \Psi = 0 \quad (\Phi, \Psi \in C_r^*(G, \omega^{-1})^\perp).$$

To see this, first note that, for all $n \in \mathbb{N}$,

$$A(G, \omega)^{(2n)} = A(G, \omega) \oplus \bigoplus_{i=0}^{n-1} [C_r^*(G, \omega^{-1})^\perp]^{(2i)},$$

where $X^{(m)}$, $m \geq 1$ implies m -th dual of a Banach space X . Thus if (3.8) holds, then both Arens products vanishes on $\bigoplus_{i=0}^{n-1} [C_r^*(G, \omega^{-1})^\perp]^{(2i)}$, which implies that $A(G, \omega)^{(2n-2)}$ is Arens regular. We will now prove (3.8). Suppose that W is the central weight on \widehat{G} defined in Definition 2.13. Let $\Phi, \Psi \in C_r^*(G, \omega^{-1})^\perp$ with norm 1. Using the identification (2.4), take two nets $\{f_\alpha\}$ and $\{g_\beta\}$ in $A(G)$ with $\|f_\alpha\|_{A(G)}, \|g_\beta\|_{A(G)} \leq 1$ such that $W^{-1}f_\alpha \rightarrow \Phi$ and $W^{-1}g_\beta \rightarrow \Psi$ in w^* -topology of $VN(G, W^{-1})$. Let $A \in VN(G)$ with $\|A\|_{VN(G)} \leq 1$. Then, by (2.7),

$$\begin{aligned} & \langle \Phi \square \Psi, AW \rangle \\ &= \lim_{\alpha} \lim_{\beta} \langle (W^{-1}f_\alpha) \cdot_{A(G, W)} (W^{-1}g_\beta), AW \rangle \\ &= \lim_{\alpha} \lim_{\beta} \langle (W^{-1} \otimes W^{-1})(f_\alpha \otimes g_\beta), \Gamma^W(AW) \rangle \\ &= \lim_{\alpha} \lim_{\beta} \langle f_\alpha \otimes g_\beta, \widetilde{\Gamma} \circ \Phi^{-1}(AW) \rangle \\ &= \lim_{\alpha} \lim_{\beta} \langle f_\alpha \otimes g_\beta, \Gamma(A)\Gamma(W)(W^{-1} \otimes W^{-1}) \rangle \\ &= \lim_{\alpha} \lim_{\beta} \sum_{\pi \in \widehat{G}} \sum_{\sigma \in \widehat{G}} d_\pi d_\sigma \text{tr}[(\widehat{f}_\alpha(\pi) \otimes \widehat{g}_\beta(\sigma))\Gamma(A)(\pi, \sigma)\Theta(\pi, \sigma)]. \end{aligned}$$

Now let $\epsilon > 0$. By hypothesis, there is a finite set E in \widehat{G} such that for every $\pi \in E^c := \widehat{G} \setminus E$, there is a finite set F (depending on ϵ and π) in \widehat{G} for

which we have:

$$(3.9) \quad \|\Theta(\pi, \sigma)\|_{M_{d_\pi} \otimes M_{d_\sigma}} \leq \epsilon \quad (\sigma \in F^c).$$

Now for every $\mathcal{A}, \mathcal{B} \subset \widehat{G}$, let

$$\Xi(\mathcal{A}, \mathcal{B}) = \sum_{\pi \in \mathcal{A}} \sum_{\sigma \in \mathcal{B}} d_\pi d_\sigma |\text{tr}[(\widehat{f}_\alpha(\pi) \otimes \widehat{g}_\beta(\sigma))\Gamma(A)(\pi, \sigma)\Theta(\pi, \sigma)]|.$$

Then, for every α and β ,

$$\begin{aligned} |\langle f_\alpha \otimes g_\beta, \Gamma(A)\Gamma(W)(W^{-1} \otimes W^{-1}) \rangle| &\leq \Xi(\widehat{G}, \widehat{G}) \\ &= \Xi(E, \widehat{G}) + \Xi(E^c, F) + \Xi(E^c, F^c). \end{aligned}$$

We will show that

$$\lim_{\alpha} \limsup_{\beta} \Xi(E, \widehat{G}) = \lim_{\alpha} \lim_{\beta} \Xi(E^c, F) = 0 \text{ and } \Xi(E^c, F^c) \leq \epsilon.$$

We have

$$\begin{aligned} \Xi(E, \widehat{G}) &\leq \sum_{\pi \in E} d_\pi \|\widehat{f}_\alpha(\pi)\|_1 \|g_\beta\|_{A(G)} \|\Gamma(A)\| \|\Theta\| \\ &\leq \sum_{\pi \in E} d_\pi \|\widehat{f}_\alpha(\pi)\|_1 \\ &\leq \sum_{\pi \in E} d_\pi^{3/2} \sum_{i,j=1}^{d_\pi} |\widehat{f}_\alpha(\pi)_{ij}|^2, \end{aligned}$$

where $\widehat{f}_\alpha(\pi) = [\widehat{f}_\alpha(\pi)_{ij}]$. Since E does not depend on α and $\Phi \in C_r^*(G, \omega^{-1})^\perp$, for all $\pi \in E$ we have

$$(3.10) \quad \lim_{\alpha} \widehat{f}_\alpha(\pi)_{ij} = \lim_{\alpha} \langle f_\alpha, \overline{\pi_{ij}} \rangle = \langle \Phi, \overline{\pi_{ij}} \rangle = 0,$$

where π_{ij} 's are the trigonometric polynomials defined in (1.1). Therefore

$$\lim_{\alpha} \limsup_{\beta} \Xi(E, \widehat{G}) = 0$$

since E is finite. For the second case, note that

$$\begin{aligned} \Xi(E^c, F) &\leq \sum_{\sigma \in F} d_\sigma \|f_\alpha\| \|\widehat{g}_\beta(\sigma)\|_1 \|\Gamma(A)\| \|\Theta\| \\ &\leq \sum_{\sigma \in F} d_\sigma \|\widehat{g}_\beta(\sigma)\|_1. \end{aligned}$$

Since F does not depend on n and $\Psi \in C_r^*(G, \omega^{-1})^\perp$, similar to (3.10), we have

$$\lim_{\beta} \widehat{g}_\beta(\sigma) = 0$$

for all $\sigma \in F$. Hence, because of finiteness of F ,

$$\lim_{\alpha} \lim_{\beta} \Xi(E^c, F) = 0.$$

Finally

$$\begin{aligned} \Xi(E^c, F^c) &\leq \sum_{\pi \in E^c} \sum_{\sigma \in F^c} d_\pi d_\sigma \|\widehat{f}_\alpha(\pi)\|_1 \|\widehat{g}_\beta(\sigma)\|_1 \|\Gamma(A)\| \|\Theta(\pi, \sigma)\| \\ &\leq \epsilon \sum_{\pi \in E^c} d_\pi \|\widehat{f}_\alpha(\pi)\|_1 \sum_{\sigma \in F^c} d_\sigma \|\widehat{g}_\beta(\sigma)\|_1, \end{aligned}$$

where the last inequality follows from (3.9). Thus, again since $\|f_\alpha\|_{A(G)}$ and $\|g_\beta\|_{A(G)} \leq 1$ we have

$$\Xi(E^c, F^c) \leq \epsilon.$$

since ϵ was arbitrary, it follows that

$$\lim_{\alpha} \lim_{\beta} \langle f_\alpha \otimes g_\beta, \Gamma(A)\Gamma(W)(W^{-1} \otimes W^{-1}) \rangle = 0.$$

Hence this shows that the first Arens product vanishes on $C_r^*(G, \omega^{-1})^\perp$ and we are done! The proof for the second Arens product is similar. \square

Corollary 3.17. *Let G be a compact, connected, simple Lie group, and let σ_a be the central weight on \widehat{G} defined in (2.14). Then $A(G, \sigma_a)$ is Arens regular if $a > 0$.*

Proof. Let Θ_a be the corresponding operator defined in Definition 3.15 associated to the weight σ_a . For every $\pi, \rho \in \widehat{G}$, we have

$$\Theta_a(\pi, \rho) = \bigoplus_{k=1}^m \frac{\sigma_a(\tau_k)}{\sigma_a(\pi)\sigma_a(\rho)} 1_{B(H_{\tau_k})},$$

where $\pi \otimes \rho \cong \bigoplus_{k=1}^m \tau_k$ is the irreducible decomposition of $\pi \otimes \rho$. Since for each $1 \leq k \leq m$, $d_{\tau_k} \leq d_\pi d_\rho$, it follows that

$$\begin{aligned} \sigma_a(\tau_k) &= (1 + \ln d_{\tau_k})^a \\ &\leq (1 + \ln d_\pi)^a + (1 + \ln d_\rho)^a. \\ &= \sigma_a(\pi) + \sigma_a(\rho). \end{aligned}$$

Thus

$$(3.11) \quad \|\Theta_a(\pi, \rho)\| \leq \frac{1}{(1 + \ln d_\pi)^a} + \frac{1}{(1 + \ln d_\rho)^a}.$$

On the other hand, as it was pointed out in the proof of Corollary 3.11, $d_\pi \rightarrow \infty$ as $\pi \rightarrow \infty$. Therefore, from (3.11), it follows that

$$\lim_{\pi \rightarrow \infty} \limsup_{\rho \rightarrow \infty} \|\Theta_a(\pi, \rho)\| = \lim_{\rho \rightarrow \infty} \limsup_{\pi \rightarrow \infty} \|\Theta_a(\pi, \rho)\| = 0.$$

Therefore $A(G, \sigma_a)$ is Arens regular by Theorem 3.16. \square

The preceding example dealt with Beurling-Fourier algebras on certain Lie groups. We can also construct Arens regular Beurling-Fourier algebra on non-abelian totally disconnected groups. We recall that, for each $n \in \mathbb{N}$, the special linear group $SL(2, 2^n)$ denotes the set of all 2×2 matrix with the determinate 1 on a finite field of 2^n elements. It is well-known that $SL(2, 2^n)$ is a finite simple group (see [3, Section 2.7]).

Corollary 3.18. *Let $G = \prod_{n=1}^{\infty} SL(2, 2^n)$, and let σ_a be the central weight on \widehat{G} defined in (2.14). Then $A(G, \sigma_a)$ is Arens regular if $a > 0$.*

Proof. Let Θ_a be the corresponding operator defined in Definition 3.15 associated to the weight σ_a . Similarly to the proof of Corollary 3.17, we have

$$\|\Theta_a(\pi, \rho)\| \leq \frac{1}{(1 + \ln d_{\pi})^a} + \frac{1}{(1 + \ln d_{\rho})^a} \quad (\pi, \rho \in \widehat{G}).$$

However it is shown in [3, Section 2.7] that every non-trivial continuous irreducible unitary representation of $SL(2, 2^n)$ has dimension at least $2^n - 1$. Thus $d_{\pi} \rightarrow \infty$ as $\pi \rightarrow \infty$ on \widehat{G} . Hence

$$\lim_{\pi \rightarrow \infty} \limsup_{\rho \rightarrow \infty} \|\Theta_a(\pi, \rho)\| = \lim_{\rho \rightarrow \infty} \limsup_{\pi \rightarrow \infty} \|\Theta_a(\pi, \rho)\| = 0.$$

Therefore $A(G, \sigma_a)$ is Arens regular by Theorem 3.16. \square

We finish this section with the following theorem that presents examples of non-Arens regular Beurling-Fourier algebras on non-abelian compact groups.

Theorem 3.19. *Let $\{G_i\}_{i \in I}$ be an infinite family of non-trivial compact groups, and let, for each $i \in I$, ω_i be a central weight on \widehat{G}_i . Let $G = \prod_{i \in I} G_i$ and $\omega = \prod_{i \in I} \omega_i$. Suppose further that ω is bounded away from zero. Then $A(G, \omega)$ is not Arens regular.*

Proof. If $J \subseteq I$, then it is clear that $A(\prod_{i \in J} G_i, \prod_{i \in J} \omega_i)$ is a quotient of $A(G, \omega)$. Thus it suffices to prove the statement of the theorem when I is infinite and countable. So we assume that $I = \mathbb{N}$. For each $i \in \mathbb{N}$, let $\pi_i \in \widehat{G}_i$ be a non-trivial representation. For each $m, n \in \mathbb{N}$, let u_m and v_n be elements of \widehat{G} defined by

$$u_m(\tilde{x}) = \pi_{2m}(x_{2m}) \quad , \quad v_n(\tilde{x}) = \pi_{2n+1}(x_{2n+1}) \quad (\tilde{x} = \{x_i\}_{i \in \mathbb{N}} \in G).$$

We have

$$(3.12) \quad u_m \otimes v_n = \begin{cases} \pi_{2m} \times \pi_{2n+1} & \text{if } 2m < 2n + 1 \\ \pi_{2n+1} \times \pi_{2m} & \text{if } 2m > 2n + 1. \end{cases}$$

In particular, $\{u_m \otimes v_n\}_{m, n \in \mathbb{N}}$ are distinct elements of \widehat{G} . Now let

$$f_m = \frac{1}{d_{u_m}} \chi_{u_m} \quad \text{and} \quad g_n = \frac{1}{d_{v_n}} \chi_{v_n},$$

where χ_{π} is the character of $\pi \in \widehat{G}$ i.e. $\chi_{\pi}(x) = \text{tr}[\pi(x)]$. Let $A \in VN(G)$ such that for every $\pi \in \widehat{G}$,

$$\mathcal{F}(A)(\pi) = \begin{cases} 1_{M_{d_{u_m}} \otimes M_{d_{v_n}}} & \text{if } \pi = u_m \otimes v_n \text{ and } 2m < 2n + 1, \\ 0 & \text{otherwise.} \end{cases}$$

Suppose that W is the central weight on \widehat{G} defined in Definition 2.13. Then, by Definition 3.15, (2.10), (2.13), and (3.4), we have

$$\begin{aligned}
\Theta(u_m, v_n) &= [\Gamma(W)(W^{-1} \otimes W^{-1})](u_m, v_n) \\
&= \bigoplus_{\sigma \in \text{supp } u_m \otimes v_n} \omega(\sigma) \omega(u_m)^{-1} \omega(v_n)^{-1} \omega(u_m \otimes v_n) 1_{M_{d\sigma}} \\
&= \omega(u_m)^{-1} \omega(v_n)^{-1} \omega(u_m \otimes v_n) 1_{M_{d_{u_m}} \otimes M_{d_{v_n}}} \\
&= 1_{M_{d_{u_m}} \otimes M_{d_{v_n}}}.
\end{aligned}$$

Thus, by (2.7), for every $m, n \in \mathbb{N}$,

$$\begin{aligned}
&\langle (W^{-1} f_m) \cdot_{A(G,W)} (W^{-1} g_n), AW \rangle \\
&= \langle (W^{-1} \otimes W^{-1})(f_m \otimes g_n), \Gamma^W(AW) \rangle \\
&= \langle f_m \otimes g_m, \widetilde{\Gamma} \circ \Phi^{-1}(AW) \rangle \\
&= \langle f_m \otimes g_n, \Gamma(A) \Gamma(W)(W^{-1} \otimes W^{-1}) \rangle \\
&= d_{u_m} d_{v_n} \text{tr}[(\widehat{f}_m(u_m) \otimes \widehat{g}_n(v_n)) \Gamma(A)(u_m, v_n) \Theta(u_m, v_n)] \\
&= \begin{cases} 1 & \text{if } 2m < 2n + 1 \\ 0 & \text{if } 2m > 2n + 1. \end{cases}
\end{aligned}$$

Therefore the repeated limit of $\langle (W^{-1} f_m) \cdot_{A(G,W)} (W^{-1} g_n), AW \rangle$ exists but they are not equal. This shows that $A(G, \omega)$ is not Arens regular. \square

Remark 3.20. We finish this section by pointing out that the proof of the preceding theorem can be adapted, with almost the same approach, to show that if $\{G_i\}_{i \in I}$ is an infinite family of non-trivial compact groups and if ω_a is the weight (2.15) on the dual of $\prod_{i \in I} G_i$, then $A(G, \omega_a)$ is not Arens regular for any $a \geq 0$.

4. THE 2×2 SPECIAL UNITARY GROUP

In this section, we apply the results of the preceding section to study explicitly the behavior of Beurling-Fourier algebras on 2×2 special unitary group:

$$SU(2) = \{A \in M_2(\mathbb{C}) \mid A \text{ is unitary, } \det A = 1\}.$$

First we make the following important observation which allows us to correspond various central weights on $\widehat{SU(2)}$ to their restriction on \mathbb{Z} .

4.1. Restriction of the weight on \mathbb{Z} . Let ω be a central weight on $\widehat{SU(2)}$. We can assume that \mathbb{T} is a closed subgroup of $SU(2)$ by the identification

$$e^{it} \mapsto \begin{bmatrix} e^{it} & 0 \\ 0 & e^{-it} \end{bmatrix} \quad (t \in [0, 2\pi]).$$

By the representation theory of $SU(2)$ [25, 29.18 and 29.20],

$$\widehat{SU(2)} = \{\pi_l \mid l = 0, 1/2, 1, 3/2, \dots\},$$

and $\dim \pi_l = 2l + 1$. Moreover,

$$\pi_l(e^{it}) = \text{diag}(e^{i2lt}, e^{i2(l-1)t}, \dots, e^{-i2lt}) \quad (t \in [0, 2\pi]).$$

Let $n \in \mathbb{Z}$, and let χ_n be the character on \mathbb{T} defined by

$$\chi_n(e^{it}) = e^{int} \quad (t \in [0, 2\pi]).$$

Then $\widehat{\pi}_l$ is an extension of χ_n if and only if $|n| \leq 2l$. Hence if we identify \mathbb{Z} with $\widehat{\mathbb{T}}$ through the Plancherel map $n \mapsto \chi_n$, we have

$$(4.1) \quad \omega_{\mathbb{T}}(n) = \inf\{\omega(\pi_l) \mid |n| \leq 2l\} \quad (n \in \mathbb{Z}).$$

This, in particular, implies that $A(\mathbb{T}, \omega_{\mathbb{T}})$ is a complete quotient of $A(SU(2), \omega)$ from Proposition 3.5. We will show in the following sections that $A(SU(2), \omega)$ behave very similarly to that of $A(\mathbb{T}, \omega_{\mathbb{T}})$.

Example 4.1. Let $a \geq 0$ and $0 \leq b \leq 1$. We define the functions $\sigma_a, \omega_a, \rho_b$ from $\widehat{SU(2)}$ into $[1, \infty)$ by

$$\begin{aligned} \sigma_a(\pi_l) &= (1 + \ln(2l + 1))^a \quad (\pi_l \in \widehat{SU(2)}), \\ \omega_a(\pi_l) &= (2l + 1)^a \quad (\pi_l \in \widehat{SU(2)}), \\ \rho_b(\pi_l) &= e^{(2l+1)^b} \quad (\pi_l \in \widehat{SU(2)}). \end{aligned}$$

Since $d_{\pi_l} = 2l + 1$, the first two weights are the one defined in Example 2.15. Also we know from [25, 29.29] that, for every $l, r = 0, 1/2, 1, 3/2, \dots$,

$$(4.2) \quad \pi_l \otimes \pi_r \cong \pi_{|l-r|} \oplus \pi_{|l-r|+1} \oplus \dots \oplus \pi_{|l+r|} = \bigoplus_{k=|l-r|}^{|l+r|} \pi_k.$$

Therefore it is routine to verify that ρ_b also defines a weight on $\widehat{SU(2)}$. Moreover, by (4.1), the restriction of the above weights on $\widehat{\mathbb{T}} = \mathbb{Z}$ corresponds to the following well-known weights on \mathbb{Z} :

$$\begin{aligned} \sigma'_a(n) &= (1 + \ln(1 + |n|))^a \quad (n \in \mathbb{Z}), \\ \omega'_a(n) &= (1 + |n|)^a \quad (n \in \mathbb{Z}), \\ \rho'_b(n) &= e^{(1+|n|)^b} \quad (n \in \mathbb{Z}). \end{aligned}$$

4.2. Operator amenability and weak amenability. Let $a \geq 0$ and $0 \leq b \leq 1$, and let σ'_a, ω'_a , and ρ'_b be the weights on \mathbb{Z} defined in Example 4.1. N. Grønbæk has characterized in [22] and [23] when either of $A(\mathbb{T}, \sigma'_a)$, $A(\mathbb{T}, \omega'_a)$, or $A(\mathbb{T}, \rho'_b)$ is amenable or weakly amenable. We summarized them below:

- (i) $A(\mathbb{T}, \sigma'_a)$ or $A(\mathbb{T}, \omega'_a)$ is amenable if and only if $a = 0$;
- (ii) $A(\mathbb{T}, \rho'_b)$ is amenable if and only if $b = 0$;
- (iii) $A(\mathbb{T}, \sigma'_a)$ is always weakly amenable;
- (iv) $A(\mathbb{T}, \omega'_a)$ has no non-zero continuous point derivation at 0 if and only if $0 \leq a < 1$;
- (v) $A(\mathbb{T}, \omega'_a)$ is weakly amenable if and only if $0 \leq a < 1/2$;
- (vi) $A(\mathbb{T}, \rho'_b)$ has non-zero continuous point derivations at 0 if $b > 0$;

(vii) $A(\mathbb{T}, \rho'_b)$ is never weakly amenable unless $b = 0$.

We will show in the following theorem that, in most cases, the analogous of these results holds for the corresponding weights on $\widehat{SU(2)}$.

Theorem 4.2. *Let $a \geq 0$ and $0 \leq b \leq 1$, and let σ_a , ω_a , and ρ_b be the weights on $\widehat{SU(2)}$ defined in Example 4.1. Then the following holds:*

- (i) $A(SU(2), \sigma_a)$ or $A(SU(2), \omega_a)$ is operator amenable if and only if $a = 0$;
- (ii) $A(SU(2), \rho_b)$ is operator amenable if and only if $b = 0$;
- (iii) $A(SU(2), \sigma_a)$ has no non-zero continuous point derivation at e ;
- (iv) $A(SU(2), \omega_a)$ has no non-zero continuous point derivation at e if $0 \leq a < 1$;
- (v) $A(SU(2), \omega_a)$ is not operator weakly amenable if $a \geq 1/2$;
- (vi) $A(SU(2), \rho_b)$ is never operator weakly amenable unless $b = 0$.

Proof. (i) and (ii). If $a = b = 0$, then these Beurling-Fourier algebras are $A(SU(2))$. Thus the result follows from [36]. On the other hand, if $a, b > 0$, then

$$\lim_{\pi_l \rightarrow \infty} \sigma_a(\pi_l) = \lim_{\pi_l \rightarrow \infty} \omega_a(\pi_l) = \lim_{\pi_l \rightarrow \infty} \rho_b(\pi_l) = \infty.$$

Therefore by Theorem 3.10, neither of $A(SU(2), \sigma_a)$, $A(SU(2), \omega_a)$, nor $A(SU(2), \rho_b)$ is operator amenable.

(iii) and (iv). It follows from the tensor formula (4.2) and Schur orthogonality relation that the conjugate of any representation π_l is itself. Moreover, for every $n \in \mathbb{N}$, and $\pi_l \in \widehat{SU(2)}$, we have

$$n(\sigma_a, \pi_l^{\otimes n}) \leq (1 + \ln n + \ln(2l + 1))^a, \quad n(\omega_a, \pi_l^{\otimes n}) \leq [n(2l + 1)]^a.$$

Therefore

$$\inf \left\{ \frac{n(\sigma_a, \pi_l^{\otimes n})}{n} \mid n \in \mathbb{N} \right\} = 0$$

for all $a \geq 0$ and

$$\inf \left\{ \frac{n(\omega_a, \pi_l^{\otimes n})}{n} \mid n \in \mathbb{N} \right\} = 0$$

when $0 \leq a < 1$. Thus the results follow from Proposition 3.13.

(v) and (vi). As it was pointed out in Example 4.1,

$$A(SU(2), \sigma_a)|_{\mathbb{T}} = A(\mathbb{T}, \sigma'_a) \quad \text{and} \quad A(SU(2), \omega_a)|_{\mathbb{T}} = A(\mathbb{T}, \omega'_a).$$

Hence operator weak amenability of $A(SU(2), \sigma_a)$ and $A(SU(2), \omega_a)$ implies the weak amenability of $A(\mathbb{T}, \sigma'_a)$ and $A(\mathbb{T}, \omega'_a)$, respectively. Thus it follows from [22] that $A(SU(2), \omega_a)$ is operator weakly amenable only if $0 \leq a < 1/2$ and $A(SU(2), \rho_b)$ is never operator weakly amenable unless $b = 0$. \square

4.3. Connection with the amenability of $A(SU(2))$. B. E. Johnson in his memoirs [26] in 1972 introduced the concept of an amenable Banach algebra and proved his famous theorem: the group algebra $L^1(G)$ is amenable if and only if G is amenable. It was believed that similar conclusion holds for the Fourier algebra $A(G)$ since $A(G)$ acts in lots of cases like a dual of $L^1(G)$. However it was Johnson himself who proved a remarkable result that

$A(SU(2))$ is not amenable [28]. Shortly after Ruan showed in [36] that the amenability of G corresponds exactly to the “operator amenability” of $A(G)$ which led to the several applications of operator spaces to harmonic analysis. Later on, Forrest and Runde [16] settled the question of amenability for the Fourier algebras: $A(G)$ is amenable if and only if G has an abelian subgroup of finite index.

We would like to analyze the non-amenability of $A(SU(2))$ and show its connection with the Beurling-Fourier algebras on $\widehat{SU(2)}$ and the classical Beurling algebras on \mathbb{Z} . Johnson used the properties of the algebra $A_\gamma(G)$ defined in (3.2) in a very clear way to obtain his result. His approach was widely generalized and studied in [17]. By [28, Theorem 3.2] (see also [17, Corollary 1.5]) the amenability of $A(G)$ implies that the maximal ideal $\{f \in A_\gamma(SU(2)) \mid f(e) = 0\}$ of $A_\gamma(G)$ has a bounded approximate identity. However $A_\gamma(G)$ is nothing but the Beurling-Fourier algebra $A(G, \omega_1)$ in (2.15) where $\omega_1(\pi_l) = 1 + 2l$ for all $\pi_l \in \widehat{SU(2)}$ (see (3.3)). Since, by Example 4.1, the restriction of $A(SU(2), \omega_1)$ on \mathbb{T} is $A(\mathbb{T}, \omega'_1)$, this implies that $\{f \in A(\mathbb{T}, \omega'_1) \mid f(e) = 0\}$ has a bounded approximate identity. Hence by Theorem 3.6, $A(\mathbb{T}, \omega'_1)$ is amenable which is impossible by [23]. This argument can also be applied to the question of weak amenability because again by a similar argument, the weak amenability of $A(SU(2))$ implies that $A(\mathbb{T}, \omega'_{1/2})$ is weakly amenable which is shown to fail in [22].

As we see, the preceding arguments shows that the (weak) amenability of the Fourier algebra $A(SU(2))$ is closely related to the (weak) amenability of a well-known Beurling algebra on \mathbb{Z} and it has inherit connection. That is why $A(SU(2))$ fails to be amenable or even weakly amenable because the Beurling algebras are known not to behave well with regard to cohomology. We believe that these connections are non-trivial and certainly worthwhile investigating more. For example, it is shown in [18] that $A(G)$ is not weakly amenable if G is compact, connected, and non-abelian. If we assume further that G is a Lie group, then again amenability or weak amenability of $A(G)$ relates closely to the behavior of certain Beurling algebras on a maximal torus of G . By investigating more this relation, we might be able to have a better understanding of the structure of Fourier algebras on compact Lie groups.

4.4. Arens regularity. Let $a \geq 0$ and $0 \leq b \leq 1$, and let σ'_a , ω'_a , and ρ'_b be the weights on \mathbb{Z} defined in Example 4.1. As it is shown in [7, Theorem 8.11], $A(\mathbb{T}, \sigma'_a)$, $A(\mathbb{T}, \omega'_a)$, or $A(\mathbb{T}, \rho'_b)$ are Arens regular for $a, b > 0$. We will show in the following theorem that the exact analogous of these results holds for the corresponding weights on $\widehat{SU(2)}$.

Theorem 4.3. *Let $a \geq 0$ and $0 \leq b \leq 1$, and let σ_a , ω_a , and ρ_b be the weights on $\widehat{SU(2)}$ defined in Example 4.1. Then:*

- (i) $A(SU(2), \sigma_a)$ or $A(SU(2), \omega_a)$ is Arens regular if and only if $a > 0$,
- (ii) $A(SU(2), \rho_b)$ is Arens regular if and only if $b > 0$.

Proof. If $a = b = 0$, then these Beurling-Foureir algebras are $A(SU(2))$. Thus the result follows from [14]. For the converse, suppose that $a > 0$. Since $SU(2)$ is a compact, connected, simple Lie group, it follows from Corollary 3.17 that $A(SU(2), \sigma_a)$ is Arens regular. For the case of ω_a , let

$$W_a = \bigoplus_{\pi \in \widehat{G}} \omega_a(\pi) 1_{B(H_\pi)},$$

and let Θ_a be the Fourier transform of $\Gamma(W_a)(W_a^{-1} \otimes W_a^{-1})$ (see Definition 3.15). For every $\pi_l, \pi_r \in \widehat{SU(2)}$, we have

$$\Theta_a(\pi_l, \pi_r) = \bigoplus_{k=|l-r|}^{|l+r|} \frac{\omega_a(\pi_k)}{\omega_a(\pi_l)\omega_a(\pi_r)} 1_{B(H_{\pi_k})},$$

where $\pi_l \otimes \pi_r \cong \bigoplus_{k=|l-r|}^{|l+r|} \pi_k$ is the irreducible decomposition of $\pi_l \otimes \pi_r$ (see the tensor formula (4.2)). Since $d_{\pi_k} \leq d_{\pi_l} + d_{\pi_r}$, it follows that

$$\omega_a(\pi_k) \leq \omega_a(\pi_l) + \omega_a(\pi_r).$$

Thus

$$\|\Theta_a(\pi_l, \pi_r)\| \leq \frac{1}{(1+2l)^a} + \frac{1}{(1+2r)^a}.$$

Hence it follows that

$$\lim_{\pi_l \rightarrow \infty} \limsup_{\pi_r \rightarrow \infty} \|\Theta_a(\pi_l, \pi_r)\| = \lim_{\pi_r \rightarrow \infty} \limsup_{\pi_l \rightarrow \infty} \|\Theta_a(\pi_l, \pi_r)\| = 0.$$

Therefore $A(G, \omega_a)$ is Arens regular by Theorem 3.16. The proof of the Arens regularity of $A(SU(2), \rho_b)$ when $b > 0$ is similar to the preceding case. \square

4.5. Arens regular subalgebras of Fourier algebras. It is shown in [21] that there are closed ideals in $L^1(\mathbb{T}^n) \cong A(\mathbb{Z}^n)$ ($n \in \mathbb{N}$) that are Arens regular. Since these ideals are non-unital, by [40], they can not have bounded approximate identity.

In this section, we show that we can construct unital, infinite-dimensional Arens regular closed subalgebras of Fourier algebras on certain products of $SU(2)$. This goes parallel to the main result of [40] since these subalgebras are not ideals. In fact, they are of the form of Beurling-Fourier algebras on $SU(2)$. This is a surprising and at the same time an interesting result since classical Beurling algebras can never be a closed subalgebra of a group algebra unless the weight is trivial. However, the relation (3.3) shows that this can happen for Beurling-Fourier algebras on certain non-abelian groups. More precisely, it is shown in [32] that for a locally compact group G , $\sup\{d_\pi \mid \pi \in \widehat{G}\}$ is finite if and only if G is almost abelian i.e. G has an abelian subgroup of finite index. Thus if G is not almost abelian, then $A_{\gamma^n}(G)$ defined in (3.2) is a Beurling-Fourier algebra with growing weight (see also (2.15) and (3.3)). We will see in the following theorem that this algebras can be Arens regular.

Theorem 4.4. *Let $n \in \mathbb{N}$, and let $G_n = SU(2) \times \cdots \times SU(2)$, 2^n -times. Then $A_{\gamma^n}(SU(2))$ is a unital, infinite-dimensional Arens regular closed subalgebra of the Fourier algebra $A(G_n)$.*

Proof. Consider the central weight

$$\omega_{2^n}(\pi_l) = d_{\pi_l}^{2^n} \quad (\pi_l \in \widehat{SU(2)}).$$

Then, by (2.15), (3.2), and (3.3), $A_{\gamma^n}(SU(2)) = A(SU(2), \omega_{2^n})$ is a unital, infinite-dimensional closed subalgebra of $A(G_n)$ and by Theorem 4.3, it is Arens regular. \square

REFERENCES

- [1] A. Beurling, *Sur les intégrales de Fourier absolument convergentes et leur application à une transformation fonctionnelle*, 9ème congrès des mathématiciens scandinaves, août 1938, Helsinki, 1939, pp. 345-366
- [2] A. Beurling, *Sur les spectres des fonctions*. Colloques internationaux du C.N.R.S., Analyse harmonique, Nancy, 1947, pp. 9-29
- [3] M. J. Collons, *Representations and characters of finite groups*, Cambridge University Press, 1990.
- [4] J. B. Conway, *A course in functional analysis*, Second edition. Graduate Texts in Mathematics, 96. Springer-Verlag, New York, 1990.
- [5] P. C. Curtis, R. J. Loy, *The structure of amenable Banach algebras*, J. London Math. Soc. (2) **40** (1989), no. 1, 89–104.
- [6] H. G. Dales, *Banach algebras and automatic continuity*, Oxford University Press, New York, 2000.
- [7] H. G. Dales and A. T.-M. Lau, *The Second Duals of Beurling Algebras*, Mem. Amer. Math. Soc, **177** n. 836 (2005).
- [8] Y. Domar, *Harmonic analysis based on certain commutative Banach algebras*, Acta Math. **96**, (1956) 166.
- [9] E. G. Effros and Z.-J. Ruan, *Operator spaces*, London Math. Soc. Monographs, New series, vol. 23, Oxford University Press, New York, 2000.
- [10] M. Enock and J.-M. Schwartz, *Kac algebras and duality of locally compact groups*. Springer-Verlag, Berlin, 1992.
- [11] P. Eymard, *L'algèbre de Fourier d'un groupe localement compact*, Bull. Soc. Math. France **92** (1964), 181–236.
- [12] G. Fendler, K. Grchenig, M. Leinert, J. Ludwig, C. Molitor-Braun, *Weighted group algebras on groups of polynomial growth*, Math. Z. **245** (2003), no. 4, 791–821.
- [13] G. B. Folland, *A course in abstract harmonic analysis*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1995.
- [14] B. E. Forrest, *Arens regularity and discrete groups*, Pacific J. Math. **151** (1991), no. 2, 217-227.
- [15] B. E. Forrest, *Arens regularity and the $A_p(G)$ algebras*, Proc. Amer. Math. Soc. **119** (1993), no. 2, 595-598.
- [16] B. E. Forrest and V. Runde, *Amenability and weak amenability of the Fourier algebra*, Math. Z. **250** (2005), 731–744.
- [17] B. E. Forrest, E. Samei and N. Spronk, *Convolutions on compact groups and Fourier algebras of coset spaces*, Studia Math, to appear, arXiv:0705.4277.
- [18] B. E. Forrest, E. Samei and N. Spronk, *Weak amenability of Fourier algebras on compact groups*, Indiana Math, J. to appear, arXiv:0808.1858.
- [19] B. E. Forrest and P. J. Wood, *Cohomology and the operator space structure of the Fourier algebra and its second dual*, Indiana Math. J. **50** (2001), 1217-1240.

- [20] M. Ghandehari, H. H. Lee and E. Samei, *Beurling-Fourier algebras of Heisenberg groups and $n \times n$ special unitary groups*, preprint.
- [21] Colin C. Graham, *Arens regularity of some familiar algebras*, preprint.
- [22] N. Grønbaek, *A characterization of weakly amenable Banach algebras*, *Studia Math.* **94** (1989), no. 2, 149-162.
- [23] N. Grønbaek, *Amenability of Weighted convolution Algebras on Locally Compact Groups*, *Trans. Amer. Math. Soc.* **319** (1990), no. 2, 765-775.
- [24] W. Hauenschild, E. Kaniuth, A. Kumar, *Ideal structure of Beurling algebras on $[FC]^+$ groups*, *J. Funct. Anal.* **51** (1983), no. 2, 213-228.
- [25] E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis*, II. Springer Verlage, 1963.
- [26] B. E. Johnson, *Cohomology in Banach algebras*, volume 127 of *Memoirs Amer. Math. Soc. American Math. Soc.*, 1972.
- [27] B.E. Johnson, *Approximate diagonals and cohomology of certain annihilator algebras*, *Amer. J. Math.* **94**, (1972) 685-698.
- [28] B. E. Johnson, *Non-amenability of the Fourier algebra of a compact group*, *J. London Math. Soc. (2)* **50** (1994) 361-374.
- [29] R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras*. Vol. I. Elementary theory. Pure and Applied Mathematics, 100. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1983.
- [30] J. Kustermans and S. Vaes, *Locally compact quantum groups in the von Neumann algebraic setting*. *Math. Scand.* 92 (2003), no. 1, 68–92.
- [31] J. Ludwig, N. Spronk, L. Turowska, *Beurling-Fourier algebras of compact groups*, preprint.
- [32] C.C. Moore, *Groups with finite dimensional irreducible representations*, *Trans. Amer. Math. Soc.* **166**, (1972) 401-410.
- [33] B. de Pagter and F. A. Sukochev, *Differentiation of operator functions in non-commutative L_p -spaces*. *J. Funct. Anal.* **212** (2004), no. 1, 28-75.
- [34] D. Rider. *Central idempotent measures on compact groups*. *Trans. Amer. Math. Soc.*, 186:459–479, 1973.
- [35] H. Reiter and J. D. Stegeman, *Classical Harmonic Analysis and Locally compact Groups*, Oxford University Press, 2000.
- [36] Zhong-Jin Ruan, *The operator amenability of $A(G)$* , *Amer. J. Math.* **117** (1995), no. 6, 1449–1474.
- [37] E. Samei, *Weak amenability and 2-weak amenability of Beurling algebras*, *J. Math. Anal. Appl.* **346** (2008), no. 2, 451-467.
- [38] N. Spronk, *Operator weak amenability of the Fourier algebra*, *Proc. Amer. Math. Soc.* **130** (2002), 3609-3617.
- [39] S. Thangavelu, *Harmonic analysis on the Heisenberg group*. Progress in Mathematics, 159. Birkhauser Boston, Inc., Boston, MA, 1998.
- [40] A. Ülger, *Arens regularity of weakly sequentially complete Banach algebras*, *Proc. Amer. Math. Soc.* **127** (1999), no. 11, 3221-3227.
- [41] E.I. Zelmanov, *On periodic compact groups*, *Israel J. Math.* **77** (1992), pp. 83-95.

DEPARTMENT OF MATHEMATICS, CHUNGBUK NATIONAL UNIVERSITY, 410 SUNGBONG-RO, HEUNGDUK-GU, CHEONGJU 361-763, KOREA

E-mail address: `hhlee@chungbuk.ac.kr`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, SASKATOON, SASKATCHEWAN, S7N 5E6, CANADA

E-mail address: `samei@math.usask.ca`