

**SOME IDENTITIES OF BERNOULLI NUMBERS AND
POLYNOMIALS ASSOCIATED WITH BERNSTEIN
POLYNOMIALS**

MIN-SOO KIM, TAEKYUN KIM, BYUNGJE LEE AND CHEON-SEOUNG RYOO

ABSTRACT. We investigate some interesting properties of the Bernstein polynomials related to the bosonic p -adic integrals on \mathbb{Z}_p .

1. INTRODUCTION

Let $C[0, 1]$ be the set of continuous functions on $[0, 1]$. Then the classical Bernstein polynomials of degree n for $f \in C[0, 1]$ are defined by

$$(1.1) \quad \mathbb{B}_n(f) = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x), \quad 0 \leq x \leq 1$$

where $\mathbb{B}_n(f)$ is called the Bernstein operator and

$$(1.2) \quad B_{k,n}(x) = \binom{n}{k} x^k (x-1)^{n-k}$$

are called the Bernstein basis polynomials (or the Bernstein polynomials of degree n) (see [10]). Recently, Acikgoz and Araci have studied the generating function for Bernstein polynomials (see [1, 2]). Their generating function for $B_{k,n}(x)$ is given by

$$(1.3) \quad F^{(k)}(t, x) = \frac{t^k e^{(1-x)t} x^k}{k!} = \sum_{n=0}^{\infty} B_{k,n}(x) \frac{t^n}{n!},$$

where $k = 0, 1, \dots$ and $x \in [0, 1]$. Note that

$$B_{k,n}(x) = \begin{cases} \binom{n}{k} x^k (1-x)^{n-k} & \text{if } n \geq k \\ 0, & \text{if } n < k \end{cases}$$

for $n = 0, 1, \dots$ (see [1, 2]).

The Bernstein polynomials can also be defined in many different ways. Thus, recently, many applications of these polynomials have been looked for by many authors. Some researchers have studied the Bernstein polynomials in the area of approximation theory (see [1, 2, 3, 7, 9, 10]). In recent years, Acikgoz and Araci [1, 2] have introduced several type Bernstein polynomials.

In the present paper, we introduce the Bernstein polynomials on the ring of p -adic integers \mathbb{Z}_p . We also investigate some interesting properties of the Bernstein polynomials related to the bosonic p -adic integrals on the ring of p -adic integers \mathbb{Z}_p .

2000 *Mathematics Subject Classification.* 11B68, 11S80.

Key words and phrases. Bernstein polynomials, Bosonic p -adic integrals.

2. BERNSTEIN POLYNOMIALS
RELATED TO THE BOSONIC p -ADIC INTEGRALS ON \mathbb{Z}_p

Let p be a fixed prime number. Throughout this paper, $\mathbb{Z}_p, \mathbb{Q}_p$ and \mathbb{C}_p will denote the ring of p -adic integers, the field of p -adic numbers and the completion of the algebraic closure of \mathbb{Q}_p , respectively. Let v_p be the normalized exponential valuation of \mathbb{C}_p with $|p|_p = p^{-1}$. For $N \geq 1$, the bosonic distribution μ_1 on \mathbb{Z}_p

$$(2.1) \quad \mu(a + p^N \mathbb{Z}_p) = \frac{1}{p^N}$$

is known as the p -adic Haar distribution μ_{Haar} , where $a + p^N \mathbb{Z}_p = \{x \in \mathbb{Q}_p \mid |x - a|_p \leq p^{-N}\}$ (cf. [4]). We shall write $d\mu_1(x)$ to remind ourselves that x is the variable of integration. Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable function on \mathbb{Z}_p . Then μ_1 yields the fermionic p -adic q -integral of a function $f \in UD(\mathbb{Z}_p)$:

$$(2.2) \quad I_1(f) = \int_{\mathbb{Z}_p} f(x) d\mu_1(x) = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x)$$

(cf. [4]). Many interesting properties of (2.2) were studied by many authors (cf. [4, 8] and the references given there). For $n \in \mathbb{N}$, write $f_n(x) = f(x + n)$. We have

$$(2.3) \quad I_1(f_n) = I_1(f) + \sum_{l=0}^{n-1} f'(l).$$

This identity is to derives interesting relationships involving Bernoulli numbers and polynomials. Indeed, we note that

$$(2.4) \quad I_1((x + y)^n) = \int_{\mathbb{Z}_p} (x + y)^n d\mu_1(y) = B_n(x),$$

where $B_n(x)$ are the Bernoulli polynomials (cf. [4]). From (1.2), we have

$$(2.5) \quad \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu_1(x) = \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^{n-k-j} B_{n-j}$$

and

$$(2.6) \quad \begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu_1(x) &= \int_{\mathbb{Z}_p} B_{n-k,n}(1-x) d\mu_1(x) \\ &= \binom{n}{k} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} \sum_{l=0}^{n-j} \binom{n-j}{l} (-1)^l B_l. \end{aligned}$$

By (2.5) and (2.6), we obtain the following proposition.

Proposition 2.1. *For $n \geq k$,*

$$\sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^{n-k-j} B_{n-j} = \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} \sum_{l=0}^{n-j} \binom{n-j}{l} (-1)^l B_l.$$

From (2.4), we note that

$$(2.7) \quad B_n(2) = (B(1) + 1)^n - n = (B + 1)^n = B_n, \quad n > 1$$

with the usual convention of replacing B^n by B_n . Thus, we have

$$\begin{aligned}
\int_{\mathbb{Z}_p} x^n d\mu_1(x) &= \int_{\mathbb{Z}_p} (x+2)^n d\mu_1(x) - n \\
(2.8) \qquad \qquad \qquad &= (-1)^n \int_{\mathbb{Z}_p} (x-1)^n d\mu_1(x) - n \\
&= \int_{\mathbb{Z}_p} (1-x)^n d\mu_1(x) - n
\end{aligned}$$

for $n > 1$, since $(-1)^n B_n(x) = B_n(1-x)$. Therefore we obtain the following theorem.

Theorem 2.2. *For $n > 1$,*

$$\int_{\mathbb{Z}_p} (1-x)^n d\mu_1(x) = \int_{\mathbb{Z}_p} x^n d\mu_1(x) + n.$$

And also we obtain

$$\begin{aligned}
\int_{\mathbb{Z}_p} B_{n-k,k}(x) d\mu_1(x) &= \int_{\mathbb{Z}_p} x^{n-k} (1-x)^k d\mu_1(x) \\
(2.9) \qquad \qquad \qquad &= \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \int_{\mathbb{Z}_p} (1-x)^{l+k} d\mu_1(x) \\
&= \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \left\{ \int_{\mathbb{Z}_p} x^{l+k} d\mu_1(x) + l + k \right\} \\
&= \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l (B_{l+k} + l + k).
\end{aligned}$$

Therefore we obtain the following result.

Corollary 2.3. *For $k > 1$,*

$$\int_{\mathbb{Z}_p} B_{n-k,k}(x) d\mu_1(x) = \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l (B_{l+k} + l + k).$$

From the property of the Bernstein polynomials of degree n , we easily see that

$$\begin{aligned}
&\int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) d\mu_1(x) \\
(2.10) \qquad \qquad \qquad &= \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} x^{2k} (1-x)^{n+m-2k} d\mu_1(x) \\
&= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l B_{2k+l}
\end{aligned}$$

and

$$\begin{aligned}
(2.11) \quad & \int_{\mathbb{Z}_p} B_{k,n}(x)B_{k,m}(x)B_{k,s}(x)d\mu_1(x) \\
&= \binom{n}{k} \binom{m}{k} \binom{s}{k} \int_{\mathbb{Z}_p} x^{3k}(1-x)^{n+m-3k}d\mu_1(x) \\
&= \binom{n}{k} \binom{m}{k} \binom{s}{k} \sum_{l=0}^{n+m+s-3k} \binom{n+m+s-3k}{l} (-1)^l B_{3k+l}.
\end{aligned}$$

Continuing this process, we obtain the following theorem.

Theorem 2.4. *The multiplication of the sequence of Bernstein polynomials*

$$B_{k,n_1}(x), B_{k,n_2}(x), \dots, B_{k,n_s}(x)$$

for $s \in \mathbb{N}$ with different degree under p -adic integral on \mathbb{Z}_p can be given as

$$\begin{aligned}
& \int_{\mathbb{Z}_p} B_{k,n_1}(x)B_{k,n_2}(x) \cdots B_{k,n_s}(x)d\mu_1(x) \\
&= \binom{n_1}{k} \binom{n_2}{k} \cdots \binom{n_s}{k} \sum_{l=0}^{n_1+n_2+\cdots+n_s-sk} \binom{n_1+n_2+\cdots+n_s-sk}{l} (-1)^l B_{sk+l}.
\end{aligned}$$

We put

$$B_{k,n}^m(x) = \underbrace{B_{k,n}(x) \times \cdots \times B_{k,n}(x)}_{m\text{-times}}.$$

Theorem 2.5. *The multiplication of*

$$B_{k,n_1}^{m_1}(x), B_{k,n_2}^{m_2}(x), \dots, B_{k,n_s}^{m_s}(x)$$

Bernstein polynomials with different degrees n_1, n_2, \dots, n_s under p -adic integral on \mathbb{Z}_p can be given as

$$\begin{aligned}
& \int_{\mathbb{Z}_p} B_{k,n_1}^{m_1}(x)B_{k,n_2}^{m_2}(x) \cdots B_{k,n_s}^{m_s}(x)d\mu_1(x) \\
&= \binom{n_1}{k}^{m_1} \binom{n_2}{k}^{m_2} \cdots \binom{n_s}{k}^{m_s} \sum_{l=0}^{n_1m_1+n_2m_2+\cdots+n_sm_s-(m_1+\cdots+m_s)k} (-1)^l \\
&\quad \times \binom{n_1m_1+n_2m_2+\cdots+n_sm_s-(m_1+\cdots+m_s)k}{l} B_{(m_1+\cdots+m_s)k+l}.
\end{aligned}$$

Theorem 2.6. *The multiplication of*

$$B_{k_1,n_1}^{m_1}(x), B_{k_2,n_2}^{m_2}(x), \dots, B_{k_s,n_s}^{m_s}(x)$$

Bernstein polynomials with different degrees n_1, n_2, \dots, n_s with different powers m_1, m_2, \dots, m_s under p -adic integral on \mathbb{Z}_p can be given as

$$\begin{aligned}
& \int_{\mathbb{Z}_p} B_{k_1,n_1}^{m_1}(x)B_{k_2,n_2}^{m_2}(x) \cdots B_{k_s,n_s}^{m_s}(x)d\mu_1(x) \\
&= \binom{n_1}{k_1}^{m_1} \binom{n_2}{k_2}^{m_2} \cdots \binom{n_s}{k_s}^{m_s} \sum_{l=0}^{n_1m_1+n_2m_2+\cdots+n_sm_s-(k_1m_1+\cdots+k_sm_s)} (-1)^l \\
&\quad \times \binom{n_1m_1+n_2m_2+\cdots+n_sm_s-(k_1m_1+\cdots+k_sm_s)}{l} B_{k_1m_1+\cdots+k_sm_s+l}.
\end{aligned}$$

Problem. Find the Witt's formula for the Bernstein polynomials in p -adic number field.

REFERENCES

- [1] M. Acikgoz and S. Araci, *A study on the integral of the product of several type Bernstein polynomials*, IST Transaction of Applied Mathematics-Modelling and Simulation, 2010.
- [2] M. Acikgoz and S. Araci, *On the generating function of the Bernstein polynomials*, Accepted to AIP on 24 March 2010 for ICNAAM 2010.
- [3] S. Bernstein, *Demonstration du theoreme de Weierstrass, fondee sur le calcul des probabilites*, Commun. Soc. Math. Kharkow (2) **13** (1912-1913), 1–2.
- [4] T. Kim, *On a q -analogue of the p -adic log gamma functions and related integrals*, J. Number Theory **76** (1999), 320–329.
- [5] T. Kim, *q -Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288–299.
- [6] T. Kim, *Barnes-type multiple q -zeta functions and q -Euler polynomials*, J. Physics A : Math. Theor., **43** (2010), 255201, 11pp.
- [7] T. Kim, L.-C. Jang and H. Yi, *Note on the modified q -Bernstein polynomials*, Discrete Dynamics in Nature and Society (in Press). arXiv 1005.4293.
- [8] T. Kim, J. Choi and Y.-H. Kim *Some identities on the q -Bernstein polynomials, q -Stirling numbers and q -Bernoulli numbers*, Adv. Stud. Contemp. Math. 20 (2010), no. 3, 335–341. arXiv 1006.2033.
- [9] G. M. Phillips, *Bernstein polynomials based on the q -integer*, Annals of Numerical Analysis **4** (1997), 511–518.
- [10] Y. Simsek and M. Acikgoz, *A new generating function of q -Bernstein-type polynomials and their interpolation function*, Abstr. Appl. Anal. ID 769095 (2010), 12 pp.

DEPARTMENT OF MATHEMATICS, KAIST, 373-1 GUSEONG-DONG, YUSEONG-GU, DAEJEON 305-701, SOUTH KOREA

E-mail address: minsookim@kaist.ac.kr

DIVISION OF GENERAL EDUCATION-MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL, 139-701, SOUTH KOREA

E-mail address: tkkim@kw.ac.kr

DEPARTMENT OF WIRELESS COMMUNICATIONS ENGINEERING, KWANGWOON UNIVERSITY, SEOUL 139-701, SOUTH KOREA

E-mail address: bj_lee@kw.ac.kr

DEPARTMENT OF MATHEMATICS, HANNAM UNIVERSITY, DAEJEON 306-791, SOUTH KOREA

E-mail address: ryooocs@hnu.kr