Constructing graphs over \mathbb{R}^n with small prescribed mean-curvature

HOLLY CARLEY City University of New York

AND

MICHAEL K.-H. KIESSLING Rutgers University

Printed: October 23, 2018

Abstract

In this paper nonlinear Hodge theory and Banach algebra estimates are employed to construct a convergent series expansion which solves the prescribed mean curvature equation $\pm \nabla \cdot (\nabla u/\sqrt{1 \pm |\nabla u|^2}) = nH$ for *n*-dimensional hypersurfaces in \mathbb{R}^{n+1} (+ sign) and $\mathbb{R}^{1,n}$ (- sign) which are graphs $\{(x, u(x)) : x \in \mathbb{R}^n\}$ of a smooth function $u : \mathbb{R}^n \to \mathbb{R}$, and whose mean curvature function H is α -Hölder continuous and integrable, with small norm. The radius of convergence is estimated explicitly from below. Our approach is inspired by, and applied to, the Maxwell–Born– Infeld theory of electromagnetism in $\mathbb{R}^{1,3}$, for which our method yields the first systematic way of explicitly computing the electrostatic potential $\phi \propto u$ for regular charge densities $\rho \propto H$ and small Born parameter, with explicit error estimates at any order of truncation of the series. In particular, our results level the ground for a controlled computation of Born–Infeld effects on the Hydrogen spectrum.

Revised version: Jan. 24, 2015; Original: Aug. 30, 2010

©2015 The authors. Reproduction of this preprint, for non-commercial purposes, is permitted.

1 Introduction

In several areas of mathematical physics one encounters the problem of ndimensional submanifolds with prescribed mean-curvature function H in an n+1 dimensional Riemannian or Lorentzian manifold. For example, the formulation of the Cauchy problem for Einstein's equations in general relativity requires a foliation of the spacetime, and a popular choice are spacelike so-called constant mean-curvature (CMC) foliations [1, 3, 27, 58, 44, 26]; in this case the mean curvature function is particularly simple, but entire CMC submanifolds do not always exist and one possible way out is to allow the mean curvature to vary in a suitable manner [27, 43]. Another, less obvious example comes from electromagnetism; more precisely, as will be explained in more detail in the next section, the graph of the electrical potential for the electrostatic solutions of the nonlinear Maxwell–Born–Infeld equations in $\mathbb{R}^{1,3}$ with prescribed charge density [8, 9, 49, 36, 16, 38, 28, 61] can be re-interpreted as an (almost) entire spacelike hypersurface in Minkowski space [39, 40, 3, 28, 38], with the charge density function playing the role of the mean curvature function. Of course, submanifolds with prescribed mean-curvature are also of interest in their own right to differential geometers.

Although the (non-)existence and regularity theory for solutions of the prescribed mean-curvature equation has evolved quite far (e.g. [30]), the nonlinearity of the equation stands in the way of writing down an explicit solution formula except in special circumstances. Thus, the problem of constructing an *n*-dimensional hypersurface with explicitly prescribed mean-curvature H in an n + 1 dimensional Euclidean or Minkowskian space has been solved in great generality only when n = 1 (by simple ODE techniques, obviously), although powerful complex variable techniques can be put to work when n = 2 (especially when $H \equiv 0$); see [20, 33, 49, 35, 41, 2, 56, 45, 21, 47, 48, 23]. However, explicit constructive results when $n \geq 3$ are typically only concerned with radially symmetric solutions (see, e.g., [4] for a more recent contribution).

In this paper we are concerned with the explicit construction of solutions to the so-called non-parametric prescribed mean-curvature problem where the hypersurfaces are graphs of some scalar function u over the entire n-dimensional base manifold when no symmetry assumption on the prescribed mean-curvature function or its solution is made. Yet to find non-special hypersurfaces without symmetry assumptions we have to settle for the neighborhood of some special solution. It is well-known that in principle one can systematically find approximate solutions by using variational methods, or by running the iterations of the fixed point map of the Schauder theory, or by directly implementing the proof of the inverse function theorem. Curiously enough, though, our literature search for *explicit formulas* of the approximate solutions at arbitrary order, with explicit estimates of the domain of convergence of such an approximation scheme, revealed nothing. We believe that ours is the first paper in which such a convergent scheme, of explicit quadratures, is presented for quite general small and regular mean curvature function; we emphasize once again that no symmetry assumptions are made.

Specifically, we will explicitly construct a certain neighborhood of the minimal / maximal hypersurfaces $(H \equiv 0)$ which consists of hypersurfaces with small and integrable mean curvature functions H which are graphs over \mathbb{R}^n , $n \geq 1$. In dimension n = 1, our "small-curvature" expansion method actually terminates already after the first term and therefore yields the solution to the problem by the usual explicit quadrature without smallness assumption. Thus our method is primarily of interest in dimensions $n \geq 2$, though for n = 2 it is perhaps only of interest as an alternative to the complex variable techniques.

Our method is motivated by a concrete problem from physics (n = 3), more precisely the problem of solving the first-order PDE system of Maxwell-Born-Infeld electrostatics. We already mentioned above that the graph of the electrostatic potential is a prescribed mean-curvature hypersurface in Minkowski space, satisfying the pertinent scalar second-order PDE problem in n = 3. Therefore in the next section we use this electrostatic analog to motivate and explain our approach, using the physicists' familiar three-dimensional vector notation for the fields. Then, in section 3, after recalling the scalar prescribed mean-curvature problem for graphs over \mathbb{R}^n in all dimensions $n \geq 1$, in both Euclidean and Minkowskian geometry (our approach treats both on an equal footing), we use nonlinear Hodge theory to give two dual reformulations of the scalar secondorder PDE problem as a first-order system for two one-forms which are the *n*-dimensional analogs of the electromagnetic vector equations in n = 3. In section 4 we state our main results in full generality: we present the solutions to the first-order Hodge systems, and by corollary to the scalar PDE, as a convergent series whose terms are given by explicit quadratures. The parameter of expansion can be seen as a measure of the smallness of H in a suitable Hölder norm. In section 5 we prove the convergence in a suitable Banach algebra for each dimension $n \in \mathbb{N}$, with n > 1, including an explicit lower estimate of the radius of convergence, obtained with the help of complex analysis; of course we also comment on the n = 1-dimensional case. In section 6 we present an algebraically simplified formulation of the three-dimensional problems which raises some interesting PDE questions. We conclude in section 7 by indicating some generalizations of our approach to related quasi-linear PDE problems in divergence form.

2 The electrostatic analog

In Maxwell's classical electromagnetic theory, Coulomb's law states that an electrostatic charge density ρ in \mathbb{R}^3 is the source for the so-called electric displacement density field D,

$$\nabla \cdot D = 4\pi\rho \qquad \text{Coulomb's law},\tag{1}$$

while Faraday's law says that the so-called electric field strength E, when stationary, is curl-free:

$$\nabla \times E = 0$$
 Faraday's law (stationary). (2)

The two fields E and D need to be linked by an "aether law," with the help of which one can eliminate either D or E and obtain a closed, generally nonlinear set of first-order PDE for the remaining field.

Note that "aether" is used here merely as a shorthand for "electromagnetic vacuum;" historically Maxwell of course thought of "the classical aether."

Of particular relevance for the prescribed mean-curvature problem is the aether law proposed in the 1930s by Max Born [8], which reads

$$D = \frac{E}{\sqrt{1 - \beta^4 |E|^2}} \qquad \text{Born's law} \tag{3}$$

which can be inverted to yield

$$E = \frac{D}{\sqrt{1 + \beta^4 |D|^2}} \qquad \text{Born's law (reverse).}$$
(4)

Here, $\beta \in (0, \infty)$ is a hypothetical new constant of nature (in the dimensionless notation of [36]). In the limit $\beta \to 0$ Born's law goes over into Maxwell's law of the "pure aether", D = E.

Remark 1. : Born's aether law (3) is only the electrostatic special case of his electromagnetic aether law (replace $|E|^2$ by $|E|^2 - |B|^2$, and add a counterpart for the magnetic induction field B and field strength H), see [8, 9], and also App. VI in [10]. Born's electromagnetic aether law was subsequently generalized by Born and Infeld [11] to a much more interesting law; the stationary case was recently treated in [37]. Also higher-dimensional generalizations of Maxwell-Born-Infeld theory with geometric significance exist; see, for instance, [28, 29, 61].

The classically fundamental problem of electrostatic fields generated by point charges, corresponding to maximal hypersurfaces with conical singularities [22], is treated in [8, 9, 49, 39, 40, 36, 16, 38]. Unfortunately, only the case with a single point charge [9, 22], or with a regular lattice of infinitely many of them [34, 28], has been solved explicitly. Even the extremely important case of two point charges still awaits solution.

In this section we make progress by presenting a convergent series solution for electrostatic finite-energy fields E in all of \mathbb{R}^3 which are generated by α -Hölder continuous charge density functions ρ which are sufficiently small in norm. This includes in particular the case of several regularized point charges, i.e. spherically symmetric and compactly supported C^{α} functions which we will use in a follow-up work on the old — but still open(!) — mathematicalphysical problem of computing Born–Infeld effects on the quantum-mechanical spectrum of atoms, especially Hydrogen. This problem is important because the high precision with which the empirical atomic spectra are known will impose strict bounds on the size of the hypothetical "new constant of nature" which enters the Born–Infeld relations — thus it will provide a crucial physical viability test of the Born–Infeld proposal for this constant, and possibly on the physical viability of their model at that. Unfortunately, conflicting results have been obtained so far with different plausible approximations to the electrostatic pair energy, see [50, 55, 32, 16, 25]. The solution obtained in this section supplies a series of better and better approximations with controlled error for smeared out point charges, which is an important stepping stone along the way to eventually settle the problem of Born–Infeld effects on the quantum-mechanical spectra of atoms with point charges. Our solution may also pave the ground for computing classical effects in plasmas, cf. [14].

2.1 Hierarchical vector series solution for small $\beta^2 \rho$

To state our solution we first stipulate some notation. As usual, $C^{k,\alpha}$ denotes the k-times continuously differentiable functions whose k-th derivatives have Hölder exponent α ; we also write C^{α} for $C^{0,\alpha}$. As for the notation C_0 , we follow the conventions of [24]. Thus, C_0 denotes the continuous functions which decay to zero at infinity, not continuous functions with compact support as in [30]; the latter are denoted by C_c in [24], and in [42]. In the same vein, bounded continuous functions will be denoted by C_b — here we chose not to follow [24], where BC is used instead. Altogether, $C_b^{k,\alpha}$, respectively $C_0^{k,\alpha}$, will denote the k-times continuously differentiable functions which together with their derivatives up to order k are bounded, respectively decay to zero at infinity, and whose k-th derivatives are in C^{α} . We equip $C_b^{k,\alpha}$ with the usual norm

$$\|f\|_{k,\alpha} = \max_{\|\ell\| \le k} \sup_{x} |D^{\ell}f(x)| + \max_{\|\ell\| = k} \sup_{x \ne y} \frac{|D^{\ell}f(x) - D^{\ell}f(y)|}{|x - y|^{\alpha}}$$
(5)

where ℓ is a multi-index. Clearly, $C_0^{k,\alpha} \subset C_b^{k,\alpha}$ is a closed subspace, i.e. if $f_n \to f$ in $\|\cdot\|_{k,\alpha}$ norm, and $f_n \in C_0^{k,\alpha}$, then $f \in C_0^{k,\alpha}$, too. For $\beta^2 \rho \in C_0^{\alpha} \cap L^1$ small in the sense stated below we now present a conver-

For $\beta^2 \rho \in C_0^{\alpha} \cap L^1$ small in the sense stated below we now present a convergent β -power series expansion of D solving the system of Maxwell–Born–Infeld equations (1), (2), (4) with vanishing condition at spatial infinity. By virtue of (4) this translates into power series solutions for E and its electric potential, ϕ , defined as $E = -\nabla \phi$ with asymptotic vanishing conditions.

Proposition 1. Let $\beta^2 \rho \in C_0^{\alpha} \cap L^1$ be small in the sense that

$$v^{(1)}(x) := -\nabla \int \frac{\beta^2 \rho(y)}{|x - y|} \, \mathrm{d}^3 y \tag{6}$$

has small $C_0^{1,\alpha}$ norm, viz.

$$\|v^{(1)}\|_{1,\alpha} < \left(2^{2/3} - 1\right)^{3/2}.$$
(7)

Then the system of PDE (1), (2), (4) with vanishing condition at spatial infinity has a solution $D = \beta^{-2} v \in C_0^{1,\alpha}$, with v given by the absolutely convergent series

$$v = \sum_{k=0}^{\infty} v^{(2k+1)},$$
(8)

with $v^{(1)}$ given in (6) and $v^{(2k+1)}$ for $k \in \mathbb{N}$ recursively given by

$$v^{(2k+1)} = \mathbf{P}V^{(2k+1)}, \qquad k \in \mathbb{N},\tag{9}$$

where $V^{(2k+1)}$ is a polynomial in the $v^{(\ell)}$ with odd $\ell < 2k+1$, viz.

$$V^{(2k+1)} = -\sum_{h=1}^{k} v^{(2(k-h)+1)} \sum_{j=1}^{h} M_{j}^{+} \sum_{|\ell|_{2j}=h-j} \prod_{i=1}^{j} v^{(2\ell_{2i-1}+1)} \cdot v^{(2\ell_{2i}+1)}, \quad (10)$$

with $|\ell|_K = \sum_{i=1}^K \ell_i$, the ℓ_i taking any non-negative integer values,

$$M_j^+ = (-1)^j \frac{(2j-1)!!}{j! 2^j} \tag{11}$$

being the j-th Maclaurin coefficient of $1/\sqrt{1+z}$ (with $M_0^+ := 1$), and where $\mathbf{P}: C_0^{1,\alpha} \to C_0^{1,\alpha}$ projects onto the solenoidal subspace of $C_0^{1,\alpha}$ -valued vector fields; in particular, $\nabla \cdot V^{(2k+1)} \in C_0^{0,\alpha} \cap L^1$, and so

$$\mathbf{P}V^{(2k+1)}(x) = V^{(2k+1)}(x) + \nabla \int \frac{\frac{1}{4\pi} \nabla \cdot V^{(2k+1)}(y)}{|x-y|} \, \mathrm{d}^3 y.$$
(12)

Remark 2. Note that the $v^{(2k+1)}$, k = 0, 1, 2, ..., depend on β in a simple manner, namely each such $v^{(2k+1)} = \beta^{2(2k+1)}D^{(k)}$, where $D^{(k)}$ is β -independent. Thus, conceptually our solution is nothing but a physicists' formal perturbative series in powers of β , treated as a formal small parameter, for which we prove convergence. Incidentally, the index (2k + 1) rather than k at the vs will be explained using perturbation theory, in section 5.

Having the solution $D = \beta^{-2}v \in C_0^{1,\alpha}$ of the PDE system (1), (2), (4) with vanishing condition at spatial infinity, we now easily obtain E and ϕ .

Corollary 1. Let $D = \beta^{-2}v \in C_0^{1,\alpha}$ with v given by (8) solve the PDE system (1), (2), (4) with vanishing condition at spatial infinity. Then $E = \beta^{-2}w \in C_0^{1,\alpha}$ with w given by

$$w = \frac{v}{\sqrt{1 + |v|^2}},$$
(13)

solves the PDE system (1), (2), (3) with vanishing condition at spatial infinity. Moreover, w has its own convergent series expansion, obtained by inserting

the series solution (8) for v into the RHS of (13) and expanding, thus

$$w(x) = \sum_{k=0}^{\infty} w^{(2k+1)}(x),$$
(14)

with

$$w^{(1)}(x) = v^{(1)}(x), (15)$$

while for $k \in \mathbb{N}$ each $w^{(2k+1)}$ is given by

$$w^{(2k+1)}(x) = \nabla \int \frac{\frac{1}{4\pi} \nabla \cdot V^{(2k+1)}(y)}{|x-y|} \, \mathrm{d}^3 y, \tag{16}$$

with $V^{(2k+1)}(x)$ given by (10) in terms of the $v^{(\ell)}$ with $\ell < 2k+1$.

Finally, having the solution $E = \beta^{-2}w$ in form of a sum of gradient fields $w^{(2k+1)}$, $k = 0, 1, 2, ..., which converges in <math>C_0^{1,\alpha}$, we can invoke $E = -\nabla \phi$ and directly read off the electric potential $\phi = \beta^{-2}u \in C_0^{2,\alpha}$, with u given as the series

$$u(x) = \int \frac{\rho(y)}{|x-y|} \, \mathrm{d}^3 y - \frac{1}{4\pi} \sum_{k \in \mathbb{N}} \int \frac{\nabla \cdot V^{(2k+1)}(y)}{|x-y|} \, \mathrm{d}^3 y, \tag{17}$$

where $V^{(2k+1)}(x)$ is given by (10) in terms of the $v^{(\ell)}$ with $\ell < 2k+1$.

Remark 3. Note that each term in the expansion of $E = \beta^{-2}w$ is a gradient field, as it should by (2).

Remark 4. The proofs of Proposition 1 and Corollary 1 are given in section 5.

Remark 5. The expansion technique given here generalizes to electromagnetostatic solutions of the Maxwell–Born–Infeld equations with regular sources, see [37]; however, the generalization comes at the price of roughly a factor 1/2smaller radius of convergence estimate.

2.2 The Helmholtz decomposition

As shown by Helmholtz, every vector field in \mathbb{R}^3 can be decomposed into a sum of a divergence-free and a curl-free field. The decomposition is not unique because a harmonic field can be transferred from one summand to the other. This non-uniqueness can be removed by imposing suitable boundary or asymptotic conditions on the individual summands; in our case asymptotic vanishing conditions. The Helmholtz decomposition sheds an interesting light on the vexpansion. For this reason we include the following elementary observations.

Writing $v = v_g + v_c$, where v_g is a gradient field (hence, curl-free) and v_c is a curl of some vector field (hence, divergence-free), and registering that $\nabla \cdot v = \nabla \cdot v_g$ and $\nabla \times v = \nabla \times v_c$, we get

$$\nabla \cdot v_g = 4\pi\beta^2 \rho \tag{18}$$

(together with $\nabla \times v_g = 0$), and

$$\nabla \times \frac{(v_g + v_c)}{\sqrt{1 \mp |(v_g + v_c)|^2}} = 0$$
(19)

(together with $\nabla \cdot v_c = 0$). We thereby have obtained a closed set of first-order vector equations for the vector field v_q separately from v_c , plus a conditionally

closed set of first-order vector equations for the vector field v_c , conditioned on v_g being given. Both sets of PDE are supplemented by the asymptotic conditions that v_g and v_c vanish at spatial infinity; with this asymptotic condition the Helmholtz decomposition becomes unique.

As announced, the Helmholtz decomposition sheds an interesting light on the v expansion. Since the curl-free component of the Helmholtz decomposition is determined completely and independently of the divergence-free Helmholtz component v_c of v — namely thusly: For $H \in C_0^{\alpha} \cap L^1$ the linear equation (18) has a unique gradient field solution $\in C_0^{1,\alpha}$ given by

$$v_g(x) := -\nabla \int \frac{\beta^2 \rho(y)}{|x - y|} \, \mathrm{d}^3 y, \tag{20}$$

with integration over \mathbb{R}^3 —, we can reinterpret our power series approach as a method to solve the v_c -equation (19) given small v_g . Indeed, note that $v_g \equiv v^{(1)}$ given earlier in (6), so we can just separate the v_g part off from our power series solution for v. The remaining power series yields the solution v_c to (19) given small v_g . The solution v_c in turn has the meaning of a Born "correction" of the Coulomb field $v^{(1)}$ caused by Born's aether law.

Put in terms of D we summarize: Our convergent series expansion (8) yields a solution D given by the β -power series

$$D = \sum_{k=0}^{\infty} \beta^{4k} D^{(k)},$$
(21)

with $D^{(k)}$ independent of β , in which $D^{(0)} = \beta^{-2}v^{(1)}$ is the Coulomb field, while the $D^{(k)} = \beta^{-2(2k+1)}v^{(2k+1)}$ with $k \in \mathbb{N}$ are the coefficient fields of its $O(\beta^{4k})$ -corrections caused by Born's law.

2.3 Prescribed mean-curvature graphs over \mathbb{R}^3

Finally, we also write $w = w_g + w_c$, where w_g is a gradient field (hence, curlfree) and w_c is a curl of some vector field (hence, divergence-free). Clearly, $\nabla \cdot w = \nabla \cdot w_g$ and $\nabla \times w = \nabla \times w_c$, and therefore the equation

$$\nabla \times w = 0 \tag{22}$$

together with the vanishing asymptotics for w_c implies right away that $w_c \equiv 0$, hence $w = w_g = -\nabla u$. Now inverting (13) to yield $v = \beta^{-2}D$ in terms of $w = -\nabla u$, and inserting the expression in (1) yields a closed second-order scalar equation for u, viz.

$$-\nabla \cdot \frac{\nabla u}{\sqrt{1 - |\nabla u|^2}} = 4\pi\beta^2\rho.$$
⁽²³⁾

This is the well-known equation of a presribed mean-curvature hypersurface in Minkowski spacetime $\mathbb{R}^{1,3}$, with height function u and mean curvature function

 $H = \frac{4\pi}{3}\beta^2\rho$. Thus our series expansion for the electrostatic Maxwell–Born(–Infeld) ϕ also provides a solution u to the three-dimensional Minkowskian prescribed mean-curvature equation.

Replacing $\beta^4 \to -\beta^4$ in (3) yields a "pseudo Born's law," in which case the electrostatic Maxwell–Born(–Infeld) equations are equivalent to the presribed mean-curvature problem in Euclidean space \mathbb{R}^4 . This equation can therefore be treated with the same expansion strategy applied to an equivalent first order system of electrostatic Maxwell–pseudo-Born–Infeld equations.

In fact, our method generalizes to arbitrary dimensions. In the ensuing sections we formulate these problems, state our results, then present their proofs. The proofs of our Maxwell–Born–Infeld results are included as special cases.

3 Formulation of the non-parametric prescribed mean-curvature problems in \mathbb{R}^{n+1} and $\mathbb{R}^{1,n}$

In this section we consider all $n \in \mathbb{N}$ equally.

3.1 The scalar problem

We recall that the mean curvature function H of a hypersurface in \mathbb{R}^{n+1} (n = 1, 2, 3, ...), which is the graph $\{(x, u(x)) : x \in \mathbb{R}^n\}$ of a real-valued C^2 function u over \mathbb{R}^n , can be computed from u in a classical manner as

$$H = \frac{1}{n} \nabla \cdot \frac{\nabla u}{\sqrt{1 + \nabla u \cdot \nabla u}},\tag{24}$$

(see, e.g., [51], and the appendix to section 14 in [30]). By replacing the Euclidean inner product operation " \cdot " by " $-\cdot$ " one obtains the analogous result for space-like hypersurfaces in Minkowski space $\mathbb{R}^{1,n}$, viz.

$$H = -\frac{1}{n}\nabla \cdot \frac{\nabla u}{\sqrt{1 - \nabla u \cdot \nabla u}},$$
(25)

(see, e.g., [1, 3, 35, 41]). The inverse problem is to find a function u, satisfying suitable asymptotic conditions, whose graph over \mathbb{R}^n describes a hypersurface in \mathbb{R}^{n+1} , respectively $\mathbb{R}^{1,n}$, for which a prescribed function $H : \mathbb{R}^n \to \mathbb{R}$ is its mean curvature w.r.t. some normal. In that case (24) and (25) become second-order quasi-linear elliptic PDEs for the unknown u, which we lump together as

$$\pm \nabla \cdot \frac{\nabla u}{\sqrt{1 \pm |\nabla u|^2}} = nH,\tag{26}$$

the upper sign for the Euclidean, the lower for the Minkowskian setting. The different signs under the square roots are of course important, while the overall sign is just a matter of convention; it could be absorbed into H.

Equation (26) needs to be supplemented by an asymptotic condition on u compatible with the choice of H. We will only consider integrable mean curvature functions $H \in L^1$, which implies that u is asymptotic to some harmonic function, and we choose to impose on u asymptotic monopole conditions, i.e.

$$u(x) \asymp \mp \frac{n}{|S^{n-1}|} \int H(y) \,\mathrm{d}^n y \,|x|^{2-n} \tag{27}$$

when n > 2, and

$$u(x) \approx \pm \frac{1}{\pi} \int H(y) \,\mathrm{d}^2 y \,\ln|x| \tag{28}$$

when n = 2. However, when n = 1, then

$$u(x) \approx \pm \frac{\frac{1}{2} \int H(y) \, \mathrm{d}y}{\sqrt{1 \mp \left|\frac{1}{2} \int H(y) \, \mathrm{d}y\right|^2}} |x|;$$
(29)

clearly, the asymptotic condition (29) with the "-" sign under the square root requires the smallness condition

$$\left|\int H(y)\,\mathrm{d}y\right| < 2.\tag{30}$$

This is special to n = 1 and unrelated to the smallness condition on H which we need to impose later on to ensure the convergence of an infinite series.

We note that when n = 1 or 2, these asymptotic conditions still allow an arbitrary constant be added to any solution u — we will fix this irrelevant freedom when stating our main results. Other options are briefly commented on in the concluding section.

3.2 The equivalent first-order Hodge systems

For all dimensions $n \in \mathbb{N}$ the second-order prescribed mean-curvature equation (26) for either sign is equivalent to a nonlinear Hodge system of first order for some 1-form. There are two mutually dual formulations.

3.2.1 The ω system

Consider a 1-form ω satisfying

$$d\omega = 0; \tag{31}$$

$$\pm \delta \frac{\omega}{\sqrt{1 \pm |\omega|^2}} = nH.$$
(32)

Then this first-order system is equivalent to the second-order equation (26) by identifying $\omega \equiv \nabla u \cdot dx$. The asymptotic conditions on ω are inherited from (27), respectively (28) or (29); thus, when n > 2,

$$\omega(x) \asymp \mp \frac{n}{|S^{n-1}|} \int H(y) \,\mathrm{d}^n y \,\nabla |x|^{2-n} \cdot \,\mathrm{d}x \tag{33}$$

and when n = 2,

$$\omega(x) \approx \pm \frac{1}{\pi} \int H(y) \,\mathrm{d}^2 y \,\nabla \ln |x| \cdot \,\mathrm{d}x \tag{34}$$

while for when n = 1,

$$\omega(x) \approx \pm \frac{\frac{1}{2} \int H(y) \,\mathrm{d}y}{\sqrt{1 \mp \left|\frac{1}{2} \int H(y) \,\mathrm{d}y\right|^2}} \mathrm{sign}(x) \,\mathrm{d}x. \tag{35}$$

3.2.2 The τ system

Dual to the above system is the following. Defining a 1-form

$$\tau = \pm \frac{\omega}{\sqrt{1 \pm |\omega|^2}},\tag{36}$$

which can be inverted to yield

$$\omega = \pm \frac{\tau}{\sqrt{1 \mp |\tau|^2}},\tag{37}$$

we see that the above first-order equations are equivalent to

$$\delta \tau = nH, \qquad (38)$$

$$d\frac{\tau}{\sqrt{1\mp |\tau|^2}} = 0.$$
 (39)

Clearly, (39) implies that there is a scalar σ such that

$$\frac{\tau}{\sqrt{1\mp|\tau|^2}} = d\sigma. \tag{40}$$

Up to a sign and an additive constant, $\sigma = u$, of course.

The asymptotic conditions on τ , inherited from (27), respectively (28) or (29), are slightly simpler now; namely, when n = 1,

$$\tau(x) \approx \frac{1}{2} \int H(y) \, \mathrm{d}y \operatorname{sign}(x) \, \mathrm{d}x \tag{41}$$

and when n = 2,

$$\tau(x) \asymp \frac{2}{|S^1|} \int H(y) \,\mathrm{d}^2 y \,\nabla \ln |x| \cdot \,\mathrm{d}x,\tag{42}$$

whereas for n > 2,

$$\tau(x) \asymp -\frac{n}{|S^{n-1}|} \int H(y) \,\mathrm{d}^n y \,\nabla |x|^{2-n} \cdot \,\mathrm{d}x.$$
(43)

3.2.3 The cubic τ system

We end this subsection with the observation that upon differentiation in (39) and multiplication with $(1 \mp |\tau|^2)^{\frac{3}{2}}$ we find a cubic version of (39),

$$0 = \left(1 \mp |\tau|^2\right) d\tau \pm d \left[\frac{1}{2} |\tau|^2\right] \wedge \tau.$$
(44)

Together with (38), and supplemented with the same asymptotic conditions, this cubically nonlinear τ system is equivalent to the τ system in the previous subsubsection. Ironically, this algebraically simplest formulation of the problem is not at all more user-friendly. We shall come back to it in section 6.

4 Statement of the main results

Henceforth we will be primarily concerned with $n \ge 2$. As mentioned in the introduction, with some minor adjustments our series solution technique produces solutions also to the n = 1-dimensional problem, but the series terminates after the first term and is identical to the conventional solution by straightforward integration. For the details, see our last remark below.

Our notation of function spaces introduced in section 2 for n = 3 carries over to arbitrary n. Now let $H \in C_0^{\alpha} \cap L^1$ and, for n > 2, define the exact 1-form

$$\tau^{(1)}(x) := -\frac{n}{|S^{n-1}|} d\int |x-y|^{2-n} H(y) \, \mathrm{d}^n y; \tag{45}$$

when n = 2 the logarithmic kernel $\ln \frac{|y|}{|x-y|}$ has to be used in (45). Note that $\tau^{(1)} \in C_0^{1,\alpha}$ for $n \ge 2$. We are now ready to state our main results.

Theorem 1. Let $H \in C_0^{\alpha} \cap L^1$ be small in the sense that

$$\|\tau^{(1)}\|_{1,\alpha} < \left(2^{2/3} - 1\right)^{3/2}.$$
(46)

Then for $n \ge 2$ the Hodge system (38), (39) with asymptotic condition (42) for n = 2 and (43) for n > 2 has an absolutely convergent series solution in $C_0^{1,\alpha}$, given by

$$\tau = \sum_{k=0}^{\infty} \tau^{(2k+1)},$$
(47)

with $\tau^{(1)}$ given by (45) for n > 2, and by its logarithmic kernel version for n = 2, while $\tau^{(2k+1)}$ for $k \in \mathbb{N}$ is recursively given by

$$\tau^{(2k+1)} = \mathbf{P}T^{(2k+1)}, \qquad k \in \mathbb{N}$$

$$\tag{48}$$

where $T^{(2k+1)}$ is a polynomial in the $\tau^{(\ell)}$ with odd $\ell < 2k+1$, viz.

$$T^{(2k+1)} = -\sum_{h=1}^{k} \tau^{(2(k-h)+1)} \sum_{j=1}^{h} M_{j}^{\mp} \sum_{|\ell|_{2j}=h-j} \prod_{i=1}^{j} \tau^{(2\ell_{2i-1}+1)} \cdot \tau^{(2\ell_{2i}+1)}, \quad (49)$$

with $|\ell|_K = \sum_{i=1}^K \ell_i$, the ℓ_i take any non-negative integer values, and with

$$M_j^{\mp} = (\pm 1)^j \frac{(2j-1)!!}{j! 2^j},\tag{50}$$

the j-th Maclaurin coefficient of $1/\sqrt{1 \mp z}$ (with $M_0^{\mp} := 1$), and with $\tau^{(a)} \cdot \tau^{(b)}$ denoting the conventional inner product of two one-forms $\tau^{(a)}$ and $\tau^{(b)}$; moreover, $\mathbf{P} : C_0^{1,\alpha} \to C_0^{1,\alpha}$ projects onto the co-closed subspace of $C_0^{1,\alpha}$ (-valued one-forms), i.e. explicitly, for n > 2:

$$\mathbf{P}T^{(2k+1)}(x) = T^{(2k+1)}(x) + \frac{1}{|S^{n-1}|} d\int \frac{\delta T^{(2k+1)}(y)}{|x-y|^{n-2}} d^n y,$$
(51)

while for n = 2 the logarithmic kernel version has to be used.

The series solution for the τ -system immediately yields:

Corollary 2. Under the conditions expressed in Theorem 1, the Hodge system (31), (32) with its pertinent asymptotic condition (33), or (34), has an absolutely convergent series solution in $C_0^{1,\alpha}$, which for n > 2 is given by

$$\omega = \mp \frac{1}{|S^{n-1}|} \sum_{k \in \mathbb{N}} d \int \frac{\delta T^{(2k-1)}(y)}{|x-y|^{n-2}} \, \mathrm{d}^n y, \tag{52}$$

while for n = 2 the logarithmic kernel $\ln \frac{|y|}{|x-y|}$ has to be used in (52). Here, for notational convenience, we have extended the list of $T^{(2k+1)}s$ defined by (49) for $k \in \mathbb{N}$ to the case k = 0, by setting

$$T^{(1)} := \tau^{(1)}. \tag{53}$$

Finally, the series solution for the ω -system now yields:

Corollary 3. Under the conditions expressed in Theorem 1, when n > 2 the scalar PDE (26) has an absolutely convergent $C_0^{2,\alpha}$ solution given by

$$u(x) = \mp \frac{1}{|S^{n-1}|} \sum_{k \in \mathbb{N}}^{\infty} \int \frac{\delta T^{(2k-1)}(y)}{|x-y|^{n-2}} \, \mathrm{d}^n y;$$
(54)

when n = 2 we use the logarithmic kernel $\ln \frac{|y|}{|x-y|}$, in which case the series converges absolutely in $C^{2,\alpha}$ on compact subsets.

Remark 6. The choice of logarithmic kernel in Corollary 3 fixes the additive constant which the asymptotic condition (28) couldn't.

Remark 7. Alternately, one can of course apply (37) to the τ -series solution (47) and map τ to ω in order to solve the ω -system. However, this is only useful in principle, whereas in practice one will work with the approximate solutions given by the pertinent partial sums of (52).

Remark 8. When n = 1, then the kernel |y| - |x - y| has to be used in (45), and we only have $\tau^{(1)} \in C_b^{1,\alpha}$. In this case the series terminates right away, i.e. $\tau^{(2k+1)} \equiv 0$ for all $k \in \mathbb{N}$, and $\tau^{(1)} = \tau$ is the solution. However, the "smallness" condition (30) has to be imposed in the Euclidean setting to map τ to ω , via (37), which is then integrated to get u.

5 Proofs of the main results

We first prove Theorem 1. Setting n = 3 in the proof proves Proposition 1.

5.1 Proof of Theorem 1

The proof consists of two parts: we first construct a formal series solution, and then prove its absolute convergence in the Banach algebra $C_0^{1,\alpha}$, for which we obtain an explicit lower estimate for the radius of convergence proportional to the "size of H."

5.1.1 The small curvature τ -hierarchy and its formal solution

To facilitate the calculations, we temporarily introduce a "smallness parameter" $\epsilon \in \mathbb{R}_+$ through the replacement of H(x) by $\epsilon H(x)$ in (38), then make the Ansatz

$$\tau = \sum_{p=1}^{\infty} \epsilon^p \tau^{(p)},\tag{55}$$

with each $\tau^{(p)}$ independent of ϵ . Inserting (55) into the pair of equations (38), (39), with H(x) replaced by $\epsilon H(x)$, and into the asymptotic conditions (42) for n = 2 and (43) for n > 2, then sorting according to powers of ϵ , we find a hierarchy of linear equations with pertinent asymptotic conditions. At each order in ϵ the linear system in question can be solved explicitly in a standard way. At the end of the procedure, we set $\epsilon = 1$ in (55) and obtain a formal solution of the τ system (38), (39).

In particular, $\tau^{(1)}$ satisfies

$$\delta \tau^{(1)} = nH, \tag{56}$$

together with

$$d\tau^{(1)} = 0. (57)$$

Consistency with our asymptotic conditions for u requires that for $H_0 \in C_0^{\alpha} \cap L^1$ we select the unique solution $\tau^{(1)} \in C_0^{1,\alpha}$ of the pair of linear equations (56), (57), given by the exact 1-form (45) for n > 2; when n = 2 our logarithmic kernel will be used.

Having the exact 1-form (45), we next find that $\tau^{(2)}$ satisfies

$$\delta \tau^{(2)} = 0 \tag{58}$$

together with

$$d\tau^{(2)} = 0,$$
 (59)

and $\tau^{(2)} \to 0$ at infinity. Clearly, the pair of linear equations (58), (59) has a unique solution $\tau^{(2)}$ which tends $\rightarrow 0$ at infinity, given by the trivial solution $\tau^{(2)} \equiv 0.$

Carrying on we now find recursively that each $\tau^{(p)}$ for even p, i.e. p = 2kfor $k \in \mathbb{N}$, satisfies

$$\delta \tau^{(2k)} = 0 \tag{60}$$

$$d\tau^{(2k)} = 0, \tag{61}$$

with $\tau^{(2k)} \to 0$ at infinity, so that $\tau^{(2k)} \equiv 0$ for general $k \in \mathbb{N}$, while for odd p = 2k + 1 with $k \in \mathbb{N}$, we find the pair of linear first-order PDE

$$\delta \tau^{(2k+1)} = 0 \tag{62}$$

$$d\tau^{(2k+1)} = dT^{(2k+1)}, \tag{63}$$

where $T^{(2k+1)}$ is the polynomial in the $\tau^{(\ell)}$ with odd $\ell < 2k+1$, given in (49), and ensuing explanations. It is easily checked that each $T^{(2k+1)}$ is in $C_0^{1,\alpha}$; namely, this follows inductively from the facts (a) that $\tau^{(1)} \in C_0^{1,\alpha}$ when $n \ge 2$, and (b) that $C_0^{1,\alpha}$ is a Banach algebra. Note also that $\delta T^{(2k+1)} \in C_0^{\alpha} \cap L^1$ for $n \geq 2$, so that (51) is well-defined. The pair of linear first-order PDE (62), (63) therefore has a unique solution $\tau^{(2k+1)} \in C_0^{1,\alpha}$ given by (48), with the projector $\mathbf{P}: C_0^{1,\alpha} \to C_0^{1,\alpha}$ given in (51) for n > 2; when n = 2 our logarithmic kernel will be used.

Thus we have formally solved our τ -problem (38), (39), with H replaced by ϵH , in terms of the formal series solution (55). We can now set $\epsilon = 1$ in (55) and obtain, for $n \ge 2$, a formal series solution (55). We can now set $\epsilon = 1$ iff (55) and obtain, for $n \ge 2$, a formal series solution of (38), (39) in $C_0^{1,\alpha}$, in the sense that each partial sum of $\tau = \sum_{k=0}^{\infty} \tau^{(2k+1)}$ is in $C_0^{1,\alpha}$. It remains to prove that the so-obtained formal series solution to the τ system, for $n \ge 2$, converges absolutely in $C_0^{1,\alpha}$ whenever (46) holds.

Absolute convergence of the formal τ series solution 5.1.2

To show that for all $n \ge 2$ the formal series $\tau = \sum_{k=0}^{\infty} \tau^{(2k+1)}$ converges ab-

To show that for all $n \geq 2$ the formal series $\tau = \sum_{k=0}^{\infty} \tau$, converges assolutely in $C_0^{1,\alpha}$ when the condition (46) holds, it remains to show that the right hand side of the norm estimate $\|\tau\|_{1,\alpha} \leq \sum_{k=0}^{\infty} \|\tau^{(2k+1)}\|_{1,\alpha}$ converges. To simplify notation we henceforth drop the subscript " $_{1,\alpha}$ " from the norm symbols. We now estimate all norms $\|\tau^{(2k+1)}\|$ for $k \in \mathbb{N}$ in terms of the 2k+1-th power of $\|\tau^{(1)}\|$. Now, $\|\tau^{(2k+1)}\| = \|\mathbf{P}T^{(2k+1)}\|$ for $k \in \mathbb{N}$, and since $\mathbf{P}: C_0^{1,\alpha} \to C_0^{1,\alpha}$, is a projector, we have the estimate: $\|\tau^{(2k+1)}\| \leq \|T^{(2k+1)}\|$. Substituting the RHS of (49) for $T^{(2k+1)}$, repeating the standard inequality $\|\sum_i \tau_i\| \leq \sum_i \|\tau_i\|$, then using the inequality $\|\tau,\tau_i\| \leq \|\tau_i\|\|\tau_i\|$ valid in Banach algebras (here $C_0^{1,\alpha}$). then using the inequality $\|\tau_i \tau_j\| \leq \|\tau_i\| \|\tau_j\|$ valid in Banach algebras (here $C_0^{1,\alpha}$), and applying repeatedly the identity $\|\tau^{(2a+1)}\| = \|\mathbf{P}T^{(2a+1)}\|$ followed by the projector estimate $\|\mathbf{P}T^{(2a+1)}\| \leq \|T^{(2a+1)}\|$ for the various pertinent values of

 $a \ge 1$ (no estimate is necessary when a = 0), and using $|M_j^{\mp}| = M_j^{-} := M_j$, for $k \geq 1$ we obtain

$$\|T^{(2k+1)}\| \leq \sum_{h=1}^{k} \|\tau^{(2(k-h)+1)}\| \sum_{j=1}^{h} |M_{j}^{\mp}| \sum_{|\ell|_{2j}=h-j} \prod_{i=1}^{2j} \|\tau^{(2\ell_{i}+1)}\|$$

$$\leq \sum_{h=1}^{k} \|T^{(2(k-h)+1)}\| \sum_{j=1}^{h} M_{j} \sum_{|\ell|_{2j}=h-j} \prod_{i=1}^{2j} \|T^{(2\ell_{i}+1)}\|.$$
(64)

Next we show that for all $k \in \mathbb{N}$ there exists some R_{2k+1} such that

$$\|T^{(2k+1)}\| \le R_{2k+1} \|\tau^{(1)}\|^{2k+1}.$$
(65)

Setting k = 1 and recalling the definition (53) we obtain the estimate

$$\|T^{(3)}\| \le \frac{1}{2} \|\tau^{(1)}\|^3.$$
(66)

Now suppose that for all $k = 1, ..., k_*$ there exists some R_{2k+1} such that (65) holds. Then the estimate (64) guarantees that (65) is true also for $k = k_* + 1$, and since $k_* \geq 1$ is arbitrary in this induction step while (66) says that the estimate is true for $k_* = 1$, it follows that (65) is true for all $k \in \mathbb{N}$.

The inductive proof that (65) holds for all $k \in \mathbb{N}$ also yields that R_{2k+1} is recursively defined for $k \in \mathbb{N}$ by

$$R_{2k+1} = \sum_{h=1}^{k} R_{2(k-h)+1} \sum_{j=1}^{h} M_j \sum_{|\ell|_{2j}=h-j} \prod_{i=1}^{2j} R_{2\ell_i+1},$$
(67)

with $R_1 := 1$.

with $R_1 := 1$. Recall that we want to show that $\sum_{k=0}^{\infty} \|\tau^{(2k+1)}\| < \infty$ for sufficiently small $\|\tau^{(1)}\|$. Since we have shown that $\|\tau^{(2k+1)}\| \le \|T^{(2k+1)}\| \le R_{2k+1}\|\tau^{(1)}\|^{2k+1}$, it suffices to show that $\sum_{k=0}^{\infty} R_{2k+1}\|\tau^{(1)}\|^{2k+1} < \infty$ for sufficiently small $\|\tau^{(1)}\|$. Setting now $\|\tau^{(1)}\| =: \xi$, we note that the formal power series $G(\xi) := \sum_{k=0}^{\infty} R_{2k+1}\xi^{2k+1}$ is nothing but the formal generating function of the R_{2k+1} , in the usual sense that, formally, $R_{2k+1} = G^{(2k+1)}(0)/(2k+1)!$. So our task is to show that the generating function is analytic about $\xi = 0$ with radius of convergence given by the RHS of (46).

With the help of the recursion relation (67) we readily find that $G(\xi)$ is the positive inverse function of $g \mapsto \xi$ given by

$$\xi = 2g - \frac{g}{\sqrt{1 - g^2}} \tag{68}$$

defined for positive ξ near $\xi = 0$, with G(0) = 0. Since the function $g \mapsto \xi$ given by (68) is analytic about g = 0 (with radius of convergence = 1) and has unit derivative at g = 0, there now exists an open neighborhood of $\xi = 0$ on which there is defined a unique inverse function $\xi \mapsto g = G(\xi)$ which vanishes at $\xi = 0$, has unit derivative at $\xi = 0$, and satisfies (68). Thus, in particular, the Maclaurin expansion of $G(\xi)$ converges to $G(\xi)$ and it generates the recursion coefficients R_{2k+1} .

We now determine the finite radius of convergence ξ_* of the power series for $G(\xi)$ about $\xi = 0$. Setting $G(\xi) = \sin \Psi(\xi)$ we see that (68) defines the function $\psi \mapsto \xi$ given by

$$\xi = 2\sin\psi - \tan\psi,\tag{69}$$

with $\xi = 0$ when $\psi = 0$. Since $\psi \mapsto \sin \psi$ is an entire function, which vanishes for $\psi = 0$ and has unit derivative there, the radius of convergence of the Maclaurin series of $\xi \mapsto G(\xi) = \sin \Psi(\xi)$ coincides with the radius of convergence of the Maclaurin series of $\xi \mapsto \Psi(\xi)$. This radius of convergence in turn is determined by those ξ value(s) closest to $\xi = 0$ at which the derivative of $\psi \mapsto \xi = 2 \sin \psi - \tan \psi$ vanishes (possibly asymptotically should $\xi \to \xi_{\infty}$ when $|\psi| \to \infty$ suitably). But this ψ derivative is $2\cos \psi - 1/\cos^2 \psi$, and it vanishes iff $2\cos^3 \psi = 1$, which gives $2^{1/3} \cos \psi \in \{1, e^{i2\pi/3}, e^{i4\pi/3}\}$. A calculation now gives the radius of convergence of $G(\xi)$ about $\xi = 0$ as

$$\xi_* = \left(2^{2/3} - 1\right)^{3/2}.\tag{70}$$

Thus we have shown that our formal series solution converges absolutely in $C_0^{1,\alpha}$ if $\|\tau^{(1)}\| < (2^{2/3} - 1)^{3/2}$. This completes our convergence proof.

Remark 9. Because of our use of the projector and Banach algebra estimates, we cannot conclude that $\|\tau^{(1)}\| < (2^{2/3} - 1)^{3/2}$ is a necessary criterion for absolute convergence of our formal power series solution. Indeed, for any radially symmetric $H \in C_0^{\alpha}$ the formal power series reduces to its first term, all other terms being identically zero, so convergence is a trivial issue and holds for any size of H, then.

5.2 Proofs of Corollaries 2 and 3

To prove Corollary 2, use (37) but insert (55) as well as the analogous Ansatz

$$\omega = \sum_{p=1}^{\infty} \epsilon^p \omega^{(p)},\tag{71}$$

then sort by powers of ϵ and easily solve for $\omega^{(p)}$ explicitly; reinsert those expressions into (71) and now set $\epsilon = 1$. The result is (52). Its absolute convergence follows verbatim as in the proof of Theorem 1.

To prove Corollary 3, integrate $\omega = \nabla u \cdot dx$, with ω from given by the absolutely convergent series (52); the constant of integration is fixed by the asymptotic conditions.

6 The cubic reformulation of the τ -problem

We now come back to our earlier remark that the nonlinearity of the τ -problem is effectively cubic. However, for technical reasons, in this section we restrict attention to n = 3. This allows us to return to the three-dimensional vector formulation for which many well-known identities of vector analysis are at our disposal. Thus the τ -problem then becomes a v-problem.

The "cubic version" of our v-problem reveals some interesting a priori differential identities which are satisfied by any solution but which are obscured by the original formulation of the v-problem.

More importantly, the algebraic simplifications dramatically reduce the combinatorial complexity of the successive approximations in the small-H hierarchy. Indeed, except for the first few low-order terms, the series solution to our original first-order v vector problem and its spin-off, the solutions to the original first-order w vector problem and the original second-order scalar u-problem, soon involves terms which look more and more unwieldy. The cubic version of the v-problem offers relief, by having to evaluate fewer integrals.

Interestingly enough, though, the solution theory of this apparently simpler cubical formulation of the small-H hierarchy seems more complicated than the "square-root formulation" and raises some challenging questions. More to the point, it will be clear from the equivalence of the original and the cubic v-problems that the cubic version has a solution v whenever v solves the original problem, and vice-versa. Moreover, upon reintroducing the ϵ parameter, it is clear from the analyticity in ϵ that a small-H expansion will produce the same series solutions as before. However, if we ignore for a moment that we already know that our formal series derived from the original version of the v-problem converges absolutely to a classical solution, from which the solvability of the cubic hierarchy of linear equations follows as a corollary, then the solvability of the cubic hierarchy of linear equations is not at all obvious but needs to be verified! We will prove the consistency ab initio, but did not succeed in proving its convergence without recourse to the square-root formulation. Perhaps some reader will feel inspired to settle this problem!

6.1 The cubically nonlinear reformulation of the *v*-problem

Carrying out the curl operation in

$$\nabla \times \frac{v}{\sqrt{1 \mp |v|^2}} = 0 \tag{72}$$

and multiplying through with $(1 \mp |v|^2)^{3/2}$ we find that, away from singularities (which occur, for instance, when $1 - |v|^2 \to 0$), any solution v of (72) satisfies

$$0 = \left(1 \mp |v|^2\right) \nabla \times v \pm \left[\frac{1}{2} \nabla |v|^2\right] \times v.$$
(73)

This equation already exhibits a cubic nonlinearity, yet it can be further manipulated into a more concise alternate format. Namely, using first the identity $\frac{1}{2}\nabla |v|^2 = (v \cdot \nabla)v + v \times (\nabla \times v)$ well-known from vector analysis, we find that (73) is equivalent to

$$0 = (1 \mp |v|^2) \nabla \times v \pm [(v \cdot \nabla)v + v \times (\nabla \times v)] \times v.$$
(74)

We next employ the identity $v \times (\nabla \times v) \times v = |v|^2 \nabla \times v - v(v \cdot \nabla \times v)$, well-known from vector algebra, which for any solution of (73) simplifies to $v \times (\nabla \times v) \times v = |v|^2 \nabla \times v$ because any solution of (73) satisfies (78). Inserting $v \times (\nabla \times v) \times v = |v|^2 \nabla \times v$ into (74) yields a cancellation and (74) becomes

$$\nabla \times v = \pm v \times (v \cdot \nabla)v. \tag{75}$$

This is perhaps the most concise cubic form of the curl equation for v. When paired with the divergence equation

$$\nabla \cdot v = 3H \tag{76}$$

we arrive at the following conclusion.

Proposition 2. The curl equation (75) paired with the divergence equation (76) forms a closed system of first-order vector PDE for v. To make them well posed the vector equations need to be supplemented by asymptotic conditions for v at spatial infinity, which as before we take to be vanishing in agreement with the monopole asymptotics of u at spatial infinity. By construction, this cubic set of equations for v, with v vanishing at infinity, is equivalent to the scalar equation (26), with u vanishing at infinity, for the appropriate choice of sign.

6.2 The Helmholtz decomposition for the cubic version

For the sake of completeness, we remark that the Helmholtz decomposition $v = v_g + v_c$ leads to analogous conclusions for the cubic version of the *v*-problem. We find the previously obtained linear equation (18) for v_g (together with $\nabla \times v_g = 0$), while the nonlinear equation for v_c (given v_q) becomes

$$\nabla \times v_c = \pm (v_g + v_c) \times (v_g + v_c) \cdot \nabla (v_g + v_c)$$
(77)

(together with $\nabla \cdot v_c = 0$). This closed set of first-order vector equations for the vector fields v_g and v_c is supplemented by the asymptotic conditions that v_q and v_c vanish at spatial infinity.

As announced, given the gradient field (20), the remaining equation (77) now becomes a closed vector equation for the solenoidal field v_c . It remains to solve equation (77) with v_q given by (20).

6.3 Spin-off: a-priori differential identities for solutions

The cubic version of the v-problem reveals two a-priori differential identities which are satisfied by any solution of the v-problem.

Proposition 3. For any solution of the v-problem,

$$v \cdot (\nabla \times v) = 0 \tag{78}$$

and

$$\nabla |v|^2 \cdot \nabla \times v = 0,\tag{79}$$

where $\nabla \times v$ is defined.

Proof of Proposition 3:

Dotting (73) with v yields (78), and dotting (73) with $\nabla |v|^2$ yields (79). Clearly, neither (78) nor (79) are generally true for arbitrary vector fields v.

6.4 The small-*H* expansion: cubic version

We now turn to the *formal* small-H solution strategy for the cubic v-problem. We begin by deriving the cubic analog of the hierarchy of linear PDE. To present a slightly different perspective, this time we take the Helmholtz point of view. Also, we take the liberty and bypass the reintroduction of ϵ and, instead of "powers of ϵ ," simply talk about "orders of smallness."

In this vein, for $H \in C_0^{\alpha} \cap L^1$ we find v_g given in (20). Now, if H is suitably small, then v_g is small in $C_0^{1,\alpha}$, and we write $v_g = v^{(1)}$. For the solenoidal part of v we now make the series Ansatz $v_c = v^{(p_1)} + v^{(p_2)} + v^{(p_3)} + \ldots$ with $1 < p_1 < p_2 < p_3 < \cdots$, and with each $v^{(p_k)}$ being of p_k -th "order of smallness," compared to $v^{(1)}$. Inserting this Ansatz into the cubic vector PDE (77) and identifying p_1 with the smallest order on the RHS, p_2 with the nextto-smallest order, and so on, we find recursively that $p_1 = 3$, then $p_2 = 5$, and $p_k = 2k + 1$ for general $k \in \mathbb{N}$. Furthermore, for $v = v_g + v_c$ with $v_g = v^{(1)}$ and $v_c = v^{(3)} + v^{(5)} + v^{(7)} + \ldots$ to be a solution of the pair of equations (76), (77), each $v^{(2k+1)}$ for $k \in \mathbb{N}$ has to satisfy

$$\nabla \times v^{(2k+1)} = \pm \sum_{h+i+j=k-1} v^{(2h+1)} \times (v^{(2i+1)} \cdot \nabla) v^{(2j+1)}$$
(80)

supplemented by the solenoidality condition

$$\nabla \cdot v^{(2k+1)} = 0. \tag{81}$$

Supposing that $v^{(2\ell+1)}$ is known for all $\ell \leq k-1$, then (80), (81) is a pair of linear first-order PDE for $v^{(2k+1)}$, with vanishing conditions at spatial infinity for $v^{(2k+1)}$. Now $v^{(1)}$ is known, and so, by induction, it follows that (80), (81) successively determine $v^{(3)}$, then $v^{(5)}$, and so on — provided that each equation in this formal linear hierarchy of equations is solvable (in $C_0^{1,\alpha}$)!

6.5 Solvability of the cubic hierarchy

Proposition 4. The infinite hierarchy of equations (80), (81), together with $v_g = v^{(1)}$ given in (20), is uniquely solvable in $C_0^{1,\alpha} \cap L^2$ at each order $k \in \mathbb{N}$,

and this solution is given by

$$v^{(2k+1)}(x) = \pm \frac{1}{4\pi} \sum_{\substack{h+i+j\\ =k-1}} \int (v^{(2h+1)} \times (v^{(2i+1)} \cdot \nabla) v^{(2j+1)})(y) \times \frac{x-y}{|x-y|^3} \,\mathrm{d}^3y.$$
(82)

Proof of Proposition 4: Since $\sum_{h+i+j=k-1} v^{(2h+1)} \times (v^{(2i+1)} \cdot \nabla) v^{(2j+1)} \in C_0^{\alpha} \cap L^1$ if $v^{(2\ell+1)} \in C_0^{1,\alpha} \cap L^2$ for all $\ell \leq k-1$, it remains to be verified that the collected terms on the RHS of (80) have vanishing divergence at each order k. With some effort one can show term by term that (82) is identical to (9), (12). For instance, based on the identity $\nabla(a \cdot b) = (a \cdot \nabla)b + (b \cdot \nabla)a + a \times \nabla \times b + b \times \nabla \times a$ for any two vector fields a and b it is readily shown that

$$-v^{(1)} \times (v^{(1)} \cdot \nabla)v^{(1)} = \frac{1}{2} \nabla |v^{(1)}|^2 \times v^{(1)} = \nabla \times (\frac{1}{2} |v^{(1)}|^2 v^{(1)}),$$
(83)

proving the equality of the RHS of (82) and that of (9), (12) when k = 1. However, the procedure of proving equality term by term soon gets very complicated, and an inductive argument is needed, instead.

Proceeding by induction, we will now show that, given $v_g = v^{(1)}$, for each $k \in \mathbb{N}$ there is a $v^{(2k+1)}$ which vanishes at infinity and solves

$$\nabla \times v^{(2k+1)} = \pm \sum_{h+i+j=k-1} v^{(2h+1)} \times (v^{(2i+1)} \cdot \nabla) v^{(2j+1)}.$$
(84)

Since the formulas become rather long, we switch to the shorter notation $v^{[k]}$ for $v^{(2k+1)}$ for all k = 0, 1, 2, 3, ...; this mildly obscures the order of smallness to which the terms belong, but shortens the length of the formulas considerably.

Step one is easily disposed of by checking explicitly that $v_q \times (v_q \cdot \nabla) v_q$ is divergence-free for any gradient field v_g . Indeed, since $\nabla \times v_g = 0$ for any gradient field v_g , first of all the well-known identity $(v \cdot \nabla)v = \frac{1}{2} \nabla |v|^2 - v \times (\nabla \times v)$ reduces to $(v_g \cdot \nabla)v_g = \frac{1}{2}\nabla |v_g|^2$, and second, $v_g \times \left(\frac{1}{2}\nabla |v_g|^2\right) = -\nabla \times \left[\frac{1}{2}|v_g|^2 v_g\right]$, so that $v_g \times (v_g \cdot \nabla) v_g = -\nabla \times \left[\frac{1}{2}|v_g|^2 v_g\right]$ is a curl, i.e. divergence-free. Hence, given $v^{[0]} = v_g$, for either sign of "±" there is a $v^{[1]}$ satisfying $\nabla \times v^{[1]} = \pm v^{[0]} \times (v^{[0]} \cdot \nabla) v^{[0]}$. Moreover, since v_g is given by (20) and $H \in C_0^{\alpha} \cap L^1$, we have that $v_g \times (v_g \cdot \nabla) v_g \in C_0^{1,\alpha} \cap L^1$, and so in particular there is a solution $v^{[1]}$ of $\nabla \times v^{[1]} = \pm v^{[0]} \times (v^{[0]} \cdot \nabla) v^{[0]}$ which vanishes at infinity.

As to the induction step, suppose that for all $k \leq m$, and either sign of "±," there is a solution $v^{[k]}$ of (84) which vanishes at spatial infinity. We now show that then also $\sum_{h+i+j=m} v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]}$ is divergence free and vanishes sufficiently rapidly at infinity.

We will need two equalities. Taking the dot product of both sides of (84)

with $v^{[l]}$ and summing for l+k=m' with $m'\leq m$ gives

$$\sum_{k+l=m'} v^{[l]} \cdot \nabla \times v^{[k]} = \pm \sum_{\substack{k+l \ =m'}} \sum_{\substack{h+i+j \ =k-1}} v^{[l]} \cdot \left[v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]} \right]$$
(85)

$$= \pm \sum_{\substack{h+i+j+l\\-m'-1}} v^{[l]} \cdot \left[v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]} \right]$$
(86)

$$= 0,$$
 (87)

where the last equality follows by the anti-symmetry of the triple product $a \cdot (b \times c)$; (87) is the order *m* expansion analogue of (78).

Similarly, dotting both sides of (84) with $(v^{[l]} \cdot \nabla)v^{[p]}$ and summing over k + l + p = m gives

$$\sum_{k+l+n=m} \nabla \times v^{[k]} \cdot \left((v^{[l]} \cdot \nabla) v^{[p]} \right) =$$
(88)

$$\sum_{\substack{k+l+n\\m=m-l+1}} \sum_{\substack{h+i+j\\m=k-1}} (v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]}) \cdot ((v^{[l]} \cdot \nabla) v^{[p]}) =$$
(89)

$$\sum_{h+i+j+l+n=m-1} (v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]}) \cdot ((v^{[l]} \cdot \nabla) v^{[p]}) = 0.$$
(90)

Now note that

$$\begin{split} \sum_{h+i+j=m} v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]} &= \\ \sum_{h+i+j=m} v^{[h]} \times \left(\frac{1}{2} \nabla (v^{[i]} \cdot v^{[j]}) - v^{[i]} \times (\nabla \times v^{[j]}) \right) = \\ \sum_{h+i+j=m} \left(-\nabla \times \left(\frac{1}{2} (v^{[i]} \cdot v^{[j]}) v^{[h]} \right) + \frac{1}{2} (v^{[i]} \cdot v^{[j]}) \nabla \times v^{[h]} \\ &- (v^{[h]} \cdot \nabla \times v^{[j]}) v^{[i]} + (v^{[h]} \cdot v^{[i]}) \nabla \times v^{[j]} \right) = \\ \sum_{h+i+j=m} \left(-\nabla \times \left(\frac{1}{2} (v^{[i]} \cdot v^{[j]}) v^{[h]} \right) + \frac{3}{2} (v^{[h]} \cdot v^{[i]}) \nabla \times v^{[j]} \right), \end{split}$$

where we used that $\sum_{h+i+j=m} (v^{[h]} \cdot \nabla \times v^{[j]}) v^{[i]} = 0$, by (87). And so,

$$\nabla \cdot \sum_{h+i+j=m} v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]} = \frac{3}{2} \sum_{h+i+j=m} \nabla (v^{[h]} \cdot v^{[i]}) \cdot \nabla \times v^{[j]} =$$
$$\sum_{h+i+j=m} \left[\left((v^{[h]} \cdot \nabla) v^{[i]} + (v^{[i]} \cdot \nabla) v^{[h]} \right) \cdot \nabla \times v^{[j]} + \left((v^{[h]} \times \nabla \times v^{[i]} + v^{[i]} \times \nabla \times v^{[h]}) \right) \cdot \nabla \times v^{[j]} \right].$$

By (90), $\sum_{h+i+j=m} \left((v^{[h]} \cdot \nabla) v^{[i]} + (v^{[i]} \cdot \nabla) v^{[h]} \right) \cdot \nabla \times v^{[j]} = 0$. Moreover, by

the anti-symmetry of the triple product $a \cdot (b \times c)$ we have

$$\sum_{h+i+j=m} (v^{[h]} \times (\nabla \times v^{[i]}) + v^{[i]} \times (\nabla \times v^{[h]})) \cdot \nabla \times v^{[j]} = 0.$$
(91)

Thus $\sum_{h+i+j=m} v^{[h]} \times (v^{[i]} \cdot \nabla) v^{[j]}$ is divergence free, as claimed.

6.6 Convergence of the cubic v series

Summing (82) over $k \in \mathbb{N}$ and adding $v^{(1)} = v_g$ given by (20) yields a formal series solution to the cubic v-problem, viz.

$$v(x) = v_g(x) \pm \frac{1}{4\pi} \sum_{\substack{k=1\\k=1}}^{\infty} \sum_{\substack{k=1\\k=1}}^{\infty} \nabla \times \int \frac{(v^{(2h+1)} \times (v^{(2i+1)} \cdot \nabla)v^{(2j+1)})(y)}{|x-y|} \,\mathrm{d}^3y.$$
(92)

We now have to address the convergence of this formal series solution.

Since the combinatorial structure of the RHS of (82) is considerably simpler than that of (9), (12), it would seem that a convergence proof is more readily forthcoming than our previous proof. Curiously, the obviously simpler series expansion for the cubic version of the v-problem has not at all yielded to our attempts of proving its convergence *directly*, i.e. without recourse to the original version of the first-order vector problem with its more complicated nonlinearity. So we are finally forced to recall the origin of the cubic version of the v-problem to conclude that (92) converges absolutely to a classical solution for small H.

So there is also a challenge: it would be good to have a simple convergence proof directly for the cubic hierarchy!

7 Related problems in divergence form

Our power series technique of solving the prescribed mean-curvature problem for graphs over \mathbb{R}^n can handle more general quasilinear problems in divergence form. In particular, let $f(s) = 1 + as + bs^2 + \cdots$ be given and analytic about s = 0, then the equation

$$\nabla \cdot \left(f(|\nabla u|) \nabla u \right) = \rho, \tag{93}$$

with $\rho \in C_0^{\alpha} \cap L^1$ small enough and asymptotic monopole condition for u, can be solved with the same solution techniques developed here for the scalar prescribed mean-curvature equation. In particular, such type of problems occur in the theory of stationary compressible fluid flows, see [6, 52, 53, 54], and other versions of nonlinear electrostatics [18].

While we here are interested in entire solutions over \mathbb{R}^n motivated by the electromagnetic Born–Infeld model, and also by some problems in the theory

of spacetime structure (e.g. [1, 3, 27, 58]), with some mild modifications our approach should be adaptable to the prescribed mean-curvature equation in bounded domains with small Dirichlet data for u (e.g. [51, 7, 31, 59, 57, 12, 19]); this then also includes the minimal / maximal hypersurface problem ($H \equiv 0$) with small Dirichlet data (e.g. [20, 45, 48, 21, 56, 46, 5]). The parametric prescribed mean-curvature problem (e.g. [60, 13, 15, 56, 62, 17]), which also captures embedded hypersurfaces which are not graphs over \mathbb{R}^n and even nonembedded hypersurfaces, has a different structure, however.

Acknowledgement This work was started while H. Carley was a Hill Assistant Professor at Rutgers University, supported in parts by the NSF through grant DMS-0406951. M. Kiessling was supported by the NSF through grants DMS-0406951 and DMS-0807705; and in the final stages of this work also by CNRS Universite de Provence through a poste rouge. Thanks go to A. Shadi Tahvildar-Zadeh and Sagun Chanillo for helpful comments on the manuscript, and to Y. Elskens for his hospitality and interest in this work.

References

- Akutagawa, K., A note on spacelike hypersurfaces with prescribed mean curvature in spatially closed globally static Lorentzian manifolds, *Memoirs Fac. Sci. Kyushu Univ.* A40, 119–123 (1986).
- [2] Akutagawa, K., and Nishikawa, S., The Gauss map and spacelike surfaces with prescribed mean curvature in Minkowski 3-space, *Tôhoku Math. J.* 42, 67–82 (1990).
- [3] Bartnik, R., Existence of maximal surfaces in asymptotically flat spacetimes Commun. Math. Phys. 94, 155–175 (1984).
- [4] Bereanu, C., Jebelean, P., and Mahwin, J., Radial solutions for some nonlinear problems involving mean curvature operators in Euclidean and Minkowskian spaces, *Proc. Amer. Math. Soc.* 137, 161-169 (2009).
- [5] Bergner, M., On the Dirichlet problem for the prescribed mean curvature equation over general domains, *Diff. Geom. Appl.*, **27**, 335-343 (2009).
- [6] Bers, L, The mathematics of subsonic and transsonic fluid dynamics, Trans. Amer. Math. Soc. 70, 465-491 (1951).
- [7] Bombieri, E., and Giusti, E., Local estimates for the gradient of nonparametric surfaces of prescribed mean curvature, *Commun. Pure Appl. Math.* 26 381–394 (1973).
- [8] Born, M., Modified field equations with a finite radius of the electron, *Nature* 132, 282 (1933).
- [9] Born, M., Théorie non-linéare du champ électromagnétique, Ann. Inst. H. Poincaré 7, 155-265 (1937).
- [10] Born, M., Atomic physics, 8th rev. ed., Blackie & Son Ltd., Glasgow (1969).
- Born, M., and Infeld, L., Foundation of the new field theory, *Nature* 132, 1004 (1933); Ditto, *Proc. Roy. Soc. London* A 144, 425-451 (1934).
- [12] Bourni, T., $C^{1,\alpha}$ theory for the Prescribed Mean Curvature Equation with Dirichlet Data, J. Geom. Anal. **21**, 982–1035 (2011).
- [13] Brezis, H., and Coron, J.M., Convergence of solutions of H-systems or how to blow bubbles, Arch. Rat. Mech. Anal. 89, 21-56 (1985).
- [14] Burton, D.A., Trines, R.M.G.M., Walton, T.J., and Wen, H., Exploring Born–Infeld electrodynamics using plasmas, *J Phys. A: Math. Theor.* 44 095501 (2011).
- [15] Caldiroli, P., and Musina, R., The Dirichlet Problem for H -Systems with Small Boundary Data: BlowUp Phenomena and Nonexistence Results, *Arch. Rat. Mech. Anal.* 181, 1-42 (2006).

- [16] Carley, H., and Kiessling, M.K.-H., Nonperturbative Calculation of Born-Infeld effects on the Schrödinger spectrum of the Hydrogen atom, *Phys. Rev. Lett.* **96**, 030402 (4pp.) (2006).
- [17] Chanillo, S., and Malchiodi, A., Asymptotic Morse theory for the equation $\Delta v = 2v_x \wedge v_y$, Comm. Anal. Geom. 13, 187–251 (2005).
- [18] Christodoulou, D., The action principle and partial differential equations, Annals of Math. Studies 146, Princeton Univ. Press (1999).
- [19] Corsato, C., Obersnel, F., Omari, P., and Rivetti, S., On the lower and upper solution method for the prescribed mean curvature equation in Minkowski space. *Disc. Cont. Dyn. Syst.* 159–169 (2013).
- [20] Courant, R., Plateau's problem and Dirichlet's principle, Annal. Math. 38, 679–724 (1937).
- [21] Dierkes, U., Hildebrandt, S., Küster, A., and Wohlrab, O., *Minimal sur-faces*, Vol I & II. Grundlehren der Mathematischen Wissenschaften 295. Springer-Verlag, Berlin (1992).
- [22] Ecker, K. Area maximizing hypersurfaces in Minkowski space having an isolated singularity, *Manuscr. Math.* 56, 375-397 (1986).
- [23] Ferraro, R., Two-dimensional solutions for Born-Infeld fields, J. High Energy Phys. 48 (2013).
- [24] Folland, G.B., Real Analysis, 2nd ed., Wiley, New York (1999).
- [25] Franklin, J., and Garon, T., Approximate calculations of Born–Infeld effects on the relativistic Hydrogen spectrum, *Phys. Lett. A* 375, 1391-1395 (2011).
- [26] Fuente, D. de la, Romero, A., and Torres, P.J. Entire spherically symmetric spacelike graphs with prescribed mean curvature function in Schwarzschild and Reissner–Nordström spacetimes, *Class. Quant. Grav.* **32** 035018 (2015).
- [27] Gerhardt, C., H-surfaces in Lorentzian manifolds, Commun. Math. Phys. 89, 523–553 (1983).
- [28] Gibbons, G.W., Born-Infeld particles and Dirichlet p-branes, Nucl. Phys. B 514, 603-639 (1998).
- [29] Gibbons, G.W., and Warner, N.P. Global structure of five-dimensional fuzzballs, *Class. Quant. Grav.* **31** 025016 (2014).
- [30] Gilbarg, D., and Trudinger, N.S., *Elliptic Partial Differential Equations of Second Order*, Second edition. Grundlehren der Mathematischen Wissenschaften 224, Springer-Verlag, Berlin New York, 1983.

- [31] Giusti, E., On the equation of surfaces of prescribed mean curvature, Inventiones Math. 46, 111–137 (1978).
- [32] Greiner, W., Müller, B., and Rafelski, J., Quantum Electrodynamics of Strong Fields, Springer, New York (1985).
- [33] Gulliver, R. D., "Regularity of Minimizing Surfaces of Prescribed Mean Curvature," Annals Math. 97, 275-305 (1973).
- [34] Hoppe, J. Some classical solutions of relativistic membrane equations in 4 spacetime dimensions, *Phys. Lett. B* **329** 1014 (1994).
- [35] Kenmotsu, K., Weierstrass formula for surfaces of prescribed mean curvature, Math. Ann. 245, 89–99 (1979).
- [36] Kiessling, M.K.-H., Electromagnetic field theory without divergence problems. 1. The Born legacy, J. Stat. Phys. 116, 1057-1122 (2004).
- [37] Kiessling, M.K.-H., Convergent perturbative power series solution of the stationary Maxwell–Born–Infeld equations with regular sources, J. Math. Phys. 52, art. 022902, 16pp. (2011).
- [38] Kiessling, M.K.-H., "On the quasi-linear elliptic PDE $-\nabla \cdot \nabla u/\sqrt{1-|\nabla u|^2} = 4\pi \sum a_k \delta_{s_k}$ in physics and geometry," Commun. Math. Phys. **314**, 509–523 (2012).
- [39] Klyachin, A.A., Solvability of the Dirichlet problem for the maximal surface equation with singularities in unbounded domains, *Dokl. Russ. Akad. Nauk* 342, 161-164 (1995); English transl.: *Dokl. Math.* 51, 340-342 (1995).
- [40] Klyachin, A.A., and Miklyukov, V.M., Existence of solutions with singularities for the maximal surface equation in Minkowski space, *Mat. Sb.* 184, 103-124 (1993); English transl.: *Russ. Acad. Sci. Sb. Math.* 80, 87-104 (1995).
- [41] Kobayashi, O., Maximal surfaces in the 3-dimensional Minkowski space L³, Tokio J. Math. 6, 297–309 (1983).
- [42] Lieb, E.H., and Loss, M., Analysis, Grad. Stud. in Math. 14, AMS (1991).
- [43] Meeks III, W.H., Pérez, J., and Ros, A., Limit leaves of an H lamination are stable. J. Diff. Geom. 84, 179-189 (2010).
- [44] Meeks, W.H., and Pérez, J., CMC foliations of closed manifolds, eprint arXiv:1404.1725v3 (2014).
- [45] Nitsche, J.C.C., Lectures on minimal surfaces, Vol. 1. (Translated from the German by Jerry M. Feinberg.) Cambridge Univ. Press, Cambridge, (1989).
- [46] Obersnel, F., and Omari, P., Positive solutions of the Dirichlet problem for the prescribed mean curvature equation, J. Diff. Eq. 249, 1674-1725 (2010).

- [47] Osserman, R., "A proof of the regularity everywhere of the classical solution to Plateau's problem," Annals Math. 91, 550-569 (1970).
- [48] Osserman, R., A survey of minimal surfaces, Van Nostrand, Holland (1969).
- [49] Pryce, M.H.L., The two-dimensional electrostatic solutions of Born's new field equations, Proc. Camb. Phil. Soc. 31, 50–68 (1935).
- [50] Rafelski, J., Fulcher, L. P., and Greiner, W., Superheavy Elements and an Upper Limit to the Electric Field Strength, *Phys. Rev. Lett.* 27, 958–961 (1971).
- [51] Serrin, J., The problem of Dirichlet for quasilinear elliptic differential equations with many independent variables, *Phil. Trans. Roy. Soc. London A* 264, 413–496 (1969).
- [52] Sibner, L.M., and Sibner, R.J., Nonlinear Hodge theory: Applications Adv. Appl. Math. 31, 1–15 (1979).
- [53] Shiffman, M., On the existence of subsonic flows of a compressible fluid, J. Rat. Mech. Anal. 1, 605–652 (1953).
- [54] Smith, P.D., Nonlinear Hodge theory on punctured Riemannian manifolds, Ind. Univ. Math. J. 31, 553–577 (1982).
- [55] Soff, G., Rafelski, J., and Greiner, W., Lower Bound to Limiting Fields in Nonlinear Electrodynamics, *Phys. Rev. A* 7, 903–907 (1973).
- [56] Struwe, M.: Plateau's problem and the Calculus of Variations. Princeton University Press (1985).
- [57] Takakuwa, S., On a parameter dependence of solvability of the Dirichlet problem for non-parametric surfaces of prescribed mean curvature, *Tokyo J. Math.* 7, 421–435 (1984).
- [58] Treibergs, A.E., Entire spacelike hypersurfaces of constant mean curvature in Minkowski space, *Invent. Math.* 66, 39–56 (1982).
- [59] Trudinger, N.S., Graphs with prescribed curvature, in: Nonlinear functional analysis and its applications, Part 2 (Berkeley, Calif., 1983), pp. 461–466, Proc. Sympos. Pure Math. 45, Part 2, AMS, Providence, RI, (1986).
- [60] Wente H., The differential equation $\Delta x = 2Hx_u \wedge x_v$ with vanishing boundary values, *Proc. Amer. Math. Soc.* **50** 131–137 (1975),
- [61] Yang, Y., Solitons in field theory and nonlinear analysis. Springer Monographs in Mathematics. Springer-Verlag, New York, (2001).
- [62] Ye, R., How to blow infinitely large soap bubbles with a fixed boundary, Annal. H. Poincarè C8, 59–78 (1991).