

Do Quenched Spinor Condensates Thermalize?

Ryan Barnett,¹ Anatoli Polkovnikov,² and Mukund Vengalattore³

¹*Joint Quantum Institute and Condensed Matter Theory Center,*

Department of Physics, University of Maryland, College Park, MD 20742, USA

²*Department of Physics, Boston University, 590 Commonwealth Ave., Boston, MA 02215, USA and*

³*Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, NY 14853, USA*

(Dated: July 18, 2022)

Motivated by recent experiments, we consider the dynamics of spin-one spinor condensates after a quantum quench from the polar to ferromagnetic state from varying the quadratic Zeeman field, q . We apply the Truncated Wigner Approximation (TWA) to the spinor system with all spatial and spin degrees of freedom. For short times, we find full agreement with the linearized Bogoliubov analysis. For longer times, where the Bogoliubov theory fails, we find that the system reaches a quasi-steady state. We compute the Bogoliubov spectrum about the ferromagnetic state with finite q and show that the resulting finite temperature correlation functions grossly disagree with the full TWA results, thus indicating that the system does not thermalize even over very long time scales. We attribute this to the presence of topological defects (i.e. spin vortices) which prevent the relaxation to the ground state over experimentally relevant time scales.

Advances in the field of ultracold atomic gases have spurred great interest in the non-equilibrium dynamics of quantum many-body systems. The ability to engineer paradigmatic model Hamiltonians, the near perfect isolation from the environment and the experimentally accessible timescales of evolution have made it possible to address fundamental questions about the dynamics of closed, interacting quantum many-body systems.

Of particular interest in this context is the study of a ‘quench’ across a quantum phase transition. Here, one or more parameters of the Hamiltonian are changed rapidly resulting in a non-equilibrium evolution of the quantum system towards the establishment of long range order. This evolution is accompanied by a spatially inhomogeneous symmetry breaking and the formation of topological defects seeded by the quench. Accurate, time-resolved studies of such quenches are of fundamental importance to a wide range of non-equilibrium phase transitions.

Recently, an instance of such a quench across a phase transition has been experimentally realized with quantum degenerate spin-1 Bose gases of ⁸⁷Rb [1, 2]. At low magnetic fields, these multicomponent fluids are characterized by a contact interaction that favors a ferromagnetic phase. At large external magnetic fields, the quadratic Zeeman energy (QZE) dominates the interaction and favors a paramagnetic (‘polar’) phase. As shown in Fig(1), these two phases are separated by a continuous phase transition. In the experiment, *in situ* images of the spin textures following a quench of the degenerate gas into the ferromagnetic phase revealed the inhomogeneous growth of transversely magnetized domains accompanied by the sporadic observation of topological defects that were characterized as spin vortices.

Motivated by this experiment, here we consider the evolution of spin textures following a quench from the polar phase to the ferromagnetic phase. While the short-time growth of magnetization during such a quench has

been analyzed [3–11], here we consider the evolution over much longer periods and study the manner in which the spin degrees of freedom thermalize following the quench. Applying the truncated Wigner approximation (TWA) to this problem (for an overview of the TWA method and its applications, see Refs. [12, 13]), we analyze the long time dynamics of the spin degrees of freedom.

We find that unless the QZE is quenched to the immediate vicinity of the critical value corresponding to the transition to the ferromagnetic phase, the system reaches a slowly evolving quasi-steady state with exponentially decaying correlations. This correlation length very slowly (logarithmically) increases in time in a manner consistent with coarsening dynamics [14]. This coarsening can be understood as a result of annihilation of vortex anti-vortex pairs created during the initial spin dynamics following the quench. Such dynamics is somewhat reminiscent to the situation in normal superfluids, where one also finds long lived metastable states following a quench [15]. We also find and interpret previously unexplored physics of the dynamics of spinor condensates such as the emergence of longitudinal magnetization.

Following the experimental situation, we consider the parameter regime characteristic of $F = 1$ spinor condensates of ⁸⁷Rb. Also, like in the experiment, the gases are confined in a quasi-2D geometry wherein the spatial extent d_y of the condensate along one dimension is less than the spin healing length. This condition implies that the spin dynamics along this axis are effectively frozen. The starting point is the Hamiltonian

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_{\text{int}} \quad (1)$$

where the free Hamiltonian is

$$\mathcal{H}_0 = \int d^2r \Psi^\dagger \left(-\frac{\hbar^2}{2m} \nabla^2 + V + qf_z^2 \right) \Psi \quad (2)$$

and $\Psi = (\psi_1, \psi_0, \psi_{-1})^T$ is a three component spinor, V is the trapping potential, $f_{x,y,z}$ are the spin-one matrices,




FIG. 1. Zero-temperature phase diagram of a spin-1 Bose condensate indicating the transition from a polar phase to a ferromagnetic phase at low quadratic Zeeman energy (QZE). The dashed line indicates the direction of the quench from an initial QZE $q \gg 2|c_2|n$ to a final QZE $q = q_f$.

and q is the quadratic Zeeman shift (the linear Zeeman shift is excluded since the spin along the magnetic field is conserved). The interaction Hamiltonian is

$$\mathcal{H}_{\text{int}} = \int d^2r \left(\frac{1}{2} c_0 n^2 + \frac{1}{2} c_2 F^2 \right). \quad (3)$$

Here we have $n = \Psi^\dagger \Psi$, $\mathbf{F} = \Psi^\dagger \mathbf{f} \Psi$, and the parameters c_0 and c_2 can be expressed in terms of the s -wave scattering lengths as $c_0 = \frac{4\pi\hbar^2}{d_y 3m} (a_0 + 2a_2)$ and $c_2 = \frac{4\pi\hbar^2}{d_y 3m} (a_2 - a_0)$. Spinor gases of ^{87}Rb are characterized by the scattering lengths $a_0(a_2) = 101.8(100.4)$ Bohr radii respectively. Because $a_2 < a_0$, the spin dependent interaction in these spinor gases favors the ferromagnetic state. Since for these parameters $c_0 \gg |c_2|$, fluctuations in the total density are suppressed. Further, due to the weak spin interaction strength, the interparticle separation at typical densities is far smaller than the spin healing length, thereby suppressing the role of quantum depletion.

As illustrated in Fig. 1, for $q > q_0 \equiv 2|c_2|n_0$, the ground state is the polar state while decreasing q below q_0 , the ground state acquires a magnetic moment and eventually reaches the fully polarized ferromagnetic state at $q = 0$. Initially, we take $q = q_i$ to be the largest energy parameter in the Hamiltonian and quench to a final state $q = q_f$ where $0 \leq q_f < q_0$. For short times, the dynamics is expected to be well-described by expanding the Hamiltonian to quadratic order around the initial polar state [3–9]. Under this expansion the Hamiltonian becomes

$$\mathcal{H} \approx \sum_{\mathbf{k}} \left((\varepsilon_{\mathbf{k}} + c_2 n_0 + q_f) (\psi_{1,\mathbf{k}}^\dagger \psi_{1,\mathbf{k}} + \psi_{-1,\mathbf{k}}^\dagger \psi_{-1,\mathbf{k}}) + c_2 n_0 (\psi_{1,\mathbf{k}}^\dagger \psi_{-1,-\mathbf{k}}^\dagger + \psi_{-1,-\mathbf{k}} \psi_{1,\mathbf{k}}) \right). \quad (4)$$

where we have dropped the term describing the stiff density fluctuations. For simplicity, we have considered the system in the continuum, in the absence of the trapping potential V . In the ferromagnetic regime this quadratic Hamiltonian gives modes with imaginary frequencies, which correspond to unstable, exponentially growing in time modes [3].

To quantify the spin dynamics after the quench, it is natural to consider transverse and longitudinal magnetization correlation functions

$$G_{\perp}(\mathbf{r}, t) = \frac{1}{n_0^2} \langle : F_{+}(\mathbf{r}) F_{-}(0) : \rangle \quad (5)$$

$$G_z(\mathbf{r}, t) = \frac{1}{n_0^2} \langle : F_z(\mathbf{r}) F_z(0) : \rangle \quad (6)$$

and the concomitant gain functions, $G_{\perp}(t)$ and $G_z(t)$, which are the above evaluated for $\mathbf{r} = 0$. For $q_0 t \gg 1$ and $q_f = 0$, the above correlations within the linearized theory give

$$G_{\perp}(t) = \frac{1}{\sqrt{8\pi q_0 t}} \frac{1}{n_0 \xi_s^2} e^{q_0 t} \quad (7)$$

$$G_z(t) = \frac{1}{64\pi q_0^2 t^2} \frac{1}{(n_0 \xi_s^2)^2} e^{2q_0 t}. \quad (8)$$

where $\xi_s = \sqrt{\hbar^2/q_0 m}$ is the spin coherence length. Interestingly, the longitudinal gain grows with twice the exponent of the transverse gain. However, the longitudinal magnetization is suppressed by a large factor $1/(n_0 \xi_s^2) \sim 10^{-4}$ for the experimental parameters of Ref. [1]. Thus, $G_z(t)$ remains much smaller than $G_{\perp}(t)$ up to the times where the latter saturates $G_{\perp} \approx 1$ and the linearized theory does not work (see Fig. 2).

As seen from the exponential growth of the gain functions, the above theory clearly fails once the transverse magnetization is of order unity. To gain a more complete understanding we turn to truncated Wigner simulations. This technique involves propagating the full spinor Gross-Pitaevskii Equations (GPEs) of motion obtained by taking the classical limit of Eq. (1) seeded with random initial conditions for canonical degrees of freedom distributed according to the Wigner transform of the initial state. For our case the initial state is taken to be the vacuum of Bogoliubov quasiparticles for a theory expanded about the polar state. For the limiting case of $q_i = \infty$ this has the particularly simple form: $W(\Psi) \approx \delta(|\psi_{0,0}|^2 - n_0) \prod_{\mathbf{k}} 8 \exp[-2 \sum_{\alpha} |\psi_{\alpha,\mathbf{k}}|^2]$, where the factor of 8 ensures the correct normalization of the Wigner function [13, 16].

Shown in Fig. 2 are the TWA results for the transverse and longitudinal gain functions in the absence of a trap compared with the analytic results based upon the above linearized theory. At short evolution times following the quench, the simulations predict the exponential growth of both the transverse and longitudinal magnetization densities in a manner that is in excellent agreement with the linearized theory. During this period, the spin textures are characterized by ferromagnetic domains that are predominantly oriented in the transverse plane.

We now move on to discuss the long-time dynamics of the quenched spinor condensate. As is seen in the gain functions in Fig. 2, the magnetization of the system




FIG. 2. The growth of transverse (red) and longitudinal (blue) magnetization following a quench to $q_f = 0$. Inset: The characteristic time constants for the growth of transverse and longitudinal magnetization densities versus the 2D interaction strength $|c_2 n|$. The dashed lines indicate the predictions of the linearized Bogoliubov treatment for the respective parameters.

eventually saturates and reaches a steady state. This motivates one to consider the prospect of thermalization in the system. More specifically, under the assumption of thermalization, the system will evolve to the ground state at $q = q_f$, with the excess energy accounted for by a superposition of elementary excitations about this configuration. We therefore will consider the theory about the state at $q = q_f$.

We define the heating (excess energy) of the system as

$$Q = \langle \mathcal{H} \rangle_i - \langle \mathcal{H} \rangle_{\text{gs}} \quad (9)$$

where the above expectation values are evaluated for the initial state immediately after the quench and the ground state for $q = q_f$. To extract the equilibrium temperature this heating can be compared with the thermal energy of the system. It is convenient to use the following parameterization of the spinor

$$\Psi = \sqrt{n} e^{i\alpha} \left(\sin \eta \cos \phi e^{i\xi}, \cos \eta e^{i\gamma}, \sin \eta \sin \phi e^{-i\xi} \right)^T. \quad (10)$$

With this, one can see that for $0 < q_f/q_0 < 1$ the classical energy is minimized for $\phi = \pi/4$, $\gamma = 0$, and $\cos(2\eta) = q_f/q_0$ [16]. One then obtains that the above heating is

$$Q = \frac{1}{4} n_0 q_0 (1 - q_f/q_0)^2. \quad (11)$$

Note that the above is based on classical energy differences. In addition to this classical contribution to the heating there is an additional quantum correction coming from the zero point fluctuations of the condensate. However this contribution is suppressed by a large factor $\sqrt{n_0 \xi_s^2} \sim 100$ and thus is not important.

The thermal energy of the final state is found from

$$Q = U(T) \equiv \text{Tr} \left(e^{-\mathcal{H}/T} \mathcal{H} \right) / Z \quad (12)$$

where $Z = \text{Tr} \left(e^{-\mathcal{H}/T} \right)$ is the partition function and the ground state energy of the system is set to be zero. To evaluate the above we develop a second Bogoliubov theory by expanding the Hamiltonian to quadratic order about the minimum at $q = q_f$. Note that within the chosen parametrization one automatically avoids subtleties related to the lack of a long range order in two dimensions at finite temperatures. The consistency of such a quadratic theory is then self-consistently checked by verifying that the resulting thermal depletion is small for the computed temperature.

The full Bogoliubov analysis of the spectrum linearized around the ferromagnetic minimum is rather cumbersome because of the phonon-magnon coupling [17] (which vanishes for the special points $q_f = 0$ and $q_f = q_0$). However, ^{87}Rb has a natural separation of energy scales since $c_0 \gg |c_2|$ which inhibits density fluctuations. This significantly simplifies the Bogoliubov analysis and effectively results in decoupling of the stiff density fluctuation mode related to the coupling c_0 and the spin modes. In this limit one finds the simplified spectrum consisting of two modes with the dispersion [16]:

$$\begin{aligned} \omega_{\mathbf{k}}^{(\perp)} &= \sqrt{\varepsilon_{\mathbf{k}}(\varepsilon_{\mathbf{k}} + q_0)} \\ \omega_{\mathbf{k}}^{(z)} &= \sqrt{(\varepsilon_{\mathbf{k}} + q_0) \left(\varepsilon_{\mathbf{k}} + \frac{q_0^2 - q_f^2}{q_0} \right)} \end{aligned} \quad (13)$$

where $\varepsilon_{\mathbf{k}}$ is the free particle dispersion. Note that these reduce to the known dispersions in the limits of $q_f = 0$ and $q_f = q_0$ [18, 19]. Furthermore, the large c_0 limit of the full dispersion [17] agrees with Eq. (13). These expressions are used to compute the thermal energy of the system. It is straightforward to verify that unless q_f is very close to q_0 , $1 - q_f/q_0 \gg 1/\sqrt{n_0 \xi_s^2}$, the dominant contribution to the thermal energy $U(T)$ comes from the quadratic dispersion of these modes: $\omega_{\mathbf{k}}^{(\perp)} \approx \omega_{\mathbf{k}}^{(z)} \approx \varepsilon_{\mathbf{k}}$. This can be justified *a posteriori* by comparing the temperature with q_0 . With these conditions, $U(T) \approx N\pi T^2/6(n_0 \xi_s^2)q_0$. Finally, equating this to the heating, one finds

$$T = \sqrt{\frac{3}{2\pi} n_0 \xi_s^2 (q_0 - q_f)}. \quad (14)$$

To test the hypothesis of thermalization, we use the developed quadratic theory to evaluate finite temperature correlation functions (evaluated at temperature Eq. (14)) and compare with long-time TWA results. The most natural correlation function to consider is G_{\perp} defined in Eq. (5). The gapless mode, Eq. (13), which leads to algebraic decay of the correlation function in two dimensions has the dominant contribution. Within the quadratic Bogoliubov theory, the correlation function is found to be [16]

$$G_{\perp}(\mathbf{r}) = \left[1 - (q_f/q_0)^2 \right] r^{-\alpha} \quad (15)$$




FIG. 3. The transverse magnetization densities for short (a) and long (b) times for $q_f = 0$. The amplitude and orientation of the transverse magnetization are indicated by the brightness and hue (color wheel shown). (c) The spin correlation function for these same times. While the correlation function at short times is in excellent agreement with Bogoliubov theory, the decay of this function at long evolution times is much more rapid than predicted by Eq. (15)

with exponent $\alpha = \sqrt{6/\pi^3} \cdot \sqrt{1/n_0\xi_s^2}$. Because of the dependence of the exponent α on the small parameter $n_0\xi_s^2$, we see that, assuming equilibrium, $G_\perp(\mathbf{r})$ will not decay by any appreciable amount over the relevant length scales of the condensate, which are typically of the order of tens or hundreds of the spin coherence length ξ_s . For the same reason in equilibrium the system should not have any vortices.

We now compare these predictions to the results of the TWA simulations for long times. As shown in Fig. 3(d), the correlation function G_\perp before saturation for short times has the functional dependence of a Bessel function $J_0(r/\xi_s)$ as previously predicted [3]. For long times, the correlation function reaches a steady state and decays by over a factor of five from its $r = 0$ value for a system size on the order of 50 spin coherence lengths. Such decay is qualitatively incompatible with the theory developed by assuming thermalization. We thus conclude that spinor condensates do not thermalize at appreciable time scales but rather reach some quasi-steady (prethermalized) regime which evolves anomalously slowly in time. Such prethermalized phases were suggested earlier in the contexts of weakly interacting fermions [20, 21], scalar superfluids [15], and dynamics of the early Universe [22]. What is perhaps unexpected is that in this system anomalously slow relaxation occurs at very high levels of the heating in the system. One possible explana-

tion for this is the presence of topological defects which persist at all time scales considered. These defects apparently have much longer life times than vortices in scalar superfluids under similar excitation levels.

In conclusion, we have examined the long time evolution of quantum degenerate spinor gases following a quench to a ferromagnetic phase. Assuming a thermalized final state, we find that the finite temperature spin correlations are characterized by an algebraic decay over length scales much larger than the relevant length scales of the condensate. In distinct contrast, numerical simulations based on the truncated Wigner approximation indicate a rapidly decaying correlation function even at extremely long evolution times. This inconsistency leads one to the conclusion that the quenched spinor condensates will not thermalize over experimentally relevant timescales. The effect of an incoherent thermal bath in contact with the spinor condensate or the role of dipolar interactions [2, 23] in the long time evolution has yet to be examined.

For valuable discussions, we thank E. Altman, A. Lamacraft, L. Mathey, J. D. Sau, and A. Turner. M. V. thanks L. M. Aycocock and S. Chakram for valuable discussions and critical comments on the manuscript. We gratefully acknowledge financial support from the JQI Physics Frontier Center and the Sherman Fairchild Foundation (R.B.), AFOSR FA9550-10-1-0110 and the Sloan Foundation (A.P.), and Cornell University (M.V.). The authors also thank the Aspen Center for Physics for hospitality, where a part of this work was completed.

-
- [1] L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn, *Nature* **443**, 312 (2006).
 - [2] M. Vengalattore, S. R. Leslie, J. Guzman, and D. M. Stamper-Kurn, *Phys. Rev. Lett.* **100**, 170403 (2008).
 - [3] A. Lamacraft, *Phys. Rev. Lett.* **98**, 160404 (2007).
 - [4] H. Saito, Y. Kawaguchi, and M. Ueda, *Phys. Rev. A* **76**, 043613 (2007).
 - [5] B. Damski and W. H. Zurek, *Phys. Rev. Lett.* **99**, 130402 (2007).
 - [6] M. Uhlmann, R. Schützhold, and U. R. Fischer, *Phys. Rev. Lett.* **99**, 120407 (2007).
 - [7] R. W. Cherng, V. Gritsev, D. M. Stamper-Kurn, and E. Demler, *Phys. Rev. Lett.* **100**, 180404 (2008).
 - [8] M. Baraban, H. F. Song, S. M. Girvin, and L. I. Glazman, *Phys. Rev. A* **78**, 033609 (2008).
 - [9] C. Klempt, O. Topic, G. Gebreyesus, M. Scherer, T. Henninger, P. Hyllus, W. Ertmer, L. Santos, and J. Arlt, *Phys. Rev. Lett.* **103**, 195302 (2009).
 - [10] H. Saito, Y. Kawaguchi, and M. Ueda, *Phys. Rev. A* **75**, 013621 (2007).
 - [11] J. D. Sau, S. R. Leslie, D. M. Stamper-Kurn, and M. L. Cohen, *Phys. Rev. A* **80**, 023622 (2009).
 - [12] P. B. Blakie, A. S. Bradley, M. J. Davis, R. J. Ballagh, and C. W. Gardiner, *Adv. in Phys.* **57**, 353 (2008).
 - [13] A. Polkovnikov, *Annals of Physics* **325**, 1790 (2010).

- [14] A. J. Bray, *Adv. in Phys.* **43**, 357 (1994).
- [15] L. Mathey and A. Polkovnikov, *Phys. Rev. A* **81**, 033605 (2010).
- [16] R. Barnett, A. Polkovnikov, and M. Vengalattore (Unpublished); See supplementary material for more information.
- [17] S. Uchino, M. Kobayashi, and M. Ueda, *Phys. Rev. A* **81**, 063632 (2010).
- [18] T.-L. Ho, *Phys. Rev. Lett.* **81**, 742 (1998).
- [19] T. Ohmi and K. Machida, *J. Phys. Soc. Japan* **67**, 1822 (1998).
- [20] M. Moeckel and S. Kehrein, *New J. of Phys.* **12**, 055016 (2010).
- [21] M. Eckstein, M. Kollar, and P. Werner, *Phys. Rev. Lett.* **103**, 056403 (2009).
- [22] J. Berges, S. Borsányi, and C. Wetterich, *Phys. Rev. Lett.* **93**, 142002 (2004).
- [23] M. Vengalattore, J. Guzman, S. Leslie, and D. Stamper-Kurn, *Phys. Rev. A* **81**, 053612 (2010).