

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

*Evidence for the Dominance of Indirect Effects in 50 Trophically-Based Ecosystem Networks*

Andria K. Salas<sup>a</sup> and Stuart R. Borrett<sup>b</sup>

<sup>a</sup>Corresponding author  
601 S. College Road  
Center for Marine Science and Dept. of Biology and Marine Biology  
University of North Carolina at Wilmington  
Wilmington, NC 28403  
616.334.4864  
aks2515@gmail.com

<sup>b</sup>601 S. College Road  
Dept. of Biology and Marine Biology and Center for Marine Science  
University of North Carolina at Wilmington  
Wilmington, NC 28403  
910.962.2411  
borretts@uncw.edu

Key words: food webs, network analysis, microbial food web, energy flow, environ analysis, cycling

45 **Abstract**

46 Indirect effects are powerful influences in ecosystems that may maintain species diversity  
47 and alter apparent relationships between species in surprising ways. Here, we applied Network  
48 Environ Analysis to 50 empirically-based trophic ecosystem models to test the hypothesis that  
49 indirect flows dominate direct flows in ecosystem networks. Further, we used Monte Carlo  
50 based perturbations to investigate the robustness of these results to potential error in the  
51 underlying data. To explain our findings, we further investigated the importance of the microbial  
52 food web in recycling energy-matter using components of the Finn Cycling Index and analysis of  
53 Environ Centrality. We found that indirect flows dominate direct flows in 37/50 (74.0%)  
54 models. This increases to 31/35 (88.5%) models when we consider only models that have  
55 cycling structure and a representation of the microbial food web. The uncertainty analysis  
56 reveals that there is less error in the *I/D* values than the  $\pm 5\%$  error introduced into the models,  
57 suggesting the results are robust to uncertainty. Our results show that the microbial food web  
58 mediates a substantial percentage of cycling in some systems (median = 30.2%), but its role is  
59 highly variable in these models, in agreement with the literature. Our results, combined with  
60 previous work, strongly suggest that indirect effects are dominant components of activity in  
61 ecosystems.

62 **1 Introduction**

63 What is an example of an isolated individual? Given that metabolism and reproduction  
64 are two criteria for living entities, no organism can be isolated (Jørgensen et al., 1992, 1999;  
65 Wilkinsin, 2006). Heterotrophs must consume carbon sources and autotrophs utilize nutrients  
66 within the soil or water that were made available by other living things. Sexual reproduction  
67 typically requires another individual of the same species. Thus, at a fundamental level, organisms  
68 must interact with their environment to grow, reach maturity, and reproduce. From these most  
69 basic interactions necessary for life develops the complex ecological community. In this paper  
70 we elucidated a part of ecosystem complexity by testing the generality of the systems ecology  
71 hypothesis that indirect effects tend to dominate direct effects in ecosystem interaction networks.

72 Indirect effects can have powerful influences on ecosystem functioning (e.g. Abrams et  
73 al., 1996; Higashi and Patten, 1989; Menge, 1997; Patten, 1983; Wootton, 1993, 1994; Yodzis,  
74 1988). For example, they have been found to help maintain diversity (Bever, 1999, 2002) and  
75 contribute to the success of a particular habitat patch (Diekötter et al., 2007). Indirect effects  
76 also can produce surprising relationships between species (Bondavalli and Ulanowicz, 1999),  
77 and can result in overall positive interactions despite local negative direct interactions (Fath and  
78 Patten, 1998; Patten, 1991; Ulanowicz and Puccia, 1990). In addition, indirect effects can  
79 develop rapidly in ecosystems (Borrett et al., 2010; Menge, 1997). Given these diverse impacts,  
80 it is clear that understanding indirect effects is necessary to understand the processes that create,  
81 constrain, and sustain ecosystems.

82 This understanding is challenging to achieve given the difficulty in empirically studying  
83 indirect effects, necessitating the need for alternative approaches such as Network Environ  
84 Analysis (NEA). NEA is applied to ecosystem models of energy-matter flow and can be used to  
85 study indirect flows of energy (e.g. Fath and Borrett, 2006; Fath and Patten, 1999; Patten et al.,  
86 1976). The ecosystem network models we used in this work are simplified representations of  
87 ecosystems that trace the path of energy-matter flux through the system. In these models, nodes  
88 represent species, groups of species, or abiotic resource pools while the directed links represent  
89 the flow of energy-matter. NEA lets us partition these flows into direct and indirect  
90 components. Indirect flows are those in which a species receives energy indirectly from another

91 species, such as when a polar bear receives energy from krill by consuming penguins that  
92 directly ate the krill. These indirect flows were the basis of our investigations.

93 We used NEA to test the hypothesis that the amount of energy flowing over indirect  
94 pathways tends to exceed the amount of energy flowing over direct pathways in ecosystems, a  
95 hypothesis that emerged in early applications of NEA (Patten, 1983, 1984, 1985). Higashi and  
96 Patten (1986, 1989) argued that the dominance of indirect flows over direct flows should be a  
97 general phenomenon by showing algebraically that indirect flow tends to increase with system  
98 size, connectance, cycling, and magnitude of direct flows. The hypotheses concerning system  
99 size and cycling are further supported by an analysis of large hypothetical ecosystem models  
100 built from a community assembly algorithm (Fath, 2004). These studies suggest that the  
101 dominance of indirect flows has the potential to occur in real ecosystems, but few studies report  
102 this phenomenon in empirically derived models (but see e.g. Borrett et al., 2006; Patten, 1983).  
103 Thus, the generality of the hypothesis in models of real systems remains to be tested. Further,  
104 little is known regarding the sensitivity of the results to uncertainty in the data used for network  
105 model construction (e.g., Borrett and Osidele, 2009; Kaufmann and Borrett, 2010).

106 There are many different types of indirect effects (Strauss, 1991 and Wootton, 1993;  
107 1994), and NEA can capture those that are and are affected by energy–matter transactions.  
108 Indirect flows are a subset of indirect effects, but the former has the ability to reflect a wider  
109 range of indirect effects than feeding relationships. Indirect effects have the ability to increase,  
110 decrease, or have no affect on indirect flows. For example, a meta-analysis by Preisser et al.  
111 (2005) suggests that the act of intimidation of predators on prey results in reduced resource  
112 acquisition by the prey species. The larger the intimidation effect of predators on prey, the  
113 smaller the flows of energy to the prey species. While NEA does not capture all types of indirect  
114 effects, the consequences of many are reflected in energy transfers. Therefore, by comparing the  
115 intensity of indirect flows to direct flows, we are estimating the magnitude of indirect effects.  
116 More importantly, this estimation should tend to be conservative given the possibility of indirect  
117 effects that minimize the strength of indirect flows.

118 To understand the variability of the relative importance of indirect effects, we further  
119 investigated the species essential to energy–matter cycling. The microbial food web plays an  
120 important role in mediating energy–matter cycling. Much of the carbon fixed by primary  
121 producers is released as particulate and dissolved organic matter. Dissolved organic carbon  
122 (DOC) is a significant source of carbon for bacteria (Larsson and Hagström, 1982) and it is this  
123 relationship that is largely the basis of the microbial food web (Hart et al., 2000), which includes  
124 benthic and pelagic bacteria, ciliates, flagellates, and microzooplankton. Particulate organic  
125 carbon (POC) from plant material and animal wastes is also a medium for microbial colonization  
126 (Fenchel, 1970). The functioning of the microbial web and loop can reintroduce the energy-  
127 matter in DOC and other detrital groups to the larger metazoan food web (Azam et al., 1983;  
128 Pomeroy, 1974). Microzooplankton also are important to the flow of energy in the system as  
129 they make available the energy in nano- and picoplankton to larger consumers (Sherr and Sherr,  
130 1988). The importance of detritus and the microbial food web has been recognized for a variety  
131 of aquatic systems, from rivers to lakes to coastal systems (Mann, 1988). However, the amount  
132 of primary production that actually passes through the microbial food web as a conduit to the  
133 metazoan food web is highly variable (as in Hart et al., 2000) and system dependent (Hart et al.,  
134 2000; Sommer et al., 2002). In summary, the microbial food web should be considered when  
135 investigating the pathways of carbon flow given its importance to the recycling process.

136 In this work we specifically tested four hypotheses. First, we tested the generality of the  
137 dominance of indirect effects hypothesis in a set of 50 empirically-derived ecosystem models.  
138 Second, we tested the idea that indirect flows through an ecosystem network tend to increase  
139 with system size, connectance, cycling, and direct flow. Third, we tested the robustness of the  
140 analytical results to potential data uncertainty in the network models using an uncertainty  
141 analysis. Finally, given the generally important role of the microbial food web in ecosystem  
142 carbon flow, we tested the hypothesis that the representation and importance of this subweb in  
143 the models helps explain why indirect flows do or do not dominate in these ecosystems.

## 144 **2 Methods and Materials**

### 145 **2.1 Model Database**

146 To test the hypothesis that indirect flows are dominant in ecosystems, we analyzed 50  
147 ecosystem models using NEA. To build our database of empirically derived ecosystem models,  
148 we started with Ulanowicz's data set (<http://www.cbl.umces.edu/~ulan/>) and then continued to  
149 search for models published in the literature. For this study, we selected ecosystems models that  
150 focused on trophic interactions that typically trace the movement of carbon or energy. We  
151 excluded models that might be considered more biogeochemically based (see Christian, et al.  
152 1996; Baird, et al. 2008; and Borrett, 2010 for more on this distinction). We also selected  
153 models that were at least minimally empirically based. These models were designed for a  
154 specific system and parameterized with some information gathered from the field. However,  
155 some literature values from related systems or best approximations may also have been used.  
156 Thus, the models range in the extent to which they were empirically derived, as some are largely  
157 field based and would be considered more empirical than those that are based solely on literature.  
158 Table 1 lists the ecosystem network models used in this study and describes some of the key  
159 features of the models.

160 This collection of 50 network models represents 35 distinct mostly freshwater and marine  
161 systems. Some of the models represent the same system at different times of the year or under  
162 different ecological conditions. Our database includes 18 of 48 models found in Ulanowicz's  
163 collection. Many of the models were initially reported at steady-state (inputs = outputs), but we  
164 balanced 15/50 of the models using the AVG2 algorithm (Allesina and Bondavalli, 2003) to  
165 meet the assumptions of NEA (see next section). These are the same models used by Borrett and  
166 Salas (2010) to test the resource homogenization systems ecology hypotheses.

### 167 **2.2 Network Environ Analysis**

#### 168 **2.2.1 Network Model Information**

169 We applied NEA to all models using a modified version of the NEA.m MATLAB®  
170 function (Fath and Borrett, 2006). Here we introduce the analysis using a hypothetical example  
171 model of an oversimplified ecosystem network consisting of detritus, bacteria, amphipods, and  
172 blue crabs (Figure 1a). In this example, the number of nodes,  $n$ , in the network is 4. The  
173 adjacency matrix ( $\mathbf{A}_{n \times n} = a_{ij}$ ) captures the structural information of the network. When there is a  
174 flow from  $j$  to  $i$ ,  $a_{ij} = 1$ , and  $a_{ij} = 0$  otherwise. Note that NEA uses a column to row orientation.  
175 In our example, bacteria decompose detritus, resulting in a 1 in the matrix at  $a_{21}$ . The amount of  
176 energy-matter flowing between the nodes is represented in the  $\mathbf{F}_{n \times n} = f_{ij}$  matrix (Figure 1a), in  
177 which the 1's of the adjacency matrix are replaced by the estimated flow values with general  
178 units of  $\text{M L}^{-2 \text{ or } -3} \text{T}^{-1}$  in mass, length, and time dimensions. Transfer of energy across the system

179 boundary is represented in  $\bar{z}$  and  $\bar{y}$ , which are vectors of the boundary inputs and outputs,  
 180 respectively (Figure 1a). The former includes processes such as immigration or transport of  
 181 organic material into the system. The latter includes processes such as fishing, emigration, and  
 182 transport of organic material out of the system. Also included in  $\bar{y}$  is respiration from all living  
 183 nodes. With this system information, we can begin our analysis.

### 184 2.2.2 Throughflow Analysis

185 Node throughflow is the amount of energy or matter that flows through each node. This  
 186 can be calculated from an input  $T_i^{in} = \sum_{j=1}^n f_{ij} + z_i$  or output  $T_j^{out} = \sum_{i=1}^n f_{ij} + y_j$  perspective.  $T^{in}$  sums  
 187 all flows into node  $i$  from nodes  $j$ , and adds to this quantity the boundary inputs into node  $i$ ,  
 188 while  $T^{out}$  sums all flows that exit node  $j$  to go to nodes  $i$ , in addition to the boundary outputs  
 189 from node  $j$ . At steady state,  $T^{in} = T^{out} = T = (T_j)$  where  $T_j$  is the total amount of energy entering  
 190 or exiting each node  $j$ . Total system throughflow ( $TST = \sum T_j$ ) is the total amount of energy or  
 191 matter that flows through the system, and its meaning can be likened to a gross domestic product  
 192 (GDP). GDP is the total monetary flow of a country's economy, and TST is the total substance  
 193 flow in an ecosystem.

194 Next, we determine the direct flow intensity,  $\mathbf{G} = (g_{ij})$ , between any two nodes. Using an  
 195 output perspective, we divide the values in each row  $i$  of the  $\mathbf{F}$  matrix by node throughflow,  
 196  $\mathbf{G} = (g_{ij}) = f_{ij} / T_j$ . Thus,  $g_{ij}$  is the percentage of each node's throughflow that goes to each  
 197 node with which it directly interacts. For example, in Figure 1b 33% of bacteria's energy flows  
 198 to detritus.

199 Both  $\mathbf{A}$  and  $\mathbf{G}$  can be raised to exponents to examine indirect pathways and flow  
 200 intensities. The matrix  $\mathbf{A}^2$ , for example, gives the numbers of pathways of length two ( $m = 2$ )  
 201 that exist between all node pairs. The matrix  $\mathbf{G}^2$  gives the flow intensities between the node pairs  
 202 over these pathways of length two. For example, there is one indirect pathway of length 2 from  
 203 amphipods to bacteria (via detritus), and 33% of the amphipod's energy flows to bacteria via this  
 204 indirect pathway (Figure 1b). The same interpretation can be given to  $\mathbf{A}^m$  and  $\mathbf{G}^m$  for any  $m$ .  
 205 Thus, the flow intensity between the nodes in the network over all indirect path lengths ( $m > 2$ ) is  
 206 equal to  $\sum_{m=2}^{\infty} \mathbf{G}^m$ .

207 If the system graph is a single strongly connected component such that each node is  
 208 reachable from every other node by a pathway of some length, then the  $\mathbf{G}^m$  matrix will be  
 209 entirely composed of non-zero numbers by some value of  $m$  (see Borrett et al., 2007, 2010).  
 210 This contrasts with the direct flow intensity matrix  $\mathbf{G}$ , which in almost all cases will have many  
 211 zeros as it is unlikely there are direct connections between all nodes. An analysis of pathways  
 212 through the  $\mathbf{A}$  and  $\mathbf{G}$  matrices with increasing  $m$  can be found in more detail in Patten (1985). In  
 213 our example (Figure 1b), we see that  $\mathbf{G}^2$  has 8 zeros, versus 10 in the  $\mathbf{G}$  matrix, and  $\mathbf{G}^{10}$  has no  
 214 zero values. This is significant because it reveals that all nodes are connected indirectly via  
 215 indirect paths. Even though the quantity of energy flowing over paths of increasing length tends  
 216 to decrease due to dissipation, longer paths are still being utilized and may significantly  
 217 contribute to TST (Borrett et al. 2010). Each node receives some contribution to its throughflow  
 218 from each and every other node from which it is reachable, further suggesting that indirect flows

219 are an important consideration. The  $\mathbf{G}^m$  matrix series can be used to determine the integral or  
 220 total flow intensity, which is captured in the matrix  $\mathbf{N}$ :

$$221 \quad \mathbf{N} \equiv \sum_{m=0}^{\infty} \mathbf{G}^m = \underbrace{\mathbf{G}^0}_{\text{Boundary}} + \underbrace{\mathbf{G}^1}_{\text{Direct}} + \underbrace{\mathbf{G}^2 + \mathbf{G}^3 + \dots + \mathbf{G}^m + \dots}_{\text{Indirect}} \quad (1)$$

222 The matrix identity,  $\mathbf{G}^0 = \mathbf{I}$ , is an  $n \times n$  matrix of zeros with ones along the principle diagonal  
 223 and is used to map in the boundary inputs,  $\mathbf{G}^1$  is the direct flow intensity, and  $\mathbf{G}^m$  is the indirect  
 224 flow intensity over paths of length  $m$ . Each element in  $\mathbf{N}$  represents the total flow intensity  
 225 between the nodes represented by that element's row and column. Multiplication by  $\bar{z}$  generates  
 226 the realized flow values (Borrett et al. 2006; Borrett and Freeze, in press). These values can be  
 227 summed to return the total system throughflow as

$$228 \quad \begin{aligned} \sum T &= \sum \mathbf{N}\bar{z} \\ &= \underbrace{\sum \mathbf{G}^0\bar{z}}_{\text{Boundary}} + \underbrace{\sum \mathbf{G}^1\bar{z}}_{\text{Direct}} + \underbrace{\sum (\mathbf{G}^2 + \mathbf{G}^3 + \dots + \mathbf{G}^m \dots)\bar{z}}_{\text{Indirect}} \end{aligned} \quad (2)$$

229 The power series in equation 1 should be convergent because ecosystems are thermodynamically  
 230 open (Jørgensen et al., 1999) and energy-matter is lost at each transfer. Thus, we can determine  
 231  $\mathbf{N}$  exactly using the identity  $\mathbf{N} = (\mathbf{I} - \mathbf{G})^{-1}$ .

232 We can use equation 2 to determine the amount of indirect flow relative to direct flow as

$$233 \quad I/D = \frac{\sum \mathbf{N}\bar{z} - \sum \mathbf{G}\bar{z} - \sum \mathbf{I}\bar{z}}{\sum \mathbf{G}\bar{z}} \quad (3)$$

234 If the  $I/D$  ratio is greater than one, it indicates that the magnitude of indirect flow is greater than  
 235 direct flow. This would be evidence to support the dominance of indirect effects hypothesis.

236 If we divide both sides of equation (2) by  $\sum T$ , we obtain the relative proportions of TST  
 237 derived from boundary, direct, and indirect flows. Indirect flow intensity (IFI) is the proportion  
 238 of total system throughflow derived from indirect pathways and direct flow intensity (DFI) is the  
 239 proportion of total system throughflow due to direct pathways. Specifically, IFI and DFI are  
 240 calculated as

$$241 \quad \text{IFI} = \frac{\sum ((\mathbf{N} - \mathbf{I} - \mathbf{G})\bar{z})}{\sum T}, \text{ and} \quad (4)$$

$$242 \quad \text{DFI} = \frac{\sum (\mathbf{G}\bar{z})}{\sum T} \quad (5)$$

243 To test the generality of the dominance of indirect flows (hypothesis 1), we calculated  $I/D$  for the  
 244 models in our data set. Further, we used a linear regression and Spearman's rank correlation test  
 245 to test the hypothesized relationship between IFI and recycling, DFI, system size, and  
 246 connectance ( $C = L/n^2$ , where  $L$  is the number of links in the network).

### 247 2.3 Contributions of nodes to system flux

248 To test the hypothesized importance of the microbial food web in energy-matter cycling  
 249 (hypothesis 4), we used two approaches: a partitioning of recycled matter into the fraction  
 250 derived from the microbial food web and other compartments, and Environ Centrality (EC).

251 2.3.1 *Recycling*

252 In contrast to the decomposition of TST described in section 2.2.2, Finn (1976)  
 253 introduced an alternative decomposition. He distinguished between flow from cyclic and acyclic  
 254 pathways:  $TST = cyclic + acyclic$ . Cyclic flow is determined as:

$$255 \quad Cyclic = \sum_{j=1}^n \left( \frac{n_{jj} - 1}{n_{jj}} \right) T_j \quad (6)$$

256 He then constructed what is known as the Finn Cycling Index (FCI) as:  $FCI = cyclic/TST$ .

257 Here, we subsetted the FCI to investigate the contribution of microbial food web to  
 258 system recycling. We first identified the nodes in each model that represent microbial  
 259 organisms. For convenience, we will conceptually reorder our network so that the first 1, 2, ...,  $k$   
 260 compartments are associated with the microbial food web. Then, we can partition the traditional  
 261 FCI (Finn 1976; 1980) into the fraction mediated by the microbial food web (MFW) and the  
 262 remainder of the nodes (other):

$$263 \quad FCI = FCI_{MFW} + FCI_{other} . \quad (7)$$

264 Therefore, the percentage of the system's cycling that is mediated by the microbial food web can  
 265 be calculated by:

$$266 \quad \%FCI_{MFW} = \frac{FCI_{MFW}}{FCI} = \frac{\sum_{j=1}^k \left( \frac{n_{jj} - 1}{n_{jj}} * T_j \right)}{\sum_{j=k+1}^n \left( \frac{n_{jj} - 1}{n_{jj}} * T_j \right)} \quad (8)$$

267 2.3.2 *Environ Centrality*

268 To further characterize the role of specific nodes, selected network models were analyzed  
 269 using environ centrality (EC; Fann, 2009). Centrality is a social science concept that “describes  
 270 the location of individuals in terms of how close they are to the center of action in the network”  
 271 (Hanneman and Riddle, 2005). Borgatti and Everett (2006) and Wasserman and Faust (1994)  
 272 provide more detail on this concept. Fann (2009) introduced an environ based centrality metric  
 273 that determines the role of each node in generating the total system throughflow. Following  
 274 Estrada (2010), it is a globally weighted walks-based metric most similar to eigenvector  
 275 centrality. This metric can be calculated for the input ( $EC^{in}$ ) or output perspective ( $EC^{out}$ ), and  
 276 the average of the two orientations was used here (average environ centrality,  $AEC$ ).

$$277 \quad EC^{in} = \frac{\sum_{j=1}^n n_{ij}}{\sum_{i=1}^n \sum_{j=1}^n n_{ij}} \quad (9)$$

$$278 \quad EC^{out} = \frac{\sum_{i=1}^n n_{ij}}{\sum_{i=1}^n \sum_{j=1}^n n_{ij}} \quad (10)$$

$$279 \quad AEC = \frac{EC_j^{in} + EC_j^{out}}{2} . \quad (11)$$

280 Thus, *AEC* gives the average proportional contribution that each node is making to total flow  
281 intensity. This metric indicates the relative significance of the nodes in generating system  
282 activity. We calculated this metric for the St. Marks seagrass system to better explain our  
283 results.

## 284 **2.4 Analytical Robustness**

285 We examined the reliability of our analytical results in two distinct ways; one addresses  
286 the potential problems caused by decisions made in the construction of the models and the other  
287 addresses the reliability of the models themselves to be reasonable representations of the  
288 underlying systems they represent.

### 289 *2.4.1 Model Structure*

290 Ecosystem network models range in their ability to reflect the actual underlying system  
291 and general overall quality. We initially choose to include all models available that have an  
292 empirical basis because the dominance of indirect flows is hypothesized to hold true for most  
293 networks. Yet all the models do not include the characteristics necessary to represent carbon  
294 flow that we believe are critical to be present for our work. Therefore, we did two analyses, one  
295 on the set of all models (Model Set 1) and another on those that meet more stringent criteria  
296 (Model Set 2). These criteria for Model Set 2 were that the model must 1) have network  
297 architecture that allows cycling, and 2) include species that are critical in mediating the process  
298 of energy cycling. These characteristics are necessary because cycles are known to be important  
299 components of ecosystem structure and function (Azam et al. 1986; Fath and Halmes 2007;  
300 Patten 1985). Species that must be included for adequate representation of the process of energy  
301 cycling include those of the microbial food web, as previously discussed. Model Set 2 is  
302 composed of 35 models representing 23 distinct systems.

### 303 *2.4.2 Uncertainty Analysis*

304 We tested the uncertainty in the *I/D* flow ratio to potential error in the model flow values  
305 (hypothesis 3) using a Monte Carlo procedure. Borrett and Osidele (2007) hypothesized that  
306 these results should be robust; here we tested this hypothesis. We constructed 10,000 models in  
307 which the original flow values were perturbed by  $\pm 5\%$ . We used the outputs for each node to  
308 rebalance the models by either subtracting or adding the amount of the change in the inputs of  
309 the node to that node's boundary output. This procedure allows the architecture, inputs, and the  
310 general distributions of flows for each network model to be maintained. Any modified model  
311 with a negative output was treated as a failed model because a negative loss does not make  
312 thermodynamic sense. We generated 10,000 successful models for each network model and  
313 applied NEA on each to estimate the analysis uncertainty.

## 314 **3 Results**

315 Our results generally confirm that indirect flows tend to dominate direct flows in  
316 empirically based networks. We present the evidence for each specific hypothesis in turn.

### 317 **3.1 Dominance of Indirect Flows**

318 Indirect flows dominate direct flows in 37 models out of 50 (74%) when the entire set of  
319 networks was considered (Figure 2, Model Set 1, all bars). When only models that met our more  
320 stringent set of criteria were considered, 31 models out of 35 (88.5%) show dominance of



321 indirect flows (Figure 2, Model Set 2, black bars only). In Model Set 2, the St. Marks seagrass  
322 sites 3 and 4 and Cypress wet and dry seasons have an  $I/D < 1$ .

323 Each of the 50 models in our data set are not necessarily independent given that some of  
324 the models reflect the same system, but under different ecological conditions or times of the  
325 year. However, they are not statistical replicates either. Given this ambiguity in the number of  
326 independent samples within our dataset, we also considered the results with respect to the  
327 number of distinct systems as opposed to models. When we do so, 27/35 systems (77.1%)  
328 exhibit dominance of indirect flows when all systems are considered. When only those systems  
329 that are used in Model Set 2 are analyzed this way, 21/23 systems (91.3%) support the  
330 hypothesis. Not included in the numerator of these calculations is the St. Marks Seagrass  
331 system, which has sites that both do and do not exhibit dominance of indirect flows.

### 332 3.2 Model Features Influencing Indirect Flows

333 System size  $n$  ( $r^2 = 0.069$ ;  $p = 0.06$ ) and connectance  $C$  ( $r^2 = 0.022$ ;  $p = 0.31$ ) do not  
334 explain much variation in IFI (Figure 3a and 3b, respectively). This is supported by Spearman's  
335 rank correlation test, as results were non-significant between IFI and both  $n$  and  $C$  ( $p = 0.078$ ;  $\rho$   
336  $= -0.25$  and  $p = 0.34$ ;  $\rho = 0.14$ , respectively). In contrast, cycling does appear to have a positive  
337 nonlinear relationship with the magnitude of IFI (Figure 3d). We did not do a linear regression  
338 analysis of the relationship between DFI and FCI on IFI because these indicators are not  
339 independent calculations, violating the assumptions of the statistical method. However, the  
340 results of Spearman's rank correlation test show a significant correlation between IFI and both  
341 DFI and FCI ( $p < 0.05$ ;  $\rho = -0.67$  and  $p < 0.05$ ;  $\rho = 0.88$ , respectively).

### 342 3.3 Analytical Robustness

343 For our uncertainty analysis, we randomly added  $\pm 5\%$  (uniform random distribution) to  
344 the observed flows. In 34 out of 50 models we generated failed perturbed models because when  
345 balancing the model one or more adjusted values became negative. The number of failures  
346 ranged from 1,256 for Ythan Estuary to 464,485 for Cypress dry season. We found it impossible  
347 to generate 10,000 models in a reasonable time period with  $\pm 5\%$  error for the Florida Bay wet  
348 and dry seasons. As these models have 14 zeros in the original output vector, we were unable to  
349 get 10,000 perturbed models that met our constraints. After generating the perturbed models, we  
350 verified that the maximum error was 5%, minimum error  $-5\%$ , and median and mean error  $\approx 0$ .

351 The error bars on Figure 2 show the full range of  $I/D$  uncertainties generated by our  
352 analysis, but here we will focus on the interquartile ranges, as this is more representative. The  
353 interquartile ranges are small indicating that the  $I/D$  values are fairly robust to  $\pm 5\%$  error in the  
354 flow values. We divided these interquartile ranges by the actual  $I/D$  values and multiplied by  
355 100 to determine the percent uncertainty in  $I/D$ . The min is 1.18%, median 2.47%, and max  
356 5.56%. This suggests that the uncertainty in  $I/D$  is less than the range of uncertainty (10%) in the  
357 data used to construct the models. Most importantly, this uncertainty in  $I/D$  does not lead to a  
358 change in the qualitative interpretation of the ratio for any of the models. However, when we  
359 consider the min and max values instead of the 1<sup>st</sup> and 3<sup>rd</sup> quartiles, we do see that the qualitative  
360 interpretation can change for the Cone Springs and Northern Benguela Estuary models from  
361 indirect flow dominance to direct flow dominance (Figure 2).

### 362 3.4 Microbial Food Web Contributions to System Behavior

363 The microbial food web mediates a large percentage of cycling in many of the models in  
364 Model Set 2 (Figure 4). The highest percentage is for the Florida Bay wet season model with

365 58.2% and the lowest is Swartkops Estuary at 0%. The median value for all models in Figure 4  
366 is 30.2% (Bothnian Bay), and in 17/35 models over a quarter of the cycling is attributed to the  
367 microbial food web components. There was no significant linear relationship between IFI and  
368 microbial food web FCI ( $r^2 = 0.014$ ,  $p = 0.50$ ) and results of Spearman's correlation test were  
369 non-significant ( $p = 0.42$ ;  $\rho = -0.14$ ).

370 Sites 1 and 2 within the St. Marks seagrass show dominance of indirect flows while sites  
371 3 and 4 do not. To explain this difference, we used EC analysis on these six models (Figure 5) to  
372 identify potential differences in the roles of microbial food web compartments. We first  
373 observed that where most of the species have a similar role in generating TST, there are a few  
374 dominant nodes that contribute to the system behavior. In sites 1 and 2 the top three nodes  
375 contributing to total system throughflow are sediment POC, benthic bacteria, and meiofauna,  
376 except for in site 2 (Jan) the third position is taken by suspended POC. Benthic bacteria mediate  
377 an average of 9.8% of TST in site 1 and 7.8 % in site 2. In site 3 the top three nodes generating  
378 system activity are sediment POC, predatory gastropods, and spot; in site 4 these nodes are  
379 sediment POC, benthic bacteria, and predatory gastropods. Benthic bacteria only mediates 2.6%  
380 of TST in site 3 and 4.5% in site 4. This analysis shows that the impact of the microbial food  
381 web is reduced in sites 3 and 4 when compared to sites 1 and 2.

## 382 4 Discussion

383 In this paper we tested the generality of the dominance of indirect flows hypothesis on 50  
384 empirically-based ecosystem models using NEA. We found that indirect flows dominate in the  
385 majority but not all of the models. When the hypothesis was tested on a subset of the models  
386 that have the model structure and species necessary to better represent energy–matter cycling, the  
387 evidence is stronger. We also tested the hypotheses relating to the magnitude of indirect flow,  
388 and we found no clear relationship between *IFI* and the magnitude of system size and  
389 connectance. There is a significant correlation, however, between *IFI* and direct flow intensity  
390 and cycling. Our analysis suggests that our NEA results are robust to model uncertainty. Lastly,  
391 the results of the node level FCI and EC support the hypothesis that the microbial food web is an  
392 important factor in the dominance of indirect effects. The amount of cycling mediated by the  
393 microbial food web cannot be used to predict the amount of indirect flow, yet the observed range  
394 in this value reflects the variation predicted by the literature (as in Hart et al., 2000). In sum, our  
395 evidence supports that indirect energy flows are just as significant, if not more so, than the  
396 observed direct feeding behavior, as seen in this study supporting the dominance of indirect  
397 flows hypothesis.

### 398 4.1 Confidence in NEA Results

399 The qualitative interpretation of the *I/D* ratios can be made with reasonable confidence,  
400 as shown by the results of the uncertainty analysis. None of the *I/D* values for any of the models  
401 change its qualitative interpretation when the error is introduced using the interquartile range and  
402 only two when using the full range. Interestingly, the uncertainty in the NEA results is less than  
403 the uncertainty introduced into the data. This is similar to the patterns found previously for  
404 network homogenization, another whole system property (Borrett and Salas 2010), and through  
405 different methods of analyzing uncertainty (Borrett and Osidele 2007). Further, this is similar to  
406 results reported for the Ascendant Perspective of Ecological Network Analysis (ENA) (Kaufman  
407 and Borrett, 2010; Kones et al., 2009).

408 While it would be ideal to test a larger percent error, doing so consistently for all models  
409 was prohibited by the thermodynamic constraints of the analysis. Therefore, our conclusions of

410 the robustness of the results of NEA to data uncertainty are limited to a smaller error range.  
411 However, we did observe that variation remained small for successful attempts using larger  
412 percent errors (data not shown).

#### 413 **4.2 Evidence for Dominance of Indirect Flows**

414 The original dominance of indirect effects hypothesis (Higashi and Patten, 1986, 1989;  
415 Patten, 1982) suggested that indirect flows should tend to dominate in the majority of ecosystem  
416 networks. Our work supports this hypothesis as 74% of all the models had  $I > D$ . However, all  
417 the models in our dataset may not all be suitable to address our specific question. While these  
418 models may have been appropriate to address the questions for which they were designed, it does  
419 not imply that they can be used broadly for all network questions. Fath (2007) provides  
420 guidelines for network construction, one of which is exhaustiveness. A model can only be  
421 exhaustive if system components are aggregated together which results in diverse and maybe  
422 heterogeneous nodes. However, ecosystem models of energy-matter should explicitly include  
423 the species that mediate the important ecosystem processes. Decomposition is one such process,  
424 which is critical in the cycling of carbon and ecosystem functioning. The explicit representation  
425 of detrital compartments has been shown to influence the results of trophic modeling (e.g.  
426 Allesina et al., 2005; Edwards, 2001; Fath and Halnes, 2007). Also vital to the adequate  
427 representation of carbon flow are the species that mediate this decomposition and their microbial  
428 consumers, and species that consume small phytoplankton. We found that models with some  
429 explicit representation of the microbial food web tended to have more recycling and therefore  
430 more indirect effects. Figure 5 shows that a significant amount of cycling is indeed being  
431 mediated by the microbial food web components in many of the models in Model Set 2. Further,  
432 10 out of the 15 models excluded from Model Set 2 had  $I/D < 1$ , while only 4/35 had  $I/D < 1$  in  
433 Model Set 2.

#### 434 **4.3 Failures of the Dominance of Indirect Flows Hypothesis in Model Set 2**

435 Despite the tendency for  $I/D$  to exceed unity, some of the models in Model Set 2 had  
436 direct flows that exceeded indirect flows. To investigate these failures we used the idea that  
437 communities that export or bury the products of primary production are typically considered  
438 autotrophic, as these products are less available for heterotrophic use (Duarte and Cedbrían,  
439 1996). We would expect autotrophic systems to have a higher tendency to fail the dominance of  
440 indirect flows hypothesis as a loss of energy-matter means there is less available for recycling,  
441 thus less indirect flow. While productivity estimates were not published for all models and we  
442 were not able to use them widely in this work, these ideas are useful in the potential explanation  
443 of the failure of the hypothesis.

444 The St. Marks seagrass, sites 3 and 4, are two of the four models that reveal an  $I/D < 1$  in  
445 Model Set 2. These, along with the other two St. Marks sites, provide an example of a system  
446 where the underlying ecology may legitimately contribute to a lack of indirect flow dominance.  
447 A combination of site location and production and cycling measures can help to explain our  
448 findings. First, the location of these sites can provide useful information. The sites that exhibit  
449  $I > D$  (Figure 3) are located within the relatively protected Goose Creek Bay (Baird et al., 1998).  
450 The sites that show  $I < D$  (Figure 3) are located in more flow-thru areas; site 3 is located within  
451 the mouth of the St. Marks River, and site 4 is located just outside of this mouth (Baird et al.,  
452 1998). Supporting this, site 3 has a low residence time compared to the other sites (Baird et al.  
453 1998). Looking at production measures, we see that site 3 has greater production and a larger  
454 production/biomass ratio (Baird et al. 1998), indicating that the products of production are likely

455 exiting the system and cannot be recycled at that site. Finally, we also see that the average  
456 system level and microbial food web FCI for sites 1 and 2 is greater than for sites 3 and 4 (Table  
457 1 and Figure 5). Thus, while the cycling in this system is generally low, there is clearly more  
458 cycling occurring at sites 1 and 2 and a significant portion of this is being mediated by bacterial  
459 decomposition.

460 Taken together, site 3 is a largely autotrophic community where the products of primary  
461 production are quickly exported from the system and cannot be utilized by the indirect pathways.  
462 Site 4 is located just outside the mouth of the St. Marks River, and while the residence time and  
463 measures of production are not notably different than sites 1 and 2, it can be reasoned that its  
464 proximity to the river, versus the bay, contribute to its likeness to site 3 versus sites 1 and 2 in  
465 regards to cycling and  $I/D$ . In both sites 3 and 4 the use of recycled carbon is not as prevalent,  
466 and therefore the relative role of bacteria in generating TST is not nearly as important. This  
467 diminished role of bacteria at sites 3 and 4 is seen in both the FCI decomposition (Figure 4) and  
468 the average EC (Figure 5). This congruence further supports the utility of the EC metric (Fann,  
469 2009). In conclusion, the failure of dominance of indirect flow in sites 3 and 4 of the St. Marks  
470 seagrass system is likely due to the ecological dynamics at these sites. In addition, the inter-site  
471 comparisons reveal the potential role of bacteria in contributing to the magnitude of indirect  
472 flows.

473 The Cypress models (wet and dry) also have  $I/D < 1$ . In this system detritus is largely  
474 buried (Ulanowicz et al., 1997), and thus the use of cycled carbon may not be significant, leading  
475 to diminished indirect flows. Therefore, this site can be considered autotrophic and like St.  
476 Marks seagrass sites 3 and 4. The representation of the microbial components in this model may  
477 also be problematic. The microbial food web is compressed into “living POC”, which includes  
478 bacteria, microprotozoans, and zooplankton, and “living sediments”, which includes  
479 bacterioplankton, microfauna, and meiofauna. This aggregation hinders the ability of the flows  
480 within the microbial food web to contribute to the sum of indirect flows, and therefore may lead  
481 to this sum being smaller than it would be if the microbial food web components were not so  
482 highly aggregated. However, this suggests that our analytical results are conservative. This idea  
483 is further supported by Fath (2004) who demonstrated that indirect flows tend to increase with  
484 system size, though our results and those of Baird et al. (2009) do not match.

485 This extreme aggregation of the microbial food web occurs in other models in which the  
486  $I/D > 1$  (e.g. Graminoids wet and dry) so it may not be the sole cause of  $I/D < 1$ . This suggests  
487 that the manner in which the microbial components are represented may matter, as opposed to  
488 just if they are or are not included in the model. The detrital fluxes may also be a problem as  
489 flows to detritus are often determined based on the balancing procedure as opposed to  
490 empirically determined (e.g. Baird and Milne, 1981; Baird et al., 1991; Miehl et al., 2009ab).  
491 Similarly, the dynamics within the microbial food web are difficult to determine empirically  
492 (Azam et al., 1983; Baird et al., 2009) and these compartments have not generally been the focus  
493 of many studies. In sum, the decomposition process and the role of the microbial food web are  
494 to some degree speculation, leading to less confidence in the results presented in Figure 5.  
495 Despite the challenges of representing the microbial food web and detritus, the degree of  
496 variation in the role of the microbial food web we found is supported by the literature (as in Hart  
497 et al., 2000; Sommer et al., 2002) and the results here reinforce the importance of its inclusion in  
498 ecosystem models.

499 **4.4 Drivers of I/D Variation**

500 An interesting result of NEA on all 50 ecosystem models is the lack of support for two of  
501 the four NEA hypotheses regarding under which conditions indirect flow intensity should  
502 increase in magnitude. There was no obvious relationship between indirect flow intensity and  
503 the size of the system and connectance but there is a significant correlation between IFI and both  
504 DFI and FCI. All four variables are predicted by the algebra of NEA to be positively correlated  
505 with the magnitude of IFI, and Fath (2004) shows a positive relationship between size of the  
506 system and indirect effects using cyber-ecosystems. However, results garnered from the use of  
507 ecosystem models do not support these hypotheses. Our results only support the predictions for  
508 DFI and FCI.

509 Our results are similar to previous studies. For example, analysis by Baird et al. (2009)  
510 also does not support the predicted relationship between  $I/D$  and  $n$ . In addition, in an analysis of  
511 the same data set used here Borrett and Salas (2010) did not find evidence of the predicted  
512 relationship between homogenization and system size and connectance, and marginal support for  
513 the relationship between homogenization and FCI. These results taken together suggest there is  
514 potentially an aspect of real ecosystems that is not being captured in the algebraic predictions  
515 and or the cyber-ecosystem community assembly model (Fath 2004; Fath and Killian 2007).  
516 Fath's original models (2004) are limited in their ability to reflect real ecosystems because they  
517 are overly simplified. There are no discernible differences in the species within the defined  
518 functional groups, which is not an accurate reflection of natural systems since we would expect  
519 some sort of niche differentiation. In addition, the flow values appear to have been randomly  
520 assigned in these networks from a uniform random distribution. Despite these simplifications,  
521 these models supported Higashi and Patten's (1989) algebraic hypotheses. Again, a key  
522 contribution of our work is that we tested the hypotheses on empirically derived ecosystem  
523 models.

524 **4.5 Methodological Challenges**

525 There are limitations of our model database that could influence our results. These  
526 potential problems largely stem from the subjective nature of model construction. For example,  
527 our models may be biased due to the limited authorship of the models in our database. The act of  
528 constructing an ecosystem model entails many decisions that could affect the results of the  
529 analysis (Abarca-Arenas and Ulanowicz, 2002; Allesina et al., 2005; Pinnegar et al., 2005), and  
530 this may bias the overall results if a large percentage of the models are constructed by the same  
531 author or group of authors. Despite this potential, there is observable variation in the  $I/D$  values  
532 for models constructed by the same author (e.g. Baird et al., 1991; Baird et al., 1998), suggesting  
533 that modeling decision bias does not necessarily obscure real ecological differences. A second  
534 issue is that the models have different aggregation schemes. Some models such as Lake Findely  
535 (Richey et al., 1978) have an extremely high degree of aggregation while others such as Lake  
536 Oneida (Miehels et al., 2009) have much finer resolution. These differences in model structure  
537 make it impossible to compare results between models (Baird et al., 1991), and might effect the  
538 overall conclusions of the study. We doubt, however, that this potential bias would change our  
539 overall results. Another limitation is our focus on trophically based models; we excluded more  
540 biogeochemically based ecosystem models from this study. Previous results, however, suggest  
541 that indirect flows are even larger in biogeochemically based ecosystems models (Borrett et al.,  
542 2006; Borrett and Osidele, 2007; Borrett et al. 2010), but this remains to be tested in a systematic  
543 fashion. Lastly, where our sample size of 50 models is the largest assembled, it is still a  
544 relatively small sample. Despite these limitations, we argue that the model database used to test

545 our hypotheses is a useful data set that is large enough to provide informative results. Despite  
546 those issues, we believe that this data is sufficient to address our hypotheses, and these results are  
547 supported by our uncertainty analyses.

## 548 **5 Conclusions**

549 In this paper we present evidence that indirect flows tend to dominant direct flows in  
550 ecosystems. We also demonstrate the importance of including the microbial food web in  
551 ecosystem energy models as this subweb is critical in mediating the flow of carbon in some  
552 ecosystems. Lastly, we found that our results were robust to error in the flow estimates,  
553 suggesting confidence in our results despite the difficulties of model construction.  
554 Our core contribution with this paper is the demonstration of the generality of the dominance of  
555 indirect effects in empirically derived and trophically based ecosystem networks. We show that  
556 when significant cycling was present in the model nearly every system examined had indirect  
557 flows dominating direct flows. These results work in concert with several recent studies to  
558 provide strong support for the hypothesis. For example, two studies suggest that relaxing the  
559 steady-state assumption of NEA would generate similar results. Borrett et al. (2006) found that  
560 indirect flows remain dominant in the Neuse River Estuary despite temporal variability, and  
561 Borrett et al. (2010) show that indirect flows develop very rapidly in ecosystems, indicating that  
562 an ecosystem does not need to remain at a certain configuration for long before indirect flows  
563 will be dominant. This latter conclusion was made using both trophic and biogeochemical  
564 networks, indicating that the results are not restricted to trophically-based ecosystem models, as  
565 is used in this work. Our findings, in combination with the initial algebra (Higashi and Patten,  
566 1986, 1989), theoretical work (Fath, 2004), and the increasingly common empirical evidence,  
567 provides strong support for the systems ecology hypothesis that indirect flows dominate direct  
568 flows in ecosystems.

## 569 **6 Acknowledgments**

570 We would like to thank M. Freeze for his help developing the methods for the uncertainty  
571 analysis, S.L. Fann for help with environ centrality, and B.C. Patten for review of the  
572 manuscript. AKS was supported by the James F. Merritt fellowship from the UNCW Center for  
573 the Marine Science and SRB was supported in part by UNCW.

## 574 **7 References**

- 575 Abarca-Arenas, L.G., Ulanowicz, R.E., 2002. The effects of taxonomic aggregation on network  
576 analysis. *Ecol. Model.* 149, 285–296.
- 577 Abrams, P.A., Menge, B.A., Mittlebach, G.G., Spiller, D., Yodzis, P., 1996. The role of indirect  
578 effects in food webs. In: Polis G, Winemiller KO, editors. *Food Webs: Dynamic and*  
579 *Structure*. New York: Chapman and Hall. p371–395.
- 580 Azam F., Fenchel T., Field J., Gray J., Meyer-Reid L., and Thingstad F., 1983. The ecological  
581 role of water-column microbes in the sea. *Mar. Ecol.: Prog. Ser.* 10, 257–263.
- 582 Allesina, S., and Bondavalli, C., 2003. Steady state of ecosystem flow networks: a comparison  
583 between balancing procedures. *Ecol. Model.* 165, 231–239.
- 584 Allesina, S., Bondavalli, C., Scharler, U.M., 2005. The consequences of the aggregation of  
585 detritus pools in ecological networks. *Ecol. Model.* 189, 221–232.
- 586 Almunia, J., Basterretxea, G., Aistegui, J., Ulanowicz, R.E., 1999. Benthic-pelagic switching in  
587 a coastal subtropical lagoon. *Estuarine, Coastal Shelf Sci.* 49, 363–384.

- 588 Baird, D., Asmus, H., Asmus, R., 2004a. Energy flow of a boreal intertidal ecosystem, the Sylt-  
589 Rømø Bight. *Mar. Ecol.: Prog. Ser.* 279, 45–61.
- 590 Baird, D., Asmus, H., Asmus, R., 2008. Nutrient dynamics in the Sylt-Romo Bight ecosystem,  
591 German Wadden Sea: an ecological network analysis approach. *Estuarine, Coastal Shelf*  
592 *Sci.* 80, 339–356.
- 593 Baird, D., Christian, R.R., Peterson, C.H., Johnson, G.A., 2004b. Consequences of hypoxia on  
594 estuarine ecosystem function: energy diversion from consumers to microbes. *Ecol. Appl.* 14,  
595 805–822.
- 596 Baird, D., Fath, B.D., Ulanowicz, R.E., Asmus, H., Asmus, R., 2009. On the consequences of  
597 aggregation and balancing of networks on system properties derived from ecological  
598 network analysis. *Ecol. Model.* 220, 3465–3471.
- 599 Baird, D., Luczkovich, J., Christian, R.R., 1998. Assessment of spatial and temporal variability  
600 in ecosystem attributes of the St. Marks National Wildlife Refuge, Apalachee Bay, Florida.  
601 *Estuarine, Coastal Shelf Sci.* 47, 329–349.
- 602 Baird, D., McGlade, J.M., and Ulanowicz, R.E., 1991. The comparative ecology of six marine  
603 ecosystems. *Philos. T. Roy. Soc. B.* 333, 15–29.
- 604 Baird, D., and Milne, H., 1981. Energy flow in the Ythan Estuary, Aberdeenshire, Scotland.  
605 *Estuarine, Coastal Shelf Sci.* 13, 455–472.
- 606 Baird, D., Ulanowicz, R.E., 1989. The seasonal dynamics of the Chesapeake Bay ecosystem.  
607 *Ecol. Monogr.* 59, 329–364.
- 608 Bever, J.D., 1999. Dynamics within mutualism and the maintenance of diversity: inference from  
609 a model of interguild frequency dependence. *Ecol. Lett.* 2, 52–62.
- 610 Bever, J.D., 2002. Negative feedback within a mutualism: host-specific growth of mycorrhiza  
611 fungi reduced plant benefit. *P. Roy. Soc. Lond. B. Bio.* 269, 2595–2601.
- 612 Borgatti, S.P., Everett, M.G., 2006. A graph-theoretic perspective on centrality. *Soc. Networks*  
613 28, 466–484.
- 614 Bondavalli, C., Ulanowicz, R.E., 1999. Unexpected effects of predators upon their prey: the case  
615 of the American alligator. *Ecosystems* 2, 49–63.
- 616 Borrett, S.R., Fath, B.D., Patten, B.C., 2007. Functional integration of ecological networks  
617 through pathway proliferation. *J. Theor. Biol.* 245, 98–111.
- 618 Borrett, S.R., Salas, A.K., 2010. Evidence for resource homogenization in 50 trophic ecosystem  
619 networks. *Ecol. Model.* 221, 1710–1716
- 620 Borrett, S.R., Osidele, O.O., 2007. Environ indicator sensitivity to flux uncertainty in a  
621 phosphorus model of Lake Sidney Lanier, USA. *Ecol. Model.* 200, 371–383.
- 622 Borrett, S.R., Whipple, S.J., Patten, B.C., Christian, R.R., 2006. Indirect effects and distributed  
623 control in ecosystems: temporal variation of indirect effects in a seven-compartment model  
624 of nitrogen flow in the Neuse River Estuary, USA–time series analysis. *Ecol. Model.* 194,  
625 178–188.
- 626 Borrett, S.R., Whipple, S. J., Patten, B.C., 2010. Rapid development of indirect effects in  
627 ecological networks. *Oikos* 119, 1136–1148.
- 628 Brylinsky, M., 1972. Steady-state sensitivity analysis of energy flow in a marine ecosystem. In  
629 B.C. Patten, editor. *Systems Analysis and Simulation in Ecology Volume 2.* Academic  
630 Press, New York. p81–101.

631 Christian, R.R., Forés, E., Comin, F., Viaroli, P., Naldi, M., Ferrari, I., 1996. Nitrogen cycling  
632 networks of coastal ecosystems: influence of trophic status and primary producer form.  
633 Ecol. Model. 87, 111–129.

634 Dame, R.F., Patten, B.C., 1981. Analysis of energy flows in an intertidal oyster reef. Mar.  
635 Ecol.: Prog. Ser. 5, 115–124.

636 Diekötter, T., Haynes, K.J., Mazeffa, D., Crist, T.O., 2007. Direct and indirect effects of habitat  
637 area and matrix composition on species interactions among flower-visiting insects. Oikos  
638 116, 1588–1598.

639 Duarte, C. M., Cebrián, J., 1996. The fate of marine autotrophic production. Limnol. Oceanogr.  
640 41, 1758–1766.

641 Edwards, A.M., 2001. Adding detritus to a nutrient-phytoplankton-zooplankton model: a  
642 dynamical-systems approach. J. Plankton Res. 23, 389–413.

643 Estrada, E., 2010. Generalized walks-based centrality measures for complex biological networks.  
644 J. Theor. Biol. 263, 556–565.

645 Fann, S.F., 2009. Environ centrality quantifies the relative roles of species in generating  
646 ecosystem activity. Honors Thesis, University of North Carolina Wilmington.

647 Fath, B.D., 2004. Network analysis applied to large-scale cyber-ecosystems. Ecol. Model. 171,  
648 329–337.

649 Fath, B.D., Borrett, S.R., 2006. A MATLAB® function for Network Environ Analysis.  
650 Environ. Modell. Softw. 21, 375–405.

651 Fath, B.D., Hales, G., 2007. Cyclic energy pathways in ecological food webs. Ecol. Model.  
652 208, 17–24.

653 Fath, B.D., Killian, M.C., 2007. The relevance of ecological pyramids in community  
654 assemblages. Ecol. Model. 208, 286–294.

655 Fath, B.D., Patten, B.C., 1998. Network synergism: emergence of positive relations in  
656 ecological systems. Ecol. Model. 107, 127–143.

657 Fath, B.D., Patten, B.C., 1991. Review of the foundations of network environ analysis.  
658 Ecosystems 2, 167–179.

659 Fath, B.D., Scharler, U.M., Ulanowicz, R.E., Hannon, B., 2007. Ecological network analysis:  
660 network construction. Ecol. Model. 208, 49–55.

661 Fenchel, T., 1970. Studies on the decomposition of organic detritus derived from the turtle grass  
662 *Thalassia testudinum*. Limnol. Oceanogr. 15, 14–20.

663 Finn, J.T., 1976. Measures of ecosystem structure and function derived from analysis of flows.  
664 J. Theor. Biol. 56, 363–380.

665 Finn, J.T., 1980. Flow analysis of models of the Hubbard brook ecosystem. Ecology 61,  
666 562–571.

667 Hanneman, R.A., Riddle, M.A. Centrality and Power *in* Introduction to social network methods.  
668 (Riverside, CA: University of California, Riverside, 2005),  
669 <http://faculty.ucr.edu/~hanneman/> (accessed October 10, 2009).

670 Hart, D.R., Stone, L., Berman, T., 2000. Seasonal dynamics of the Lake Kinneret food web: The  
671 importance of the microbial loop. Limnol. Oceanogr. 45, 350–361.

672 Heymans, J.J., Baird, D., 2000. A carbon flow model and network analysis of the Northern  
673 Benguela Upwelling system, Namibia. Ecol. Model. 126, 9–32.

674 Higashi, M., Patten, B.C., 1986. Further aspects of the analysis of indirect effects in ecosystems.  
675 Ecol. Model. 31, 69–77.



- 676 Higashi, M., Patten, B.C., 1989. Dominance of indirect causality in ecosystems. *Am. Nat.* 133,  
677 288–302.
- 678 Jørgensen, S.E., Fath, B.D., Bastianoni, S., Marques, J., Muller, F., Nielsen, S.N., Patten, V.,  
679 Tiezzi, E., Ulanowicz, R., 2007. *A New Ecology: Systems Perspective*. Netherlands:  
680 Elsevier. 288p.
- 681 Jørgensen, S.E., Patten, B.C., Straskraba, M., 1992. Ecosystems emerging - toward an ecology  
682 of complex-systems in a complex future. *Ecol. Model.* 62, 1–27.
- 683 Jørgensen, S.E., Patten, B.C., Straskraba, M., 1999. Ecosystems emerging: 3. Openness. *Ecol.*  
684 *Model.* 114, 41–64.
- 685 Kaufman, A.G., Borrett, S.R., 2010. Ecosystem network analysis indicators are generally robust  
686 to parameter uncertainty in a phosphorus model of Lake Sidney Lanier, USA. *Ecol. Model.*  
687 221, 1230–1238
- 688 Kones, J.K., Soetaert, K., van Oevelen, D., Owino, J.O., 2009. Are network indices robust  
689 indicators of food web functioning? A Monte Carlo approach. *Ecol. Model.* 220, 370–382.
- 690 Larsson, U., Hagström, Å., 1982. Fractionated phytoplankton primary production, exudates  
691 release, and bacterial production in a Baltic eutrophication gradient. *Mar. Biol.* 67, 57–70.
- 692 Link, J., Overholtz, W., O'Reilly, J., Green, J., Dow, D., Palka, D., Legault, C., Vitaliano, J.,  
693 Guida, V., Fogarty, M., Brodziak, J., Methratta, L., Stockhausen, W., Col, L., Griswol, C.,  
694 2008. The northeast U.S. continental shelf energy modeling and analysis exercise (EMAX):  
695 Ecological network model development and basic ecosystem metrics. *J. Marine Syst.* 74,  
696 453–474.
- 697 Mann, K.H., 1988. Production and use of detritus in various freshwater, estuarine, and coastal  
698 marine ecosystems. *Limnol. Oceanogr.* 33, 910–930.
- 699 Menge, B.A., 1997. Detection of direct versus indirect effects: were experiments long enough?  
700 *Am. Nat.* 149, 801–823.
- 701 Miehls, A.L.J., Mason, D.M., Frank, K.A., Krause, A.E., Peacor, S.D., Taylor, W.W., 2009a.  
702 Invasive species impacts on ecosystem structure and function: A comparison of Oneida  
703 Lake, New York, USA, before and after zebra mussel invasion. *Ecol. Model.* 220,  
704 3194–3209.
- 705 Miehls, A.L.J., Mason, D.M., Frank, K.A., Krause, A.E., Peacor, S.D., Taylor, W.W., 2009b.  
706 Invasive species impacts on ecosystem structure and function: A comparison of the Bay of  
707 Quinte, Canada and Oneida Lake, USA, before and after zebra mussel invasion. *Ecol.*  
708 *Model.* 220, 3182–3193.
- 709 Monaco, M.E., Ulanowicz, R.E., 1997. Comparative ecosystem trophic structure of three U.S.  
710 Mid-Atlantic estuaries. *Mar. Ecol.: Prog. Ser.* 161, 239–254.
- 711 Odum, H.T., 1957. Trophic structure and productivity of Silver Springs, Florida. *Ecol. Monogr.*  
712 27, 55–112.
- 713 Patten, B.C., 1982. Environs: Relativistic elementary particles for ecology. *Am. Nat.* 119,  
714 179–219.
- 715 Patten, B.C., 1983. On the quantitative dominance of indirect effects in ecosystems. In W.K.  
716 Lauenroth, G.V. Skogerboe, and M. Flug, editors. *Analysis of Ecological Systems: State-*  
717 *of-the-Art in Ecological Modelling*. Elsevier, Amsterdam. p27–37.
- 718 Patten, B.C., 1984. Further developments toward a theory of the quantitative importance of  
719 indirect effects in ecosystem. *Verh. Gesellschaft für Ökologie* 13, 271–284.
- 720 Patten, B.C., 1985. Energy cycling in the ecosystem. *Ecol. Model.* 28, 1–71.

- 721 Patten B.C., Bosserman, R.W., Finn, J.T., Cale, W.G., 1976. Propagation of cause in  
722 ecosystems. In Patten, B.C., editor. *Systems Analysis and Simulation in Ecology*, Volume  
723 IV. Academic Press, New York. p457–579.
- 724 Patten, B.C., Higashi, M., Burns, T., 1990. Trophic dynamics in ecosystem networks:  
725 significance of cycles and storage. *Ecol. Model.* 51, 1–28.
- 726 Patten, B.C., 1991. Network ecology: indirect determination of the life-environment relationship  
727 in ecosystems. Higashi, M., Burns, T.P., editors. *Theoretical Studies of Ecosystems: The*  
728 *Network Perspective*. Cambridge: Cambridge University Press. p289–351
- 729 Pinnegar, J.K., Blanchard, J.L., Mackinson, S., Scott, R.D., Duplisea, D.E., 2005. Aggregation  
730 and removal of weak-links in food web models: system stability and recovery from  
731 disturbance. *Ecol. Model.* 184, 229–248.
- 732 Pomeroy, L.R., 1974. The ocean's food web, a changing paradigm. *BioScience* 24, 499–504.
- 733 Preisser, E.L., Bolnick, D.I., Benard, M.F., 2005. Scared to death? The effects of intimidation  
734 and consumption in predator-prey interactions. *Ecology* 86, 501–509.
- 735 Richey, J.E., Richey, J.E., Wissmar, R.C., Devol, A.H., Likens, G.E., Eaton, J.S., Wetzel, R.G.,  
736 Odum, W.E., Johnson, N.M., Loucks, O.L., Prentki, R.T., Rich, P.H., 1978. Carbon flow in  
737 four lake ecosystems: a structural approach. *Science* 202, 1183–1186.
- 738 Rybarczyk, H., Elkaim, B., Ochs, L., Loquet, N., 2003. Analysis of the trophic network of a  
739 macrotidal ecosystem: the Bay of Somme (Eastern Channel). *Estuarine, Coastal Shelf Sci.*  
740 58, 405–421.
- 741 Sandberg, J., Elmgren, R., Wulff, F., 2000. Carbon flows in Baltic sea food webs-a re-  
742 evaluation using a mass balance approach. *J. Marine. Syst.* 25, 249–260.
- 743 Sherr, E., Sherr, B., 1988. Role of microbes in pelagic food webs: a revised concept. *Limnol.*  
744 *Oceanogr.* 33, 1225–1227.
- 745 Sommer, U., Stibor, H., Katechakis, A., Sommer, F., Hansen, T., 2002. Pelagic food web  
746 configurations at different levels of nutrient richness and their implications for the ratio fish  
747 production: primary production. *Hydrobiologia* 484, 11–20.
- 748 Strauss, S.Y., 1991. Indirect effects in community ecology: their definition, study and  
749 importance. *Trends Ecol. Evol.* 6, 206–210.
- 750 Tilly, L.J., 1968. The structure and dynamics of Cone Spring. *Ecol. Monogr.* 38, 169–197.
- 751 Turner, A., Mittelbach, G., 1990. Predator avoidance and community structure: interactions  
752 among piscivores, planktivores, and plankton. *Ecology* 71, 2241–2254.
- 753 Ulanowicz, R.E., 1986. *Growth and Development: Ecosystems Phenomenology*. toExcel Press,  
754 Lincoln.
- 755 Ulanowicz, R.E., 1995. Trophic flow networks as indicators of ecosystem stress. In Polis,  
756 G.A. and Winemiller, K.O., editors. *Food Webs: Integration of Patterns and Dynamics*.  
757 Chapman and Hall, NY. p358–368.
- 758 Ulanowicz, R.E., Bondavalli, C., Egnotovitch, M.S., 1997. Network analysis of trophic dynamics  
759 in South Florida ecosystem, FY 96: The Cypress wetland ecosystem. Annual Report to the  
760 United States Geological Service Biological Resources Division Ref. No. [UMCES]CBL  
761 97-075, Chesapeake Biological Laboratory, University of Maryland.
- 762 Ulanowicz, R.E., Bondavalli, C., Egnotovitch, M.S., 1998. Network analysis of trophic dynamics  
763 in South Florida ecosystem, FY 97: The Florida Bay ecosystem. Annual Report to the  
764 United States Geological Service Biological Resources Division Ref. No. [UMCES] CBL  
765 98-123, Chesapeake Biological Laboratory, University of Maryland.

766 Ulanowicz, R.E., Bondavalli, C., Heymans, J.J., Egnotovitch, M.S., 1999. Network analysis of  
767 trophic dynamics in South Florida ecosystem, FY 98: The mangrove ecosystem. Annual  
768 Report to the United States Geological Service Biological Resources Division Ref. No.  
769 [UMCES] CBL 99-0073. Technical Report Series No. TS-191-99, Chesapeake Biological  
770 Laboratory, University of Maryland.

771 Ulanowicz, R.E., Bondavalli, C., Heymans, J.J., Egnotovitch, M.S., 2000. Network analysis of  
772 trophic dynamics in South Florida ecosystem, FY 99: The graminoid ecosystem. Annual  
773 Report to the United States Geological Service Biological Resources Division Ref. No.  
774 [UMCES] CBL 00-0176, Chesapeake Biological Laboratory, University of Maryland.

775 Ulanowicz, R.E., Puccia, C.J., 1990. Mixed tropic impacts in ecosystems. *Coenoses* 5:7–16.

776 Wasserman, S., Faust, K., 1994. *Social Network Analysis: Methods and Applications*.  
777 Cambridge University Press, Cambridge, U.K. 857p.

778 Wilkinson, D.M., 2006. *Fundamental Processes in Ecology an Earth Systems Approach*.  
779 Oxford University Press, New York. 200p.

780 Wootton, J.T., 1993. Indirect effects and habitat use in an intertidal community: interaction  
781 chains and interaction modifications. *Am. Nat.* 141, 71–89.

782 Wootton, J.T., 1994. The nature and consequences of indirect effects in ecological communities.  
783 *Annu. Rev. Ecol. Syst.* 25, 443–466.

784 Yodzis, P., 1988. The indeterminacy of ecological interactions as perceived through perturbation  
785 experiments. *Ecology* 69, 508–515.

786  
787  
788  
789  
790  
791  
792  
793

793 Table 1: 50 empirically derived and trophically-based ecosystem models. The complete list is  
 794 Model Set 1 and models excluded from Model Set 2 are indicated with superscript 1 (no cycling)  
 795 and/or superscript 2 (no microbial food web representation).  
 796  
 797

Model	Units	$n^a$	$C^a$	$TST^a$	$FCI^a$	$I/D^a$	Source
Lake Findley <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	4	0.38	51	0.30	1.72	Richey et al. 1978
Mirror Lake <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	5	0.36	218	0.32	1.86	Richey et al. 1978
Lake Wingra <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	5	0.40	1517	0.40	1.86	Richey et al. 1978
Marion Lake <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	5	0.36	243	0.31	2.18	Richey et al. 1978
Cone Springs <sup>2</sup>	kcal m <sup>-2</sup> yr <sup>-1</sup>	5	0.32	30626	0.09	1.02	Tilly 1968
Silver Springs <sup>1,2</sup>	kcal m <sup>-2</sup> yr <sup>-1</sup>	5	0.28	29175	0.00	0.20	Odum 1957
English Channel <sup>1,2</sup>	kcal m <sup>-2</sup> yr <sup>-1</sup>	6	0.25	2280	0.00	0.44	Brylinsky 1972
Oyster Reef	Kcal m <sup>-2</sup> yr <sup>-1</sup>	6	0.33	84	0.11	1.58	Dame and Patten 1981
Bay of Somme <sup>2</sup>	mgC m <sup>-2</sup> d <sup>-1</sup>	9	0.30	2035	0.14	0.74	Rybarczyk et al. 2003
Bothnian Bay	gC m <sup>-2</sup> yr <sup>-1</sup>	12	0.22	130	0.18	1.97	Sandberg et al. 2000
Bothnian Sea	gC m <sup>-2</sup> yr <sup>-1</sup>	12	0.24	458	0.27	2.52	Sandberg et al. 2000
Ythan Estuary <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	13	0.23	4181	0.24	1.89	Baird and Milne 1981
Baltic Sea	mgC m <sup>-2</sup> d <sup>-1</sup>	15	0.17	1974	0.13	1.93	Baird et al. 1991
Ems Estuary	mgC m <sup>-2</sup> d <sup>-1</sup>	15	0.19	1019	0.32	2.95	Baird et al. 1991
Swarkops Estuary	mgC m <sup>-2</sup> d <sup>-1</sup>	15	0.17	13996	0.47	2.73	Baird et al. 1991
Southern Benguela Upwelling	mgC m <sup>-2</sup> d <sup>-1</sup>	16	0.23	1774	0.19	1.24	Baird et al. 1991
Peruvian Upwelling	mgC m <sup>-2</sup> d <sup>-1</sup>	16	0.22	33496	0.04	1.16	Baird et al. 1991
Crystal River (control) <sup>2</sup>	mgC m <sup>-2</sup> d <sup>-1</sup>	21	0.19	15063	0.07	0.67	Ulanowicz 1986, 1995
Crystal River (thermal) <sup>2</sup>	mgC m <sup>-2</sup> d <sup>-1</sup>	21	0.14	12032	0.09	0.71	Ulanowicz 1986, 1995
Charca de Maspalomas Lagoon	mgC m <sup>-2</sup> d <sup>-1</sup>	21	0.13	6010331	0.18	2.69	Alumunia et al. 1999
Northern Benguela Upwelling	mgC m <sup>-2</sup> d <sup>-1</sup>	24	0.21	6608	0.05	1.04	Heymans and Baird 2000
Neuse Estuary (early summer 1997)	mgC m <sup>-2</sup> d <sup>-1</sup>	30	0.09	13826	0.12	1.48	Baird et al. 2004b
Neuse Estuary (late summer 1997)	mgC m <sup>-2</sup> d <sup>-1</sup>	30	0.11	13038	0.13	1.31	Baird et al. 2004b
Neuse Estuary (early summer 1998)	mgC m <sup>-2</sup> d <sup>-1</sup>	30	0.09	14025	0.12	1.41	Baird et al. 2004b
Neuse Estuary (late summer 1998)	mgC m <sup>-2</sup> d <sup>-1</sup>	30	0.10	15031	0.11	1.26	Baird et al. 2004b
Gulf of Maine	g ww m <sup>-2</sup> yr <sup>-1</sup>	31	0.35	18382	0.15	2.11	Link et al. 2008
Georges Bank	g ww m <sup>-2</sup> yr <sup>-1</sup>	31	0.35	16890	0.18	2.58	Link et al. 2008
Middle Atlantic Bight	g ww m <sup>-2</sup> yr <sup>-1</sup>	32	0.37	17917	0.18	2.78	Link et al. 2008
Narragansett Bay	mgC m <sup>-2</sup> yr <sup>-1</sup>	32	0.15	3917246	0.51	6.93	Monaco and Ulanowicz 1997
Southern New England Bight	g ww m <sup>-2</sup> yr <sup>-1</sup>	33	0.03	17597	0.16	2.56	Link et al. 2008
Chesapeake Bay	mgC m <sup>-2</sup> yr <sup>-1</sup>	36	0.09	3227453	0.19	2.76	Baird and Ulanowicz 1989
St. Marks Seagrass, site 1 (Jan)	mgC m <sup>-2</sup> d <sup>-1</sup>	51	0.08	1316	0.13	1.70	Baird et al. 1998
St. Marks Seagrass, site 1 (Feb)	mgC m <sup>-2</sup> d <sup>-1</sup>	51	0.08	1591	0.11	1.66	Baird et al. 1998
St. Marks Seagrass, site 2 (Jan)	mgC m <sup>-2</sup> d <sup>-1</sup>	51	0.07	1383	0.09	1.24	Baird et al. 1998
St. Marks Seagrass, site 2 (Feb)	mgC m <sup>-2</sup> d <sup>-1</sup>	51	0.08	1921	0.08	1.30	Baird et al. 1998
St. Marks Seagrass, site 3 (Jan)	mgC m <sup>-2</sup> d <sup>-1</sup>	51	0.05	12651	0.01	0.12	Baird et al. 1998
St. Marks Seagrass, site 4 (Feb)	mgC m <sup>-2</sup> d <sup>-1</sup>	51	0.08	2865	0.04	0.71	Baird et al. 1998
Sylt Rømø Bight	mgC m <sup>-2</sup> d <sup>-1</sup>	59	0.08	1353406	0.09	1.54	Baird et al. 2004a
Graminoids (wet)	gC m <sup>-2</sup> yr <sup>-1</sup>	66	0.18	13677	0.02	1.20	Ulanowicz et al. 2000
Graminoids (dry)	gC m <sup>-2</sup> yr <sup>-1</sup>	66	0.18	7520	0.04	1.41	Ulanowicz et al. 2000
Cypress (wet)	gC m <sup>-2</sup> yr <sup>-1</sup>	68	0.12	2572	0.04	0.62	Ulanowicz et al. 1997
Cypress (dry)	gC m <sup>-2</sup> yr <sup>-1</sup>	68	0.12	1918	0.04	0.70	Ulanowicz et al. 1997
Lake Oneida (pre-ZM) <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	74	0.22	1638	<0.01	0.17	Miehls et al. 2009a
Lake Quinte (pre-ZM) <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	74	0.21	1467	<0.01	0.15	Miehls et al. 2009b
Lake Oneida (post-ZM) <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	76	0.22	1365	<0.01	0.41	Miehls et al. 2009a
Lake Quinte (post-ZM) <sup>2</sup>	gC m <sup>-2</sup> yr <sup>-1</sup>	80	0.21	1925	0.01	0.30	Miehls et al. 2009b
Mangroves (wet)	gC m <sup>-2</sup> yr <sup>-1</sup>	94	0.15	3272	0.10	1.69	Ulanowicz et al. 1999
Mangroves (dry)	gC m <sup>-2</sup> yr <sup>-1</sup>	94	0.15	3266	0.10	1.70	Ulanowicz et al. 1999
Florida Bay (wet)	mgC m <sup>-2</sup> yr <sup>-1</sup>	125	0.12	2721	0.14	1.73	Ulanowicz et al. 1998
Florida Bay (dry)	mgC m <sup>-2</sup> yr <sup>-1</sup>	125	0.13	1779	0.08	1.29	Ulanowicz et al. 1998

<sup>a</sup>  $n$  is the number of nodes in the network model,  $C = L/n^2$  is the model connectance when  $L$  is the number of direct links or energy-matter transfers,  $TST = \sum \sum f_{i,j} + \sum z_i$  is the total system throughflow,  $FCI = 20$  the Finn Cycling Index, and  $I/D$  is the indirect/direct flow ratio.

797 **8 Figure Legends**

798

799 **Figure 1:** Hypothetical example ecosystem network with  $n = 4$ . A) Information regarding who  
800 is connected to whom in the network is in the adjacency matrix, **A**. The actual amount of energy  
801 flowing between the nodes is found in the flow matrix, **F**. Energy being transported across the  
802 system boundary into the nodes is represented in  $\bar{z}$ , and energy exiting the system across the  
803 boundary is in  $\bar{y}$ . B) Direct flow intensity is represented in **G**, and is the percentage of energy  
804 each node receives directly from each other node. When **G** is raised to a power  $m$ , the resulting  
805 matrices represent the indirect flow intensity between the nodes over all the paths of length  $m$ .

806

807 **Figure 2:** Indirect/direct (*I/D*) flow ratio for 50 empirically derived and trophically-based  
808 ecosystem models. Black bars are networks composing Model Set 2, those models that satisfy  
809 the second set of criteria of cycling structure and representation of the microbial food web.  
810 Extension of bars beyond  $I/D = 1$  indicates indirect flow dominance. The range of uncertainty  
811 between maximum and minimum values generated from our Monte-Carlo type uncertainty  
812 analysis is indicated for each model by the error bars. A few of these values have been  
813 previously published using an alternate calculation: <sup>1</sup>Jørgensen and Fath 2007; <sup>3</sup>Baird et al. 2007;  
814 <sup>4</sup>Borrett et al (2010). Only one value has been published using the calculation we use here:  
815 <sup>2</sup>Borrett et al. (2006).

816

817 **Figure 3:** Test of the relationship between indirect flow intensity (IFI) and (a) system size,  $n$ , (b)  
818 connectance,  $C$ , (c) direct flow intensity, DFI, and (d) Finn Cycling Index, FCI. There is no  
819 significant linear relationship between IFI and system size ( $r^2 = 0.069$ ;  $p = 0.066$ ) and  
820 connectance ( $r^2 = 0.022$ ;  $p = 0.31$ ).

821

822 **Figure 4:** Rank order percentage of Finn Cycling Index (FCI) that is mediated by components of  
823 the microbial food web in the ecosystem models in Model Set 2.

824


825 **Figure 5:** Average Environ Centrality for St. Marks seagrass system, site 1 (a, b), site 2 (c, d),  
826 site 3 (e), and site 4 (f). Average Environ Centrality is the percentage of TST mediated by each  
827 node. The nodes are ranked on the x-axis from highest to lowest Average Environ Centrality,  
828 with the three highest ranking nodes indicated.

829

829 Figure 1  
830

831

A Model



$$\mathbf{A} = \begin{bmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$\mathbf{F} = \begin{bmatrix} 0 & 5 & 3 & 3 \\ 15 & 0 & 0 & 0 \\ 0 & 9 & 0 & 0 \\ 0 & 0 & 5 & 0 \end{bmatrix}$$

$$\vec{z} = \begin{bmatrix} 4 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$


$$\vec{y} = \begin{bmatrix} 0 & 1 & 1 & 2 \end{bmatrix}$$

B Analysis


$$\mathbf{G} = \begin{bmatrix} 0 & .33 & .33 & .60 \\ 1 & 0 & 0 & 0 \\ 0 & .60 & 0 & 0 \\ 0 & 0 & .56 & 0 \end{bmatrix}$$

$$\mathbf{G}^2 = \begin{bmatrix} .33 & .20 & .33 & 0 \\ 0 & .33 & .33 & .60 \\ .60 & 0 & 0 & 0 \\ 0 & .33 & 0 & 0 \end{bmatrix}$$

$$\mathbf{G}^{10} = \begin{bmatrix} .12 & .10 & .09 & .07 \\ .12 & .12 & .10 & .10 \\ .10 & .07 & .07 & .05 \\ .05 & .05 & .04 & .04 \end{bmatrix}$$




832 Figure 3  
833



834  
835  
836  
837



837 Figure 4



838

