

Classification of 3D consistent quad-equations

Raphael Boll*

July 21, 2022

Abstract

We consider 3D consistent systems of six independent quad-equations assigned to the faces of a cube. The well-known classification of 3D consistent quad-equations, the so-called ABS-list, is included in this situation. The extension of these equations to the whole lattice \mathbb{Z}^3 is possible by reflecting the cubes. For every quad-equation we will give at least one system included leading to one of its zero-curvature representations.

PACS number: 02.30.Ik

1 Introduction

One of the definitions of integrability of lattice equations, which becomes increasingly popular in the recent years, is based on the notion of multidimensional consistency. For two-dimensional lattices, this notion was clearly formulated first in [NW01], and it was proposed to use as a synonym of integrability in [BS02, Nij02]. The outstanding importance of 3D consistency in the theory of discrete integrable systems became evident no later than with the appearance of the well-known ABS-classification of integrable equations in [ABS03]. In that article Adler, Bobenko and Suris used a definition of 3D consistency only allowing equations on the faces of a cube, which differ only by the parameter values assigned to the edges of a cube. In [Atk08] and [BS10] appeared a lot of systems of quad-equations not satisfying this strict definition of 3D consistency. However, these systems can also be seen as families of Bäcklund transformations and

*Institut für Mathematik, MA 7-2, Technische Universität Berlin, Str. des 17. Juni 136, 10623 Berlin, Germany; e-mail: boll@math.tu-berlin.de

they lead to zero curvature representations of participating quad-equations in the same way (see [BS02]).

As already done in [ABS09] the definition of 3D consistency can be extended: In contrast to the restriction, that all faces of a cube must carry the same equation up to parameters assigned to edges of the cube, we will allow different equations on all faces of a cube. The classification in that article is restricted to so-called equations of type Q, i.e. those whose biquadratics are all non-degenerate (we will give a precise definition in the next section). The present paper is devoted to systems containing equations which are not necessarily of type Q. Our classification will cover systems appearing in [Atk08] and all systems in [BS10], as well as equations which are equivalent to the Hietarinta equation [Hie04], to the “new” equation in [HV10] and to the equation in [LY09]. Moreover, it will contain also many novel systems.

In addition, every system in this classification can be extended to the whole net \mathbb{Z}^3 by reflecting the cubes.

The outline of our approach is the following: In Section 2 we will present a complete classification of a single quad-equation modulo Möbius transformations acting independently on the fields at the four vertices of an elementary quadrilateral. In Section 3 we will give a classification of 3D consistent systems of quad-equations possessing the so-called tetrahedron property modulo Möbius transformations acting independently on the fields at the eight vertices of an elementary cube.

2 Quad-Equations on Single Quadrilaterals

At the beginning we will introduce some objects and notations. We will start with the most important one, the *quad-equation* $Q(x_1, x_2, x_3, x_4) = 0$, where $Q \in \mathbb{C}[x_1, x_2, x_3, x_4]$ is an irreducible multi-affine polynomial.

Very useful tools characterizing quad-equations, are the *biquadratics*. We define them for every permutation (i, j, k, ℓ) of $(1, 2, 3, 4)$ as follows

$$Q^{i,j} = Q^{i,j}(x_i, x_j) = Q_{x_k} Q_{x_\ell} - Q Q_{x_k, x_\ell}.$$

One can show that $Q^{i,j} \neq 0$. A biquadratic $Q^{1,2}$ is called *non-degenerate* if no polynomial in its equivalence class with respect to Möbius transformations in x_1 and x_2 is divisible by a factor $x_1 - \alpha_1$ or $x_2 - \alpha_2$ (with $\alpha_i \in \mathbb{C}$). Otherwise, $Q^{1,2}$ is called *degenerate*. The following Lemma gives some informations about the relation between non-degenerate biquadratics of a quad-equation:

Lemma 2.1. *Biquadratics on opposite edges (we consider the two diagonals as opposite edges, too) are either both degenerate or both non-degenerate.*

Moreover, it follows that the number of non-degenerate biquadratics of a quad-equation is even. Another restriction for biquadratics of a quad-equation comes along with the next Lemma:

Lemma 2.2. *There do not exist any quad-equations with exactly two degenerate biquadratics.*

A multi-affine polynomial Q is of *type Q* if all its biquadratics are non-degenerate. Otherwise it is of *type H1* if two out of six biquadratics are non-degenerate and of *type H2* if all six biquadratics are degenerate.

For every permutation (i, j, k, ℓ) of $(1, 2, 3, 4)$ the quartic polynomial

$$r^i = r^i(x_i) = \left(Q_{x_\ell}^{j,k}\right)^2 - 2Q^{j,k}Q_{x_\ell, x_\ell}^{j,k}$$

is called a corresponding *discriminant*. This polynomial turns out to be independent on permutations of (j, k, ℓ)

The Theorem 2 in [ABS09] and independently also [IR02] gives a complete classification of biquadratics up to Möbius transformations.

The group $(\text{Möb})^4$ acts on quad-equations by Möbius transformations on all fields independently.

2.1 Quad-Equations of Type Q

Quad-equations of type Q were already classified in [ABS09]. Every quad-equation of type Q is equivalent modulo $(\text{Möb})^4$ to one of the following quad-equations characterized by the quadruples of discriminants:

- $(\delta, \delta, \delta, \delta)$:

$$Q = \alpha_2(x_1 - x_2)(x_3 - x_4) - \alpha_1(x_1 - x_4)(x_2 - x_3) + \delta\alpha_1\alpha_2(\alpha_1 + \alpha_2) \quad (Q_1)$$

- (x_1, x_2, x_3, x_4) :

$$Q = \alpha_2(x_1 - x_2)(x_3 - x_4) - \alpha_1(x_1 - x_4)(x_2 - x_3) + \alpha_1\alpha_2(\alpha_1 + \alpha_2)(x_1 + x_2 + x_3 + x_4) - \alpha_1\alpha_2(\alpha_1 + \alpha_2)(\alpha_1^2 + \alpha_1\alpha_2 + \alpha_2^2) \quad (Q_2)$$

- $(x_1^2 - \delta, x_2^2 - \delta, x_3^2 - \delta, x_4^2 - \delta)$:

$$Q = (\alpha_1 - \alpha_1^{-1})(x_1x_2 + x_3x_4) + (\alpha_2 - \alpha_2^{-1})(x_1x_4 + x_2x_3) - (\alpha_1\alpha_2 - \alpha_1^{-1}\alpha_2^{-1})(x_1x_3 + x_2x_4) + \frac{\delta}{4}(\alpha_1 - \alpha_1^{-1})(\alpha_2 - \alpha_2^{-1})(\alpha_1\alpha_2 - \alpha_1^{-1}\alpha_2^{-1}) \quad (Q_3)$$

- $((x_1 - 1)(k^2x_1^2 - 1), (x_2 - 1)(k^2x_2^2 - 1), (x_3 - 1)(k^2x_3^2 - 1), (x_4 - 1)(k^2x_4^2 - 1))$:

$$Q = \text{sn}(\alpha_1)\text{sn}(\alpha_2)\text{sn}(\alpha_1 + \alpha_2)(k^2x_1x_2x_3x_4 + 1) - \text{sn}(\alpha_1)(x_1x_2 + x_3x_4) - \text{sn}(\alpha_2)(x_1x_4 + x_2x_3) + \text{sn}(\alpha_1 + \alpha_2)(x_1x_3 + x_2x_4) \quad (Q_4)$$

2.2 Quad-Equations of Type H1

A complete classification of type H1 quad-equations was not known before. Every quad-equation of type H1 is equivalent modulo $(\text{Möb})^4$ to one of the following quad-equations characterized by the quadruples of discriminants:

- $(\epsilon, 0, \epsilon, 0)$:

$$Q = (x_1 - x_3)(x_2 - x_4) + (\alpha_2 - \alpha_1)(1 + \epsilon x_2 x_4) \quad (H_1^\epsilon)$$

- $(\epsilon x_1 + 1, 1, \epsilon x_3 + 1, 1)$:

$$Q = (x_1 - x_3)(x_2 - x_4) + (\alpha_2 - \alpha_1)(x_1 + x_2 + x_3 + x_4) + \alpha_2^2 - \alpha_1^2 \\ + \epsilon(\alpha_2 - \alpha_1)(2x_2 + \alpha_1 + \alpha_2)(2x_4 + \alpha_1 + \alpha_2) + \epsilon(\alpha_2 - \alpha_1)^3 \quad (H_2^\epsilon)$$

- $(x_1^2 + \delta\epsilon, x_2^2, x_3^2 + \delta\epsilon, x_4^2)$:

$$Q = \alpha_1(x_1 x_2 + x_3 x_4) - \alpha_2(x_1 x_4 + x_2 x_3) + (\alpha^2 - \beta^2) \left(\delta + \frac{\epsilon x_2 x_4}{\alpha_1 \alpha_2} \right) \quad (H_3^\epsilon)$$

Remark. All these equations were already mentioned in [ABS09].

2.3 Quad-Equations of Type H2

A complete classification of type H2 quad-equations was not known before. Every quad-equation of type H2 is equivalent modulo $(\text{Möb})^4$ to one of the following quad-equations characterized by the quadruples of discriminants:

- $(0, 0, 0, 0)$:

$$Q = x_1 + x_2 + x_3 + x_4$$

- $(1, 0, 1, \delta)$:

$$Q = x_1 + x_3 + x_2(x_4 + \delta x_1)$$

- $(x_1^2, x_2^2, x_3^2, x_4^2)$:

$$Q = x_1 x_3 + x_2 x_4 + \delta_1 x_2 x_3 + \delta_2 x_3 x_4$$

3 Quad-Equations on the Faces of a Cube

We will now consider systems of the type

$$\begin{aligned} A(x, x_1, x_2, x_{12}) &= 0, & \bar{A}(x_3, x_{13}, x_{23}, x_{123}) &= 0, \\ B(x, x_2, x_3, x_{23}) &= 0, & \bar{B}(x_1, x_{12}, x_{13}, x_{123}) &= 0, \\ C(x, x_1, x_3, x_{13}) &= 0, & \bar{C}(x_2, x_{12}, x_{23}, x_{123}) &= 0 \end{aligned} \quad (1)$$




Figure 1: Equations on a Cube

where the equations A, \dots, \bar{C} are quad-equations assigned to the faces of a cube in the manner demonstrated in Figure 1. Such a system is *3D consistent* if the three values for x_{123} (calculated by using $\bar{A} = 0$, $\bar{B} = 0$ or $\bar{C} = 0$) coincide for arbitrary initial data x, x_1, x_2 and x_3 . It possesses the *tetrahedron property* if two equations

$$K(x, x_{12}, x_{13}, x_{23}) = 0 \text{ and } \bar{K}(x_1, x_2, x_3, x_{123}) = 0$$

are satisfied for every solution of the system. It can be shown that the polynomials K and \bar{K} are multi-affine and irreducible.

The group $(\text{Möb})^8$ acts on such a system by Möbius transformations on all vertex fields independently.

We classify all 3D consistent systems 1 with a tetrahedron property, whereas the classification of the case, that all quad-equations are of type Q, was already done in [ABS09]. The proofs are omitted because they are too long for this article. Note, that in many cases the tetrahedron property is a consequence of the other assumptions.

3.1 Two Equations of Type Q and Four Equations of Type H1

In this section we consider systems (1) with

- B and \bar{B} of type Q,
- A, \bar{A}, C and \bar{C} of type H1 and
- the non-degenerate biquadratics of \bar{A}, C and \bar{C} on edges neighboring B or \bar{B} .

Below is the list of all 3D consistent systems modulo $(\text{Möb})^8$ with these properties. It turns out that their tetrahedron property follows from the above assumptions.

Theorem 3.1. *Every 3D consistent system (1) satisfying the properties of this section is equivalent modulo (Möb)⁸ to one of the following systems which can be characterized by the quadruples of discriminants of A:*

- $(\epsilon, 0, \epsilon, 0)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}; \alpha, \beta) &= (x - x_2)(x_1 - x_{12}) - (\alpha - \beta)(1 + \epsilon x_1 x_{12}), \\ B(x, x_2, x_3, x_{23}; \epsilon) &= (\beta - \gamma)(x - x_2)(x_3 - x_{23}) - (\alpha - \beta)(x - x_{23})(x_2 - x_3) \\ &\quad - \epsilon(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma) \end{aligned}$$

- $(\epsilon x + 1, 1, \epsilon x_2 + 1, 1)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}; \alpha, \beta) &= (x - x_2)(x_1 - x_{12}) - (\alpha - \beta)(x + x_1 + x_2 + x_{12}) \\ &\quad - \alpha^2 + \beta^2 - \epsilon(\alpha - \beta)(2x_1 + \alpha + \beta)(2x_{12} + \alpha + \beta) - \epsilon(\alpha - \beta)^3, \\ B(x, x_2, x_3, x_{23}; \epsilon) &= (\beta - \gamma)(x - x_2)(x_3 - x_{23}) - (\alpha - \beta)(x - x_{23})(x_2 - x_3) \\ &\quad + (\alpha - \beta)(\beta - \gamma)(\alpha - \gamma) - 2\epsilon(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma)(x + x_2 + x_3 + x_{23}) \\ &\quad - 4\epsilon^2(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma) \left((\alpha - \beta)^2 + (\alpha - \beta)(\beta - \gamma) + (\beta - \gamma)^2 \right) \end{aligned}$$

- $(x^2 + \delta\epsilon, x_1^2, x_2^2 + \delta\epsilon, x_{12}^2)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}; \alpha, \beta) &= \alpha(xx_1 + x_2x_{12}) - \beta(xx_{12} + x_1x_2) \\ &\quad + (\alpha^2 - \beta^2) \left(\delta + \frac{\epsilon x_1 x_{12}}{\alpha\beta} \right), \\ B(x, x_2, x_3, x_{23}; \epsilon) &= \gamma(\alpha^2 - \beta^2)(xx_2 + x_3x_{23}) - \beta(\alpha^2 - \gamma^2)(xx_3 + x_2x_{23}) \\ &\quad + \alpha(\beta^2 - \gamma^2)(xx_{23} + x_2x_3) + \frac{\delta\epsilon(\alpha^2 - \beta^2)(\alpha^2 - \gamma^2)(\beta^2 - \gamma^2)}{\alpha\beta\gamma} \end{aligned}$$

with

$$\begin{aligned} \bar{A} &= A(x_3, x_{13}, x_{23}, x_{123}; \alpha, \beta), & \bar{B} &= B(x_1, x_{12}, x_{13}, x_{123}; 0), \\ \bar{C} &= A(x, x_1, x_3, x_{13}; \alpha, \gamma), & \bar{C} &= A(x_2, x_{12}, x_{23}, x_{123}; \alpha, \gamma), \\ \bar{K} &= A(x, x_{12}, x_{23}, x_{13}; \beta, \gamma), & \bar{K} &= A(x_2, x_1, x_3, x_{123}; \beta, \gamma). \end{aligned}$$

Remark. Equivalent systems appeared in [ABS09, Atk08, BS10].

3.2 Six Equations of Type H1, First Case

In this section we consider systems (1) with

- $A, \bar{A}, \dots, \bar{C}$ of type H1 and

- all non-degenerate biquadratics on diagonals of faces.

Below is the list of all 3D consistent systems modulo (Möb)⁸ with these properties and with the tetrahedron property. It turns out that their tetrahedron property follows from the above assumptions except for the system characterized by the quadruple $(\epsilon, 0, 0, \epsilon)$.

Theorem 3.2. *Every 3D consistent system (1) satisfying the properties of this section and possessing the tetrahedron property is equivalent modulo (Möb)⁸ to one of the following systems which can be characterized by the quadruples of discriminants of A :*

- $(\epsilon, 0, 0, \epsilon)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}; \alpha, \beta) &= (x - x_{12})(x_1 - x_2) - (\alpha - \beta)(1 + \epsilon x_1 x_2), \\ K(x, x_{12}, x_{13}, x_{23}; \epsilon) &= (\beta - \gamma)(x - x_{12})(x_{13} - x_{23}) - (\alpha - \beta)(x - x_{23})(x_{12} - x_{13}) \\ &\quad - \epsilon(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma) \end{aligned}$$

- $(\epsilon x + 1, 1, 1, \epsilon x_{12} + 1)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}; \alpha, \beta) &= (x - x_{12})(x_1 - x_2) - (\alpha - \beta)(x + x_1 + x_2 + x_{12}) - \alpha^2 \\ &\quad + \beta^2 - \epsilon(\alpha - \beta)(2x_1 + \alpha + \beta)(2x_2 + \alpha + \beta) - \epsilon(\alpha - \beta)^3, \\ K(x, x_{12}, x_{13}, x_{23}; \epsilon) &= (\beta - \gamma)(x - x_{12})(x_{13} - x_{23}) - (\alpha - \beta)(x - x_{23})(x_{12} - x_{13}) \\ &\quad + (\alpha - \beta)(\beta - \gamma)(\alpha - \gamma) - 2\epsilon(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma)(x + x_{12} + x_{13} + x_{23}) \\ &\quad - 4\epsilon^2(\alpha - \beta)(\beta - \gamma)(\alpha - \gamma)\left((\alpha - \beta)^2 + (\alpha - \beta)(\beta - \gamma) + (\beta - \gamma)^2\right) \end{aligned}$$

- $(x^2 + \delta\epsilon, x_1^2, x_2^2, x_{12}^2 + \delta\epsilon)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}; \alpha, \beta) &= \alpha(xx_1 + x_2x_{12}) - \beta(xx_2 + x_1x_{12}) \\ &\quad + (\alpha^2 - \beta^2)\left(\delta + \frac{\epsilon x_1 x_2}{\alpha\beta}\right), \\ K(x, x_{12}, x_{13}, x_{23}; \epsilon) &= \gamma(\alpha^2 - \beta^2)(xx_{12} + x_{13}x_{23}) - \beta(\alpha^2 - \gamma^2)(xx_{13} + x_{12}x_{23}) \\ &\quad + \alpha(\beta^2 - \gamma^2)(xx_{23} + x_{12}x_{13}) + \frac{\delta\epsilon(\alpha^2 - \beta^2)(\alpha^2 - \gamma^2)(\beta^2 - \gamma^2)}{\alpha\beta\gamma} \end{aligned}$$

with

$$\begin{aligned} \bar{A} &= A(x_{13}, x_3, x_{123}, x_{23}; \alpha, \beta), & B &= A(x, x_2, x_3, x_{23}; \beta, \gamma), \\ \bar{B} &= A(x_{12}, x_1, x_{123}, x_{13}; \beta, \gamma), & C &= A(x, x_1, x_3, x_{13}; \alpha, \gamma), \\ \bar{C} &= A(x_{12}, x_2, x_{123}, x_{23}; \alpha, \gamma), & \bar{K} &= K(x_1, x_2, x_3, x_{123}; 0). \end{aligned}$$

Remark. Equivalent systems appeared in [ABS03, ABS09].

3.3 Six Equations of Type H1, Second Case

In this section we consider systems (1) with

- $A, \bar{A}, \dots, \bar{C}$ of type H1,
- the non-degenerate biquadratics of A and \bar{A} on diagonals of faces and
- the non-degenerate biquadratics of B, \bar{B}, C and \bar{C} on edges not neighboring A and \bar{A} .

Below is the list of all 3D consistent systems modulo (Möb)⁸ with these properties. It turns out that their tetrahedron property follows from the above assumptions.

Theorem 3.3. *Every 3D consistent system (1) satisfying the properties of this section is equivalent modulo (Möb)⁸ to one of the following systems which can be characterized by the quadruples of discriminants of A :*

- $(\epsilon, 0, 0, \epsilon)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}) &= (x - x_{12})(x_1 - x_2) - (\alpha - \beta)(1 + \epsilon x_1 x_2), \\ B(x, x_2, x_3, x_{23}; \beta) &= (x - x_3)(x_2 - x_{23}) + \gamma(1 + \epsilon x_2 x_{23}), \\ K(x, x_{12}, x_{13}, x_{23}) &= (x - x_{12})(x_{13} - x_{23}) - (\alpha - \beta)(1 + \epsilon x_{13} x_{23}) \end{aligned}$$

- $(\epsilon x + 1, 1, 1, \epsilon x_{12} + 1)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}) &= (x - x_{12})(x_1 - x_2) - (\alpha - \beta)(x + x_1 + x_2 + x_{12}) - \alpha^2 + \beta^2 \\ &\quad - \epsilon(\alpha - \beta)(2x_1 + \alpha + \beta)(2x_2 + \alpha + \beta) - \epsilon(\alpha - \beta)^3, \\ B(x, x_2, x_3, x_{23}; \beta) &= (x - x_3)(x_2 - x_{23}) + \gamma(x + x_2 + x_3 + x_{23}) + \gamma^2 + 2\beta\gamma \\ &\quad + \epsilon\gamma(2x_2 + 2\beta + \gamma)(2x_{23} + 2\beta + \gamma) + \gamma^3\epsilon, \\ K(x, x_{12}, x_{13}, x_{23}) &= (x - x_{12})(x_{13} - x_{23}) - (\alpha - \beta)(x + x_{12} + x_{13} + x_{23}) - \alpha^2 \\ &\quad + \beta^2 - 2\gamma(\alpha - \beta) - \epsilon(\alpha - \beta)(2x_{13} + \alpha + \beta + 2\gamma)(2x_{23} + \alpha + \beta + 2\gamma) \\ &\quad - \epsilon(\alpha - \beta)^3 \end{aligned}$$

- $(x^2 + \delta\epsilon, x_1^2, x_2^2, x_{12}^2 + \delta\epsilon)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}) &= \alpha(xx_1 + x_2x_{12}) - \beta(xx_2 + x_1x_{12}) + (\alpha^2 - \beta^2) \left(\delta + \frac{\epsilon x_1 x_2}{\alpha\beta} \right), \\ B(x, x_2, x_3, x_{23}; \beta) &= xx_2 + x_3x_{23} - \gamma(xx_{23} + x_2x_3) - (\gamma^2 - 1) \left(\delta\beta + \frac{\epsilon x_2 x_{23}}{\beta\gamma} \right), \\ K(x, x_{12}, x_{13}, x_{23}) &= \alpha(xx_{13} + x_{12}x_{23}) - \beta(xx_{23} + x_{12}x_{13}) \\ &\quad + (\alpha^2 - \beta^2) \left(\delta\gamma + \frac{\epsilon x_{13} x_{23}}{\alpha\beta\gamma} \right) \end{aligned}$$

with

$$\begin{aligned} \bar{A} &= A(x_3, x_{13}, x_{23}, x_{123}), & \bar{B} &= B(x_{12}, x_1, x_{123}, x_{13}; \beta), & C &= B(x, x_1, x_3, x_{13}; \alpha), \\ \bar{C} &= B(x_{12}, x_2, x_{123}, x_{23}; \alpha), & \bar{K} &= K(x_3, x_{123}, x_1, x_2). \end{aligned}$$

Remark. These systems were not known before.

3.4 Four Equations of Type H1 and Two Equations of Type H2, First Case

Theorem 3.4. *No 3D consistent system (1) exists with the tetrahedron property and with*

- A and \bar{A} of type H2,
- B, \bar{B}, C and \bar{C} of type H1,
- the non-degenerate biquadratics of B, \bar{B}, C and \bar{C} on diagonals.

3.5 Four Equations of Type H1 and Two Equations of Type H2, Second Case

In this section we consider systems (1) with

- A and \bar{A} of type H2,
- B, \bar{B}, C and \bar{C} of type H1 and
- the non-degenerate biquadratics of B, \bar{B}, C and \bar{C} on edges not neighboring A and \bar{A} .

Below is the list of all 3D consistent systems modulo (Möb)⁸ with these properties and with the tetrahedron property. It turns out that in the last case the tetrahedron property follows from the above assumptions.

Theorem 3.5. *Every 3D consistent system (1) satisfying the properties of this section and possessing the tetrahedron property is equivalent modulo (Möb)⁸ to one of the following systems which can be characterized by the quadruples of discriminants of A :*

- $(0, 0, 0, 0)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}) &= x + x_1 + x_2 + x_{12}, \\ B(x, x_2, x_3, x_{23}; \alpha) &= (x - x_3)(x_2 - x_{23}) + \alpha \end{aligned}$$

with

$$\begin{aligned} \bar{A} &= A(x_3, x_{13}, x_{23}, x_{123}), & \bar{B} &= B(x_1, x_{12}, x_{13}, x_{123}; \alpha), \\ C &= B(x, x_1, x_3, x_{13}; -\alpha), & \bar{C} &= B(x_2, x_{12}, x_{23}, x_{123}; -\alpha), \\ K &= A(x, x_{12}, x_{13}, x_{23}), & \bar{K} &= A(x_1, x_2, x_3, x_{123}). \end{aligned}$$

- $(1, 1, 0, \delta_1 \delta_2)$:

$$B(x, x_2, x_3, x_{23}; \delta_2) = (x - x_3)(x_2 - x_{23}) + \delta_1 \alpha (x_2 + x_{23}) + \delta_2 \alpha (x + x_3) + \delta_1 \delta_2 \alpha^2,$$

$$\begin{aligned} \bar{B}(x_1, x_{12}, x_{13}, x_{123}; \delta_1) &= (x_1 - x_{13})(x_{12} - x_{123}) + 2(\delta_1 - 1)\alpha \\ &\quad + (\delta_1 \delta_2 + \delta_1 - \delta_2)\alpha(x_{12} + x_{123}) + 2\delta_1 \delta_2 \alpha x_{12} x_{123}, \end{aligned}$$

$$K(x, x_{13}, x_{23}, x_{12}; \alpha) = x + x_{13} + (\delta_1 - 1)x_{23} + \delta_1 \alpha + x_{12}(\delta_2 x + \delta_1 x_{23} + \delta_1 \delta_2 \alpha)$$

with

$$\begin{aligned} A &= K(x, x_1, x_2, x_{12}; 0), & \bar{A} &= K(x_3, x_{13}, x_{23}, x_{123}), \\ C &= B(x, x_1, x_3, x_{13}; \delta_1 + \delta_2 - \delta_1 \delta_2), & \bar{C} &= \bar{B}(x_2, x_{12}, x_{23}, x_{123}; 0), \\ \bar{K} &= K(x_3, x_1, x_2, x_{123}; \alpha). \end{aligned}$$

Remark. If $\delta_1 = \delta_2 = 0$, A is reducible.

- $(x^2, x_1^2, x_{12}^2, x_2^2)$:

$$A(x, x_1, x_2, x_{12}; \delta_1, \delta_2) = xx_{12} + x_1 x_2 + \delta_1 x_1 x_{12} + \delta_2 x_2 x_{12},$$

$$B(x, x_2, x_3, x_{23}; \alpha) = xx_{23} + x_2 x_3 - \alpha(xx_2 + x_3 x_{23}) - \delta_2(\alpha^2 - 1)x_2 x_{23}$$

with

$$\begin{aligned} \bar{A} &= A(x_3, x_{13}, x_{23}, x_{123}; \delta_1, \delta_2), & \bar{B} &= B(x_1, x_{12}, x_{13}, x_{123}; \alpha), \\ C &= B(x, x_1, x_3, x_{13}; \alpha^{-1}), & \bar{C} &= B(x_2, x_{12}, x_{23}, x_{123}; \alpha^{-1}), \\ K &= A(x, x_{13}, x_{23}, x_{12}; \delta_1 \alpha^{-1}, \delta_2 \alpha), & \bar{K} &= A(x_3, x_1, x_2, x_{123}; \delta_1 \alpha^{-1}, \delta_2 \alpha) \end{aligned}$$

or

$$A(x, x_1, x_2, x_{12}; \delta_1) = xx_2 + x_1 x_{12} + \delta_1 x_1 x_2 + \delta_2 x_2 x_{12},$$

$$B(x, x_2, x_3, x_{23}; \delta_1 \delta_2) = xx_{23} + x_2 x_3 - \alpha(xx_2 + x_3 x_{23}) + \delta_1 \delta_2(\alpha^2 - 1)x_2 x_{23}$$

with

$$\begin{aligned} \bar{A} &= A(x_3, x_{13}, x_{23}, x_{123}; \delta_1), & \bar{B} &= B(x_{123}, x_{13}, x_{12}, x_1; 0), \\ C &= B(x, x_1, x_3, x_{13}; -\delta_1), & \bar{C} &= B(x_{123}, x_{23}, x_{12}, x_2; -\delta_1), \\ K &= A(x, x_{13}, x_{23}, x_{12}; \delta_1 \alpha), & \bar{K} &= A(x_3, x_1, x_2, x_{123}; \delta_1 \alpha). \end{aligned}$$

Remark. These systems were not known before. Special cases of equations $A=0$ in the last two systems are equivalent modulo (Möb)⁴ to the equations whose integrability was shown in [Hie04, LY09] and the “new” integrable equation from [HV10].

3.6 Two Equations of Type H1 and Four Equations of Type H2

In this section we consider systems (1) with

- A, \bar{A}, C and \bar{C} of type H2,
- B and \bar{B} of type H1 and
- the non-degenerate biquadratics of B and \bar{B} on diagonals.

Below is the list of all 3D consistent systems modulo (Möb)⁸ with these properties and with the tetrahedron property.

Theorem 3.6. *Every 3D consistent system (1) satisfying the properties of this section is equivalent modulo (Möb)⁸ to one of the following systems which can be characterized by the quadruples of discriminants of A :*

- $(0, 0, 0, 0)$:

$$\begin{aligned} A(x, x_1, x_2, x_{12}) &= x + x_1 + x_2 + x_{12}, \\ B(x, x_2, x_3, x_{23}; \alpha) &= (x - x_{23})(x_2 - x_3) + \alpha \end{aligned}$$

with

$$\begin{aligned} \bar{A} &= A(x_3, x_{13}, x_{23}, x_{123}), & \bar{B} &= B(x_1, x_{12}, x_{13}, x_{123}; \alpha), \\ C &= A(x, x_1, x_3, x_{13}), & \bar{C} &= A(x_2, x_{12}, x_{23}, x_{123}), \\ K &= B(x, x_{12}, x_{13}, x_{23}; -\alpha), & \bar{K} &= B(x_1, x_2, x_3, x_{123}; -\alpha). \end{aligned}$$

- $(1, 1, 0, \delta_1\delta_2)$:

$$\begin{aligned} B(x, x_2, x_3, x_{23}; \delta_2) &= (x - x_{23})(x_2 - x_3) + \delta_1\alpha(x_2 + x_3) + \delta_2\alpha(x + x_{23}) + \delta_1\delta_2\alpha^2, \\ \bar{B}(x_1, x_{12}, x_{13}, x_{123}; \delta_1) &= (x_1 - x_{123})(x_{12} - x_{13}) + 2(\delta_1 - 1)\alpha \\ &\quad + (\delta_1\delta_2 + \delta_1 - \delta_2)\alpha(x_1 + x_{123}) + 2\delta_1\delta_2\alpha x_1 x_{123}, \\ C(x, x_1, x_3, x_{13}; \alpha) &= x + x_{13} + (\delta_1 - 1)x_3 + \delta_1\alpha + x_1(\delta_2x + \delta_1x_3 + \delta_1\delta_2\alpha) \end{aligned}$$

with

$$\begin{aligned} A &= C(x, x_1, x_2, x_{12}; 0), & \bar{A} &= C(x_{23}, x_{123}, x_3, x_{13}; 0), \\ \bar{C} &= C(x_{23}, x_{123}, x_2, x_{12}; \alpha), & K &= B(x, x_{12}, x_{13}, x_{23}; \delta_1 + \delta_2 - \delta_1\delta_2), \\ \bar{K} &= \bar{B}(x_1, x_2, x_3, x_{123}; 0) \end{aligned}$$

or

$$\begin{aligned}
 C(x, x_1, x_3, x_{13}; \alpha) &= x + x_3 + (\delta_1 - 1)x_{13} + \delta_1\alpha + x_1(\delta_2x + \delta_1x_{13} + \delta_1\delta_2\alpha), \\
 K(x, x_{12}, x_{13}, x_{23}; \delta_2) &= (x - x_{23})(x_{12} - x_{13}) + \delta_1\alpha(x_{12} + x_{13}) + \delta_2\alpha(x + x_{23}) \\
 &\quad + \delta_1\delta_2\alpha^2, \\
 \bar{K}(x_1, x_2, x_3, x_{123}; \delta_1) &= (x_1 - x_{123})(x_2 - x_3) + 2(\delta_1 - 1)\alpha \\
 &\quad + (\delta_1\delta_2 + \delta_1 - \delta_2)\alpha(x_1 + x_{123}) + 2\delta_1\delta_2\alpha x_1x_{123}
 \end{aligned}$$

with

$$\begin{aligned}
 A &= C(x, x_1, x_2, x_{12}; 0), & \bar{A} &= C(x_{23}, x_{123}, x_3, x_{13}; 0), \\
 B &= K(x, x_2, x_3, x_{23}; \delta_1 + \delta_2 - \delta_1\delta_2), & \bar{B} &= \bar{K}(x_1, x_{12}, x_{13}, x_{123}; 0), \\
 \bar{C} &= C(x_{23}, x_{123}, x_2, x_{12}; \alpha).
 \end{aligned}$$

Remark. If $\delta_1 = \delta_2 = 0$, A is reducible.

- $(x^2, x_1^2, x_{12}^2, x_2^2)$:

$$\begin{aligned}
 A(x, x_1, x_2, x_{12}; \delta_1, \delta_2) &= xx_1 + x_2x_{12} + \delta_1x_1x_{12} + \delta_2x_1x_2, \\
 B(x, x_2, x_3, x_{23}; \alpha) &= xx_3 + x_2x_{23} - \alpha(xx_2 + x_3x_{23}) - \delta_2(\alpha^2 - 1)x_2x_3
 \end{aligned}$$

with

$$\begin{aligned}
 \bar{A} &= A(x_{23}, x_{123}, x_3, x_{13}; \delta_1, \delta_2), & \bar{B} &= B(x_{12}, x_1, x_{123}, x_{13}; \alpha), \\
 C &= A(x, x_1, x_3, x_{13}; \delta_1\alpha^{-1}, \delta_2\alpha), & \bar{C} &= A(x_{23}, x_{123}, x_2, x_{12}; \delta_1\alpha^{-1}, \delta_2\alpha), \\
 K &= B(x, x_{12}, x_{13}, x_{23}; \alpha^{-1}), & \bar{K} &= B(x_2, x_1, x_{123}, x_3; \alpha^{-1})
 \end{aligned}$$

or

$$\begin{aligned}
 A(x, x_1, x_2, x_{12}; \delta_1) &= xx_2 + x_1x_{12} + \delta_1x_2x_{12} + \delta_2x_1x_2, \\
 B(x, x_2, x_3, x_{23}; \delta_1\delta_2) &= xx_3 + x_2x_{23} - \alpha(xx_2 + x_3x_{23}) + \delta_1\delta_2(\alpha^2 - 1)x_2x_3
 \end{aligned}$$

with

$$\begin{aligned}
 \bar{A} &= A(x_{23}, x_{123}, x_3, x_{13}; \delta_1), & \bar{B} &= B(x_1, x_{12}, x_{13}, x_{123}; 0), \\
 C &= A(x, x_1, x_3, x_{13}; \delta_1\alpha), & \bar{C} &= A(x_{23}, x_{123}, x_2, x_{12}; \delta_1\alpha), \\
 K &= B(x, x_{12}, x_{13}, x_{23}; -\delta_1), & \bar{K} &= B(x_1, x_2, x_3, x_{123}; -\delta_1)
 \end{aligned}$$

or

$$\begin{aligned}
 A(x, x_1, x_2, x_{12}; \delta_1) &= xx_{12} + x_1x_2 + \delta_1x_2x_{12} + \delta_2x_1x_{12}, \\
 B(x, x_2, x_3, x_{23}; -\delta_1) &= xx_3 + x_2x_{23} - \alpha(xx_2 + x_3x_{23}) - \delta_1(\alpha^2 - 1)x_2x_3
 \end{aligned}$$

with

$$\begin{aligned}\bar{A} &= A(x_{23}, x_{123}, x_3, x_{13}; \delta_1), & \bar{B} &= B(x_1, x_{12}, x_{13}, x_{123}; -\delta_1), \\ C &= A(x, x_1, x_3, x_{13}; \delta_1\alpha), & \bar{C} &= A(x_{23}, x_{123}, x_2, x_{12}; \delta_1\alpha), \\ K &= B(x, x_{12}, x_{13}, x_{23}; \delta_1\delta_2), & \bar{K} &= B(x_1, x_2, x_3, x_{123}; 0).\end{aligned}$$

Remark. Except two special cases (see [Atk08]) these systems were not known before.

3.7 Six Equations of Type H2

In this section we consider systems (1) with $A, \bar{A}, \dots, \bar{C}$ of type H2. Below is the list of all those systems possessing the tetrahedron property.

Theorem 3.7. *Every 3D consistent system (1) satisfying the properties of this section is equivalent modulo $(M\ddot{o}b)^8$ to one of the following systems which can be characterized by the quadruples of discriminants of A :*

- $(0, 0, 0, 0)$:

$$K(x, x_{12}, x_{13}, x_{23}; \alpha, \beta) = (\alpha - \alpha\beta + \beta)x - \alpha\beta x_{12} - \alpha x_{13} + \beta x_{23}$$

with

$$\begin{aligned}A &= K(x, x_1, x_2, x_{12}; -1, 1), & \bar{A} &= K(x_{13}, x_3, x_{123}, x_{23}; \alpha\beta - \alpha - \beta, 1), \\ B &= K(x, x_2, x_3, x_{23}; -\alpha, 1), & \bar{B} &= K(x_{12}, x_1, x_{123}, x_{13}; (\alpha\beta - \alpha - \beta)/\alpha, 1), \\ C &= K(x, x_1, x_{13}, x_3; 1, -\beta), & \bar{C} &= K(x_{12}, x_2, x_{23}, x_{123}; 1, (\alpha\beta - \alpha - \beta)/\beta), \\ \bar{K} &= K(x_{123}, x_3, x_2, x_1; \alpha, \beta).\end{aligned}$$

- $(0, 0, 1, 1)$:

$$A(x, x_1, x_2, x_{12}) = xx_1 - x_2 - x_{12}, \quad \bar{A}(x_3, x_{13}, x_{23}, x_{123}) = (x_3 + x_{13})x_{23} + x_{123}$$

with

$$\begin{aligned}B &= A(x, x_{23}, x_2, x_3), & \bar{B} &= \bar{A}(x_{12}, x_{13}, x_1, x_{123}), & C &= A(-x, x_1, x_3, x_{13}), \\ \bar{C} &= \bar{A}(x_2, x_{12}, x_{23}, -x_{123}), & K &= A(-x, x_{23}, x_{12}, x_{13}), & \bar{K} &= \bar{A}(x_2, x_3, x_1, -x_{123})\end{aligned}$$

or

$$\begin{aligned}A(x, x_1, x_2, x_{12}) &= xx_1 - x_2 - x_{12}, \\ \bar{A}(x_3, x_{13}, x_{23}, x_{123}) &= (x_3 + x_{13})x_{123} + x_3 - x_{23}\end{aligned}$$

with

$$\begin{aligned} B &= \bar{A}(x_2, x_3, -x_{23}, \alpha^{-1}x), & \bar{B} &= A(\alpha x_1, x_{123} + 1, -x_{12}, -x_{13}), \\ C &= A(x + \alpha, x_1, -x_3, -x_{13}), & \bar{C} &= \bar{A}(x_2, x_{12}, -x_{23}, x_{123}), \\ K &= \bar{A}(x_{12}, x_{13}, x_{23}, \alpha^{-1}x) & \bar{K} &= A(\alpha x_1, x_{123}, x_2, x_3). \end{aligned}$$

- $(x^2, x_1^2, x_{12}^2, x_2^2)$:

$$A(x, x_1, x_2, x_{12}; \delta_1, \delta_2) = xx_{12} + x_1x_2 + \delta_1x_1x_{12} + \delta_2x_2x_{12}$$

with

$$\begin{aligned} \bar{A} &= A(x_{13}, x_3, x_{123}, x_{23}; \delta_1, \delta_2), & B &= A(x, x_2, x_3, x_{23}; \delta_2, 0), \\ \bar{B} &= A(x_{13}, x_1, x_{123}, x_{12}; \delta_1\delta_3, \delta_2), & C &= A(x_{13}, x_3, x, x_1, \delta_1, -\delta_3), \\ \bar{C} &= A(x_{123}, x_{23}, x_2, x_{12}, \delta_1, -\delta_3), & K &= A(x_{13}, x, x_{23}, x_{12}; -\delta_3, -\delta_1\delta_2), \\ \bar{K} &= A(x_3, x_1, x_{123}, x_2; -\delta_3, 0) \end{aligned}$$

or

$$A(x, x_1, x_2, x_{12}; \delta_1, \delta_2) = xx_1 + x_2x_{12} + \delta_1x_1x_{12} + \delta_2x_1x_2$$

with

$$\begin{aligned} \bar{A} &= A(x_{13}, x_3, x_{123}, x_{23}; \delta_1, \delta_2), & B &= A(x, x_3, x_2, x_{23}; \delta_2, 0), \\ \bar{B} &= A(x_{13}, x_1, x_{123}, x_{12}; \delta_1\delta_3, \delta_2), & C &= A(x_{12}, x_1, x_3, x; -\delta_3, -\delta_1\delta_2), \\ \bar{C} &= A(x_{123}, x_{12}, x_{23}, x_2; -\delta_3, 0), & K &= A(x_{13}, x_{12}, x, x_{23}; \delta_1, -\delta_3), \\ \bar{K} &= A(x_{123}, x_1, x_2, x_3; \delta_1, -\delta_3). \end{aligned}$$

Remark. All systems except for the one characterized by the quadruple $(0, 0, 0, 0)$ are new. The latter is equivalent to a special case of the one of the systems from [Atk09] (in that paper the tetrahedron property is not assumed).

4 Concluding Remarks

The Sections 3.2, 3.3 and 3.5 gives us a Bäcklund transformation for every quad-equation of type H1 and of type H2 . Due to [BS02] this leads us to a zero-curvature representation for every quad-equation of type H1 and type H2.

Moreover, our classification includes all known 3D consistent systems except two systems mentioned in [Atk08], the system mentioned in [Hie04] and two systems only containing linear equations (see [Atk09, ABS09]). Of course, all these system do not possess the tetrahedron property.

Acknowledgments

The author is supported by the Berlin Mathematical School and is indebted to Yuri B. Suris for his continued guidance.

References

- [ABS03] Vsevolod E. Adler, Alexander I. Bobenko, and Yuri B. Suris, *Classification of Integrable Equations on Quad-Graphs. The Consistency Approach*, Comm. Math. Phys. **233** (2003), pp. 513–543.
- [ABS09] ———, *Discrete nonlinear hyperbolic equations. Classification of integrable cases*, Funct. Anal. Appl. **43** (2009), pp. 3–17.
- [Atk08] James Atkinson, *Bäcklund transformations for integrable lattice equations*, J. Phys. A: Math. Theor. **41** (2008), no. 135202, 8pp.
- [Atk09] ———, *Linear quadrilateral lattice equations and multidimensional consistency*, J. Phys. A: Math. Theor. **42** (2009), no. 454005, 7pp.
- [BS02] Alexander I. Bobenko and Yuri B. Suris, *Integrable systems on quad-graphs*, Intern. Math. Research Notices **11** (2002), pp. 573–611.
- [BS10] Raphael Boll and Yuri B. Suris, *Non-symmetric discrete Toda systems from quad-graphs*, Applicable Analysis **89** (2010), no. 4, pp. 547–569.
- [Hie04] Jarmo Hietarinta, *A new two-dimensional lattice model that is 'consistent around a cube'*, J. Phys. A: Math. Theor. **37** (2004), pp. 67–73.
- [HV10] Peter E. Hydon and Claude-M. Viallet, *Asymmetric integrable quad-graph equations*, Applicable Analysis **89** (2010), no. 4, pp. 493–506.
- [IR02] Apostolos Iatrou and John A. G. Roberts, *Integrable mappings of the plane preserving biquadratic invariant curves II*, Nonlinearity **14** (2002), pp. 459–489.
- [LY09] Decio Levi and Ravil I. Yamilov, *On a linear integrable difference equation on the square*, Ufa Math. J. **1** (2009), no. 2, pp. 101–105.
- [Nij02] Frank W. Nijhoff, *Lax pair for Adler (lattice Krichever-Novikov) system*, Phys. Lett. A **297** (2002), pp. 49–58.
- [NW01] Frank W. Nijhoff and Alan J. Walker, *The discrete and continuous Painlevé VI hierarchy and the Garnier systems*, Glasg. Math. J. **43A** (2001), pp. 109–123.