Spontaneous Parity–Time Symmetry Breaking and Stability of Solitons in Bose-Einstein Condensates

Zhenya Yan¹, Bo Xiong²,^{*} and W. M. Liu²

¹Key Laboratory of Mathematics Mechanization, Institute of Systems Science,

AMSS, Chinese Academy of Sciences, Beijing 100190, China

²Beijing National Laboratory for Condensed Matter Physics,

Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

(Dated: November 12, 2021)

We report explicitly a novel family of exact \mathcal{PT} -symmetric solitons and further study their spontaneous \mathcal{PT} symmetry breaking, stabilities and collisions in Bose-Einstein condensates trapped in a \mathcal{PT} -symmetric harmonic trap and a Hermite-Gaussian gain/loss potential. We observe the significant effects of mean-field interaction by modifying the threshold point of spontaneous \mathcal{PT} symmetry breaking in Bose-Einstein condensates. Our scenario provides a promising approach to study \mathcal{PT} -related universal behaviors in non-Hermitian quantum system based on the manipulation of gain/loss potential in Bose-Einstein condensates.

PACS numbers: 11.30.Er, 05.45.Yv, 03.75.Lm

Parity (\mathcal{P}) and time-reversal (\mathcal{T}) symmetries are fundamental notions in physics. There has already been much attention to these systems which admire the combined \mathcal{PT} symmetry but do not obey \mathcal{P} and \mathcal{T} symmetries separately. Despite the fact that \mathcal{PT} -Hamiltonian can, in general, be non-Hermitian, the pioneering work of Bender and Boettcher [1] showed that a family of non-Hermitian Hamiltonian with \mathcal{PT} symmetry, \hat{H} = $\hat{p} + \hat{x}^2 (i\hat{x})^\epsilon$, can still show entirely real spectra whereas the generated dynamics is (pseudo) unitary corresponding to the *unbroken* \mathcal{PT} symmetry. Their results have inspired considerable scientific attention devoted to study of non-Hermitian \mathcal{PT} -symmetric Hamiltonian in many aspects, such as \mathcal{PT} -symmetric classical-mechanics theory [2], \mathcal{PT} -symmetric quantum mechanics [3], \mathcal{PT} symmetric quantum field theory [4], pseudo-Hermitian quantum mechanics [5], Lie algebras [6], complex crystal [7] and \mathcal{PT} -symmetric wave chaos [8] etc. Some review papers ([9], for example) present an overview of the theoretical and experimental progress on the study of non-Hermitian \mathcal{PT} -symmetric Hamiltonian.

For the case considered here, given that the action of the parity \mathcal{P} is linear and has the effect $\hat{p} \to -\hat{p}$ and $\hat{x} \to$ $-\hat{x}$ whereas the time operator \mathcal{T} is antilinear and has the effect $\hat{p} \to -\hat{p}$, $\hat{x} \to \hat{x}$, and $i \to -i$, it then follows that a necessary (but not sufficient) condition for a Hamiltonian to be \mathcal{PT} symmetric is $V(\hat{x}) = V^*(-\hat{x})$. In other words, \mathcal{PT} symmetry implies that the real and imaginary parts of the \mathcal{PT} -symmetric potential $V(\hat{x}) \equiv V_{\mathrm{R}}(\hat{x}) + iV_{\mathrm{I}}(\hat{x})$ should be spatially symmetric and antisymmetric, respectively, i.e., $V_{\mathrm{R}}(\hat{x}) = V_{\mathrm{R}}(-\hat{x})$ and $V_{\mathrm{I}}(\hat{x}) = -V_{\mathrm{I}}(-\hat{x})$. It turns out that the \mathcal{PT} -symmetric condition is sufficient to guarantee that the energy spectrum is real and time evolution is unitary, the condition of Dirac Hermiticity is not necessary [9].

There have been several experimental and theoretical studies on one-dimensional Schrödinger equation with different \mathcal{PT} -symmetric potentials such as the periodic potentials [10], non-periodic potentials [11] and \mathcal{PT} symmetric lattices [12]. Some experiments in the framework of wave optics based on $\operatorname{Al}_x \operatorname{Ga}_{1-x}$ [13] and Fe-doped LiNbO₃ [14] provide experimental observation of the behaviors in the both passive and active \mathcal{PT} -symmetric optical coupled two-channel systems. More recently, Musslimani *et al.* [15] have shown that the optical nonlinearity can shift the \mathcal{PT} -symmetric threshold and in turn allow nonlinear eigenmodes with real eigenvalues to exist in the Scarff II potential [16] and periodic potentials [17].

Very interestingly, by using the recently realized experimental techniques [18], \mathcal{PT} -symmetric gain/loss potential is very promising to be experimentally realizable with BECs in the near future. Moreover, a dissipative source (non- \mathcal{PT} -symmetric potentials, i.e., the gain/loss distribution chosen as a Gaussian function) have been used to generate nonlinear coherent excitations in Bose-Einstein condensates (BECs) [18]. As is well known that, for ultra-cold alkali atoms, the confining potential is usually well approximated with the quadratic form [19]. Thus, the natural question appeared is whether the spontaneous \mathcal{PT} symmetry breaking and stability of solitons are universal. The goal of this letter is to give the positive answer to this question by showing that \mathcal{PT} -symmetric soliton-type solutions are pretty natural in BECs trapped by a harmonic potential and synthetic \mathcal{PT} -symmetric gain/loss potential.

In this Letter, we find a new class of exact \mathcal{PT} -symmetric solitons and further study their spontaneous \mathcal{PT} symmetry breaking, stabilities and collisions in BECs with a \mathcal{PT} -symmetric harmonic trap and Hermite-Gaussian gain/loss potential. In particular, the mean-field interaction drastically modifies the threshold point of spontaneous \mathcal{PT} symmetry breaking of solitons in BECs. These results provide a promising approach towards observation of other \mathcal{PT} -related universal features

in new classes of \mathcal{PT} -synthetic system.

We concentrate on a quasi-one-dimensional (1D) BECs in the mean-field regime described by the 1D Gross-Pitaevskii (GP) equation with complex \mathcal{PT} -symmetric potential

$$i\hbar\frac{\partial\Psi}{\partial t} = \left[-\frac{\hbar^2}{2m}\frac{\partial^2}{\partial z^2} + V_{\rm R}(z) + iV_{\rm I}(z) + g_{\rm 1D}|\Psi|^2\right]\Psi,\quad(1)$$

here the complex amplitude distribution $\Psi \equiv \Psi(z,t)$ is macroscopic order parameter, the nonlinearity $g_{1D} = 2a_s \hbar \omega_{\perp}$ is the effective 1D coupling strength, where a_s stands for the *s*-wave scattering length which can be changed through Feshbach resonance, and ω_{\perp} denotes the transverse confining frequency. The external potential is chosen as a harmonic trap $V_{\rm R}(z) = (1/2)m\omega_z^2 z^2$ with *m* being the atomic mass and ω_z is the longitudinal confining frequency [19]. The gain/loss potential $V_{\rm I}(z)$ is phenomenologically incorporated to account for the manipulation of gain/loss potential in BECs by a novel experimental microscopy technique [18].

After the dimensionless transformations of the density, length, time and energy measured in units of $g/(2a_s)$, $a_{\perp} = \sqrt{\hbar/(m\omega_{\perp})}$, ω_{\perp}^{-1} , and $\hbar\omega_{\perp}$, we arrive at an effective 1D GP equation with non-Hermitian \mathcal{PT} -symmetric potential in the dimensionless variables

$$i\frac{\partial\psi}{\partial t} = \left[-\frac{1}{2}\frac{\partial^2}{\partial z^2} + v_{\rm R}(z) + iv_{\rm I}(z) + g|\psi|^2\right]\psi \quad (2)$$

which is associated with $\delta \mathcal{L}/\delta \psi^* = 0$ in which the Lagrangian density can be written as $\mathcal{L} = i(\psi \psi_t^* - \psi^* \psi_t) + |\psi_z|^2 + 2[v_R(z) + iv_I(z)]|\psi|^2 + g|\psi|^4$, where $v_R(z) = (1/2)\omega^2 z^2$ with the trap parameter $\omega = \omega_z/\omega_{\perp}$, $v_I(z) = \hbar\omega_{\perp}V_I(z)$, and g is a interaction parameter. Based on the previous discussion, the real and the imaginary components of the \mathcal{PT} -symmetric potential should satisfy the relations: $v_R(z) = v_R(-z)$ and $v_I(-z) = -v_I(-z)$, respectively. We here concentrate on the attractive interaction g < 0. The repulsive interaction g > 0 does not pose new changes and will be considered elsewhere.

We here focus on spatially localized soliton solutions of Eq. (2) for which $\lim_{|z|\to\infty} \psi(z,t) = 0$. Our goal is to seek a class of soliton-type stationary solutions in the form $\psi(z,t) = \phi(z) \exp(-i\mu t)$ with μ being the chemical potential and the *complex function* $\phi(z) \in \mathbb{C}[z]$ obeying the stationary GP equation with \mathcal{PT} -symmetric potential

$$\mu\phi = \left[-\frac{1}{2}\frac{d^2}{dz^2} + v_{\rm R}(z) + iv_{\rm I}(z) + g|\phi|^2\right]\phi.$$
 (3)

Let us now consider the more interesting and physically relevant \mathcal{PT} -symmetric potential with the external trapping potential $v_{\rm R}(z)$ and the gain/loss potential $v_{\rm I}(z)$ being of the harmonic trap $v_{\rm R}(z) = (1/2)\omega^2 z^2$ and a family of the Hermite-Gaussian distributions

$$v_{\mathrm{I},n}(z) = w_0 \sqrt{\omega} \left[2nH_{n-1}(\sqrt{\omega}z) - \sqrt{\omega}zH_n(\sqrt{\omega}z) \right] e^{-\frac{\omega z^2}{2}}, \quad (4)$$

where w_0 is a constant parameter which can be controlled experimentally, the even integer index n (i.e., n = 0, 2, 4, ...) is necessary to make sure $v_{I,n}(z)$ be the odd function for the \mathcal{PT} -symmetric potential and determines the number of nodes across the Gaussian envelope, $H_n(z) = (-1)^n e^{z^2} d^n e^{-z^2}/dz^n$ stands for the Hermite polynomial. For the odd integer index n, the function $v_{I,n}(z)$ given by Eq. (4) does not obey the \mathcal{PT} -symmetric condition, so we do not considered these cases here.

As a consequence, we find that there exists a family of exact \mathcal{PT} -symmetric soliton-type solutions as

$$\phi_n(z) = \frac{\sqrt{2}w_0}{3\sqrt{|g|}} H_n(\sqrt{\omega}z) e^{-\frac{\omega z^2}{2}} \exp[i\varphi_n(z)], \quad (5)$$

where n = 0, 2, 4, ..., g < 0, the chemical potential satisfies the condition $\mu = \omega[n + 1/2]$, and the phase $\varphi_n(z)$ is given by $\varphi_n(z) = (2/3)w_0 \int_0^z [H_n(\sqrt{\omega}s) \exp(-\omega s^2/2)] ds$. It is easy to show that these solutions is well localized since $\phi_n(z)|_{|z|\to\infty} \to 0$.

In the following, without loss of generality, we will study the cases of n = 0 and n = 2, respectively, for their spontaneous \mathcal{PT} symmetry breaking, stabilities and collisions properties. Thus, we can write the gain/loss terms in the forms of

$$v_{\mathrm{I},0}(z) = -w_0 \omega z \exp\left(-\omega z^2/2\right),\tag{6a}$$

$$v_{\rm I,2}(z) = w_0 \omega (10z - 4\omega z^3) \exp\left(-\omega z^2/2\right),$$
 (6b)

for n = 0 and n = 2, respectively. The corresponding exact \mathcal{PT} -symmetric solitons should be written as

$$\phi_{0}(z) = \frac{\sqrt{2}w_{0}}{3\sqrt{|g|}} e^{-\frac{\omega z^{2}}{2}} \exp\left[\frac{iw_{0}}{3}\sqrt{\frac{2\pi}{\omega}}\operatorname{erf}\left(\sqrt{\omega/2}z\right)\right] (7a)$$

$$\phi_{2}(z) = \frac{2\sqrt{2}w_{0}}{3\sqrt{|g|}} (2\omega z^{2} - 1)e^{-\frac{\omega z^{2}}{2}} \exp[i\varphi_{2}(z)], \quad (7b)$$

for n = 0 and n = 2, respectively, where erf (x) denotes error function and $\varphi_2(z) = (2/3)w_0[\sqrt{2\pi/\omega}\operatorname{erf}(\sqrt{\omega/2}z) - 4z\exp(-\omega z^2/2)].$

As is well known that, for our studied \mathcal{PT} -symmetric potentials, the linear spectrum can be entirely real as long as the w_0 is operated below the phase transition point (unbroken \mathcal{PT} symmetry) for a fixed value of ω . Above this so-called \mathcal{PT} -symmetric threshold, a phase transition occurs and the linear spectrum enters the complex domain.

Notice that in our studied system, the corresponding linear spectrum problem associated with the \mathcal{PT} symmetric potential (6a), exhibits an entirely real spectrum provided that,

$$|w_0| \le 1.945,$$
 (8)

for the case of g = -1, $\omega = 1$ and n = 0. A single-peak soliton-type stationary state (7a) in the \mathcal{PT} -symmetric



FIG. 1: (color online). Intensity evolution of a single-peak soliton state in a \mathcal{PT} -symmetric potential (6a) for g = -1 and $\omega = 1$. (a) $w_0 = 0.8$. (b) $w_0 = 1.0$. The inset depicts the real (top) and imaginary (bottom) component of \mathcal{PT} -symmetric potential.

potential (6a) for the case of g = -1, $\omega = 1$ and $w_0 = 0.8$ is shown in Fig. 1(a). The stability of this \mathcal{PT} -symmetric soliton has been confirmed by numerical simulation of Eq. (2) using beam propagation methods with adding random noise on both amplitude and phase of exact \mathcal{PT} -symmetric soliton state. Thus for this given w_0 , one can see that \mathcal{PT} -symmetric soliton states can be found with real energy in this system. Very interestingly, one finds that even if the potential (6a) has below the phase transition point ($w_0 = 1.0$, its single particle spectrum is real), single-peak soliton has already been unstable with breaking \mathcal{PT} symmetry and imaginary energy. Fig. 1(b) shows the numerical simulation of Eq. (2) for the case of g = -1, $\omega = 1$ and $w_0 = 1.0$ by using the same method as in Fig. 1(a). This means that, the single-peak soliton itself can modify the real part of potential through the mean field interaction term in Eq. (2). Thus, this new effective trapping potential shifts the \mathcal{PT} threshold w_0 of Eq. (8) towards a smaller value and in turn damages this stationary soliton state against breaking \mathcal{PT} symmetry. For the obtained singlepeak \mathcal{PT} -symmetric soliton (7a) we find that the quantity $S = (i/2)(\phi \phi_x^* - \phi_x \phi^*) = 4\omega_0^3/(9|g|) \exp(-3\omega z^2/2)$ associated with the transverse power flow density is positive with $\omega_0 > 0$, which means that the power always flows in one direction, i.e., from the gain to loss region.

We next investigate multi-peak soliton-type stationary state (7b) and their dynamics supported with the \mathcal{PT} symmetric potential (6b). The linear spectrum properties of such a \mathcal{PT} -symmetric potential can be understood by examining the corresponding linear spectrum problem of Eq. (2), And one finds that purely real spectrums are



FIG. 2: (color online). Intensity evolution of a multi-peak soliton state in a \mathcal{PT} -symmetric potential (6b) for g = -1 and $\omega = 1$. (a) $w_0 = 0.1$. (b) $w_0 = 0.2$. The inset depicts the imaginary component of \mathcal{PT} - symmetric potential.

possible in the range

$$|w_0| \le 0.00002,\tag{9}$$

for the case of $g = -1, \omega = 1$ and n = 2. Since the similar idea holds for such \mathcal{PT} -symmetric potential (6b), by numerical simulation of Eq. (2) using beam propagation methods, however, one finds a stable \mathcal{PT} symmetric multi-peak soliton-type state even for the parameters of $q = -1, \omega = 1$ and $w_0 = 0.1$ as is shown in Fig. 2(a). Thus, very interestingly, in contrast to the stable \mathcal{PT} -symmetric single-peak soliton, here, the multipeak soliton itself eventually, shifts the \mathcal{PT} threshold w_0 of Eq. (9) towards a rather larger value and makes soliton-type state preserve \mathcal{PT} symmetry very well by drastically modifying the real part of potential through the mean field interaction term. With increasing the w_0 further, the multi-peak soliton will be unstable with breaking \mathcal{PT} symmetry and imaginary energy. Fig. 2(b) shows the numerical simulation of Eq. (2) for the case of $g = -1, \omega = 1$ and $w_0 = 0.2$ by using the same method as former cases.

Finally, we need to check if this \mathcal{PT} -symmetric soliton indeed behave as solitons. To this end, we take initial conditions in the form of a superposition of two singlepeak (7a) or multi-peak (7b) solitons with symmetric central positions. In such a case, and given that the single-peak or multi-peak solitons were found above to be robust objects behaving similarly to solitons of an integrable system, one may expect that the solitons would perform harmonic oscillations in the presence of the sufficiently large parabolic trap even with \mathcal{PT} -symmetric potentials. As shown in the Fig. 3(a) for the case of g = -1, $\omega = 1$ and $w_0 = 0.3$, two single-peak solitons propagate in opposite directions and undergo a pretty



FIG. 3: (color online). Dynamic evolution of collisions of two single-peak and multi-peak solitons in a \mathcal{PT} -symmetric potentials (6a) and (6b) for g = -1 and $\omega = 1$. Here (a) $w_0 = 0.3$ and (b) $w_0 = 0.8$ for two single-peak solitons. (c) $w_0 = 0.2$ and (d) $w_0 = 0.25$ for two multi-peak solitons.

elastic collision and the two solitons remain unscathed after the collision. The stability of this superposition of two single-peak solitons has been confirmed by numerical simulation of Eq. (2) using beam propagation methods as before. But when the gain/loss parameter w_0 becomes larger (for example, $w_0 = 0.8$), two single-peak solitons undergo a inelastic collision as shown in the Fig. 3(b). A superposition of two multi-peak solitons (7b) with symmetric central positions is also found to be robust objects behaving similarly to above and undergos a elastic collision as shown in the Fig. 3(c) for the case of g = -1, $\omega = 1$ and $w_0 = 0.2$. But when the gain/loss parameter w_0 becomes a little larger (for example, $w_0 = 0.25$), two multi-peak solitons become unstable clearly as shown in the Fig. 3(d). All these findings certainly indicate that many kinds of robust stable nonlinear modes can be supported by \mathcal{PT} -symmetric potential in BECs.

Experimental realization of the ideas presented here is promising. A condensate in a dilute vapor of sodium or rubidium atoms is placed in a cigar-shaped potential may provide an accessible laboratory to experimentally observe spontaneous \mathcal{PT} symmetry breaking and stability of solitons. Such an experiment would require precise control over the rates of gain/loss process in the condensate where atoms are injected continuously into the condensate in left side, whereas they are removed simultaneously from the condensate in right side. The combination of high spatial resolution with *in situ* detection [18] will hopefully provide new possibilities for the preparation, the manipulation and the characterization of non-Hermitian \mathcal{PT} symmetry condensate.

In summary, we have demonstrated that a novel class of exact \mathcal{PT} -symmetric solitons can be well supported

in BECs with \mathcal{PT} -symmetric gain/loss potential. Their spontaneous \mathcal{PT} symmetry breaking, stability, as well as collision dynamics of such \mathcal{PT} -symmetric solitons were examined in detail. We observe the significant effects of mean-field potential by modifying the threshold point of spontaneous \mathcal{PT} symmetry breaking. Our results may intrigue a new class of \mathcal{PT} -synthetic universal properties study that relies on manipulations of macroscopic matter-wave field with BECs.

This work was supported by the NSFC under grants Nos. 60821001/F02, 10874235, 10934010, 60978019, the NKBRSFC under grants Nos. 2009CB930701, 2010CB922904 and 2011CB921500.

- * Electronic address: stevenxiongbo@gmail.com
- C. M. Bender and S. Boettcher, Phys. Rev. Lett. 80, 5243 (1998).
- C. M. Bender, S. Boettcher, and P. N. Meisinger, J. Math. Phys. (N.Y.) 40, 2201 (1999); C. M. Bender, D. D. Holm, and D. W. Hook, J. Phys. A 40, F81 (2007).
- [3] C. M. Bender *et al.*, Phys. Rev. Lett. **98**, 040403 (2007);
 C. M. Bender *et al.*, *ibid.* **104**, 061601 (2010).
- [4] C. M. Bender, D. C. Brody, and H. F. Jones, Phys. Rev. Lett. 89, 270401 (2002); *ibid.* 92, 119902(E) (2004); C.
 M. Bender, and S. P. Klevansky, Phys. Rev. Lett. 105, 031601 (2010).
- [5] A. Mostafazadeh and A. Batal, J. Phys. A 37, 11645 (2004); A. Mostafazadeh, Phys. Rev. Lett. 99, 130502 (2007).
- [6] B. Bagchi and C. Quesne, Phys. Lett. A 273, 285 (2000).
- M. Znojil, Phys. Lett. A 285, 7 (2001); Z. Ahmed, Phys. Rev. A 64, 042716 (2001); A. Khare and U. Sukhatme, J. Math. Phys. (N.Y.) 46, 082106 (2005); S. Longhi, Phys. Rev. Lett. 103, 123601 (2009); S. Longhi, Phys. Rev. A 81, 022102 (2010).
- [8] C. T. Carl et al., Phys. Rev. Lett. 104, 054102 (2010).
- [9] C. M. Bender, Rep. Prog. Phys. 70, 947 (2007); A. Mostafazadeh, arXiv:0810.5643.
- [10] C. M. Bender, G. V. Dunne, and P. N. Meisinger, Phys. Lett. A **252**, 272 (1999); K. G. Makris *et al.*, Phys. Rev. Lett. **100**, 103904 (2008).
- [11] C. M. Bender and S. Boettcher, J. Phys. A **31**, L273 (1998); M. Znojil, *ibid.* **32**, 4563 (1999); S. Klaiman, U. Günther, and N. Moiseyev, Phys. Rev. Lett. **100**, 080402 (2008).
- [12] O. Bendix *et al.*, Phys. Rev. Lett. **103**, 030402 (2009);
 K. G. Makris *et al.*, Phys. Rev. A **81**, 063807 (2010).
- [13] A. Guo et al., Phys. Rev. Lett. 103, 093902 (2009).
- [14] C. E. Rüter *et al.*, Nature Phys. 6, 192 (2010); T. Kottos, Nature Phys. 6, 166 (2010).
- [15] Z. H. Musslimani *et al.*, Phys. Rev. Lett. **100**, 030402 (2008); J. Phys. A **41**, 244019 (2008).
- [16] Z. Ahmed, Phys. Lett. A 282, 343 (2001).
- [17] H. F. Jones, Phys. Lett. A 262, 242 (1999); J. K. Boyd,
 J. Math. Phys. 42, 15 (2001); E. Narevicius, P. Serra,
 and N. Moiseyev, Europhys. Lett. 62, 789 (2003).
- [18] T. Gericke *et al.*, Nature Phys. **4**, 949 (2008); V. A. Brazhnyi *et al.*, Phys. Rev. Lett. **102**, 144101 (2009).
- [19] F. Dalfovo et al., Rev. Mod. Phys. 71, 463 (1999); O.

Morsch and M. Oberthaler, Rev. Mod. Phys. **78**, 179 (2006); R. Carretero-González, D. J. Frantzeskakis, and

P. G. Kevrekidis, Nonlinearity **21**, R139 (2008).