Chiral extrapolation of SU(3) amplitudes

Gerhard Ecker

University of Vienna, Faculty of Physics, Boltzmanngasse 5, A-1090 Wien, Austria

Abstract. Approximations of chiral SU(3) amplitudes at NNLO are proposed to facilitate the extrapolation of lattice data to the physical meson masses. Inclusion of NNLO terms is essential for investigating convergence properties of chiral SU(3) and for determining low-energy constants in a controllable fashion. The approximations are tested with recent lattice data for the ratio of decay constants F_K/F_{π} .

Keywords: NNLO CHPT, lattice QCD PACS: 12.38.Gc,12.39.Fe,14.40.Be

MOTIVATION

Lattice QCD has made enormous progress in the light quark sector. Since most lattice studies employ quark masses with $m_q^{\text{lattice}} > m_q^{\text{phys}}$ an extrapolation in meson masses is needed. Although chiral perturbation theory (CHPT) amplitudes [1] are available to NNLO for most quantities of interest [2] the NNLO contributions are usually quite involved. In addition, they are mostly available only in numerical form for chiral SU(3).

In the lattice community, the convergence properties of chiral SU(3) are generally considered problematic. For a thorough discussion of this issue, NLO amplitudes are not sufficient. Moreover, using quark masses with $m_s^{\text{lattice}} < m_s^{\text{phys}}$ would be very useful for assessing the convergence properties of chiral SU(3) [3].

Tuning the quark masses offers a new environment for extracting chiral low-energy constants (LECs). This is especially welcome for LECs that are difficult to access phenomenologically, both at NLO and at NNLO.

For all these reasons, analytic approximations of NNLO CHPT amplitudes are expected to be very useful. In this talk, I describe recent attempts to derive such approximations [4] that are

- more sophisticated than the double-log approximation [5],
- yet much simpler than the full numerical expressions.

CHPT AT NNLO

A compact representation of CHPT is in terms of the generating functional of Green functions [1]. The NNLO functional Z_6 of $O(p^6)$ is itself a sum of different contributions shown pictorially in Fig. 1. I concentrate here on

the so-called irreducible contributions [6, 4]

$$Z_{6}^{a+b+d+g} = \int d^{4}x \{ [C_{a}^{r}(\mu) + \frac{1}{4F_{0}^{2}} \left(4\Gamma_{a}^{(1)}L - \Gamma_{a}^{(2)}L^{2} + 2\Gamma_{a}^{(L)}(\mu)L \right) \right] O_{a}(x) + \frac{1}{(4\pi)^{2}} \left[L_{i}^{r}(\mu) - \frac{\Gamma_{i}}{2}L \right] H_{i}(x;M) + \frac{1}{(4\pi)^{4}}K(x;M) \} .$$
(1)

The C_a^r are LECs of $O(p^6)$, $O_a(x)$ the associated monomials of the chiral Lagrangian [7]. The chiral log *L* is given by $L = 1/(4\pi)^2 \ln M^2/\mu^2$, L_i^r are LECs of $O(p^4)$ and F_0 is the pseudoscalar decay constant in the chiral SU(3) limit. The coefficients $\Gamma_a^{(1)}$, $\Gamma_a^{(2)}$ and $\Gamma_a^{(L)}$ are listed in Ref. [6]. $H_i(x;M)$ (1-loop) and K(x;M) (2-loop) are nonlocal functionals. $Z_6^{a+b+d+g}$ and Z_6^{total} are independent of both scales μ, M .

APPROXIMATIONS

The drawback of the explicit form of the generating functional of $O(p^6)$ for chiral SU(3) is that the 2-loop contribution K(x;M) is in general only numerically available. The approximation suggested in Ref. [4] (referred to as Approximation I from now on) has the following properties:

- *K*(*x*;*M*) (and the reducible 2-loop contributions) are dropped;
- All chiral logs are kept;
- *Z*₆^(appr.I) is still invariant with respect to the renormalization scale μ;
- The residual dependence on the scale *M* (appearing only in the chiral log *L*) is the only vestige of the 2-loop part;



FIGURE 1. Skeleton diagrams for the generating functional Z_6 of $O(p^6)$. Simple dots, crossed circles, black box denote vertices from LO, NLO, NNLO Lagrangians, respectively. Propagators and vertices carry the full tree structure associated with the lowest-order Lagrangian.

- Approximation I respects the large- N_c hierarchy of $O(p^6)$ contributions:
 - $C_a, L_i L_j \longrightarrow L_i \times \text{loop} \longrightarrow 2\text{-loop}$
- Only tree and 1-loop amplitudes are needed;
- Unlike the double-log approximation [5], it allows for the extraction of LECs with the correct scale dependence.

One important question remains to be answered: how trustworthy is this approximation? As the following examples will show, the answer depends on the quantity under consideration.

APPLICATIONS

F_K/F_{π}

The ratio of pseudoscalar decay constants F_K/F_{π} exhibits a picture-book type of chiral expansion as shown in Table 1. The separately scale-dependent contributions of $O(p^6)$ are given for $\mu = 770$ MeV. The entries for "numerical results" were provided by Bernard and Passemar (Ref. [8] and private communication).

For an SU(3) quantity normalized to one at LO, successive terms in the chiral expansion are expected to show the following generic behaviour:

$$\begin{array}{ll} O(p^4) & O(p^6) & O(p^8) \\ \lesssim 0.3 & \lesssim 0.3^2 = 0.09 & \lesssim 0.3^3 = 0.027 \end{array}$$

This suggests as criterion for an acceptable NNLO approximation that the accuracy should not be worse than 3 %, the typical size of a term of $O(p^8)$. Table 1 shows that

this criterion is only barely met by Approximation I. A possible modification consists in ignoring the strict large- N_c counting to include also products of 1-loop functions originating from diagrams a,c in Fig. 1. As Table 1 indicates, this Approximation II [9] satisfies the criterion for an acceptable approximation.

In addition to two combinations of LECs of $O(p^6)$, also various L_i appear in F_K/F_{π} . At $O(p^4)$, only L_5 enters. We have therefore extracted L_5 , $C_{14} + C_{15}$ and $C_{15} + 2C_{17}$ from a fit of the recent lattice data of the BMW Collaboration [10], using for the remaining L_i (appearing only at $O(p^6)$) the values of fit 10 of Ref. [11]. The results [4, 9] are displayed in Table 2. The fitted values of F_K/F_{π} agree with the detailed analysis of Ref. [10]. For both F_K/F_{π} and L_5 , there is practically no difference between the two approximations but the LECs of $O(p^6)$ show a bigger spread. The results are shown for $M = M_K$. For a discussion of the *M*-dependence, I refer to Refs. [4, 9]. The fit also demonstrates that NNLO terms are absolutely essential. While the NNLO fit (Approximation II) is well behaved (χ^2 /dof = 1.2, statistical errors only), the NLO fit with the single parameter L_5 is a catastrophe ($\chi^2/dof = 4$).

Other applications

 F_{π}/F_0 in chiral SU(3). From the chiral point of view, the interest is not so much in "deriving" F_{π} from lattice data but rather to extract the still badly known F_0 and the NLO LEC L_4 . F_0 sets the scale of the chiral SU(3) expansion and is therefore essential for an appraisal of convergence properties. The comparison of Approximation

$O(p^4)$ $O(p^6)$ 2-loop $L_i \times loop$ tree numerical results [11, 8] 0.14 0.002 0.051 0.008 Appr. I ($M = M_K$) - 0.030 - 0.011 - 0.011 TABLE 2. Fit results for F_K/F_{π} and LECs (statistical errors only) F_K/F_{π} $10^3L_5^r$ $10^3(C_{14}^r + C_{15}^r)$ GeV ² $10^3(C_1^r)$ Appr. I ($M = M_K$) 1.198(5) 0.76(9) 0.37(8) Appr. II ($M = M_K$) 1.200(5) 0.75(9) 0.20(7) BMW [10] 1.192(7)_{stat}(6)_{syst} 0.20(7)	-						
2-loop $L_i \times loop$ tree numerical results [11, 8] 0.14 0.002 0.051 0.008 Appr. I ($M = M_K$) - 0.030 Appr. II ($M = M_K$) - 0.011 TABLE 2. Fit results for F_K/F_{π} and LECs (statistical errors only) F _K /F _π 0.03($C_{14}^r + C_{15}^r$) GeV ² Appr. I ($M = M_K$) Appr. I ($M = M_K$) Appr. II ($M = M_K$)			$O(p^4)$		$O(p^6)$		-
numerical results [11, 8] 0.14 0.002 0.051 0.008 Appr. I $(M = M_K)$ - 0.030 Appr. II $(M = M_K)$ - 0.011 CABLE 2. Fit results for F_K/F_{π} and LECs (statistical errors only) F_K/F_{π} $10^3 L_5^r$ $10^3 (C_{14}^r + C_{15}^r)$ GeV ² $10^3 (C_1^r$ Appr. I $(M = M_K)$ 1.198(5) 0.76(9) 0.37(8) Appr. II $(M = M_K)$ 1.200(5) 0.75(9) 0.20(7) BMW [10] 1.192(7)_{stat}(6)_{syst} 0.75(9) 0.20(7)				2-loop	$L_i \times \text{loop}$	tree	
Appr. I $(M = M_K)$ - 0.030 Appr. II $(M = M_K)$ - 0.011 CABLE 2. Fit results for F_K/F_{π} and LECs (statistical errors only) F_K/F_{π} $10^3 L_5^r$ $10^3 (C_{14}^r + C_{15}^r)$ GeV ² $10^3 (C_1^r$ Appr. I $(M = M_K)$ 1.198(5) $0.76(9)$ $0.37(8)$ Appr. II $(M = M_K)$ 1.200(5) $0.75(9)$ $0.20(7)$ BMW [10] 1.192(7) _{stat} (6) _{syst} $0.75(9)$ $0.20(7)$		numerical results [11, 8]	0.14	0.002	0.051	0.008	
Appr. II $(M = M_K)$ - 0.011 ABLE 2. Fit results for F_K/F_{π} and LECs (statistical errors only) F_K/F_{π} $10^3 L_5^r$ $10^3 (C_{14}^r + C_{15}^r)$ GeV ² $10^3 (C_1^r$ Appr. I $(M = M_K)$ 1.198(5) 0.76(9) 0.37(8) Appr. II $(M = M_K)$ 1.200(5) 0.75(9) 0.20(7) BMW [10] 1.192(7)_{stat}(6)_{syst}		Appr. I $(M = M_K)$		- 0.030			
EABLE 2. Fit results for F_K/F_{π} and LECs (statistical errors only) F_K/F_{π} $10^3 L_5^r$ $10^3 (C_{14}^r + C_{15}^r)$ GeV ² $10^3 (C_1^r$ Appr. I ($M = M_K$) 1.198(5) 0.76(9) 0.37(8) Appr. II ($M = M_K$) 1.200(5) 0.75(9) 0.20(7) BMW [10] 1.192(7) _{stat} (6) _{syst}		Appr. II $(M = M_K)$		- 0.011			
Appr. I $(M = M_K)$ 1.198(5) 0.76(9) 0.37(8) Appr. II $(M = M_K)$ 1.200(5) 0.75(9) 0.20(7) BMW [10] 1.192(7) _{stat} (6) _{syst}	ABLE 2. Fit rest						
Appr. II $(M = M_K)$ 1.200(5)0.75(9)0.20(7)BMW [10]1.192(7) _{stat} (6) _{syst}		ults for F_K/F_{π} and LECs (st F_K/F_{π}	tatistical err $10^3 L_5^r$	fors only) $10^3 (C_{14}^r)$	$+C_{15}^r$) GeV ²	10 ³	$C_{15}^r + 2$
BMW [10] $1.192(7)_{\text{stat}}(6)_{\text{syst}}$	Appr. I $(M = M_K)$	ults for F_K/F_{π} and LECs (st F_K/F_{π} 1.198(5)	tatistical err $10^{3}L_{5}^{r}$ $0.76(9)$	$\frac{10^3 (C_{14}^r)}{0}$	$+C_{15}^r) \text{ GeV}^2$	- 10 ³ ($C_{15}^r + 2$ 1.29
	Appr. I $(M = M_K)$ Appr. II $(M = M_K)$	ults for F_K/F_{π} and LECs (so F_K/F_{π}) 1.198(5)) 1.200(5)	$ \frac{10^3 L_5^r}{0.76(9)} \\ 0.75(9) $	$\frac{10^3 (C_{14}^r)}{0.000}$	$+C_{15}^r) \text{ GeV}^2$.37(8) .20(7)	2 10 ³ ($C_{15}^r + 2$ 1.29 0.71

TABLE 1. Chiral expansion of $F_K/F_{\pi} - 1$

I with available numerical results (Ref. [8] and private communication) is very encouraging in this case [9].

 K_{l3} vector form factor $f_+(t)$. A crucial quantity for a precision determination of the CKM element V_{us} is the value of $f_+(0)$. For at least two reasons [9], both approximations discussed here do not appear very promising in this case.

- The chiral expansion of $f_+(0)$ shows a rather atypical behaviour.
- The approximations considered so far do not match with available numerical results [12, 13, 8].

CONCLUSIONS

- 1. We have provided user-friendly extrapolation formulas [4, 9] for $N_f = 2 + 1$ lattice data.
- 2. The proposed approximations are superior to NLO and chiral double-log approximations and should be useful for investigating the convergence properties of chiral SU(3).
- 3. They allow for the extraction of NLO and NNLO LECs with the correct scale dependence.
- 4. Analysis of the BMW data [10] for F_K/F_{π} provides the best available determination of L_5 , assuming the large- N_c hierarchy $|L_4|, |L_6| \ll L_5$.
- 5. The proposed approximations are intended to be used by lattice groups: they offer considerably more insight than, e.g., polynomial fits.

ACKNOWLEDGMENTS

The results presented here have grown out of a very pleasant collaboration with Pere Masjuan and Helmut Neufeld. I also thank Véronique Bernard, Steve Gottlieb and Emilie Passemar for helpful discussions.

REFERENCES

- J. Gasser and H. Leutwyler, Annals Phys. 158 (1984) 142; Nucl. Phys. B 250 (1985) 465.
- J. Bijnens, Prog. Part. Nucl. Phys. 58 (2007) 521 [arXiv:hep-ph/0604043].
- A. Bazavov, C. Bernard, C. E. DeTar *et al.*, Rev. Mod. Phys. 82 (2010) 1349. [arXiv:0903.3598 [hep-lat]].
- 4. G. Ecker, P. Masjuan and H. Neufeld, Phys. Lett. B **692** (2010) 184. [arXiv:1004.3422 [hep-ph]].
- 5. J. Bijnens, G. Colangelo and G. Ecker, Phys. Lett. B 441 (1998) 437 [arXiv:hep-ph/9808421].
- 6. J. Bijnens, G. Colangelo and G. Ecker, Annals Phys. **280** (2000) 100 [arXiv:hep-ph/9907333].
- J. Bijnens, G. Colangelo and G. Ecker, JHEP 9902 (1999) 020 [arXiv:hep-ph/9902437].
- V. Bernard and E. Passemar, JHEP **1004** (2010) 001 [arXiv:0912.3792 [hep-ph]].
- 9. G. Ecker, P. Masjuan and H. Neufeld, in preparation.
- S. Dürr *et al.*, Phys. Rev. D **81** (2010) 054507 [arXiv:1001.4692 [hep-lat]].
- 11. G. Amorós, J. Bijnens and P. Talavera, Nucl. Phys. B **568** (2000) 319 [arXiv:hep-ph/9907264].
- 12. P. Post and K. Schilcher, Eur. Phys. J. C **25** (2002) 427. [hep-ph/0112352].
- J. Bijnens and P. Talavera, Nucl. Phys. B 669 (2003) 341. [hep-ph/0303103].