

Quantum theory of fermion preheating

Jürgen Berges, Daniil Gelfand, and Jens Pruschke

Institute for Nuclear Physics, Darmstadt University of Technology, Schlossgartenstr. 9, 64285 Darmstadt, Germany

We show that quantum effects dramatically enhance the production of fermions during preheating after inflation in the early universe. As a consequence fermions rapidly approach a quasi-stationary distribution with a thermal occupancy in the infrared, while the inflaton enters a turbulent scaling regime. The failure of standard semi-classical descriptions based on the Dirac equation with a homogeneous background field is caused by non-perturbatively high boson occupation numbers. We comment on related phenomena in heavy-ion collisions for the production of quark matter fields from highly occupied gauge bosons.

PACS numbers: 11.10.Wx, 98.80.Cq, 12.38.Mh

The precise understanding of phenomena out of equilibrium plays a crucial role for our knowledge about the primordial universe. Important examples are the generation of density fluctuations, nucleosynthesis or baryogenesis – the latter being responsible for our own existence.

Most dramatic processes such as far-from-equilibrium particle production at the end of inflation pose a particular theoretical challenge [1]. The corresponding non-perturbative phenomenon of preheating is conventionally described using classical approximations for the bosonic inflaton [2]. Their validity for macroscopic occupation numbers has been verified explicitly in quantum field theory [3]. Much less is known about preheating dynamics with fermions. Since identical fermions cannot occupy the same state, their quantum nature is highly relevant and a consistent quantum theory of fermion production after inflation is of crucial importance. Related questions arise in the context of heavy-ion collision experiments, where the production of quark matter fields from highly occupied gauge bosons can occur far from equilibrium [4].

So far, preheating dynamics with fermions has been mainly investigated based on semi-classical descriptions using the Dirac equation with coupling to a homogeneous inflaton field [5]. The findings exhibit fermion number densities which vary periodically in time, governed by resonant excitations. Also backreaction of fermions onto inflaton dynamics has been included. A similar semi-classical prescription has been used to compute quark production from classical gluons in the context of heavy-ion collisions [6]. Further inclusion of quantum corrections is complicated by a secular perturbative time evolution which becomes rapidly invalid. Suitable approximations may be based on the two-particle irreducible (2PI) generating functional, which are also known to be able to describe the approach to thermal equilibrium [7]. Simplified discussions with these techniques, which evaluate the dynamics for vanishing inflaton field amplitude, exhibit new nonlinear phenomena such as instability-induced fermion production [8].

In this work we present a quantum theory of fermion preheating. For the first time we consistently include quantum corrections to next-to-leading order (NLO) in

the Yukawa coupling between the inflaton field and massless fermions. Even for weak couplings this turns out to change semi-classical or leading-order (LO) results so dramatically that we consider a complementary non-perturbative method for comparison. It is based on lattice simulations following the techniques of Ref. [9]. This treats the fermions exactly, but the inflaton dynamics becomes classical-statistical. In $3 + 1$ dimensions this is computationally expensive and becomes feasible with the implementation of 'low-cost' fermion algorithms [10]. Remarkably, higher-order corrections turn out to leave the NLO results practically unchanged for the considered range of weak couplings. We show that the NLO corrections introduce a dynamical mechanism for the enhancement of quantum effects in the presence of large inflaton occupation numbers. In the theory of preheating the inflaton occupation number depends on time, thus leading to a dynamic 'switching on' of the quantum enhancement. As a consequence fermions rapidly approach a quasi-stationary distribution with a thermal occupancy in the infrared, while the inflaton enters the scaling regime characteristic for strong turbulence.

In inflationary cosmology, the universe at early times expands quasi-exponentially in a vacuum-like state. During this stage of inflation, all energy is contained in a slowly evolving inflaton field. Eventually the inflaton has to transfer its energy to particles, thereby starting the thermal history of the hot Friedmann universe. If the creation of particles is sufficiently slow (for instance, if the inflaton is coupled only gravitationally to the matter fields) this transfer may be described by perturbative decay. However, for a wide range of couplings the particle production is governed by a nonequilibrium instability [1–3]. These instabilities are known to lead to exponential growth of inflaton occupation numbers in long wavelength modes on time scales much shorter than the asymptotic thermal equilibration time. This is followed by a turbulent phase with different universal scaling regimes for non-perturbative long wavelength modes [11] and perturbative higher momenta [12]. Despite the differences between early-universe dynamics and the 'Little Bang' generated in a heavy-ion collision, there

are remarkable parallels concerning the role of instabilities and turbulence for the process of thermalization [14].

To reveal the production of matter fields from inflaton decay requires the inclusion of fermions. We consider a quantum field theory with interaction of the Yukawa type

$$-\frac{g}{N_f} \bar{\psi}_i \left((1 - \gamma^5) \Phi_{ij}^\dagger + (1 + \gamma^5) \Phi_{ij} \right) \psi_j, \quad (1)$$

for $i, j = 1, \dots, N_f$ massless Dirac fermions, where summation over repeated indices is implied. (We use natural units with $\hbar = c = k_B = 1$.) The interaction is constructed to couple the left- and right-handed fermion field components $\psi_L = (1 - \gamma^5)\psi/2$, $\psi_R = (1 + \gamma^5)\psi/2$ for a non-vanishing expectation value

$$\phi(t) = \langle \text{Tr } \Phi(t, \mathbf{x}) \rangle \quad (2)$$

of the inflaton field. This describes the generation of an effective fermion mass by spontaneous symmetry breaking. Together with the scattering or decay processes implied by (1), the interaction contains the major building blocks required to discuss fermion production.

To be more specific we consider two flavors, $N_f = 2$, with symmetry group $SU_L(2) \times SU_R(2) \sim O(N_s = 4)$, where $\Phi = (\sigma + i\vec{\tau}\vec{\pi})/2$ and an inflaton potential $V = m^2(\sigma^2 + \vec{\pi}^2)/2 + \lambda(\sigma^2 + \vec{\pi}^2)^2/4!N_s$ characteristic for a linear sigma model. (Here $\vec{\tau}$ denote Pauli matrices.) In chaotic inflationary models, after the end of inflation the inflaton field coherently oscillates around the minimum of its potential with a very large amplitude on the order of the Planck mass. The coupling of the field to its own quantum fluctuations leads to the phenomenon of parametric resonance [1]. For our purposes it will be sufficient to consider fermion preheating without expansion. Though our results will be for parametric resonance in chaotic inflation scenarios, we concentrate on characteristic properties which are independent of the underlying instability type and details of the model.

In the corresponding quantum field theory fermion production can be computed from correlation functions

$$F(x, y) = \frac{1}{2} \langle [\psi(x), \bar{\psi}(y)] \rangle, \quad (3)$$

$$\rho(x, y) = i \langle \{\psi(x), \bar{\psi}(y)\} \rangle. \quad (4)$$

Here $F(x, y)$ denotes the commutator or statistical two-point function, and $\rho(x, y)$ is the anti-commutator of two fermion fields or the spectral function. In thermal equilibrium these would be translation invariant and related by a fluctuation-dissipation relation in Fourier space: $F_{\text{eq}}(\omega, \mathbf{p}) = -i[1/2 - n_{\text{FD}}(\omega)]\rho_{\text{eq}}(\omega, \mathbf{p})$ with the Fermi-Dirac distribution $n_{\text{FD}}(\omega) = 1/[\exp(\omega/T) + 1]$ for zero net fermion charge [7]. The question of fermion production out of equilibrium may be discussed in terms of a time-dependent occupation number $n_\psi(t, \mathbf{p})$, which starts out at $n_\psi(t = 0, \mathbf{p}) \simeq 0$ after inflation and approaches a constant distribution in equilibrium. Calling




FIG. 1: Time-dependent fermion occupation number.

$M_\psi(t) = g\phi(t)/N_f$ we consider a standard definition [5]

$$n_\psi(t, \mathbf{p}) = \frac{1}{2} - \frac{|\mathbf{p}| F_V(t, t; \mathbf{p}) + M_\psi(t) F_S(t, t; \mathbf{p})}{\sqrt{\mathbf{p}^2 + M_\psi^2(t)}}, \quad (5)$$

where $F_S = \text{tr}(F/4)$, $F_V = \text{tr}(\vec{p} \cdot \vec{\gamma} F/4)$ with the trace acting in Dirac space. These are the scalar and vector components of the flavor-diagonal F , which represent the only nonvanishing components at initial time consistent with symmetries.

Fig. 1 compares results for the fermion occupation number obtained from three different approximations, for $\xi \equiv g^2/\lambda = 0.1$ in units of the rescaled initial inflaton field amplitude $\phi_0 = \phi(t = 0)/\sqrt{6N_s/\lambda}$. For the semi-classical approximation (LO), the correlator (3) obeys the evolution equation

$$\left[i\gamma^\mu \partial_\mu - \frac{g}{N_f} \phi(t) \right] F(x, y) = 0. \quad (6)$$

Since in this approximation the result shows strongly oscillatory behavior, a curve at fixed time $t = 250/\phi_0$ (dotted) is given together with a time-average over a period of about $20/\phi_0$ (dashed). For the NLO result (solid curve) the right-hand side of (6) is replaced by a quantum correction, which reads diagrammatically:

(7)

Here the dashed lines denote self-consistently dressed fermion propagators and the solid line represents the inflaton propagator. The computation of the NLO correction is computationally expensive due to the appearance of memory integrals in the coupled evolution equations for the fermion and the inflaton degrees of freedom.

We observe that the NLO quantum correction, which by naive power counting is $O(g^2)$, changes the LO result both qualitatively and quantitatively. This happens




FIG. 2: Total number density of fermions.

despite the fact that the employed coupling, $g = 0.1$, is small and the semi-classical equation (6) takes already into account $O(g)$ contributions. For both LO and NLO results we use the same approximation for the bosonic sector of our model with $\lambda = 0.1$, which employs the well-tested 2PI $1/N_s$ expansion [13]. Before we turn to the lattice simulation results, which treat the fermions to all orders in g , we discuss the mechanism responsible for the dramatically enhanced NLO quantum correction.

For this we consider in Fig. 2 the total number density $N_\psi(t) = 8 \int d^3p n_\psi(t, \mathbf{p}) / (2\pi)^3$ as a function of time. Both LO (dashed) and NLO (solid) results agree rather well at early times at which parametric resonance leads to an exponential increase of the inflaton two-point function, $F_\phi(t, t; \mathbf{p}_0) \sim A_0 e^{2\gamma_0 t}$. Here γ_0 denotes the growth rate of the maximally amplified momentum mode $\mathbf{p}_0$ of the inflaton correlation function [1–3]. We find that significant deviations between the LO and NLO fermion production occur at the time

$$t_\phi - \frac{1}{4\gamma_0} \ln(t_\phi \phi_0) \simeq \frac{1}{2\gamma_0} \ln \left(\frac{12N_s \sqrt{\pi^3 |\gamma_0''| \phi_0^3}}{(N_s + 1)A_0 |\mathbf{p}_0|^2 \lambda} \right), \quad (8)$$

where parametric resonance ends. (Here γ_0'' is the second derivative of the momentum dependent growth rate evaluated at $|\mathbf{p}_0|$ [1–3].) At this time the inflaton two-point function starts to become non-perturbatively large, i.e. parametrically $F_\phi(t, t; \mathbf{p}_0) \sim A_0/\lambda$. At the same characteristic time, the inflaton field amplitude $\phi(t)$ starts to decrease as can be seen from Fig. 3, which diminishes the $g\phi(t)$ contribution to the LO equation of motion (6).

The non-perturbative dependence of F_ϕ on the inverse of the coupling for $t \gtrsim t_\phi$ is the reason for the amplified fermion production by quantum corrections. The mechanism goes as follows. The loop correction (7) $\sim g^2$ contains a boson line with a contribution F_ϕ of order $1/\lambda$, which together lead to corrections of order $\xi = g^2/\lambda$. Since inflation typically requires small λ , we




FIG. 3: Inflaton field amplitude.

observe strong enhancement with $\xi \gg g^2$. We have explicitly verified that the fermion production rate scales proportional to $\sim \xi$ in this regime. We emphasize that the observed phenomenon of dynamic enhancement of quantum corrections is not restricted to chaotic inflation and parametric resonance. The non-perturbatively large correlations lead to universal behavior [11], and the discussion for tachyonic preheating or spinodal instabilities would follow along the same lines.

More generally, for any boson field with non-perturbatively large occupation number and Yukawa-type coupling to fermions a similar enhancement of quantum corrections may be observed. This concerns, in particular, also quantum chromodynamics. In the context of heavy-ion collisions large gluon occupation numbers $\sim 1/\alpha_s$, where the running gauge coupling α_s is small at sufficiently high energies, would enhance the diagrammatic contribution (7) to order one if the boson line is associated to gluons and the fermion lines to quarks.

In view of the dramatic differences between LO and NLO results, it remains to demonstrate that higher-order quantum corrections do not change much our understanding. Since the interaction term (1) is quadratic in the fermion fields, we 'integrate out' the fermions on a lattice and study the remaining bosonic effective theory by numerical integration of the equations of motion and Monte Carlo sampling [9]. This requires numerical integration of the fermion propagator, which becomes feasible in $3 + 1$ dimensions using 'low-cost' fermion algorithms suggested in Ref. [10]. The approach treats the fermions exactly. However, the inflaton dynamics becomes classical-statistical, which has been shown in the past to be an accurate description for sufficiently early times we are considering here [3].

Remarkably, the lattice simulation results shown in Fig. 1 (symbols) agree well with the NLO approximation up to momenta of about $1.3/\phi_0$ [15] for the employed




FIG. 4: Inflaton occupation number of transverse modes.

small ξ . We conclude that the distribution indeed exhibits a thermal value $n_{\text{eq}}(0) = 1/2$ for massless fermions at low momenta, which drops at a characteristic scale $\sim \phi_0$. We emphasize that the inflaton is still far from equilibrium at this stage. Fig. 4 shows the inflaton occupation number as a function of momentum at different times, following standard conventions of Refs. [1–3]. In particular, at time $t = 250/\phi_0$ the inflaton clearly approaches a non-thermal power-law distribution $\sim 1/|\mathbf{p}|^4$ characteristic for strong turbulence [11].

A potentially important application of fermion preheating is the possibility to produce very heavy fermions. Superheavy fermions may be interesting for the dark matter problem and for the problem of ultra-high energy cosmic rays. Out of equilibrium this question can be addressed using the dynamical spectral function (4). For the discussion of $\rho(x, y)$ we introduce Wigner coordinates, $X = (x + y)/2$ and $x - y$. Fourier transformation with respect to the latter gives the dependence on frequency-momenta. For the spatially homogeneous system the spectral function then depends in addition only on time X^0 . For the temporal component $\rho_V^0 = \text{tr}(\gamma^0 \rho)/4$ the sum over all frequencies is normalized to one,

$$-i \int \frac{d\omega}{2\pi} \rho_V^0(X^0; \omega, \mathbf{p}) = 1, \quad (9)$$

which corresponds to the equal-time anti-commutation relation for fermions. Fig. 5 shows NLO results at different times and $|\mathbf{p}|/\phi_0 \simeq 0$. Initially (dashed-dotted curve) we find a double peak at $\omega \simeq 0$, corresponding to massless fermions, and $\omega \simeq g\phi(t=0)/N_f$ characterizing very heavy fermion states. At later times these peaks become much broader and lower, which is characteristic for the presence of strong correlations without quasi-particle interpretations.

We thank Denes Sexty for collaboration on related topics. This work is supported by the BMBF grant 06DA9018.




FIG. 5: Nonequilibrium fermion spectral function.

-
- [1] J. H. Traschen and R. H. Brandenberger, Phys. Rev. D **42** (1990) 2491; L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. Lett. **73** (1994) 3195.
 - [2] S. Y. Khlebnikov and I. I. Tkachev, Phys. Rev. Lett. **77** (1996) 219.
 - [3] J. Berges and J. Serreau, Phys. Rev. Lett. **91** (2003) 111601.
 - [4] J. Adams et. al., Nucl. Phys. A **757** (2005) 102; K. Adcox et. al., *ibid.* **757** (2005) 184; I. Arsene et. al., *ibid.* **757** (2005) 1; B. B. Back et. al., *ibid.* **757** (2005) 28.
 - [5] J. Baacke, K. Heitmann and C. Patzold, Phys. Rev. D **58** (1998) 125013; P. B. Greene and L. Kofman, Phys. Lett. B **448** (1999) 6; G. F. Giudice, M. Peloso, A. Riotto and I. Tkachev, JHEP **9908** (1999) 014; J. Garcia-Bellido, S. Mollerach and E. Roulet, JHEP **0002** (2000) 034; J. Baacke, N. Kevlishvili and J. Pruschke, JCAP **0706** (2007) 004.
 - [6] F. Gelis, K. Kajantie and T. Lappi, Phys. Rev. Lett. **96** (2006) 032304.
 - [7] J. Berges, S. Borsanyi and J. Serreau, Nucl. Phys. B **660** (2003) 51.
 - [8] J. Berges, J. Pruschke and A. Rothkopf, Phys. Rev. D **80** (2009) 023522.
 - [9] G. Aarts and J. Smit, Nucl. Phys. B **555** (1999) 355.
 - [10] S. Borsanyi and M. Hindmarsh, Phys. Rev. D **79** (2009) 065010.
 - [11] J. Berges, A. Rothkopf and J. Schmidt, Phys. Rev. Lett. **101** (2008) 041603.
 - [12] R. Micha and I. I. Tkachev, Phys. Rev. Lett. **90** (2003) 121301.
 - [13] J. Berges, Nucl. Phys. A **699** (2002) 847. G. Aarts et al, Phys. Rev. D **66** (2002) 045008.
 - [14] J. Berges, Nucl. Phys. A **820** (2009) 65C.
 - [15] For the 64^3 lattice with spacing $a_s = 1.5/\phi_0$ the observed differences at somewhat larger $|\mathbf{p}|$ are found to depend on the spatial Wilson term, which we have to add to suppress doublers for the employed Wilson fermions. A possible better agreement on even larger lattices and a systematic study of cutoff dependencies also for those momenta is beyond the scope of the present work.