# Convexity of quantum  $\chi^2$ -divergence

Frank Hansen

February 15 2011

#### Abstract

The quantum  $\chi^2$ -divergence has recently been introduced and applied to quantum channels (quantum Markov processes). In contrast to the classical setting the quantum  $\chi^2$ -divergence is not unique but depends on the choice of quantum statistics. In the reference [\[13\]](#page-6-0) a special one-parameter family of quantum  $\chi^2_{\alpha}(\rho, \sigma)$ -divergences for density matrices were studied, and it was established that they are convex functions in  $(\rho, \sigma)$  for parameter values  $\alpha \in [0, 1]$ , thus mirroring the classical theorem for the  $\chi^2(p,q)$ -divergence for probability distributions  $(p, q)$ . We prove that any quantum  $\chi^2$ -divergence is a convex function in its two arguments.

Key words and phrases: Quantum  $\chi^2$ -divergence, quantum statistics, monotone metric, convexity, operator monotone function.

# 1 Introduction

The geometrical formulation of quantum statistics originates in a study by Chentsov of the classical Fisher information. Chentsov proved [\[3\]](#page-5-0) that the Fisher-Rao metric is the only Riemannian metric, defined on the tangent space, that is decreasing under Markov morphisms. Since Markov morphisms represent coarse graining or randomization, it means that the Fisher information is the only Riemannian metric possessing the attractive property that distinguishability of probability distributions becomes more difficult when they are observed through a noisy channel.

Morozova [\[11\]](#page-6-1) extended the analysis to quantum mechanics by replacing Riemannian metrics defined on the tangent space of the simplex of probability distributions with positive definite sesquilinear (originally bilinear) forms  $K_{\rho}$  defined on the tangent space of a quantum system, where  $\rho$  is a positive definite state. Customarily,  $K_{\rho}$  is extended to all operators (matrices) supported by the underlying Hilbert space, cf. [\[12,](#page-6-2) [5\]](#page-6-3) for details. Noisy channels are in this setting represented by stochastic (completely positive and trace preserving) mappings, and the contraction property is replaced by the monotonicity requirement

$$
K_{T(\rho)}(T(A), T(A)) \le K_{\rho}(A, A)
$$

for every stochastic mapping  $T : M_n(\mathbb{C}) \to M_m(\mathbb{C})$ . Unlike the classical situation, these requirements no longer uniquely determine the metric.

We consider the following class of functions which is used to characterize monotone metrics.

**Definition 1.1.**  $\mathcal{F}_{op}$  is the class of functions  $f : (0, +\infty) \to (0, +\infty)$  such that

- $(i)$  f is operator monotone,
- (*ii*)  $f(t) = tf(t^{-1})$  for  $t > 0$ ,

(*iii*) 
$$
f(1) = 1
$$
.

By the combined efforts of Chentsov, Morozova and Petz [\[12\]](#page-6-2) it was established that a monotone metric is given on the canonical form

(1) 
$$
K_{\rho}(A,B) = \text{Tr} A^* c(L_{\rho}, R_{\rho})(B),
$$

where the so-called Morozova-Chentsov function  $c$  is of the form

<span id="page-1-0"></span>
$$
c(x,y) = \frac{1}{yf(xy^{-1})} \qquad f: \mathbf{R}_{+} \to \mathbf{R}_{+}
$$

for a function  $f \in \mathcal{F}_{op}$  and c is taken in the two commuting positive definite (super) operators  $L_{\rho}$  and  $R_{\rho}$  defined by setting

$$
L_{\rho}A = \rho A
$$
 and  $R_{\rho}A = A\rho$ .

It is condition  $(ii)$  in the definition above that ensures symmetry of the metric in the sense that  $K_{\rho}(A, B) = K_{\rho}(B, A)$  for self-adjoint A and B.

Lesniewski and Ruskai [\[9\]](#page-6-4) gave equivalent descriptions in terms of operator convex functions and operator monotone decreasing functions. In particular

$$
K_{\rho}(A, B) = \text{Tr} \, A^* R_{\rho}^{-1} k (L_{\rho} R_{\rho}^{-1}) B,
$$

where  $k: \mathbf{R}_{+} \to \mathbf{R}_{+}$  is an operator monotone decreasing function satisfying  $k(t^{-1}) = tk(t)$  for  $t > 0$  and  $k(1) = 1$ . The Morozova-Chentsov-Petz formalism is then recovered by inserting the operator monotone function  $f(t) = 1/k(t)$  in equation [\(1\)](#page-1-0).

**Definition 1.2.** The  $\chi_f^2$ -divergence (relative to a choice of monotone metric) is given by

$$
\chi_f^2(\rho,\sigma) = K_\sigma^c(\rho - \sigma, \rho - \sigma),
$$

where

$$
c(x, y) = \frac{1}{yf(xy^{-1})}
$$
  $x, y > 0$ 

is the Morozova-Chentsov function specified by a function  $f \in \mathcal{F}_{op}$ .

The functions

$$
f_{\alpha}(t) = \frac{2t^{\alpha}}{t^{2\alpha - 1} + 1} \qquad t > 0
$$

with parameter  $\alpha \in [0, 1]$  are elements in  $\mathcal{F}_{op}$  and correspond to the functions

$$
k(t) = \frac{1}{2}(t^{-\alpha} + t^{\alpha - 1}) \qquad t > 0
$$

in the Ruskai-Lesniewski formalism. The associated Morozova-Chentsov functions are given by

$$
c_{\alpha}(x,y) = \frac{1}{y f_{\alpha}(xy^{-1})} = \frac{x^{\alpha-1} y^{-\alpha} + x^{-\alpha} y^{\alpha-1}}{2}
$$

and the divergences for positive definite density matrices are

$$
\chi_{\alpha}^{2}(\rho,\sigma) = \text{Tr}(\rho - \sigma)\sigma^{-\alpha}(\rho - \sigma)\sigma^{\alpha-1} = \text{Tr}\,\rho\sigma^{-\alpha}\rho\sigma^{\alpha-1} - 1.
$$

This expression is convex in  $(\rho, \sigma)$  as pointed out in [\[13\]](#page-6-0).

## 2 Convexity

Consider a function  $f: \mathbf{R}_{+} \to \mathbf{R}_{+}$ . The perspective (function) of f is the function  $g$  of two variables defined by setting

$$
g(t,s) = sf(ts^{-1}) \qquad t,s > 0.
$$

Effros showed [\[4\]](#page-6-5) that if f is operator convex then, whenever meaningful, the perspective is a convex operator function of two variables. In particular, it becomes operator convex in the sense of Korányi [\[8\]](#page-6-6). It is also convex in functions of commuting operators. In particular, the function

$$
(\rho, \sigma) \to g(L_{\rho}, R_{\sigma})
$$

is convex in pairs of positive definite  $n \times n$  matrices, equivalent to the statement that the function

$$
(\rho, \sigma) \to \text{Tr} \, A^* g(L_\rho, R_\sigma) A
$$

is convex for any  $n \times n$  matrix A. Similar statements are valid also for operator concave functions.

**Theorem 2.1.** The  $\chi^2_f(\rho, \sigma)$ -divergence is convex in  $(\rho, \sigma)$  for any f in  $\mathcal{F}_{op}$ . *Proof.* Let us consider a function f in  $\mathcal{F}_{op}$  with Morozova-Chensov function

$$
c(x,y) = \frac{1}{yf(xy^{-1})} = F(x,y)^{-1} \qquad x, y > 0,
$$

where  $F(x, y) = yf(xy^{-1})$  is the perspective of f. Since f is operator concave we obtain that  $F(x, y)$  is operator concave as a function of two variables. Inversion (of super operators) is decreasing. By using linearity of the mappings  $\sigma \to L_{\sigma}$  and  $\sigma \to R_{\sigma}$  we therefore obtain

$$
F(L_{\lambda\sigma_1+(1-\lambda)\sigma_2}, R_{\lambda_1\sigma_1+(1-\lambda)\sigma_2})^{-1}
$$
  
=  $F(\lambda L_{\sigma_1}+(1-\lambda)L_{\sigma_2}, \lambda R_{\sigma_1}+(1-\lambda)R_{\sigma_2})^{-1}$   
 $\leq (\lambda F(L_{\sigma_1}, R_{\sigma_1})+(1-\lambda)F(L_{\sigma_2}, R_{\sigma_2}))^{-1}$ 

for states (density matrices)  $\rho_1, \rho_2$  and  $\sigma_1, \sigma_2$  and real numbers  $\lambda \in [0, 1]$ . The divergence  $\chi_f^2(\rho, \sigma)$  is given on the form

$$
\chi_f^2(\rho,\sigma) = \text{Tr}(\rho-\sigma)c(L_\sigma,R_\sigma)(\rho-\sigma),
$$

and by setting

 $\rho = \lambda \rho_1 + (1 - \lambda) \rho_2$  and  $\sigma = \lambda \sigma_1 + (1 - \lambda) \sigma_2$ 

we obtain the inequality

$$
\chi_f^2(\rho, \sigma) = \text{Tr}(\rho - \sigma) F(L_{\sigma}, R_{\sigma})^{-1}(\rho - \sigma)
$$
  
\n
$$
= \text{Tr}(\rho - \sigma) F(L_{\lambda\sigma_1 + (1-\lambda)\sigma_2}, R_{\lambda\sigma_1 + (1-\lambda)\sigma_2})^{-1}(\rho - \sigma),
$$
  
\n
$$
\leq \text{Tr}(\rho - \sigma) (\lambda F(L_{\sigma_1}, R_{\sigma_1}) + (1 - \lambda) F(L_{\sigma_2}, R_{\sigma_2}))^{-1}(\rho - \sigma)
$$
  
\n
$$
= \text{Tr}(\lambda(\rho_1 - \sigma_1) + (1 - \lambda)(\rho_2 - \sigma_2)) (\lambda F(L_{\sigma_1}, R_{\sigma_1}) + (1 - \lambda) F(L_{\sigma_2}, R_{\sigma_2}))^{-1}
$$
  
\n
$$
(\lambda(\rho_1 - \sigma_1) + (1 - \lambda)(\rho_2 - \sigma_2))
$$
  
\n
$$
\leq \lambda \text{Tr}(\rho_1 - \sigma_1) F(L_{\sigma_1}, R_{\sigma_1})^{-1}(\rho_1 - \sigma_1) + (1 - \lambda) \text{Tr}(\rho_2 - \sigma_2) F(L_{\sigma_2}, R_{\sigma_2})^{-1}(\rho_2 - \sigma_2)
$$
  
\n
$$
= \lambda \chi_f^2(\rho_1, \sigma_1) + (1 - \lambda) \chi_f^2(\rho_2, \sigma_2),
$$

where we in the second inequality, applied on super operators, used that the mapping

$$
(A,\xi)\to(\xi\mid A^{-1}\xi)
$$

is jointly convex for positive invertible operators  $A$  on a Hilbert space  $H$ , and vectors  $\xi \in H$ , cf. [\[6,](#page-6-7) Proposition 4.3]. This is also a direct consequence of convexity of the mapping

$$
(A, B) \to B^* A^{-1} B
$$

for B arbitrary and A positive definite, cf. [\[10,](#page-6-8) Remark after Theorem 1] and [\[6,](#page-6-7) Remark 4.5]. Furthermore, it is related to  $[1,$  Theorem 3.1]. **QED** 

Any function f in  $\mathcal{F}_{op}$  satisfies the inequalities

$$
\frac{2t}{t+1} \le f(t) \le \frac{t+1}{2} \qquad t > 0,
$$

where the smallest function in  $\mathcal{F}_{op}$  corresponds to the Bures metric. We mention the following characterization [\[5,](#page-6-3) [7,](#page-6-9) [2\]](#page-5-2) of the functions in  $\mathcal{F}_{op}$ .

**Theorem 2.2.** A function f in  $\mathcal{F}_{op}$  admits a canonical representation

<span id="page-4-0"></span>(2) 
$$
f(t) = \frac{1+t}{2} \exp \left[ - \int_0^1 \frac{(1-\lambda^2)(1-t)^2}{(\lambda+t)(1+\lambda t)(1+\lambda)^2} h(\lambda) d\lambda \right],
$$

where the weight function  $h : [0,1] \rightarrow [0,1]$  is measurable. The equivalence class containing h is uniquely determined by f. Any function on the given form is in  $\mathcal{F}_{op}$ .

Notice that the integral kernel is non-negative for every  $t > 0$ . The representation induces an order relation  $\preceq$  in  $\mathcal{F}_{op}$  stronger than the point-wise order by setting  $f \preceq g$  if the representing weight functions  $h_f$  and  $h_g$  satisfies  $h_f \geq h_g$  almost everywhere. With this order relation  $(\mathcal{F}_{op}, \preceq)$  becomes a lattice, inducing a lattice structure on the set of quantum  $\chi^2$  divergences. It is compatible with the parametrization of the Wigner-Yanase-Dyson metrics, cf. [\[2,](#page-5-2) Theorem 2.8].

The representation in [\(2\)](#page-4-0) may be used to construct families of metrics that increase monotonously from the smallest (Bures) metric to the largest. In fact any family of weight functions that decreases monotonously from the constant 1 to the zero function will induce this property. In [\[5,](#page-6-3) Proposition 3 we considered the constant weight functions  $h_{\alpha} = \alpha$  for  $0 \le \alpha \le 1$  and obtained in this way a family of metrics

$$
f_{\alpha}(t) = t^{\alpha} \left(\frac{1+t}{2}\right)^{1-2\alpha} \qquad t > 0,
$$

that decreases monotonously from the largest monotone metric down to the Bures metric for  $\alpha$  increasing from 0 to 1. In the Lesniewski-Ruskai representation that corresponds to the functions

$$
k_{\alpha}(\omega) = w^{-\alpha} \left(\frac{1+\omega}{2}\right)^{2\alpha-1}
$$

as mentioned in [\[13,](#page-6-0) Appendix A].

## <span id="page-5-1"></span>References

- [1] T. Ando and F. Hiai. Operator log-convex functions and operator means. Math. Ann., DOI 10.1107/s00208-010-0577-4, 2010.
- <span id="page-5-2"></span>[2] K. Audenaert, L. Cai, and F. Hansen. Inequalities for quantum skew information. Lett Math Phys, 85:135–146, 2008.
- <span id="page-5-0"></span>[3] N.N. Censov. Statistical Decision Rules and Optimal Inferences, Transl. Math. Monogr., volume 53. Amer. Math. Soc., Providence, 1982.
- <span id="page-6-5"></span><span id="page-6-3"></span>[4] E.G. Effros. A matrix convexity approach to some celebrated quantum inequalities. Proc. Natl. Acad. Sci. USA, 106:1006–1008, 2009.
- [5] F. Hansen. Characterizations of symmetric monotone metrics on the state space of quantum systems. Quantum Information and Computation, 6:597–605, 2006.
- <span id="page-6-9"></span><span id="page-6-7"></span>[6] F. Hansen. Extensions of lieb's concavity theorem. Journal of Statistical Physics, 124:87–101, 2006.
- <span id="page-6-6"></span>[7] F. Hansen. Metric adjusted skew information. Proc. Natl. Acad. Sci. USA, 105:9909–9916, 2008.
- <span id="page-6-4"></span>[8] A. Korányi. On some classes of analytic functions of several variables. Trans. Amer. Math. Soc., 101:520–554, 1961.
- [9] A. Lesniewski and M.B. Ruskai. Monotone Riemannian metrics and relative entropy on non-commutative probability spaces. J. Math. Phys., 40:5702–5724, 1999.
- <span id="page-6-8"></span><span id="page-6-1"></span>[10] E.H. Lieb and M.B. Ruskai. Some operator inequalities of the Schwarz type. Adv. in Math., 12:269–273, 1974.
- [11] E.A. Morozova and N.N. Chentsov. Markov invariant geometry on state manifolds (Russian). Itogi Nauki i Techniki, 36:69–102, 1990. Translated in J. Soviet Math. 56:2648-2669, 1991.
- <span id="page-6-2"></span><span id="page-6-0"></span>[12] D. Petz. Monotone metrics on matrix spaces. Linear Algebra Appl., 244:81–96, 1996.
- [13] K. Temme, M.J. Kastoryano, M.B. Ruskai, M.M. Wolf, and F. Verstrate. The  $\chi^2$ -divergence and mixing times of quantum Markov processes. J. Math. Phys., 51:122201, 2010.

Frank Hansen: Institute for International Education, Tohoku University, Japan. Email: frank.hansen@m.tohoku.ac.jp.