ON THE TRANSPOSITION ANTI-INVOLUTION IN REAL CLIFFORD ALGEBRAS III: THE AUTOMORPHISM GROUP OF THE TRANSPOSITION SCALAR PRODUCT ON SPINOR SPACES

RAFAL ABLAMOWICZ AND BERTFRIED FAUSER

ABSTRACT. A signature $\varepsilon = (p,q)$ dependent transposition anti-involution $T_{\varepsilon}^{\varepsilon}$ of real Clifford algebras $C_{p,q}$ for non-degenerate quadratic forms was introduced in [\[1\]](#page-22-0). In [\[2\]](#page-22-1) we showed that, depending on the value of $(p - q)$ mod 8, the map T_{ε} gives rise to transposition, complex Hermitian, or quaternionic Hermitian conjugation of representation matrices in spinor representation. The resulting scalar product is in general different from the two known standard scalar products [\[12\]](#page-22-2). We provide a full signature (p, q) dependent classification of the invariance groups $G_{p,q}^{\varepsilon}$ of this product for $p+q \leq 9$. The map T_{ε}^{\sim} is identified as the "star" map known [\[14\]](#page-23-0) from the theory of (twisted) group algebras, where the Clifford algebra $C\ell_{p,q}$ is seen as a twisted group ring $\mathbb{R}^t[(\mathbb{Z}_2)^n]$, $n = p + q$. We discuss and list important subgroups of stabilizer groups $G_{p,q}(f)$ and their transversals in relation to generators of spinor spaces.

1. INTRODUCTION

This article concludes the developments of [\[1,](#page-22-0) [2\]](#page-22-1). Hence, all notation and definitions are the same. Recall that in addition to reversion, grade involution and conjugation, any universal real Clifford algebra $C\ell(V,Q)$ of a non-degenerate quadratic real vector space possesses a transposition anti-involution T_{ε} ^{*} as a unique extension of a certain orthogonal map $t_{\varepsilon}: V \to V^{\flat}$. Using the identification $V^{\flat} \cong V^*$, the orthogonal map t_{ε} was shown to be a symmetric non-degenerate correlation on V. For the properties of t_{ε} see [\[1\]](#page-22-0).

Let $C\ell_{p,q}$ be a simple Clifford algebra $(p - q \neq 1 \mod 4)$ and let $S = C\ell_{p,q} f$ be a left spinor ideal generated by a primitive idempotent f. As usual, we let $\mathbb{K} = f\mathcal{C}\ell_{p,q}f$. Then, T_{ε}^{\sim} allows one to define [\[2,](#page-22-1) Prop. 3] a K-valued transposition spinor scalar product $S \times S \to \mathbb{K}$ $as¹$ $as¹$ $as¹$

$$
\forall \psi, \phi \in S, \quad \exists! \lambda \in \mathbb{K}, \quad \text{s.t.} \quad (\psi, \phi) \mapsto T_{\varepsilon}(\psi)\phi = \lambda f = f\lambda. \tag{1}
$$

¹⁹⁹¹ Mathematics Subject Classification. 15A66, 16S35, 20C05, 68W30.

Key words and phrases. grade involution, group ring, indecomposable module, involution, minimal left ideal, monomial order, primitive idempotent, reversion, semisimple algebra, spinor, stabilizer, transversal, twisted group ring, universal Clifford algebra.

¹Recall that elements λ of $\mathbb{K} = fC\ell_{p,q}f$ commute with f because $\lambda = fuf$ for some $u \in C\ell_{p,q}$. The notion ∃! stands for 'there exists a unique'.

The group $G_{p,q}^{\varepsilon}$ was defined in [\[2\]](#page-22-1) as the invariance group of this transposition scalar product:

$$
G_{p,q}^{\varepsilon} = \{ g \in C\ell_{p,q} \, | \, T_{\varepsilon}^{\sim}(g)g = 1 \}. \tag{2}
$$

In particular, the product is invariant under two of its subgroups: Salingaros' [\[17,](#page-23-1) [18,](#page-23-2) [19\]](#page-23-3) finite vee group $G_{p,q} < G_{p,q}^{\varepsilon}$ and the stabilizer group $G_{p,q}(f) \triangleleft G_{p,q}$ of a primitive idempotent f under the conjugate action of $G_{p,q}$ on $Cl_{p,q}$. Since the stabilizer group $G_{p,q}(f)$ plays an important role in constructing and understanding spinor representation of Clifford algebras, we classified these groups in [\[2\]](#page-22-1).

In this article, in Section [2,](#page-1-0) we discuss important subgroups of $G_{p,q}(f)$ which are related to the idempotent f and to the division ring K . Furthermore, we show how transversals of these subgroups in the vee group $G_{p,q}$ are in fact generators of the (left) spinor spaces $C_{\mathcal{C}}S_{\mathbb{K}}$ viewed as real vector spaces or as right modules over K.

Then, in Section [3,](#page-7-0) we undertake a systematic study of the transposition scalar product $T_{\varepsilon}(\psi)\phi$ and its automorphism group $G_{p,q}^{\varepsilon}$ for all signatures (p,q) , $p+q\leq 9$, in simple and semisimple Clifford algebras.

In Section [4](#page-18-0) we conclude that our transposition anti-involution T_{ε}^{\sim} of $C\ell_{p,q}$ is related to a certain anti-involution $* : \mathbb{R}^t[(\mathbb{Z}_2)^n] \to \mathbb{R}^t[(\mathbb{Z}_2)^n]$ when we identify the Clifford algebra $C\ell_{p,q}$ with a twisted group ring $\mathbb{R}^t[(\mathbb{Z}_2)^n]$ of an abelian group $(\mathbb{Z}_2)^n = \mathbb{Z}_2 \times \cdots \times \mathbb{Z}_2$, n times where $n = p + q$, in the sense of [\[6\]](#page-22-3).

Finally, in Section [5,](#page-20-0) we summarize the results obtained in all three papers and provide some ideas how they can be extended and generalized.

2. Subgroups of the stabilizer group

Let $C\ell_{p,q}$ be a real universal Clifford algebra with Grassmann basis β sorted by the admissible order InvLex [\[1\]](#page-22-0). We begin by recalling the definition of Salingaros' vee group [\[17\]](#page-23-1).

Definition 1. A vee group $G_{p,q}$ is defined as the set $G_{p,q} = \{\pm m \mid m \in \mathcal{B}\}\$ in $C\ell_{p,q}$ together with the Clifford product as the group binary operation.

This finite group is of order $|G_{p,q}| = 2^{1+p+q}$ and its commutator subgroup $G'_{p,q} = \{\pm 1\}.$

We know that any polynomial f in $C\ell_{p,q}$ expressible as a product of idempotents $\frac{1}{2}(1 \pm \mathbf{e}_{\underline{i}_s})$ namely

$$
f = \frac{1}{2} (1 \pm \mathbf{e}_{\underline{i}_1}) \frac{1}{2} (1 \pm \mathbf{e}_{\underline{i}_2}) \cdots \frac{1}{2} (1 \pm \mathbf{e}_{\underline{i}_k}),
$$
(3)

where $\mathbf{e}_{i_1}, \ldots, \mathbf{e}_{i_k}, k = q - r_{q-p}$, are commuting basis monomials in \mathcal{B} with square 1, is a primitive idempotent in $\mathcal{C}\ell_{p,q}$ ^{[3](#page-1-2)} Furthermore, $\mathcal{C}\ell_{p,q}$ has a complete set of 2^k such primitive mutually annihilating idempotents which add up to the identity 1. Moreover, the basis

²Here, r_i is Radon-Hurwitz number defined by recursion as $r_{i+8} = r_i + 4$ and these initial values: $r_0 = 0, r_1 = 1, r_2 = r_3 = 2, r_4 = r_5 = r_6 = r_7 = 3$ [\[8,](#page-22-4) [12\]](#page-22-2).

³In the following, for f we can pick any of the 2^k primitive idempotents. In examples, our default choice for f will always be the one in which all signs in the factorization (3) are plus.

monomials which define f generate a group $\langle \mathbf{e}_{i_1}, \ldots, \mathbf{e}_{i_k} \rangle \cong (\mathbb{Z}_2)^k$, see [\[2\]](#page-22-1) and references therein.

With any primitive idempotent f we associate three groups:

(i) The *stabilizer* $G_{p,q}(f)$ of f defined as

$$
G_{p,q}(f) = \{ m \in G_{p,q} \mid mfm^{-1} = f \} < G_{p,q}.\tag{4}
$$

The stabilizer $G_{p,q}(f)$ is a normal subgroup of $G_{p,q}$. These groups were classified, depending on the signature (p, q) , in [\[2\]](#page-22-1). In particular, recall that

$$
|G_{p,q}(f)| = \begin{cases} 2^{1+p+r_{q-p}}, & p-q \neq 1 \text{ mod } 4; \\ 2^{2+p+r_{q-p}}, & p-q = 1 \text{ mod } 4. \end{cases}
$$
(5)

(ii) A new abelian *idempotent group* $T_{p,q}(f)$ of f, a subgroup of $G_{p,q}(f)$, generated by the commuting basis monomials e_{i_1}, \ldots, e_{i_k} , and their negatives, which appear in the factorization (3) of f .

(iii) A field group $K_{p,q}(f)$ of f, a subgroup of $G_{p,q}(f)$, related to the (skew double) field $\mathbb{K} \cong fC\ell_{p,q}f.$

Definition 2. Let f be a primitive idempotent in $C\ell_{p,q}$ defined as in [\(3\)](#page-1-3). Then, the *idempotent group* $T_{p,q}(f)$ *of* f is defined as

$$
T_{p,q}(f) = \langle \pm 1, \mathbf{e}_{\underline{i}_1}, \dots, \mathbf{e}_{\underline{i}_k} \rangle < G_{p,q}(f),\tag{6}
$$

where $k = q - r_{q-p}$.

Thus, immediately from this definition we obtain the following

Lemma 1. $T_{p,q}(f)$ has the following structure:

$$
T_{p,q}(f) \cong G'_{p,q} \times \langle \mathbf{e}_{\underline{i}_1}, \dots, \mathbf{e}_{\underline{i}_k} \rangle \cong G'_{p,q} \times (\mathbb{Z}_2)^k \quad and \quad |T_{p,q}(f)| = 2^{1+k} \tag{7}
$$

where $k = q - r_{q-p}$.

Proof. This is straightforward since $G'_{p,q}$ and $\langle \mathbf{e}_{\underline{i}_1}, \ldots, \mathbf{e}_{\underline{i}_k} \rangle$ are normal subgroups of $T_{p,q}(f)$ with trivial intersection, $T_{p,q}(f) = \langle \mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_k} \rangle \tilde{G}'_{p,q}$, and $k = q - r_{q-p}$ as in [\(3\)](#page-1-3).

Our third group, the field group $K_{p,q}(f)$, is related to the division ring $\mathbb{K} = fC\ell_{p,q}f$. Recall that for simple Clifford algebras, left spinor ideals $S = Cl_{p,q}f$ are right K-modules where $\mathbb{K} \simeq \mathbb{R}$, (respectively, $\mathbb{K} \simeq \mathbb{C}$, or $\mathbb{K} \simeq \mathbb{H}$,) when $p - q = 0, 1, 2 \mod 8$ (respectively, $p - q = 3, 7 \mod 8$, or $p - q = 4, 5, 6 \mod 8$ [\[2\]](#page-22-1).

Definition 3. Let f be a primitive idempotent in $C\ell_{p,q}$ defined as in [\(3\)](#page-1-3) and let $\mathbb{K} =$ $f\mathcal{C}\ell_{p,q}f$. Let K be a set of Grassmann monomials in B which span K as a real algebra, that is, $\mathbb{K} = \text{span}_{\mathbb{R}}\{m \mid m \in \mathcal{K}\}.$ Then, the field group of f is defined as

$$
K_{p,q}(f) = \langle \pm 1, m \mid m \in \mathcal{K} \rangle < G_{p,q}(f). \tag{8}
$$

Therefore, immediately from the definition, we get that

$$
|K_{p,q}(f)| = \begin{cases} 2, & p-q = 0, 1, 2 \text{ mod } 8; \\ 4, & p-q = 3, 7 \text{ mod } 8; \\ 8, & p-q = 4, 5, 6 \text{ mod } 8. \end{cases}
$$
(9)

Before we state our main theorem which summarizes relations between the groups $G_{p,q}$, $G_{p,q}(f)$, $T_{p,q}(f)$, $K_{p,q}(f)$, and $G'_{p,q}$, we recall the definition of a *(left)* transversal [\[16,](#page-23-4) Sect. 5.6].

Definition 4. Let K be a subgroup of a group G. A transversal ℓ of K in G is a subset of G consisting of exactly one element $\ell(bK)$ from every (left) coset bK, and with $\ell(K) = 1$.

A transversal of a normal subgroup $K \triangleleft G$ is the same as the image of a *lifting* function $\ell: Q \to G$ in the extension of K by Q [\[16,](#page-23-4) Sect. 10.2] (using Rotman's notation ℓ for the lifting too).

Theorem 1 (Main Theorem). Let f be a primitive idempotent in a simple or semisimple Clifford algebra $C\ell_{p,q}$ and let $G_{p,q}, G_{p,q}(f), T_{p,q}(f), K_{p,q}(f)$, and $G'_{p,q}$ be the groups defined above. Furthermore, let $S = Cl_{p,q}f$ and $\mathbb{K} = fCl_{p,q}f$.

- (i) Elements of $T_{p,q}(f)$ and $K_{p,q}(f)$ commute.
- (ii) $T_{p,q}(f) \cap K_{p,q}(f) = G'_{p,q} = {\pm 1}.$
- (iii) $G_{p,q}(f) = T_{p,q}(f)K_{p,q}(f) = K_{p,q}(f)T_{p,q}(f).$
- $(iv) |G_{p,q}(f)| = |T_{p,q}(f)K_{p,q}(f)| = \frac{1}{2}$ $\frac{1}{2}|T_{p,q}(f)||K_{p,q}(f)|.$
- (v) $G_{p,q}(f) \triangleleft G_{p,q}$, $T_{p,q}(f) \triangleleft G_{p,q}$, and $K_{p,q}(f) \triangleleft G_{p,q}$. In particular, $T_{p,q}(f)$ and $K_{p,q}(f)$ are normal subgroups of $G_{p,q}(f)$.
- (vi) We have:

$$
G_{p,q}(f)/K_{p,q}(f) \cong T_{p,q}(f)/G'_{p,q},\tag{10}
$$

$$
G_{p,q}(f)/T_{p,q}(f) \cong K_{p,q}(f)/G'_{p,q}.\tag{11}
$$

(vii) We have:

$$
(G_{p,q}(f)/G'_{p,q})/(T_{p,q}(f)/G'_{p,q}) \cong G_{p,q}(f)/T_{p,q}(f) \cong K_{p,q}(f)/\{\pm 1\}
$$
 (12)

and the transversal of $T_{p,q}(f)$ in $G_{p,q}(f)$ spans $\mathbb K$ over $\mathbb R$ modulo f.

- (viii) The transversal of $G_{p,q}(f)$ in $G_{p,q}$ spans S over K modulo f.
- (ix) We have $(G_{p,q}(f)/T_{p,q}(f)) \triangleleft (G_{p,q}/T_{p,q}(f))$ and

$$
(G_{p,q}/T_{p,q}(f))/(G_{p,q}(f)/T_{p,q}(f)) \cong G_{p,q}/G_{p,q}(f)
$$
\n(13)

and the transversal of $T_{p,q}(f)$ in $G_{p,q}$ spans S over R modulo f. (x) The stabilizer $G_{p,q}(f)$ can be viewed as

$$
G_{p,q}(f) = \bigcap_{x \in T_{p,q}(f)} C_{G_{p,q}}(x) = C_{G_{p,q}}(T_{p,q}(f))
$$
\n(14)

where $C_{G_{p,q}}(x)$ is the centralizer of x in $G_{p,q}$ and $C_{G_{p,q}}(T_{p,q}(f))$ is the centralizer of $T_{p,q}(f)$ in $G_{p,q}$.

Proof. All of the following statements have been verified additionally by explicit computations in CLIFFORD for all signatures (p, q) , with $p + q \leq 9$.

- (i) Let $m \in K_{p,q}(f)$ where $K_{p,q}(f)$ is defined in [\(8\)](#page-2-0). To show that m commutes with every element in the group $T_{p,q}(f)$ defined in [\(6\)](#page-2-1), it is enough to show that $m\mathbf{e}_{\underline{i}_j}=\mathbf{e}_{\underline{i}_j}m$ for every $\mathbf{e}_{\underline{i}_j}\in\mathcal{T}$ where $\mathcal{T}=\{\mathbf{e}_{\underline{i}_1},\ldots,\mathbf{e}_{\underline{i}_k}\}$ is the set of $k=q-r_{q-p}$ basis monomials which define any primitive idempotent f as in [\(3\)](#page-1-3). Suppose, to the contrary, that there exists $e_{i_j} \in \mathcal{T}$ which does not commute with m. Thus, e_{i_j} and m anti-commute, or $\mathbf{e}_{\underline{i}}$, $m = -m\mathbf{e}_{\underline{i}}$. Let $P_j^{\pm} = \frac{1}{2}$ $\frac{1}{2}(1 \pm \mathbf{e}_{i_j})$. Without any loss of generality, let us assume that P_j^+ is a factor of f so that $f = f P_j^+ = P_j^+ f$. On the other hand, $mf = fm$ since $m \in K_{p,q}(f) \subset \mathbb{K}$ and all elements of K commute with f. Thus, $fm = fP_j^+m = fmP_j^- = m(fP_j^-) = 0$ since $P_j^+P_j^- = P_j^-P_j^+ = 0$, or, since m is invertible, $f = 0$ which is false. Therefore, m commutes with every element of $T_{p,q}(f)$ which is generated by the elements of $\mathcal T$ and ± 1 .
- (ii) Obviously, $\{\pm 1\} \subset T_{p,q}(f) \cap K_{p,q}(f)$. Let $m \in T_{p,q}(f) \cap K_{p,q}(f)$. Then, $m^2 = 1$ since every element of $T_{p,q}(f)$, except the identity, is of order 2. Thus, since also $m \in K_{p,q}(f)$, we conclude that $m = \pm 1$, or, $T_{p,q}(f) \cap K_{p,q}(f) \subset {\pm 1}$.
- (iii) Obviously, $T_{p,q}(f), K_{p,q}(f) < G_{p,q}(f)$ because elements of $T_{p,q}(f)$ and $K_{p,q}(f)$ commute with f, hence, they stabilize f. Therefore, we have $T_{p,q}(f)K_{p,q}(f)$ = $K_{p,q}(f)T_{p,q}(f) < G_{p,q}(f)$ where the equality comes from (i). To show that $T_{p,q}(f)K_{p,q}(f) = G_{p,q}(f)$, we use the following order argument: By the Product Formula [\[16,](#page-23-4) Prop. 2.73], we have that

$$
|T_{p,q}(f)K_{p,q}(f)| = \frac{|T_{p,q}(f)||K_{p,q}(f)|}{|T_{p,q}(f) \cap K_{p,q}(f)|} = 2^k |K_{p,q}(f)| = |G_{p,q}(f)| \tag{15}
$$

thanks to Lemma [1;](#page-2-2) part (ii) of this Theorem; and the formula for $|G_{p,q}(f)|$ given in [\(5\)](#page-2-3).

- (iv) This follows from (ii), (iii), and the Product Formula [\[16,](#page-23-4) Prop. 2.73].
- (v) The fact that $G_{p,q}(f) \triangleleft G_{p,q}$ was proven in [\[2\]](#page-22-1). Then, $T_{p,q}(f) \triangleleft G_{p,q}$ and $K_{p,q}(f) \triangleleft G_{p,q}$ has been verified with CLIFFORD in Clifford algebras $C\ell_{p,q}$ for $p+q \leq 9$. Of course, when $K_{p,q}(f) = {\pm 1}$ then $T_{p,q}(f) = G_{p,q}(f) \triangleleft G_{p,q}$ and $K_{p,q}(f)$ is trivially normal in $G_{p,q}$. The rest follows immediately from (i) and (iii).
- (vi) This follows from (iii), (v), and the Second Isomorphism Theorem [\[16,](#page-23-4) Thm. 2.74].
- (vii) The first isomorphism in [\(11\)](#page-3-0) follows from the fact that $G'_{p,q} \lhd G_{p,q}$ and the Third Isomorphism Theorem [\[16,](#page-23-4) Thm. 2.75]. The second isomorphism in [\(11\)](#page-3-0) follows from (ii), (iii), and the Second Isomorphism Theorem [\[16,](#page-23-4) Thm. 2.74]. The fact that the transversal of $T_{p,q}(f)$ in $G_{p,q}(f)$ spans K over R modulo f has been verified with CLIFFORD in Clifford algebras $C\ell_{p,q}$ for $p+q \leq 9$.
- (viii) The fact that the transversal of $G_{p,q}(f)$ in $G_{p,q}$ spans S over K modulo f has been verified with CLIFFORD in Clifford algebras $C\ell_{p,q}$ for $p+q \leq 9$.
- (ix) The isomorphism in [\(13\)](#page-3-1) follows from (iii) and the Third Isomorphism Theorem. The remaining statement has been verified with CLIFFORD in Clifford algebras $Cl_{p,q}$ for $p+q \leq 9$.

(x) This follows from basic group theory and definitions of centralizers.

 \Box

Corollary 1. We have two normal series in $G_{p,q}$:

$$
G_{p,q} \ge G_{p,q}(f) \ge T_{p,q}(f) \ge G'_{p,q} \ge \{1\},\tag{16}
$$

$$
G_{p,q} \ge G_{p,q}(f) \ge K_{p,q}(f) \ge G'_{p,q} \ge \{1\}.
$$
\n(17)

Recall that in CLIFFORD information about each Clifford algebra $C\ell_{p,q}$ for $p+q \leq 9$ is stored in a built-in data file. This information can be retrieved in the form of a sevenelement list with a command clidata([p,q]). For example, for $C\ell_{3,0}$ we find:

$$
\mathtt{data} = [complex, 2, simple, \frac{1}{2}\mathrm{Id} + \frac{1}{2}\mathbf{e}_1, [\mathrm{Id}, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_{23}], [\mathrm{Id}, \mathbf{e}_2], [\mathrm{Id}, \mathbf{e}_2]] \tag{18}
$$

where Id denotes the identity element of the algebra. In particular, from the above we find that: (i) $Cl_{3,0}$ is a simple algebra isomorphic to $Mat(2,\mathbb{C})$; (data[1], data[2], data[3]) (ii) Expression $\frac{1}{2}$ Id + $\frac{1}{2}$ **e**₁ (data[4]) is a primitive idempotent f which may be used to generate a spinor ideal $S = C\ell_{3,0}f$; (iii) The fifth entry data^[5] provides, modulo f, a real basis for S, that is, $S = \text{span}_{\mathbb{R}}\{f, e_2f, e_3f, e_{23}f\}$; (iv) The sixth entry data [6] provides, modulo f, a real basis for $\mathbb{K} = fC\ell_{3,0}f \cong \mathbb{C}$, that is, $\mathbb{K} = \text{span}_{\mathbb{R}}\{f, e_{23}f\}$; and, (v) The seventh entry data[7] provides, modulo f, a basis for S over K, that is, $S = \text{span}_{\mathbb{K}}\{f, e_2f\}$.

Thus, based on the above theorem, we have the following corollary:

Corollary 2. Let data be the list of data returned by the procedure clidata in CLIFFORD. Then, data[5] is a transversal of $T_{p,q}(f)$ in $G_{p,q}$; data[6] is a transversal of $T_{p,q}(f)$ in $G_{p,q}(f);$ and $\tt{data}[7]$ is a transversal of $G_{p,q}(f)$ in $G_{p,q}.$ Therefore, $|\tt{data}[5]|=$ $|\texttt{data}[6]| |\texttt{data}[7]|$. This is equivalent to $\frac{G_{p,q}}{T_{p,q}}$ $\left|\frac{G_{p,q}}{T_{p,q}(f)}\right|=\left|\frac{G_{p,q}(f)}{T_{p,q}(f)}\right|$ $\frac{G_{p,q}(f)}{T_{p,q}(f)}\Big|\Big|\frac{G_{p,q}}{G_{p,q}(f)}\Big|.$

Proof. From Thm. [1](#page-3-2) part (ix) we find that data [5] is a transversal of $T_{p,q}(f)$ in $G_{p,q}$; from part (vii) we find that $data[6]$ is a transversal of $T_{p,q}(f)$ in $G_{p,q}(f)$; and from part (viii) we find that $data[7]$ is a transversal of $G_{p,q}(f)$ in $G_{p,q}$. Therefore, the relation $|\text{data}[5]| = |\text{data}[6]| |\text{data}[7]|$ follows from [\(13\)](#page-3-1) and Lagrange's Theorem.

Remark. Before we proceed, we summarize first some definitions cf. [\[12,](#page-22-2) Sect. 17.2]:

- $G_{p,q}^{\varepsilon} = \{ g \in \mathbb{C}\ell_{p,q} \mid T_{\varepsilon}^{\varepsilon}(g)g = 1 \}$ 'Signed vee group' $G_{p,q}^{\varepsilon}$ defined in [\(2\)](#page-1-4).
- $G_{p,q} = \{m \in \mathcal{B} \mid \pm m\}$ Salingaros' vee group.
- $\text{Lip}(p,q) = \{s \in Cl_{p,q} \mid \forall x \in \mathbb{R}^{p,q}, sxs^{-1} \in \mathbb{R}^{p,q}\}\$ Lipschitz group.
- $\text{Pin}(p,q) = \{s \in \text{Lip}(p,q) \mid s\tilde{s} = \pm 1\} < \text{Lip}(p,q)$ Pin group.
- $\text{Spin}(p,q) = \text{Pin}(p,q) \cap C\ell_{p,q}^+ < \text{Pin}(p,q) < \text{Lip}(p,q) \text{Spin} \text{ group}.$
- $\text{Spin}_{+}(p,q) = \{s \in \text{Spin}(p,q) \mid s\tilde{s} = 1\}$ for $p \neq 0$ and $q \neq 0$, orthochronous isometries.

Now we collect a few facts relating the above defined groups to the groups commonly used. These facts are easy to prove and some are known:

• $G_{p,q} < G_{p,q}^{\varepsilon}$ because $\forall m \in G_{p,q}$ we have $T_{\varepsilon}^{\sim}(m) = m^{-1}$ or $T_{\varepsilon}^{\sim}(m)m = 1$.

 4 See [\[3,](#page-22-5) [4\]](#page-22-6) how to use CLIFFORD.

• $\forall m \in G_{p,q}$ we have $T_{\varepsilon}^{\sim}(m)m=1$ where $T_{\varepsilon}^{\sim}=t_{\varepsilon} \circ \tilde{\ }$. Thus, $\tilde{\ }=T_{\varepsilon} \circ t_{\varepsilon}$ (for t_{ε} see [\[1,](#page-22-0) Disp. (21)] and so $\forall m \in G_{p,q}$ we have $m\tilde{m} = mT_{\varepsilon}^{\prime}(t_{\varepsilon}(m)) = \pm 1$. Therefore,

$$
G_{p,q} < \mathbf{Pin}(p,q) < \mathbf{Lip}(p,q),
$$

thus $G_{p,q}$ is a discrete subgroup of $\text{Pin}(p,q)$. Together with $G_{p,q} < G_{p,q}^{\varepsilon}$, we get

$$
G_{p,q} < \mathbf{Pin}(p,q) \cap G_{p,q}^{\varepsilon}.
$$

Thus, the above connects $G_{p,q}$, $\text{Pin}(p,q)$ and $G_{p,q}^{\varepsilon}$.

We illustrate the above Thm. [1](#page-3-2) with the following three examples.

Example 1. We consider $C\ell_{1,1} \cong \text{Mat}(2,\mathbb{R})$. In CLIFFORD, we find the following data for $C\ell_{1,1}$:

$$
\mathtt{data} = [real, 2, simple, \frac{1}{2}\mathrm{Id} + \frac{1}{2}\mathbf{e}_{12}, [\mathrm{Id}, \mathbf{e}_1], [\mathrm{Id}], [\mathrm{Id}, \mathbf{e}_1]]. \tag{19}
$$

Let $f=\frac{1}{2}$ $\frac{1}{2}(1 + \mathbf{e}_{12})$. Then, the groups are as follows:

$$
G_{1,1} = {\pm 1, \pm \mathbf{e}_1, \pm \mathbf{e}_2, \pm \mathbf{e}_{12}},
$$
\n(20)

$$
G_{1,1}(f) = {\pm 1, \pm \mathbf{e}_{12}}, \tag{21}
$$

$$
T_{1,1}(f) = {\pm 1, \pm \mathbf{e}_{12}}, \tag{22}
$$

$$
K_{1,1}(f) = \{\pm 1\}.\tag{23}
$$

Thus, from (20) , (21) , (22) the quotient groups and their transversals^{[5](#page-6-3)} are:

$$
G_{1,1}/G_{1,1}(f) = \{G_{1,1}(f), \mathbf{e}_1 G_{1,1}(f)\} \stackrel{\ell}{\to} \{1, \mathbf{e}_1\} = \text{data}[7],\tag{24}
$$

$$
G_{1,1}(f)/T_{1,1}(f) = \{T_{1,1}(f)\} \stackrel{\ell}{\to} \{1\} = \text{data}[6],\tag{25}
$$

$$
G_{1,1}/T_{1,1}(f) = \{T_{1,1}(f), \mathbf{e}_1 T_{1,1}(f)\} \stackrel{\ell}{\to} \{1, \mathbf{e}_1\} = \text{data}[5].
$$
 (26)

Example 2. We consider $C\ell_{1,2} \cong \text{Mat}(2, \mathbb{C})$. In CLIFFORD, we find the following data for $C\ell_{1,2}$:

$$
\mathtt{data} = [complex, 2, simple, \frac{1}{2}\mathrm{Id} + \frac{1}{2}\mathbf{e}_{13}, [\mathrm{Id}, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_{12}], [\mathrm{Id}, \mathbf{e}_2], [\mathrm{Id}, \mathbf{e}_1]]. \tag{27}
$$

Let $f=\frac{1}{2}$ $\frac{1}{2}(1 + e_{13})$. Then, the groups are as follows:

$$
G_{1,2} = {\pm 1, \pm \mathbf{e}_1, \pm \mathbf{e}_2, \pm \mathbf{e}_3, \pm \mathbf{e}_{12}, \pm \mathbf{e}_{13}, \pm \mathbf{e}_{23}, \pm \mathbf{e}_{123}},
$$
\n(28)

$$
G_{1,2}(f) = {\pm 1, \pm \mathbf{e}_2, \pm \mathbf{e}_{13}, \pm \mathbf{e}_{123}},
$$
\n(29)

$$
T_{1,2}(f) = {\pm 1, \pm \mathbf{e}_{13}},\tag{30}
$$

$$
K_{1,2}(f) = \{\pm 1, \pm \mathbf{e}_2\}.
$$
\n(31)

⁵Note that transversal sets, which just provide coset representatives, are not unique: For the given (left) coset aK of K, a normal subgroup of G, we may assign $\ell(aK) = a$ or $\ell(aK) = b$ as long as $aK = bK$ or, equivalently, $ab^{-1} \in K$.

Thus, the quotient groups and their transversals are:

$$
G_{1,2}/G_{1,2}(f) = \{G_{1,2}(f), \mathbf{e}_1 G_{1,2}(f)\} \stackrel{\ell}{\to} \{1, \mathbf{e}_1\} = \text{data}[7],\tag{32}
$$

$$
G_{1,2}(f)/T_{1,2}(f) = \{T_{1,2}(f), \mathbf{e}_2 T_{1,2}(f)\} \stackrel{\ell}{\to} \{1, \mathbf{e}_2\} = \text{data}[6],
$$
\n
$$
G_{1,2}/T_{1,2}(f) = \{T_{1,2}(f), \mathbf{e}_1 T_{1,2}(f), \mathbf{e}_2 T_{1,2}(f), \mathbf{e}_{12} T_{1,2}(f)\} \stackrel{\ell}{\to} \tag{33}
$$

$$
\{1,{\bf e}_1,{\bf e}_2,{\bf e}_{12}\}={\tt data}[5].~~(34)
$$

Example 3. We consider $C\ell_{1,3} \cong \text{Mat}(2, \mathbb{H})$. In CLIFFORD, we find the following data for $C\ell_{1,3}$:

data =
$$
[quaternionic, 2, simple, \frac{1}{2}Id + \frac{1}{2}e_{14}, \quad [Id, e_1, e_2, e_3, e_{12}, e_{13}, e_{23}, e_{123}], [Id, e_2, e_3, e_{23}], [Id, e_1]]. \quad (35)
$$

Let $f=\frac{1}{2}$ $\frac{1}{2}(1 + e_{14})$. Then, the groups are as follows:

$$
G_{1,4} = \{\pm 1, \pm \mathbf{e}_1, \pm \mathbf{e}_2, \pm \mathbf{e}_3, \pm \mathbf{e}_4, \pm \mathbf{e}_{12}, \pm \mathbf{e}_{13}, \pm \mathbf{e}_{14}, \pm \mathbf{e}_{23},\pm \mathbf{e}_{24}, \pm \mathbf{e}_{34}, \pm \mathbf{e}_{123}, \pm \mathbf{e}_{124}, \pm \mathbf{e}_{134}, \pm \mathbf{e}_{1234}\},
$$
 (36)

$$
G_{1,4}(f) = \{ \pm 1, \pm \mathbf{e}_2, \pm \mathbf{e}_3, \pm \mathbf{e}_{14}, \pm \mathbf{e}_{23}, \pm \mathbf{e}_{124}, \pm \mathbf{e}_{134}, \pm \mathbf{e}_{1234} \},
$$
(37)

$$
T_{1,4}(f) = {\pm 1, \pm \mathbf{e}_{14}}, \tag{38}
$$

$$
K_{1,4}(f) = \{ \pm 1, \pm \mathbf{e}_2, \pm \mathbf{e}_3, \pm \mathbf{e}_{23} \}.
$$
 (39)

Thus, the quotient groups and their transversals are:

$$
G_{1,4}/G_{1,4}(f) = \{G_{1,4}(f), \mathbf{e}_1 G_{1,4}(f)\} \stackrel{\ell}{\to} \{1, \mathbf{e}_1\} = \text{data}[7],\tag{40}
$$

$$
G_{1,4}(f)/T_{1,4}(f) = \{T_{1,4}(f), \mathbf{e}_2 T_{1,4}(f), \mathbf{e}_3 T_{1,4}(f), \mathbf{e}_{23} T_{1,4}(f)\} \xrightarrow{\ell} \{1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_{23}\} = \text{data}[6],
$$
\n
$$
G_{1,4}/T_{1,4}(f) = \{T_{1,4}(f), \mathbf{e}_1 T_{1,4}(f), \mathbf{e}_2 T_{1,4}(f), \mathbf{e}_3 T_{1,4}(f),
$$
\n
$$
\mathbf{e}_{12} T_{1,4}(f), \mathbf{e}_{13} T_{1,4}(f), \mathbf{e}_{23} T_{1,4}(f), \mathbf{e}_{123} T_{1,4}(f)\} \xrightarrow{\ell} \{1, \mathbf{e}_1 T_{1,4}(f), \mathbf{e}_2 T_{1,4}(f), \mathbf{e}_3 T_{1,4}(f)\} \xrightarrow{\ell} \{1, \mathbf{e}_2 T_{1,4}(f), \mathbf{e}_3 T_{1,4}(f), \mathbf{e}_4 T_{1,4}(f)\}.
$$
\n(41)

$$
\{1, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_{12}, \mathbf{e}_{13}, \mathbf{e}_{23}, \mathbf{e}_{123}\} = \text{data}[5]. \tag{42}
$$

The transversal of $G_{p,q}(f)$ in $G_{p,q}$ and the transversal of $T_{p,q}(f)$ in $G_{p,q}(f)$ are listed in Tables 1–4 in [\[4,](#page-22-6) Appendix A]. The stabilizer groups $G_{p,q}(f)$ are listed in Tables 1–5 in [\[2,](#page-22-1) Appendix A].

3. Transposition scalar product on spinor spaces

In [\[12,](#page-22-2) Ch. 18], Lounesto discusses scalar products on $S = \mathbb{C}\ell_{p,q}f$ for simple Clifford algebras and on $\check{S} = S \oplus \hat{S} = C\ell_{p,q}e$, $e = f + \hat{f}$, for semisimple Clifford algebras. Recall that \hat{f} denotes the grade involution of f. It is well known that in each case the spinor representation is faithful. Following Lounesto, we let

 $\check{\mathbb{K}}$ be either \mathbb{K} or $\mathbb{K} \oplus \hat{\mathbb{K}}$. \check{S} be either S or $S \oplus \hat{S}$

when $C\ell_{p,q}$ is simple or semisimple, respectively. Then, in the simple algebras, the two scalar products are

$$
S \times S \to \mathbb{K}, \quad (\psi, \phi) \mapsto \begin{cases} \beta_+(\psi, \phi) = s_1 \tilde{\psi} \phi \\ \beta_-(\psi, \phi) = s_2 \bar{\psi} \phi \end{cases}
$$
(43)

whereas in the semisimple algebras they are

$$
\check{S} \times \check{S} \to \check{\mathbb{K}}, \quad (\check{\psi}, \check{\phi}) \mapsto \begin{cases} (\beta_+(\psi, \phi), \beta_+(\psi_g, \phi_g)) = (s_1 \check{\psi} \phi, s_1 \check{\psi}_g \phi_g) \\ (\beta_-(\psi, \phi), \beta_-(\psi_g, \phi_g)) = (s_2 \check{\psi} \phi, s_2 \check{\psi}_g \phi_g) \end{cases}
$$
(44)

when $\check{\psi} = \psi + \psi_g$ and $\check{\phi} = \phi + \phi_g$, $\psi, \phi \in S$, $\psi_g, \phi_g \in \hat{S}$, and where $\tilde{\psi}, \tilde{\psi}_g$ (resp. $\bar{\psi}, \bar{\psi}_g$) denotes reversion (resp. Clifford conjugation) of ψ, ψ_q and s_1, s_2 are special monomials in the Clifford algebra basis which guarantee that the products $s_1\tilde{\psi}\phi$, $s_2\bar{\psi}\phi$, hence also $s_1\tilde{\psi}_g\phi_g$, $s_2\bar{\psi}_g\phi_g$, belong to K ≅ \mathbb{R}^6 \mathbb{R}^6 . The automorphism groups of β_+ and β_- are defined in the simple case as, respectively, $G_+ = \{s \in \mathbb{C}\ell_{p,q} \mid s\tilde{s} = 1\}$ and $G_- = \{s \in \mathbb{C}\ell_{p,q} \mid s\bar{s} = 1\},$ and as ${}^{2}G_{-}$ and ${}^{2}G_{+}$ in the semisimple case. They are shown in [\[12,](#page-22-2) Tables 1 and 2, p. 236].

3.1. Simple Clifford algebras. In Example 3 in [\[2\]](#page-22-1) it was shown that the transposition scalar product in $S = C\ell_{2,2}f$ is different from each of the two Lounesto's products whereas Example 4 showed that the transposition product in $S = C\ell_{3,0}f$ coincided with β_+ . Furthermore, it was remarked that $T_{\varepsilon}(\psi)\phi$ always equaled β_{+} for Euclidean signatures $(p,0)$ and β for anti-Euclidean signatures $(0, q)$. We formalize this in the following.

Proposition 1. Let $\psi, \phi \in S = \mathcal{C}\ell_{p,q}f$ and $(\psi, \phi) \mapsto T_{\varepsilon}(\psi)\phi = \lambda f, \lambda \in \mathbb{K}$, be the transposition scalar product. Let β_+ and β_- be the scalar products on S shown in [\(43\)](#page-8-1). Then, there exist monomials s_1, s_2 in the transversal ℓ of $G_{p,q}(f)$ in $G_{p,q}$ such that

$$
T_{\varepsilon}(\psi)\phi = \begin{cases} \beta_{+}(\psi,\phi) = s_{1}\tilde{\psi}\phi, & \forall\psi,\phi \in C\ell_{p,0}f, \\ \beta_{-}(\psi,\phi) = s_{2}\bar{\psi}\phi, & \forall\psi,\phi \in C\ell_{0,q}f. \end{cases}
$$
(45)

Proof. In [\[1,](#page-22-0) Cor. 1] it was shown that T_{ε} reduces to reversion (resp. conjugation) antiinvolution in $Cl_{p,0}$ for the Euclidean signature $(p,0)$ (resp. for the anti-Euclidean signature $(0, q)$).

⁶In simple Clifford algebras, the monomials s_1 and s_2 also satisfy: (i) $\tilde{f} = s_1 fs_1^{-1}$ and (ii) $\bar{f} = s_2 fs_2^{-1}$. The identity (i) (resp. (ii)) is also valid in the semisimple algebras provided $\beta_+ \not\equiv 0$ (resp. $\beta_- \not\equiv 0$).

Table 1 (Part 1): Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in simple Clifford algebras $C\ell_{p,q} \cong \mathrm{Mat}(2^k,{\mathbb R})$

		$\mathbf{u} = \mathbf{u} - \mathbf{u}$		
		(p,q) $(0,0)$ $(1,1)$ $(2,0)$ $(2,2)$ $(3,1)$ $(3,3)$ $(0,6)$		
				$G_{p,q}$ $O(1)$ $O(2)$ $\overline{O(2)}$ $O(4)$ $O(4)$ $O(8)$ $\overline{O(8)}$

 $k = q - r_{q-p}, p - q \neq 1 \mod 4, p - q = 0, 1, 2 \mod 8$

Table 1 (Part 2): Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in simple Clifford algebras $C\ell_{p,q} \cong \mathrm{Mat}(2^k, \mathbb{R})$

$k = q - r_{q-p}, p - q \neq 1 \mod 4, p - q = 0, 1, 2 \mod 8$									
				(p,q) $(4,2)$ $(5,3)$ $(1,7)$ $(0,8)$	$(4,4)$ $(8,0)$				
				$G_{p,q}$ $O(8)$ $O(16)$ $O(16)$ $\overline{O(16)}$ $O(16)$ $\overline{O(16)}$					

Let $u \in \mathcal{C}\ell_{p,q}$ and let $[u]$ be a matrix of u in the spinor representation π_S of $\mathcal{C}\ell_{p,q}$ realized in the spinor $(C\ell_{p,q}, \mathbb{K})$ -bimodule ${}_{C\ell}S_{\mathbb{K}} \cong C\ell_{p,q}f\mathbb{K}$. Then, by [\[2,](#page-22-1) Prop. 5],

$$
[T_{\varepsilon}^{\sim}(u)] = \begin{cases} [u]^T & \text{if } p - q = 0, 1, 2 \text{ mod } 8; \\ [u]^{\dagger} & \text{if } p - q = 3, 7 \text{ mod } 8; \\ [u]^{\dagger} & \text{if } p - q = 4, 5, 6 \text{ mod } 8; \end{cases}
$$
(46)

where T , \dagger , and \dagger denote, respectively, transposition, complex Hermitian conjugation, and quaternionic Hermitian conjugation. Thus, we immediately have:

Proposition 2. Let $G_{p,q}^{\varepsilon} \subset C\ell_{p,q}$ where $C\ell_{p,q}$ is a simple Clifford algebra. Then, $G_{p,q}^{\varepsilon}$ is: The orthogonal group $O(N)$ when $K \cong \mathbb{R}$; the complex unitary group $U(N)$ when $K \cong \mathbb{C}$; or, the compact symplectic group $Sp(N) = U_{\mathbb{H}}(N)$ when $\mathbb{K} \cong \mathbb{H}$.^{[7](#page-9-0)} That is,

$$
G_{p,q}^{\varepsilon} = \begin{cases} O(N) & \text{if } p - q = 0, 1, 2 \text{ mod } 8; \\ U(N) & \text{if } p - q = 3, 7 \text{ mod } 8; \\ Sp(N) & \text{if } p - q = 4, 5, 6 \text{ mod } 8; \end{cases}
$$
(47)

where $N = 2^k$ and $k = q - r_{q-p}$.

Proof. The isomorphism $C\ell_{p,q} \cong \text{Mat}(N,\mathbb{K})$, where $\mathbb{K} \cong fC\ell_{p,q}f$ for a primitive idempotent f, implies that for every $u \in G_{p,q}^{\varepsilon}$ the matrix $[T_{\varepsilon}^{\varepsilon}(u)u] = [T_{\varepsilon}^{\varepsilon}(u)][u] = 1$. Then, [\(46\)](#page-9-1) implies that $[u]$ ^T[u] = 1, [u][†][u] = 1, or [u][‡][u] = 1 depending on the value of p−q mod 8. $□$

The scalar product $T_{\varepsilon}(\psi)\phi$ was computed with CLIFFORD for all signatures (p, q) , $p+q \leq$ 9. Observe that as expected, in Euclidean (resp. anti-Euclidean) signatures $(p, 0)$ (resp.

⁷See Fulton and Harris [\[7\]](#page-22-7) for a definition of the quaternionic unitary group $U_{\mathbb{H}}(N)$. In our notation we follow loc. cit. page 100, 'Remark on Notations'.

ON THE TRANSPOSITION ANTI-INVOLUTION III 11

Table 2 (Part 1): Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in simple Clifford algebras $C\ell_{p,q} \cong \mathrm{Mat}(2^k,\mathbb{C})$

		$\frac{1}{2}$ $\frac{1}{4}$ $\frac{1}{2}$		
		$ (p,q) (0,1) (1,2) (3,0) (2,3) (0,5) (4,1) (1,6) (7,0)$		
		$\overline{G_{p,q}}$ $U(1)$ $U(2)$ $U(2)$ $U(4)$ $U(4)$ $U(4)$ $U(5)$ $U(8)$ $U(8)$		

 $k = q - r_{q-n}, p-q \neq 1 \mod 4, p-q = 3, 7 \mod 8$

Table 2 (Part 2): Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in simple Clifford algebras $C\ell_{p,q} \cong \mathrm{Mat}(2^k,\mathbb{C})$

$k = q - r_{q-p}, p - q \neq 1 \mod 4, p - q = 3, 7 \mod 8$									
						$ (p,q) (5,2) (3,4) (4,5) (6,3) (2,7) (0,9) (8,1)$			
						$\begin{array}{ c c c c c c c c } \hline G_{p,q} & U(8) & U(8) & U(16) & U(16) & U(16) & U(16) \hline \end{array}$			

Table 3 (Part 1): Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in simple Clifford algebras $C\ell_{p,q} \cong \mathrm{Mat}(2^k,\mathbb{H})$

	\mathbf{u} \mathbf{v} \mathbf{u}				
(p,q)	(0, 2)	(0,4)		$(4,0)$ $(1,3)$ $(2,4)$ $(6,0)$	
	$G_{p,q}$ $\left \overline{\left \overline{Sp(1)} \right } \right \left \overline{\left \overline{Sp(2)} \right } \right \left \overline{Sp(2)} \right $ $Sp(2)$ $\overline{\left \overline{Sp(4)} \right }$				

 $k = q - r_{q-n}, p-q \neq 1 \mod 4, p-q = 4, 5, 6 \mod 8$

Table 3 (Part 2): Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in simple Clifford algebras $C\ell_{p,q} \cong \mathrm{Mat}(2^k,\mathbb{H})$

	$\kappa - q - r_{q-p}, p - q + 1$ mod 4, $p - q = 4, 0, 0$ mod 8									
(p,q)	(1, 5)									
$G_{p,q}$		$Sp(4)$ $Sp(4)$ $Sp(8)$ $Sp(8)$ $Sp(8)$ $Sp(8)$								

 $(0, q)$) the group $G_{p,0}^{\varepsilon}$ (resp. $G_{0,q}^{\varepsilon}$) coincides with the corresponding automorphism group of the scalar product β_+ (resp. β_-) listed in [\[12,](#page-22-2) Table 1, p. 236] (resp. [12, Table 2, p. 236]. This is indicated by a single (resp. double) box around the group symbol in Tables 4 and 5. For example, for the Euclidean signatures $(5,0)$, we show $G_{5,0}^{\varepsilon}$ as $\left[{}^{2}Sp(2)\right]$ whereas for the anti-Euclidean signatures $(0, 7)$, we show $G_{0,7}^{\varepsilon}$ as $||^{2}O(8)||$.

For simple Clifford algebras, the automorphism groups $\overline{G_{p,q}^{\varepsilon}}$ are displayed in Tables 1, 2, and 3 (for semisimple algebras, see Tables 4 and 5). In each case the form is positive definite and non-degenerate. Also, unlike in the case of the forms β_+ and β_- , there is

no need for the extra monomial factor like s_1, s_2 in [\(43\)](#page-8-1) (and [\(44\)](#page-8-2)) to guarantee that the product $T_{\varepsilon}(\psi)\phi$ belongs to K since this is always the case [\[1,](#page-22-0) [2\]](#page-22-1). Recall that the only role of the monomials s_1 and s_2 is to permute entries of spinors $\psi\phi$ and $\psi\phi$ to assure that $\beta_+(\psi, \phi)$ and $\beta_-(\psi, \phi)$ belong to the (skew) field K. That is, more precisely, that $\beta_+(\psi, \phi)$ and $\beta_-(\psi, \phi)$ have the form $\lambda f = f\lambda$ for some λ in K. The idempotent f in the spinor basis in S corresponds uniquely to the identity coset $G_{p,q}(f)$ in the quotient group $G_{p,q}/G_{p,q}(f)$. Based on [\[2,](#page-22-1) Prop. 2] we know that since the vee group $G_{p,q}$ permutes entries of any spinor ψ , the monomials s_1 and s_2 belong to the transversal of the stabilizer $G_{p,q}(f) \triangleleft G_{p,q}$ [\[2,](#page-22-1) Cor. 2].^{[8](#page-11-0)}

One more difference between the scalar products β_+ and β_- , and the transposition product $T_{\varepsilon}(\psi)\phi$ is that in some signatures one of the former products may be identically zero whereas the transposition product is never identically zero. The signatures (p, q) in which one of the products β_+ or β_- is identically zero can be easily found in [\[12,](#page-22-2) Tables 1] and 2, p. 236] as the automorphism group of the product is then a general linear group.

We summarize our findings. Let $T_{\varepsilon}(\psi)\phi$ be the scalar product on the right K-linear spinor space S in a simple Clifford algebra $Cl_{p,q}$.

• In the real case $(p - q = 0, 1, 2 \mod 8)$ when $\dim_{\mathbb{R}} S = N$, the scalar product (modulo f) is just the symmetric non-degenerate and positive definite form

$$
T_{\varepsilon}(\psi)\phi = T_{\varepsilon}(\phi)\psi = \psi_1\phi_1 + \ldots + \psi_N\phi_N.
$$
 (48)

• In the complex case $(p - q = 4, 5, 6 \mod 8)$ when $\dim_{\mathbb{C}} S = N$, the scalar product (modulo f) is just the standard (complex) Hermitian non-degenerate and positive definite form

$$
T_{\varepsilon}(\psi)\phi = \overline{T_{\varepsilon}(\phi)\psi} = \overline{\psi}_1\phi_1 + \ldots + \overline{\psi}_N\phi_N
$$
 (49)

where $\overline{T_{\varepsilon}(\phi)\psi}$ denotes "complex" conjugation in K.

• In the quaternionic case $(p-q=3, 7 \mod 8)$ when $\dim_{\mathbb{H}} S = N$, the scalar product (modulo f) is just the standard (quaternionic) Hermitian (or, "symplectic scalar product") which is also non-degenerate and positive definite.^{[9](#page-11-1)}

$$
T_{\varepsilon}(\psi)\phi = \overline{T_{\varepsilon}(\phi)\psi} = \overline{\psi}_1\phi_1 + \ldots + \overline{\psi}_N\phi_N
$$
 (50)

where $\overline{T_{\varepsilon}(\phi)\psi}$ denotes "quaternionic" conjugation in K.

Here we give a few low-dimensional examples of the transposition scalar product $T_{\varepsilon}(\psi)\phi$.

Example 4. Consider $C\ell_{2,2} \cong \text{Mat}(4,\mathbb{R})$ with $f = \frac{1}{4}$ $\frac{1}{4}(1+\mathbf{e}_{13})(1+\mathbf{e}_{24})$. Then, $\mathbb{K} = fC\ell_{2,2}f \cong$ $\text{span}_{\mathbb{R}}\{1\} \cong \mathbb{R}$ and a transversal of the stabilizer $G_{2,2}^{(1)}(f) \triangleleft G_{2,2}$ is $\mathcal{M} = \{1, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_{12}\}.$ Let

 ${}^{8}\text{In}$ [\[12,](#page-22-2) Page 233], Lounesto states correctly that "the element s can be chosen from the standard basis of $C\ell_{p,q}$." In fact, one can restrict the search for s to the transversal of the stabilizer $G_{p,q}(f)$ in $G_{p,q}$ which has a much smaller size $2^{q-r_{q-p}}$ compared to the size 2^{p+q} of the Clifford basis.

⁹An R-bilinear form $K: V \times V \to \mathbb{K}$ on a real, complex, or quaternionic vector space V is non-degenerate if whenever $K(v, w) = 0, \forall v$ then $w = 0$. The form is *positive definite* if $K(v, v) > 0$ for $v \neq 0$ [\[7\]](#page-22-7).

 $\psi, \phi \in S = C\ell_{2,2}f$. Hence,

$$
\psi = \psi_1 f + \psi_2 \mathbf{e}_1 f + \psi_3 \mathbf{e}_2 f + \psi_4 \mathbf{e}_{12} f, \tag{51}
$$

$$
\phi = \phi_1 f + \phi_2 \mathbf{e}_1 f + \phi_3 \mathbf{e}_2 f + \phi_4 \mathbf{e}_{12} f,\tag{52}
$$

where $\psi_i, \phi_i \in \mathbb{R}, i = 1, \ldots, 4$. It is then easy to check that the form

$$
T_{\varepsilon}(\psi)\phi = (\psi_1\phi_1 + \psi_2\phi_2 + \psi_3\phi_3 + \psi_4\phi_4)f
$$
\n
$$
(53)
$$

is invariant under $O(4)$. Of course, in spinor representation, we have

$$
[T_{\varepsilon}(\psi)] = \begin{bmatrix} \psi_1 & \psi_2 & \psi_3 & \psi_4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ and } [\phi] = \begin{bmatrix} \phi_1 & 0 & 0 & 0 \\ \phi_2 & 0 & 0 & 0 \\ \phi_3 & 0 & 0 & 0 \\ \phi_4 & 0 & 0 & 0 \end{bmatrix}.
$$
 (54)

Notice that the scalar products β_+ and β_- modulo f are:

$$
\beta_{+}(\psi,\phi) = s_{1}\tilde{\psi}\phi = (\psi_{3}\phi_{2} - \psi_{2}\phi_{3} + \psi_{4}\phi_{1} - \psi_{1}\phi_{4})f
$$
\n
$$
= tr([\psi_{1} \quad \psi_{2} \quad \psi_{3} \quad \psi_{4}] \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \phi_{1} \\ \phi_{2} \\ \phi_{3} \\ \phi_{4} \end{bmatrix})f
$$
\n(55)

$$
\beta_{-}(\psi,\phi) = s_{2}\bar{\psi}\phi = (\psi_{4}\phi_{1} + \psi_{2}\phi_{3} - \psi_{1}\phi_{4} - \psi_{3}\phi_{2})f
$$

= tr([ψ_{1} ψ_{2} ψ_{3} ψ_{4}]]
$$
\begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \phi_{1} \\ \phi_{2} \\ \phi_{3} \\ \phi_{4} \end{bmatrix} f.
$$
 (56)

Each product is invariant under $Sp(4, \mathbb{R})$. (cf. [\[12,](#page-22-2) Tables 1 and 2, p. 236]) The monomials $s_1 = s_2 = e_{12}$ belong to M.^{[10](#page-12-0)} For example, the monomial s_1 permutes entries in the first column as seen below for β_{+} :

$$
[\tilde{\psi}\phi] = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \lambda & 0 & 0 & 0 \end{bmatrix} \quad \text{whereas} \quad [s_1\tilde{\psi}\phi] = \begin{bmatrix} -\lambda & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}
$$
 (57)

where $\lambda = \psi_2 \phi_3 - \psi_3 \phi_2 + \psi_1 \phi_4 - \psi_4 \phi_1 \in \mathbb{R}$. A similar role is played by the monomial s_2 in $\beta_-\$.

The above example shows that, in general, $\tilde{\psi}$ and $\bar{\psi}$, do not belong to the dual S^* of the spinor space S. On the other hand, $T_{\varepsilon}(\psi)$ always belongs to S^* for every $\psi \in S$ [\[1,](#page-22-0) [2\]](#page-22-1). We have classified all automorphism groups of the transposition scalar product for real simple Clifford algebras. The results are summarized in Table 1.

¹⁰The choice of each of the monomials s_1 and s_2 is of course not unique. In general, there are $2^{p+q}/N$ such choices modulo the commutator subgroup $\{\pm 1\}$ of $G_{p,q}$.

Example 5. Consider $C\ell_{1,2} \cong \text{Mat}(2,\mathbb{C})$ with $f = \frac{1}{2}$ $\frac{1}{2}(1 + e_{13})$. Then, $\mathbb{K} = fC\ell_{1,2}f \cong$ span_R{1, e₂} $\cong \mathbb{C}$ and a transversal of the stabilizer $\bar{G}_{1,2}(f) \triangleleft G_{1,2}$ is $\mathcal{M} = \{1, e_1\}$. Let $\psi, \phi \in S = C\ell_{1,2}f$. Recalling that the left minimal ideal S is a right K-module, we write "complex" coefficients on the right:

$$
\psi = f\psi_1 + \mathbf{e}_1 f \psi_2 = f(\psi_{11} + \psi_{12} \mathbf{e}_2) + \mathbf{e}_1 f(\psi_{21} + \psi_{22} \mathbf{e}_2), \tag{58}
$$

$$
\phi = f\phi_1 + \mathbf{e}_1 f\phi_2 = f(\phi_{11} + \phi_{12}\mathbf{e}_2) + \mathbf{e}_1 f(\phi_{21} + \phi_{22}\mathbf{e}_2),
$$
\n(59)

where $\psi_{ij}, \phi_{ij} \in \mathbb{R}, \psi_i, \phi_i \in \mathbb{K}, i, j = 1, 2$. It is then easy to check that

$$
T_{\varepsilon}(\psi)\phi = \overline{\psi}_1\phi_1 + \overline{\psi}_2\phi_2 = (\psi_{11}\phi_{11} + \psi_{22}\phi_{22} + \psi_{21}\phi_{21} + \psi_{12}\phi_{12})f +
$$

$$
(\psi_{21}\phi_{22} - \psi_{22}\phi_{21} + \psi_{11}\phi_{12} - \psi_{12}\phi_{11})\mathbf{e}_2f.
$$
 (60)

Thus, $T_{\varepsilon}(\psi)\phi$ is invariant under the complex unitary group $U(2)$. Then,

$$
\beta_{+}(\psi,\phi) = s_{1}\tilde{\psi}\phi = (\psi_{11}\phi_{21} + \psi_{22}\phi_{12} + \psi_{21}\phi_{11} + \psi_{12}\phi_{22})f +
$$

$$
(\psi_{21}\phi_{12} - \psi_{22}\phi_{11} + \psi_{11}\phi_{22} - \psi_{12}\phi_{21})\mathbf{e}_{2}f
$$

$$
= (\overline{\psi}_{1}\phi_{2} + \overline{\psi}_{2}\phi_{1})f = \text{tr}([\overline{\psi}_{1} \quad \overline{\psi}_{2}] \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \phi_{1} \\ \phi_{2} \end{bmatrix})f
$$
(61)

where $s_1 = e_1 \in \mathcal{M}$ and ψ_1, ψ_2 denotes complex conjugation. Since the matrix is similar to a diagonal matrix diag $(1, -1)$, the automorphism group of this form is the complex unitary group $U(1, 1)$. Then,

$$
\beta_{-}(\psi,\phi) = s_{2}\bar{\psi}\phi = (\psi_{11}\phi_{21} + \psi_{22}\phi_{12} - \psi_{21}\phi_{11} - \psi_{12}\phi_{22})f +
$$

$$
(-\psi_{21}\phi_{12} - \psi_{22}\phi_{11} + \psi_{11}\phi_{22} + \psi_{12}\phi_{21})\mathbf{e}_{2}f
$$

$$
= (\psi_{1}\phi_{2} - \psi_{2}\phi_{1})f = \text{tr}([\psi_{1} \quad \psi_{2}] \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} \phi_{1} \\ \phi_{2} \end{bmatrix})f
$$
(62)

where $s_2 = e_1 \in \mathcal{M}$. Thus, β_- is invariant under $Sp(2, \mathbb{C})$. See also [\[12,](#page-22-2) Tables 1 and 2, p. 236].^{[11](#page-13-0)}

We have classified all automorphism groups of the transposition scalar product for complex simple Clifford algebras. The results are summarized in Table 2. In our next example we consider a quaternionic simple Clifford algebra.

Example 6. Consider $C\ell_{1,3} \cong \text{Mat}(2, \mathbb{H})$ with $f = \frac{1}{2}$ $\frac{1}{2}(1 + e_{14})$. Then, $\mathbb{K} = fC\ell_{1,3}f \cong$ span_R{1, e₂, e₃, e₂₃} $\cong \mathbb{H}$ and a transversal of the stabilizer $G_{1,3}(f) \triangleleft G_{1,3}$ is $\mathcal{M} = \{1, e_1\}.$ Let $\psi, \phi \in S = C\ell_{1,3}f$. Hence,

$$
\psi = f\psi_1 + \mathbf{e}_1 f\psi_2 \quad \text{and} \quad \phi = f\phi_1 + \mathbf{e}_1 f\phi_2 \tag{63}
$$

where

$$
\psi_i = \psi_{i1} + \psi_{i2}\mathbf{e}_2 + \psi_{i3}\mathbf{e}_3 + \psi_{i4}\mathbf{e}_{23} \in \mathbb{K},\tag{64}
$$

$$
\phi_i = \phi_{i1} + \phi_{i2} \mathbf{e}_2 + \phi_{i3} \mathbf{e}_3 + \phi_{i4} \mathbf{e}_{23} \in \mathbb{K}
$$
\n(65)

¹¹For the definitions of $U(1, 1)$ and $Sp(2, \mathbb{C})$, see [\[12,](#page-22-2) Ch. 18].

for $\psi_{ij}, \phi_{ij} \in \mathbb{R}, i = 1, 2; j = 1, ..., 4$. Then we get:

$$
T_{\varepsilon}(\psi,\phi) = (\overline{\psi}_{1}\phi_{1} + \overline{\psi}_{2}\phi_{2})f = \left(\sum_{i=1}^{2} \sum_{j=1}^{4} \psi_{ij}\phi_{ij}\right)f +
$$

\n
$$
(\psi_{[11}\phi_{12]} - \psi_{[13}\phi_{14]} + \psi_{[21}\phi_{22]} - \psi_{[23}\phi_{24]})\mathbf{e}_{2}f +
$$

\n
$$
(\psi_{[11}\phi_{13]} + \psi_{[12}\phi_{14]} + \psi_{[21}\phi_{23]} + \psi_{[22}\phi_{24]})\mathbf{e}_{3}f +
$$

\n
$$
(\psi_{[11}\phi_{14]} - \psi_{[12}\phi_{13]} + \psi_{[21}\phi_{24]} - \psi_{[22}\phi_{23]})\mathbf{e}_{23}f,
$$
\n(66)

which is invariant under $U_{\mathbb{H}}(2) = Sp(2)$. Here, $\overline{\psi}_1, \overline{\psi}_2$ denotes quaternionic conjugation.^{[12](#page-14-0)} The other two scalar products are as follows:

$$
\beta_{+}(\psi,\phi) = s_{1}\tilde{\psi}\phi = (\psi_{(11}\phi_{21)} + \psi_{(12}\phi_{22)} + \psi_{(13}\phi_{23)} + \psi_{(14}\phi_{24)})f +
$$

\n
$$
(\psi_{[11}\phi_{22]} - \psi_{[12}\phi_{21]} - \psi_{[13}\phi_{24]} + \psi_{[14}\phi_{23]})\mathbf{e}_{2}f +
$$

\n
$$
(\psi_{[11}\phi_{23]} + \psi_{[12}\phi_{24]} - \psi_{[13}\phi_{21]} - \psi_{[14}\phi_{22]})\mathbf{e}_{3}f +
$$

\n
$$
(\psi_{[11}\phi_{24]} - \psi_{[12}\phi_{23]} + \psi_{[13}\phi_{22]} + \psi_{[21}\phi_{14]})\mathbf{e}_{23}f
$$

\n
$$
= (\overline{\psi}_{1}\phi_{2} + \overline{\psi}_{2}\phi_{1})f = \text{tr}([\overline{\psi}_{1} \ \overline{\psi}_{2}] \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \phi_{1} \\ \phi_{2} \end{bmatrix})f
$$
 (67)

where $s_1 = e_1 \in \mathcal{M}$. Since the matrix is similar to a diagonal matrix diag(1, -1), the automorphism group of this form is the quaternionic unitary group $U_{1,1} \mathbb{H}$. Then,

$$
\beta_{-}(\psi,\phi) = s_{2}\bar{\psi}\phi = (\psi_{[11}\phi_{21]} - \psi_{[12}\phi_{22]} - \psi_{[13}\phi_{23]} + \psi_{[14}\phi_{24]})f +
$$

\n
$$
(\psi_{[11}\phi_{22]} + \psi_{[12}\phi_{21]} + \psi_{[13}\phi_{24]} + \psi_{[14}\phi_{23]})\mathbf{e}_{2}f +
$$

\n
$$
(\psi_{[11}\phi_{23]} - \psi_{[12}\phi_{24]} + \psi_{[13}\phi_{21]} - \psi_{[14}\phi_{22]})\mathbf{e}_{3}f +
$$

\n
$$
(\psi_{(11}\phi_{24)} + \psi_{(12}\phi_{23)} - \psi_{(13}\phi_{22)} - \psi_{(14}\phi_{21}))\mathbf{e}_{23}f
$$

\n
$$
= (\psi_{1}^{*}\phi_{2} - \psi_{2}^{*}\phi_{1})f = \text{tr}([\psi_{1}^{*} \ \psi_{2}^{*}] \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} \phi_{1} \\ \phi_{2} \end{bmatrix})f
$$
(68)

where $s_2 = e_1 \in \mathcal{M}$ and ψ_1^*, ψ_2^* denotes an anti-involution on $\mathbb{K} \cong \mathbb{H}$ which sends ψ_{i1} + ψ_{i2} **e**₂ + ψ_{i3} **e**₃ + ψ_{i4} **e**₂₃ $\mapsto \psi_{i1} + \psi_{i2}$ **e**₂ + ψ_{i3} **e**₃ - ψ_{i4} **e**₂₃. In [\[12,](#page-22-2) Tables 1 and 2, p. 236] we find that β_+ and β_- scalar products are invariant under $Sp(2, 2)$.^{[13](#page-14-1)}

Remark. If we define quaternions as pairs of complex numbers $\mathbb{H} = \mathbb{C} \oplus \mathbb{C} j$ via the Cayley-Dickson doubling process [\[12,](#page-22-2) Sect. 23.2] where $\mathbb{C} = \mathbb{R} \oplus \mathbb{R}i$, that is, write quaternions as $q = z + wj$ or as $q = (z, w)$ where $z, w \in \mathbb{C}$, then the star anti-involution in [\(68\)](#page-14-2) is an R-linear map of H sending $z + w_j \mapsto z + \bar{w}j$ or $(z, w) \mapsto (z, \bar{w})$ where \bar{w} is the complex conjugate of w. Observe that the R-linear map of H sending $z + wj \mapsto \overline{z} - wj$ or $(z, w) \mapsto (\overline{z}, -w)$ is the quaternionic conjugation of H.

We have classified all automorphism groups of the transposition scalar product for quaternionic simple Clifford algebras. The results are summarized in Table 3.

¹²To shorten display, we set $\psi_{[11}\phi_{12]} = \psi_{11}\phi_{12} - \psi_{12}\phi_{11}$ and $\psi_{(11}\phi_{21)} = \psi_{11}\phi_{21} + \psi_{21}\phi_{11}$, etc.

 $1^3Sp(2,2) = U(2,2) \cap Sp(4,\mathbb{C})$ and $Sp(2,2)/\{\pm 1\} \cong SO_+(4,1)$.

3.2. Semisimple Clifford algebras. Faithful spinor representation of a semisimple Clifford algebra $C\ell_{p,q}$ $(p - q = 1 \mod 4)$ is realized in a left ideal $\check{S} = S \oplus \hat{S} = C\ell_{p,q}e$ where $e = f + \hat{f}$ for any primitive idempotent f. Recall that $\hat{ }$ denotes grade involution. We refer to [\[12,](#page-22-2) pp. 232–236] for some of the concepts. In particular, $S = C\ell_{p,q}f$ and $\hat{S} = C\ell_{p,q}\hat{f}$. Thus, every spinor $\check{\psi} \in \check{S}$ has unique components $\psi \in S$ and $\psi_g \in \hat{S}$. We refer to the elements $\check{\psi} \in \check{S}$ as "spinors" whereas to its components $\psi \in S$ and $\psi_{\sigma} \in \hat{S}$ we refer as $\frac{\pi}{2}$ $\frac{1}{2}$ -spinors".

For the semisimple Clifford algebras $C\ell_{p,q}$, we will view spinors $\check{\psi} \in \check{S} = S \oplus \hat{S}$ as ordered pairs $(\psi, \psi_g) \in S \times \hat{S}$ when $\check{\psi} = \psi + \psi_g$. Likewise, we will view elements $\check{\lambda}$ in the double fields $\check{\mathbb{K}} = \mathbb{K} \oplus \hat{\mathbb{K}}$ as ordered pairs $(\lambda, \lambda_g) \in \mathbb{K} \times \hat{\mathbb{K}}$ when $\check{\lambda} = \lambda + \lambda_g$. As before, $\mathbb{K} = fC\ell_{p,q}f$ while $\hat{\mathbb{K}} = \hat{f} C \ell_{p,q} \hat{f}$. Recall that $\check{\mathbb{K}} \cong {}^2\mathbb{R} \stackrel{\text{def}}{=} \mathbb{R} \oplus \mathbb{R}$ or $\check{\mathbb{K}} \cong {}^2\mathbb{H} \stackrel{\text{def}}{=} \mathbb{H} \oplus \mathbb{H}$ when, respectively, $p - q = 1 \mod 8$, or $p - q = 5 \mod 8$.

In this section we classify automorphism groups of the transposition scalar product

$$
\check{S} \times \check{S} \to \check{\mathbb{K}}, \quad (\check{\psi}, \check{\phi}) \mapsto T_{\varepsilon}(\check{\psi}, \check{\phi}) \stackrel{\text{def}}{=} (T_{\varepsilon}(\psi)\phi, T_{\varepsilon}(\psi_g)\phi_g) \in \check{\mathbb{K}} \tag{69}
$$

when $\check{\psi} = \psi + \psi_q$ and $\check{\phi} = \phi + \phi_q$.

Proposition 3. Let $G_{p,q}^{\varepsilon} \subset C\ell_{p,q}$ where $C\ell_{p,q}$ is a semisimple Clifford algebra. Then, $G_{p,q}^{\varepsilon}$ is: The double orthogonal group ${}^{2}O(N) \stackrel{def}{=} O(N) \times O(N)$ when $\check{\mathbb{K}} \cong {}^{2} \mathbb{R}$ or the double compact symplectic group ${}^{2}Sp(N) \stackrel{def}{=} Sp(N) \times Sp(N)$ when $\mathbb{K} \cong {}^{2} \mathbb{H}.^{14}$ $\mathbb{K} \cong {}^{2} \mathbb{H}.^{14}$ $\mathbb{K} \cong {}^{2} \mathbb{H}.^{14}$ That is,

$$
G_{p,q}^{\varepsilon} = \begin{cases} 2O(N) = O(N) \times O(N) & \text{when } p - q = 1 \text{ mod } 8; \\ 2Sp(N) = Sp(N) \times Sp(N) & \text{when } p - q = 5 \text{ mod } 8; \end{cases}
$$
(70)

where $N = 2^{k-1}$ and $k = q - r_{q-p}$.

Proof. This result follows directly when we recall that $C\ell_{p,q} \cong {}^{2}\text{Mat}(N,\mathbb{R})$ or $C\ell_{p,q} \cong$ ²Mat(N, H) in the semisimple case depending whether $p-q=1 \mod 8$ or $p-q=5 \mod 8$, respectively. Then, apply Proposition [2](#page-9-2) to each non universal component of $C\ell_{p,q}$ while remembering to reduce N by a factor of 2.

The automorphism groups $G_{p,q}^{\varepsilon}$ for semisimple Clifford algebras $C\ell_{p,q}$ for $p+q\leq 9$ are shown in Tables 4 and 5. All results in these tables, like in Tables 1, 2, and 3, have been verified with CLIFFORD.

We show two examples.

Example 7. Consider $C\ell_{2,1} \cong {}^2 \text{Mat}(2,\mathbb{R})$ with $f = \frac{1}{4}$ $\frac{1}{4}(1 + \mathbf{e}_1)(1 + \mathbf{e}_{23})$ and $\hat{f} = \frac{1}{4}$ $rac{1}{4}(1 \mathbf{e}_1$)(1+ \mathbf{e}_{23}). Then, a transversal of the stabilizer $G_{2,1}(f) = G_{2,1}(f) \triangleleft G_{2,1}$ is $\mathcal{M} = \{1, \mathbf{e}_2\}.$ Let $\psi, \phi \in S = C\ell_{2,1}f$ and $\psi_g, \phi_g \in \hat{S} = C\ell_{2,1}\hat{f}$. Hence,

$$
\psi = \psi_{11}f + \psi_{21}\mathbf{e}_2f, \qquad \phi = \phi_{11}f + \phi_{21}\mathbf{e}_2f,\tag{71}
$$

$$
\psi_g = \psi_{11}\hat{f} + \psi_{21}\mathbf{e}_2\hat{f}, \qquad \phi_g = \phi_{11}\hat{f} + \phi_{21}\mathbf{e}_2\hat{f} \tag{72}
$$

¹⁴Recall that $Sp(N) = U_{\mathbb{H}}(N)$ where $U_{\mathbb{H}}(N)$ is the quaternionic unitary group [\[7\]](#page-22-7).

ON THE TRANSPOSITION ANTI-INVOLUTION III 17

Table 4: Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in semisimple Clifford algebras $C\ell_{p,q} \cong {}^2\mathrm{Mat}(2^{k-1},\mathbb{R})$

$k = q - r_{q-p}, p - q = 1 \mod 4, p - q = 1 \mod 8$										
				$ (p,q) (1,0) (2,1) (3,2) (0,7) (4,3) (1,8) (5,4)$						
								$\Big \, G^{\varepsilon}_{p,q} \, \left \,\left \, {^2O(1)}\right \, {^2O(2)}\right \, {^2O(4)}\Big \, \Big \, {^2O(8)}\, \Big \, {^2O(8)}\, \Big \, {^2O(16)}\, \Big \, {^2O(16)}\, \Big \, \left {\,^2O(16)}\right \, \Big \,$		

Table 5: Automorphism group $G_{p,q}^{\varepsilon}$ of $T_{\varepsilon}(\psi)\phi$ in semisimple Clifford algebras $C\ell_{p,q} \cong {}^2\mathrm{Mat}(2^{k-1},\mathbb{H})$

where $\psi_{i1}, \phi_{i1} \in \mathbb{R}, i = 1, 2^{15}$ $\psi_{i1}, \phi_{i1} \in \mathbb{R}, i = 1, 2^{15}$ $\psi_{i1}, \phi_{i1} \in \mathbb{R}, i = 1, 2^{15}$ It is then easy to see that the transposition scalar product on S:

$$
T_{\varepsilon}(\psi)\phi = (\psi_{11}\phi_{11} + \psi_{21}\phi_{21})f\tag{73}
$$

is invariant under $O(2)$. Likewise, the transposition scalar product on \hat{S}

$$
T_{\varepsilon}(\psi_g)\phi_g = (\psi_{11}\phi_{11} + \psi_{21}\phi_{21})\hat{f}
$$
\n(74)

is also invariant under $O(2)$. Thus, on $\check{S} = S \oplus \hat{S}$, we get

$$
T_{\varepsilon}(\check{\psi})\check{\phi} = (T_{\varepsilon}(\psi)\phi, T_{\varepsilon}(\psi_g)\phi_g) = (\psi_{11}\phi_{11} + \psi_{21}\phi_{21})(f, \hat{f}) \in \check{K}
$$
\n(75)

where $\check{K} \cong {}^{2} \mathbb{R}$ and $G_{2,1}^{\varepsilon} = {}^{2}O(2)$. Of course, in the faithful spinor representation in $\check{S} = S \oplus \hat{S}$ over the double field ²R, we have

$$
[T_{\varepsilon}(\check{\psi})] = \begin{bmatrix} (\psi_{11}, \psi_{11}) & (\psi_{21}, \psi_{21}) \\ (0, 0) & (0, 0) \end{bmatrix} \text{ and } [\check{\phi}] = \begin{bmatrix} (\phi_{11}, \phi_{11}) & (0, 0) \\ (\phi_{21}, \phi_{21}) & (0, 0) \end{bmatrix}
$$
(76)

which gives the same result in matrix form:

$$
[T_{\varepsilon}(\check{\psi})][\check{\phi}] = \begin{bmatrix} (\lambda, \lambda_g) & (0, 0) \\ (0, 0) & (0, 0) \end{bmatrix}
$$
\n(77)

¹⁵By a small abuse of notation, we write the same numerical components for ψ and ψ_g , also for ϕ and ϕ_q , although in general they may be different. In order to identify the automorphism groups what matters is really the form of the polynomial in the numerical components but not the actual components.

where $\lambda_g = \lambda = \psi_{11}\phi_{11} + \psi_{21}\phi_{21} \in \mathbb{R}$. For a comparison, notice that the scalar products β_+ and β_- on \dot{S} are

$$
\beta_+(\check{\psi}, \check{\phi}) = (0, 0), \tag{78}
$$

$$
\beta_{-}(\check{\psi},\check{\phi}) = (\psi_{11}\phi_{21} - \psi_{21}\phi_{11})(f,\hat{f})
$$
\n(79)

and their automorphism groups are, respectively, ${}^2GL(2,\mathbb{R})$ and ${}^2Sp(2,\mathbb{R})$.^{[16](#page-17-0)}

Example 8. Consider $C\ell_{1,4} \cong {}^2\text{Mat}(2,\mathbb{H})$ with $f = \frac{1}{4}$ $\frac{1}{4}(1 + \mathbf{e}_{234})(1 + \mathbf{e}_{15})$ and $\hat{f} = \frac{1}{4}$ $rac{1}{4}(1$ $e_{234})(1+e_{15})$. Then, a transversal of the stabilizer $G_{1,4}(f) = G_{1,4}(f) \triangleleft G_{1,4}$ is $\mathcal{M} = \{1, e_1\}$. Let $\psi, \phi \in S = \mathcal{C}\ell_{1,4}f$ and $\psi_g, \phi_g \in \hat{S} = \mathcal{C}\ell_{1,4}\hat{f}$. Let's assign, modulo f, quaternionic components to $\frac{1}{2}$ -spinors as follows:

$$
\psi_i := \psi_{11} + \psi_{12}\mathbf{e}_2 + \psi_{13}\mathbf{e}_3 + \psi_{14}\mathbf{e}_{23}, \quad \phi_i := \phi_{11} + \phi_{12}\mathbf{e}_2 + \phi_{13}\mathbf{e}_3 + \phi_{14}\mathbf{e}_{23},
$$

for $i = 1, 2$. Thus,

$$
\psi = f\psi_1 + \mathbf{e}_1 f\psi_2, \qquad \phi = f\phi_1 + \mathbf{e}_1 f\phi_2,\tag{80}
$$

$$
\psi_g = \hat{f}\psi_1 + \mathbf{e}_1 \hat{f}\psi_2, \qquad \phi_g = \hat{f}\phi_1 + \mathbf{e}_1 \hat{f}\phi_2.
$$
\n(81)

Then, modulo f , we find the transposition scalar product on S

$$
T_{\varepsilon}(\psi)\phi = (\overline{\psi}_1\phi_1 + \overline{\psi}_2\phi_2)f\tag{82}
$$

where $\overline{\psi}_i$ denotes "quaternionic" conjugation. It is easy to see that this product is invariant under $Sp(2) = U_{\mathbb{H}}(2)$. Furthermore, modulo \hat{f} , we find the transposition scalar product on \tilde{S}

$$
T_{\varepsilon}(\psi_g)\phi_g = (\overline{\psi}_1\phi_1 + \overline{\psi}_2\phi_2)\hat{f}
$$
\n(83)

hence it is also invariant under $Sp(2)$. Therefore, on $\check{S} = S \oplus \hat{S}$, we get

$$
T_{\varepsilon}(\check{\psi})\check{\phi} = (T_{\varepsilon}(\psi)\phi, T_{\varepsilon}(\psi_g)\phi_g) = (\overline{\psi}_1\phi_1 + \overline{\psi}_2\phi_2)(f, \hat{f}) \in \check{\mathbb{K}} \cong {}^2\mathbb{H}
$$
(84)

and $G_{1,4}^{\varepsilon} = {}^2Sp(2)$. We can also verify this result using the faithful spinor representation in $\check{S} = S \oplus \hat{S}$ over the double (skew) field ²H:

$$
[T_{\varepsilon}(\check{\psi})] = \begin{bmatrix} (\overline{\psi}_1, \overline{\psi}_1) & (\overline{\psi}_2, \overline{\psi}_2) \\ (0, 0) & (0, 0) \end{bmatrix} \text{ and } [\check{\phi}] = \begin{bmatrix} (\phi_1, \phi_1) & (0, 0) \\ (\phi_2, \phi_2) & (0, 0) \end{bmatrix}
$$
(85)

which gives the same result in matrix form:

$$
[T_{\varepsilon}(\check{\psi})][\check{\phi}] = \begin{bmatrix} (\lambda, \lambda_g) & (0, 0) \\ (0, 0) & (0, 0) \end{bmatrix}
$$
 (86)

¹⁶In [\[12,](#page-22-2) Table 1, page 236], Lounesto shows for short only $GL(2,\mathbb{R})$ whereas in [12, Table 2, page 236], he also shows ${}^2Sp(2,\mathbb{R})$. Thus, our definition of the symplectic group $Sp(N,\mathbb{R})$ is the same as his.

where $\lambda_g = \lambda = \overline{\psi}_1 \phi_1 + \overline{\psi}_2 \phi_2 \in \mathbb{H}$. We again compare this product with β_+ and β_- on \check{S} which are:

$$
\beta_+(\check{\psi}, \check{\phi}) = (\overline{\psi}_1 \phi_2 + \overline{\psi}_2 \phi_1)(f, \hat{f}), \tag{87}
$$

$$
\beta_{-}(\check{\psi},\check{\phi}) = (0,0),\tag{88}
$$

and their automorphism groups are, respectively, $^2U_{1,1}\mathbb{H} = U_{1,1}\mathbb{H} \times U_{1,1}\mathbb{H}$ and $^2GL(2,\mathbb{H})$.^{[17](#page-18-1)}

4. Clifford algebra as a twisted group ring

For any group G , its commutator subgroup G' is a normal subgroup of G , and the quotient group G/G' is abelian. [\[16,](#page-23-4) Prop. 5.57] Hence, since the vee group $G_{p,q}$ is not abelian (for $n = p + q \ge 2$) of order $2 \cdot 2^{p+q}$ and its commutator subgroup $G'_{p,q} = {\pm 1}$, the abelian quotient group is just

$$
G_{p,q}/G'_{p,q} \cong (\mathbb{Z}_2)^n = \underbrace{\mathbb{Z}_2 \times \cdots \times \mathbb{Z}_2}_{n} \quad \text{with} \quad n = p + q.
$$

Thus, we can view the universal Clifford algebra $C\ell_{p,q}$ of a non-degenerate quadratic form Q of signature (p, q) as a twisted group ring $C\ell_{p,q} = \mathbb{R}^t[(\mathbb{Z}_2)^n]$ over the abelian group $(\mathbb{Z}_2)^n$.^{[18](#page-18-2)} Let e_1, \ldots, e_n be generators for *n* isomorphic copies of \mathbb{Z}_2 , that is,

$$
\mathbb{Z}_2 \cong \langle e_1 \rangle \cong \langle e_2 \rangle \cong \cdots \cong \langle e_n \rangle
$$

where the cyclic groups $\langle e_i \rangle = \{e_i^{a_i}\}, a_i = 0, 1, i = 1, \ldots, n$, are written multiplicatively.^{[19](#page-18-3)} Then, the *twisted group ring* $\mathbb{R}^{t}[(\mathbb{Z}_{2})^{n}]$ [\[14,](#page-23-0) Sect. 2] is an associative R-algebra with basis $\{\bar{x} \mid x \in (\mathbb{Z}_2)^n\}$ and multiplication defined distributively for all $x, y \in (\mathbb{Z}_2)^n$ as

$$
\bar{x}\bar{y} = \gamma(x, y)\overline{xy}, \qquad \gamma(x, y) \in \mathbb{R}^* = \mathbb{R} \setminus \{0\}.
$$

Due to the required associativity of the algebra $(\bar{x}\bar{y})\bar{z} = \bar{x}(\bar{y}\bar{z})$ for any $x, y, z \in (\mathbb{Z}_2)^n$, the function $\gamma : (\mathbb{Z}_2)^n \times (\mathbb{Z}_2)^n \to \mathbb{R}^*$ must satisfy the relation

$$
\gamma(x, y)\gamma(xy, z) = \gamma(y, z)\gamma(x, yz), \qquad \forall z, y, z \in (\mathbb{Z}_2)^n
$$

which implies that γ is a 2-cocycle. Furthermore, it is shown [\[14,](#page-23-0) Lemma 2.1] that a general twisted group ring $K^t[G]$ has an identity $\gamma(1,1)^{-1}\overline{1}$ and that the elements $\overline{x} \in K^t[G]$ are all units. The inverse of \bar{x} is:

$$
\bar{x}^{-1} = \gamma(x, x^{-1})\gamma(1, 1)^{-1}\overline{x^{-1}} = \gamma(x^{-1}, x)\gamma(1, 1)^{-1}\overline{x^{-1}}.
$$

¹⁷The group $U_{p,q}$ is defined in [\[7,](#page-22-7) Page 99] as the group of automorphisms of a *Hermitian form* of signature (p, q) on a quaternionic vector space V of dimension $p + q$ thus having the standard expression $\sum_{i=1}^p \overline{v}_i w_i - \sum_{i=p+1}^{p+q} \overline{v}_i w_i$. In [\[12,](#page-22-2) Table 1, page 236] we find ${}^2Sp(2, 2)$ and in [12, Table 2, page 236] we find $GL(2, \mathbb{H})$, respectively.

¹⁸For a general theory of group rings see [\[14\]](#page-23-0).

¹⁹We could simplify notation by identifying an ordered *n*-tuple $(e_1^{a_1}, e_2^{a_2}, \ldots, e_n^{a_n})$ from $(\mathbb{Z}_2)^n$ with a monomial $e_1^{a_1}e_2^{a_2}\cdots e_n^{a_n}$ in $\mathbb{R}^t[(\mathbb{Z}_2)^n]$.

One can easily show that the formula $(xy)^{-1} = y^{-1}x^{-1}, \forall x, y \in G$, carries over to $(\bar{x}\bar{y})^{-1} = \bar{y}^{-1}\bar{x}^{-1}, \forall \bar{x}, \bar{y} \in K^t[G]$:

$$
(\bar{x}\bar{y})(\bar{y}^{-1}\bar{x}^{-1}) = \bar{x}(\bar{y}\bar{y}^{-1})\bar{x}^{-1} = \bar{x}\gamma(1,1)^{-1}\bar{1}\bar{x}^{-1} = \bar{x}\bar{x}^{-1}\gamma(1,1)^{-1}\bar{1}
$$

$$
= \gamma(1,1)^{-1}\bar{1}\gamma(1,1)^{-1}\bar{1} = \gamma(1,1)^{-1}\gamma(1,1)^{-1}\bar{1}\bar{1}
$$

$$
= \gamma(1,1)^{-1}\gamma(1,1)^{-1}\gamma(1,1)\bar{1} = \gamma(1,1)^{-1}\bar{1}
$$
(89)

In [\[14,](#page-23-0) Page 5], one finds this definition of a map $\ast: K[G] \to K[G]$ in a general group ring $K[G]$:

$$
\left(\sum a_x x\right)^* = \sum a_x x^{-1}
$$

where the summation is over $x \in G$. It follows easily that, for any $\alpha, \beta \in K[G]$, we have: (i) $(\alpha + \beta)^* = \alpha^* + \beta^*$, (ii) $(\alpha\beta)^* = \beta^*\alpha^*$, (iii) $\alpha^{**} = \alpha$, (iv) $1^* = 1$ for the unity. Hence, * is an anti-involution of the group ring.

Notice that a similar map can be defined in any *twisted* group ring $* : K^t[G] \to K^t[G]$ namely

$$
\left(\sum a_x \bar{x}\right)^* = \sum a_x \bar{x}^{-1}.
$$

Furthermore, this map is an anti-involution of $K^t[G]$ as it satisfies the same four properties (i)–(iv). In particular, property (ii) follows from the identity $(\bar{x}\bar{y})^{-1} = \bar{y}^{-1}\bar{x}^{-1}$ shown above.

We recall from [\[2,](#page-22-1) Lemma 1] some properties of the transposition anti-involution T_{ϵ}^{γ} , and, in particular, its action $T_{\varepsilon}^{\sim}(m) = m^{-1}$ on a monomial m in the Grassmann basis \mathcal{B} which is, as we see now, identical to the action $*(m) = m^{-1}$ on every $m \in \mathcal{B}$. We formulate our concluding result.

Theorem 2. The anti-involution T_{ε}^{\sim} on the Clifford algebra $C_{p,q}$ is the anti-involution $*$ of $C\ell_{p,q}$ viewed as the twisted group ring $\mathbb{R}^t[(\mathbb{Z}_2)^n]$.

For a Hopf algebraic discussion of Clifford algebras as twisted group algebras, see [\[6,](#page-22-3) [13\]](#page-22-8) and references therein.

Remark. Following [\[12,](#page-22-2) Chapt. 21], consider *n*-tuples $\underline{a} = a_1 a_2 \dots a_n$ of binary digits $a_i = 0, 1$, with addition $\underline{a} \oplus \underline{b} = \underline{c}$ defined by term wise addition modulo 2. This is the natural component wise product group structure on $(\mathbb{Z}_2)^n$. Thus, with this operation, the set of n-tuples is a group and the group characters are Walsh functions

$$
w_{\underline{a}}(\underline{b}) = (-1)^{\sum_{i=1}^{n} a_i b_i}
$$

As group characters, the Walsh functions satisfy $w_k(a \oplus b) = w_k(a)w_k(b)$. For any binary *n*-tuple $\underline{a} = a_1 a_2 \dots a_n$, let's adopt notation

$$
e_{\underline{a}} \stackrel{\text{def}}{=} e_1^{a_1} e_2^{a_2} \cdots e_n^{a_n}
$$

for any element $(e_1^{a_1})$ $\binom{a_1}{1}, \binom{a_2}{2}, \ldots, \binom{a_n}{n} \in (\mathbb{Z}_2)^n$. Then, Lounesto shows how one can define the Clifford product in the algebra $C\ell_{p,q}$ on the "basis monomials" (later extended by linearity and associativity to all algebra elements) with the help of Walsh functions, namely,

$$
e_{\underline{a}}e_{\underline{b}} \stackrel{\text{def}}{=} (-1)^{\sum_{i=1}^{p} a_i b_i} w_{\underline{a}}(h(\underline{b})) e_{\underline{a} \oplus \underline{b}} \tag{90}
$$

where h , the inverse of the so called *Gray code*, is defined as

$$
h(\underline{a})_i = \sum_{j=1}^i a_j \bmod 2.
$$

Thus, we have a close connection between viewing Clifford algebra $C\ell_{p,q}$ as a twisted group ring $\mathbb{R}^t[(\mathbb{Z}_2)^n]$ and as a real associative algebra with the monomial basis $\{e_a\}$ for all binary n -tuples a and multiplication defined with the help of the Walsh function and inverse Gray code as in [\(90\)](#page-19-0). In particular, one should be able to express the twisting 2-cocycle γ needed to define the product in $\mathbb{R}^t[(\mathbb{Z}_2)^n]$ (see [\[6,](#page-22-3) [13\]](#page-22-8)) in terms of the Walsh functions and the Gray code.

5. Conclusions

We summarize and conclude our work on the transposition involution in real Clifford algebras, including the developments of [\[1,](#page-22-0) [2\]](#page-22-1). The present set of three papers develops in a concise and explicit manner all important aspects of the transposition map, which we have lifted from the matrix representations to an abstract map in real Clifford algebras. We did this in an exhaustive manner for all signatures $\varepsilon = (p, q)$ for $p + q \leq 9$ well aware of the fact that with mod 8 periodicity of real Clifford algebras, we have a complete treatment for any signature. Our goal was to explicitly verify all our claims with CLIFFORD [\[5\]](#page-22-9). This has revealed some corrections necessary to results already published in the literature. We also want to mention that despite an extensive search, we were unable to find our transposition map in the literature, contrary to the immediate feeling that this map 'should already have been known'. However, see its relation to the '*-map' in the twisted group rings given in Theorem [2.](#page-19-1) Of course, the transposition map for Euclidean and anti-Euclidean signatures coincides with, respectively, the reversion and the conjugation anti-involutions of $C\ell_{p,q}$ used in the standard constructions of Spin and Pin groups, but for non-definite signatures it differs.

The first paper [\[1\]](#page-22-0) is concerned with a general setup of the transposition map. This includes a careful choice of sorted bases, with an admissible order, allowing for further generalizations. Our starting point was to find the abstract map in abstract real Clifford algebras induced by the linear algebra map of transposition in the left (right) regular matrix representations of the real Clifford algebras. We do not reiterate the points given in the conclusions of [\[1\]](#page-22-0) but we focus here on the general outline of our work.

The transposition map explains the block structure (grading) of the regular representation matrices. Moreover, we exhibited an explicit decomposition of the transposition map with respect to the graded decomposition of the Clifford algebras $Cl_{p,q} = Cl_{p,0} \otimes Cl_{0,q}$ for arbitrary signatures. This fact, depending on the symmetry of the employed non singular polar bilinear form $B_{p,q}$, emerges from the decomposition of the related quadratic form $Q_{p,q} = Q'_{p,0} \perp Q''_{0,q}$ into a positive and a negative definite form. On the technical side we have extended the duality result of Lounesto

$$
\langle u \mathsf{I}_B v, w \rangle = \langle v, \tilde{u} \wedge v \rangle \tag{91}
$$

relating the contraction and the Grassmann wedge product, to the general Clifford case

$$
\langle u, vw \rangle = \langle \tilde{v}u, v \rangle \qquad \text{provided } \det Q_{p,q} \neq 0. \tag{92}
$$

Choosing a sorted basis \mathcal{B} which diagonalizes $Q_{p,q}$, we have considered two canonical isomorphisms $V \to V^*$ identifying the vector space V with its dual V^* . These maps are given by: (i) the dual basis \mathcal{B}^* defined as $e_i^*(e_j) = \delta_{i,j}$ and (ii) the reciprocal basis \mathcal{B}^{\flat} defined as $\mathbf{e}_i^{\flat} =$ $\overline{\mathbf{e}_i}$ ε_i . This leads to the universal dual (or *reciprocal*) Clifford algebra Cl_n^* . The isomorphism $\flat: V \to V^*$ depends on the quadratic form $Q_{p,q}$ or, equivalently, on its symmetric polar bilinear form $B_{p,q}$. The consequences are somewhat straightforward and we gave a few explicit examples.

In general, the regular representation is reducible. Thus, it is much more interesting to explore how the transposition map is related to the irreducible spinor representations. This is done in [\[2\]](#page-22-1). Algebraic spinor spaces S, seen as left $C\ell$ -modules, are minimal left ideals. These ideals can be generated by a primitive idempotent f such that $S(f) = C\ell_{p,q}f$. As real Clifford algebras are simple or are direct sums of two simple subalgebras, such spinor ideals are right modules over K which is either a (skew) field or a double (skew) field. The spinor ideal $_{C\ell}S(f)_{\mathbb{K}}$ is hence a left $C\ell$ and right K-module. The abstract transposition map factors into a transposition map on the left action by $C\ell$ and an (anti) involution on K. It was shown, for simple and semisimple algebras, that this involution, depending on the signature mod 8, amounts to the identity on \mathbb{R} , complex conjugation on \mathbb{C} , and quaternionic conjugation on H. An important role in these constructions is played by the vee group $G_{p,q}$ and the stabilizer group $G_{p,q}(f)$ of the primitive idempotent element f generating the spinor space $S(f)$. We have explicitly constructed left transversals of cosets which provide bases for the spinor spaces. A full classification of the stabilizer groups $G_{p,q}(f)$ was given in [\[2\]](#page-22-1) for all signatures $p + q = n \leq 9$, which is complete when taken mod 8 in both simple and semisimple algebras. These stabilizer groups do play an important role in quantum computing too. The latter, however, usually employs only complex Clifford algebras which posses a much simpler mod 2 structure.

We showed that the transposition map allows one to form a new transposition scalar product on spinor spaces which we have studied in the present paper. Only in the Euclidean and anti-Euclidean cases, this scalar product is identical to the two known spinor scalar products, see for example [\[12\]](#page-22-2), and it is different in all other signatures. The present paper provides a complete classification of the invariance groups $G_{p,q}^{\varepsilon}$ of the transposition scalar product. We also gave a detailed exposition on $T_{p,q}(f)$, the idempotent group $T_{p,q}(f)$ of f, and the field group $K_{p,q}(f)$ of f, as normal subgroups in the stabilizer group $G_{p,q}(f)$ of the primitive idempotent f, and on their coset spaces $G_{p,q}/T_{p,q}(f)$, $G_{p,q}(f)/T_{p,q}(f)$, and $G_{p,q}/G_{p,q}(f)$. These subgroups allow to construct very effectively non-canonical transversals and hence basis elements of the spinor spaces and the (skew double) field underlying the spinor space. This leads to a description of the situation tabulated for $p + q = n \leq 9$ which is complete and sufficient due to the mod 8 periodicity. A further aspect which we have emphasized is the fact that the Clifford algebras can be seen as twisted group rings $\mathbb{R}^t[(\mathbb{Z}_2)^n]$. In particular, we have observed that our transposition is then mapped to the 'star

map' in such rings [\[14\]](#page-23-0). This observation relates our work to recent approaches to Clifford algebras using Hopf algebraic methods applied to the twisted group ring $\mathbb{R}^t[(\mathbb{Z}_2)^n]$ [\[6,](#page-22-3) [13\]](#page-22-8). In particular, the 'star' map of a twisted group ring is actually the antipode map.

Let β be an automorphism of K. The right module $S_{\mathbb{K}}$ can be given a new module structure $S_{\mathcal{K}}$ by $\psi \cdot \lambda = \psi \beta(\lambda)$. Similarly we can twist the left action of Cl seen as the matrix algebra over K. As an example, the complex left module $\mathbb{C}M$ can receive an antilinear complex vector space structure $\mathbb{C}M$, such that $z \cdot m = \overline{z}m$. Seen as representations, these modules are in general not isomorphic.

As a final remark, observe that the transposition anti-involution connects left and right spinor modules $(T_{\varepsilon}^{\sim} : c_{\ell} S_{\mathbb{K}} \to \mathbb{K} S_{\ell} c_{\ell})$ or, equivalently, regular left and right representations, and it implements an *isomorphism* between the left and the right modules. Such a map introduces a Frobenius algebra structure [\[11\]](#page-22-10), in general twisted (or of the second kind), on the Clifford algebra and its respective representations. This is expected due to the fact that Clifford algebras can be seen as twisted group rings. Our work on the stabilizer and invariance groups shows that only in the case when $K \cong \mathbb{R}$ this identification is straightforward. In all other cases, the left and the right modules have different actions of the base field K due to the action of the transposition involution on the base field, which induces the twisting in the sense of the twisted Frobenius structures [\[11,](#page-22-10) Ch. 7.1]. The clarifications brought forward in this set of three articles will help to investigate this fact.

REFERENCES

- [1] R. Ab lamowicz and B. Fauser, On the Transposition Anti-Involution in Real Clifford Algebras I: The Transposition Map (to appear in Linear and Multilinear Algebra)
- [2] R. Ab lamowicz and B. Fauser, On the Transposition Anti-Involution in Real Clifford Algebras II: Stabilizer Groups of Primitive Idempotents (to appear in Linear and Multilinear Algebra)
- [3] R. Ab lamowicz, *Clifford Algebra Computations with Maple*, in "Clifford (Geometric) Algebras with Applications in Physics, Mathematics, and Engineering", W. E. Baylis, (Ed.) (Birkhäuser, Boston, 1996) 463–502
- [4] R. Ablamowicz, Spinor Representations of Clifford algebras: A Symbolic Approach, Computer Physics Communications Thematic Issue - Computer Algebra in Physics Research 115, No. 2–3 (1998) 510– 535
- [5] R. Ablamowicz and B. Fauser, CLIFFORD for Maple, $http://math.html, `interch.edu/rafal/` (C1996-2011)$
- [6] H. Albuquerque and S. Majid, *Clifford algebras obtained by twisting of group algebras*, J. of Pure and Appl. Algebra 171, (2002) 133–148
- [7] W. Fulton and J. Harris, Representation Theory: A First Course, (Springer, New York, 1991)
- [8] A. J. Hahn, Quadratic Algebras, Clifford Algebras, and Arithmetic Witt Groups, (Undergraduate Texts in Mathematics) (Springer-Verlag, New York, 1994)
- [9] J. Helmstetter, Groupes de Clifford pour de formes quadratiques de rang quelconque, C. R. Acad. Sci. Paris 285 (1977) 175–177
- [10] J. Helmstetter and A. Micali, *Quadratic Mappings and Clifford Algebra*, (Birkhäuser, Basel, 2008)
- [11] L. Kadison, New examples of Frobenius extensions, Univ. Lecture Series Vol. 14, (Amer. Math. Soc. Providence Rhode Island, 1999)
- [12] P. Lounesto, Clifford Algebras and Spinors, 2nd ed. (Cambridge University Press, Cambridge, 2001)
- [13] S. Morier-Genoud and V. Ovsienko, *Simple graded commutative algebras*, J. of Algebra 323 (2010) 1649–1664
- [14] D. S. Passman, The Algebraic Structure of Group Rings, (Robert E. Krieger Publishing Company, 1985)
- [15] I. R. Porteous, Clifford Algebras and the Classical Groups, (Cambridge University Press, Cambridge, 1995)
- [16] J.J. Rotman, Advanced Modern Algebra, Revised Printing, (Prentice Hall, Upper Saddle River, 2002)
- [17] N. Salingaros, Realization, extension, and classification of certain physically important groups and algebras, J. Math. Phys. 22 (1981) 226–232
- [18] N. Salingaros, On the classification of Clifford algebras and their relation to spinors in n dimensions, J. Math. Phys. 23 (1) (1982) 1–7
- [19] N. Salingaros, The relationship between finite groups and Clifford algebras, J. Math. Phys. 25 (1984) 738–742
- [20] V. V. Varlamov, Universal Coverings of the Orthogonal Groups, Adv. in Applied Clifford Algebras 14, No. 1 (2004) 81–168

E-mail address: rablamowicz@tntech.edu

Department of Mathematics, Box 5054, Tennessee Technological University, Cookeville, TN 38505, USA

E-mail address: B.Fauser@cs.bham.ac.uk

SCHOOL OF COMPUTER SCIENCE, THE UNIVERSITY OF BIRMINGHAM, EDGBASTON-BIRMINGHAM, W. Midlands, B15 2TT, United Kingdom