

# Anomalous random walk of a greedy traveling salesman

Adam Lipowski<sup>1</sup> and Dorota Lipowska<sup>2</sup>

<sup>1</sup>*Faculty of Physics, Adam Mickiewicz University, Poznań, Poland*

<sup>2</sup>*Faculty of Modern Languages and Literature, Adam Mickiewicz University, Poznań, Poland*

Using Monte Carlo simulations we examine the diffusive properties of the greedy algorithm in the  $d$ -dimensional traveling salesman problem. Our results show that the average squared distance from the origin  $\langle r^2 \rangle$  after  $t$  steps scales as  $t^\alpha$ . While for  $d = 3$  and  $4$ ,  $\alpha$  is very close to unity, for  $d = 2$  it is greater than unity ( $\alpha \sim 1.08(1)$ ). The distribution of lengths also shows marked differences between  $d = 2$  and  $d > 2$  versions. A simple strategy adopted by the salesman might resemble strategies chosen by some foraging and hunting animals, for which anomalous diffusive behaviour has recently been reported but interpreted in terms of Lévy flights. Our results suggest that anomalous diffusive behaviour in such systems might appear due to the dimension-dependent properties of a search space.

Keywords: traveling salesman, greedy algorithm, anomalous random walk

Anomalous diffusion appears whenever the typical square displacement of a tracer particle  $R^2(t)$  does not obey the normal diffusion law  $R^2(t) \sim t^\alpha$  with  $\alpha = 1$ . Deviations include numerous examples of subdiffusion ( $\alpha < 1$ ) such as, for example, photoconductivity of amorphous materials [1], diffusion in convective rolls [2] or a random walk on percolation cluster [3]. On the other side, there is much rarer superdiffusive ( $\alpha > 1$ ) behaviour, of which the Richardson diffusion in turbulent fluids is a classical example [4]. Some macromolecules or biological systems also exhibit anomalous dynamics with superdiffusive behaviour reported for micelles [5] or living cell migrations [6].

Recently, a number of studies have indicated that foraging or hunting strategies of some animals also show superdiffusive behaviour and examples include ants [7], fruit flies [8], or wandering albatrosses [9]. The prevailing understanding is that while searching for food these animals move in steps of different lengths with fat power-law distributions consistent with Lévy flights. However, some other works show that this hypothesis should be taken with care. Indeed, re-analysing data for the wandering albatrosses, Edwards *et al.* concluded that previously suggested Lévy-flight scenario appeared due to spurious inclusion of very long flights [10]. There are also claims that the idea of animals using stochastic but essentially blind search strategies does not seem to be viable since it neglects the role of animals' intelligence and experience in guiding them [11]. Nevertheless, more extensive recent studies for 14 open-ocean predator fish species seem to support Lévy-flight scenario [12].

In the present paper we show that a simple search strategy, i.e., go to the nearest (intelligence) and not yet visited (experience) place, implies in a sufficiently low dimension ( $d \leq 2$ ) anomalous diffusive behaviour. For  $d > 2$ , most likely normal diffusion takes place. The distribution of distances does not have fat power-law tails, however, when the search exploits all available places, much broader and possibly Lévy type distributions can




FIG. 1. An exemplary configuration of the path of the greedy traveling salesman for  $N = 2 \cdot 10^4$  cities (dots) and of duration  $t = 10^3$ .

be seen.

More specifically, we consider a traveling salesman [13] that has to visit cities that are uniformly distributed in the  $d$ -dimensional space of linear size  $L$ . We assume a unit density of cities, i.e.,  $N/L^d = 1$ , where  $N$  is the number of cities. The salesman is not interested in finding the minimal tour but visits cities according to the following greedy algorithm: in each step, from the last visited city it moves to the nearest not yet visited city. For most simulations, the starting point is the city placed in the center of the space, i.e., of the cartesian coordinates  $(L/2, L/2, \dots, L/2)$ . The quantity of main interest is the average square distance  $\langle r^2 \rangle$  from the starting point and especially its asymptotic form after many steps  $t$ . An example of a typical trajectory in a two-dimensional case is shown in Fig. 1. Although qualitatively it looks like an ordinary random-walk trajectory, more detailed analysis shows that it is actually superdiffusive.

To examine the asymptotic behaviour of  $\langle r^2 \rangle$ , it is essential to average over many configurations with large




FIG. 2. Sectorization of the two-dimensional space. For a trajectory that at a certain moment arrives at the sector  $s_0$ , the program searches for available cities within  $s_0$ . (i) If it finds some, then to find the nearest city it is sufficient to examine  $s_0$  and its neighbouring sectors as indicated in (a). This is because any city outside these sectors is further from any point in  $s_0$  than the length of the diagonal of  $s_0$  (which is the largest distance within  $s_0$ ). (ii) If there are no available cities in  $s_0$ , the program searches for them in a larger area (bold square in (b)). If it finds some, then to find the nearest city it is sufficient to examine sectors as indicated in (b). (iii) If no cities are found, the program examines even larger areas (until some cities are found).

number of cities  $N$  so that a relatively large part of the trajectory of the salesman was not affected by the boundaries. Because of these finite-size effects, we terminate the generation of trajectories after number of steps that is much smaller than  $N$ . As a result, the  $t$ -th city on a trajectory is typically a close neighbour of the  $(t-1)$ -th city. Instead of searching for the nearest not yet visited city in the entire configuration, it would be much faster to restrict the search to the cities which are in the vicinity of the last visited city. Such a modification can be implemented by means of the sectorization of the space. In addition to coordinates, we store the information about a sector to which a given city belongs. Then, to find the  $t$ -th city, we search the sectors surrounding the  $(t-1)$ -th city (see Fig. 2). Let us emphasize that the above described sectorization does not introduce any approximation as for a given configuration of cities the algorithm generates exactly the same trajectory as the one without sectorization. In our simulations we used such mesh that each sector contained approximately 10 cities. (In the two-dimensional case, where the longest trajectories had to be generated, the sectorization reduced the simulation time by the factor  $\sim 10^3$ .)

Using the above algorithm, we simulated  $d=2, 3$ , and 4 systems containing up to  $3 \cdot 10^8$  cities. Results for the two-dimensional case are shown in Fig. 3. Since the asymptotic slope seems to be bigger than unity, these data show slightly superdiffusive behaviour, but more quantitative analysis of the slope would be desirable. One should take into account that the asymptotic regime in




FIG. 3. The logarithmic plot of the time dependence of  $\langle r^2 \rangle$  for several values of  $N$  indicated in the right bottom. The straight line corresponds to the linear increase  $\langle r^2 \rangle \sim t$ . For each  $N$  the plotted results are averages over at least  $10^5$  independent configurations.




FIG. 4. The time-dependent exponent  $\alpha(t)$  calculated in the two-dimensional case for several values of  $N$ .

the numerical data will be visible only in an intermediate interval since both the initial transit as well as large- $t$  data should be neglected (for long trajectories  $\langle r^2 \rangle$  is strongly affected by finite-size effects). To estimate  $\alpha$ , we first examine the time-dependent quantity  $\alpha(t)$  defined as the local exponent estimated from data spanning over one decade and centered at  $t$  (i.e., to estimate  $\alpha(t)$  we used  $\langle r^2 \rangle$  calculated for  $t'$  such that  $\log_{10}(t) - 0.5 < \log_{10}(t') < \log_{10}(t) + 0.5$ ).

The obtained results are presented in Fig. 4. As expected, for large  $t$  all curves bent downward and in the middle range a plateau emerges. Our estimate based on data around  $t = 10^5$  suggests that  $\alpha = 1.08(1)$ .

We used the same technique in the three- and four-dimensional versions, and their final results are shown in Fig. 5 and Fig. 6, respectively. In these higher-dimensional cases, the linear size  $L$  increases slower with the number of cities  $N$ , and that is why simulations bent




FIG. 5. The time-dependent exponent  $\alpha(t)$  calculated in the three-dimensional case for several values of  $N$ .




FIG. 6. The time dependent exponent  $\alpha(t)$  calculated in the four-dimensional case for several values of  $N$ .

downward already for smaller value of  $t$ . The emerging plateaux indicate that  $\alpha$  is very close to unity both for  $d = 3$  and 4. Let us also notice that in the  $d = 1$  case the salesman, possibly after some initial switches, will stick to one side and then move ballistically in the same direction, trivially implying  $\alpha = 2$ .

We also analysed distributions of length  $l$  between subsequent (in a trajectory) cities in the  $d = 2$  and 3 cases. One should take into account, however, that these distributions are not entirely stationary and to some extent vary in time  $t$  (in general, in the late-time evolution very long distances appear and in the final stage they could be of the order of the system size  $L$ ). To reduce this variability, we generated trajectories that were relatively short (most likely none of the runs reached the boundary of the system).

The logarithmic plot of these distributions (Fig. 7) shows that in the  $d = 2$  version the distribution is much broader than in the  $d = 3$  version, and perhaps of a qualitatively different large- $l$  tail. Let us notice that in both cases  $\langle l \rangle$  is similar and of the order of unity (inset




FIG. 7. The plot of the logarithm of the distribution  $P(l)$  of intervals  $l$  calculated for paths of length  $t = 5000$ . The inset shows that the variability of the average interval  $\langle l \rangle$  (at time  $t$ ) during such runs is rather small.




FIG. 8. The same distribution  $P(l)$  as in Fig. 7 but in the log-log scale. The inset shows the log-log plot of  $P(l)$  for runs where all cities are visited ( $N = 10^5$ ). The dotted line has the slope  $-3$  ( $\sim l^{-3}$ ).

in Fig. 7) but the recorded fluctuations of  $l$  are several times larger in the  $d = 2$  case. Let us also observe that large- $l$  tails of these distributions do not seem to have a power-law form. This indicates that superdiffusive behaviour in the  $d = 2$  case is not due to sufficiently broad Lévy-like distributions of lengths  $l$  but rather due to the development of long-time correlations. We will return to this point in the final part of the paper.

The log-log plot of these distributions confirms the absence of power-law tails for large  $l$  for these distributions (Fig. 8). It turns out that the other side of these distributions, namely small- $l$  tails can be understood with some simple arguments. Indeed, these tails are not much different than the small- $l$  tails in the initial distribution of cities. For simplicity, let us examine first the  $d = 1$  case, where  $N-1$  cities are uniformly dis-

tributed on the interval of length  $L = N$  and one city is placed in the center of the interval (i.e., it has coordinate  $L/2$ ). Calculating the probability density  $P_1(x)$  that exactly one city is at the distance  $x$  from the central city (to the right or left) and  $N-2$  cities are further than  $x$  from it, in the limit  $N \rightarrow \infty$  we obtain  $P_1(x) = \frac{2(N-1)}{L}(1 - \frac{2x}{L})^{N-2} = 2 \exp(-2x)$ . Generalizing the above calculation to the  $d = 2$  and  $d = 3$  cases, we obtain  $P_2(x) \sim x \exp(-\pi x^2)$  and  $P_3(x) \sim x^2 \exp(-\frac{4}{3}\pi x^3)$ , respectively. As can be seen in Fig. 8, the small- $l$  tails are indeed close to the tails of the initial distributions both for  $d = 2$  and 3. One can also notice that for large  $x$  the initial distributions  $P_2(x)$  and  $P_3(x)$  decay very fast (faster than exponentially) while the corresponding distributions  $P(l)$  decay slower (see Fig. 7). Thus, the traveling salesman generates paths where short distances appear with approximately the same distribution as in the initial distribution but long distances are generated much more frequently.

It is perhaps interesting, especially in the context of modeling animal search strategies, that the distributions  $P(l)$  become much broader when calculations are made on paths that visit all the cities (inset in Fig. 8). This is because in the final part of the tour the salesman has to visit all the "leftovers" that are typically quite distant from each other. These numerical data suggest that in two-dimensional data the tail of the resulting distribution might even have the power-law decay,  $P(l) \sim l^{-\mu}$  and  $\mu \sim 3$ , and thus could be consistent with the limiting Lévy distribution. Let us notice that there are some claims that Lévy distributions in the foraging data appeared most likely due to a spurious inclusion of late-stage flights [10]. These flights, e.g., of albatrosses returning to their nests, were obviously very long, but should not be considered as part of the searching pattern.

Finally, we would like to comment on the origin of the anomalous diffusion in our model. It is not caused by broad, power-law distributions of lengths, but rather by long-time correlations. More specifically, this is the memory of the salesman that restricts its movements and effectively pushes the salesman out of regions that have already been visited. While in high-dimensional systems ( $d > 2$ ) it has only a minor effect, in low-dimensional systems ( $d = 1$  and 2) this mechanism generates superdiffusive behaviour. Let us notice that a similar mechanism generates superdiffusive behaviour in the so-called self-avoiding walk problem [14]. However, this model of a polymer is known to be superdiffusive for  $d = 1$  ( $\alpha = 2$ ),

$d = 2$  ( $\alpha = 1.5$ ), and  $d = 3$  ( $\alpha = 1.2$ ), with normal diffusion resuming for  $d > 3$ . Our model might thus belong to a novel universality class, where excluded-volume effects play an important role, at least in a sufficiently low dimension. Its direct applicability to search strategies is rather limited, but the model might indicate that the very dimension of the search space play an important role. In particular, it suggests that much broader distributions should appear in two- rather than in three-dimensional search spaces. Since a horizontal span of the search space for albatrosses or sharks is much larger than vertical, perhaps such space could be considered as effectively two-dimensional and as our work suggests broad distributions might in such a case appear. It would be thus interesting to examine species for which search space is less anisotropic.

We acknowledge access to the computing facilities at Poznań Supercomputing and Networking Center.

- 
- [1] H. Pfister and H. Scher, *Adv. Phys.* **27**, 747 (1978).
  - [2] Y. Pomeau, A. Pumir, and W. R. Young, *Phys. Fluids A* **1**, 462 (1989).
  - [3] S. Havlin and D. ben Avram, *Adv. Phys.* **36**, 695 (1987).
  - [4] L. F. Richardson, *Proc. Roy. Soc. London A* **110**, 709 (1926). R. Metzler and J. Klafter, *Phys. Rep.* **339**, 1 (2000).
  - [5] A. Ott, J. P. Bouchaud, D. Langevin, and W. Urbach, *Phys. Rev. Lett.* **65**, 2201 (1990). R. Ganapathy, A. K. Sood, and S. Ramaswamy, *Europhys. Lett.* **77**, 18007 (2007).
  - [6] P. Dieterich, R. Klages, R. Preuss, and A. Schwab, *Proc. Nat. Acad. Sci.* **105**, 459 (2008).
  - [7] M. F. Shlesinger, in *On growth and Form* (eds. H. E. Stanley and N. Ostrowsky) (Nijhoff, Dordrecht, 1986).
  - [8] B. Cole, *J. Anim. Behav.* **50**, 1317 (1995).
  - [9] G. M. Viswanathan, V. Afanasyev, S. V. Buldyrev, E. J. Murphy, P. A. Prince, and H. E. Stanley, *Nature* **381**, 413 (1996).
  - [10] A. M. Edwards *et al.*, *Nature* **449**, 1044 (2007).
  - [11] J. Travis, *Science* **318**, 742 (2007).
  - [12] N. E. Humphries *et al.*, *Nature* **465**, 1066 (2010).
  - [13] D. L. Applegate, R. E. Bixby, V. Chvátal and W. J. Cook, *The traveling salesman problem: a computational study*, Princeton Series in Applied Mathematics. (Princeton University Press 2007).
  - [14] B. Li, N. Madras, and A. D. Sokal, *J. Stat. Phys.* **80**, 661 (1995).