

EVOLUTION OF THE MILKY WAY HALO BY ACCRETION OF DWARF SATELLITE GALAXIES

Mykola Petrov¹ and Gerhard Hensler¹

Abstract. Within the Cold Dark Matter scenario the hierarchical merging paradigm is the natural result to form massive galactic halos by the minor mergers of sub-halos and, by this, inherently their stellar halo. Although this must be also invoked for the Milky Way, the context of chemical and kinematic coherence of halo stars and dwarf spheroidal galaxies is yet unsolved a focus of present-day research. To examine this issue we model the chemo-dynamical evolution of the system of satellites selected from the cosmological Via Lactea II simulations to be similar for the Milky Way environment but at an early epoch.

1 Introduction

From the hierarchical merging paradigm and from numerical simulations of Λ CDM cosmology numerous Dark Matter subhalos are expected to assemble around a massive halo and to accumulate its mass. If they have already experienced star formation (SF), their stars should be merged into a spheroid and be identifiable by their kinematics and chemical abundances. Although these low-mass sub-halos and their baryonic content as dwarf spheroidal galaxies (dSphs) therefore serve as the key to pinpoint this cosmological Λ CDM description, observational detections of stellar streams within the Milky Way (MW) halo are rare though requested from accretion models (Johnston *et al.* 2008), the stellar abundances in present-day dSphs deviate clearly from the halo (Tolstoy *et al.* 2009), and evolutionary models of dSph SF, chemistry, and gas expulsion are still too simplistic. Not only that their accretion epoch occurred continuously over the Hubble time instead of a short event (Prantzos (2008)), and their gas is not only removed by tidal and ram-pressure stripping (Mayer *et al.* 2007) but is instead re-accreted while orbiting

This work was supported by the Initiative Kolleg (IK) 'The Cosmic Matter Circuit' I033-N program and computational time on the Vienna Scientific, Grape & Astro Clusters.

¹ Institute of Astronomy, University of Vienna, Austria; e-mail: mykola.petrov@univie.ac.at & e-mail: gerhard.hensler@univie.ac.at

the MW (Bouchard *et al.* 2009), also their consideration as isolated systems lacks reality (Revaz *et al.* 2009).

2 Modeling

To model the evolution of the system of dSphs in the gravitational field of the MW for which the accretion by the host galaxy is probable within the Hubble time, with the highest possible mass resolution the Via Lactea II (Diemand *et al.* 2008) simulation was used to follow 250 sub-haloes as DM progenitors of dSphs in the DM mass range of $10^6 M_\odot < M_{sat} < 6 \cdot 10^8 M_\odot$ from $z = 4.56$ for the first Gyr. Because the production and chemical evolution of 11 elements is intended to be treated within a proper computational time the number of gas particles is limited to 10^6 as also the DM. These facts limit the radius of consideration to within a radius of 40 kpc of the MW center of mass. For the simulations an advanced version of the single-gas chemo-dynamical SPH/N-body code (Petrov & Hensler 2010) is applied.

Starting in virial equilibrium of a 10^4 K warm gas, because re-ionization is improbable to have affected the Local Group dSphs (Grebel *et al.* 2004), cooling allows the gas particles to achieve SF conditions in all satellites (Fig. 1), but its efficiency directly depends on the mass of a satellite and its dynamical history (merging with other satellites or disruption by the MW potential). The stellar feedback by supernovae type II (SNeII) releases sufficient energy to expel hot gas from the main bodies of less massive dSphs facilitated by tidal interactions. This gas accumulates in the MW halo while massive dSphs merge and continue SF. In Fig. 1 (right panel) SF epoch is shown as function of the oxygen abundance of newly formed stars. For the first 10^8 yr of the simulation there is a considerable variance of stellar oxygen abundance in the whole system ($-5 \leq [O/H] \leq -0.5$) reflecting the very inhomogeneous production and distribution of enriched gas. After 10^8 yr the merger of sub-satellites ISM promotes the mixing of heavy elements. Finally, almost complete recycling of the gas erases the abundance inhomogeneities so that O in stars converges to $-1 \leq [O/H] \leq 0$ with a small dispersion.

In a comprehensive paper (Petrov & Hensler 2010) we will present detailed analyses of the SF history, gas exchange, stellar abundance evolution of dSphs and the MW halo in the early universe and will discuss their implications for our cosmological picture.

References

- Bouchard, A. *et al.* 2009, AJ, 137, 3038
 Diemand, J. *et al.* 2008, Nature, 454, 735
 Grebel, E.K. & Gallagher, J.S. 2004, ApJ, 610, L89
 Johnston, K. *et al.* 2008, ApJ, 689, 936
 Mayer, L. *et al.* 2007, Nature, 445, 738
 Petrov, M. & Hensler, G. 2010, ApJ, submitted
 Prantzos, N. 2008, A&A, 489, 525




Fig. 1. *Left:* SFR of the first 1Gyr as function of time and total mass of satellites. *Right:* Formation epoch t of stars with stellar oxygen abundance, $[O/H]$, colour-coded for the newly formed stellar mass (on logarithmic scales).

Revaz, Y. *et al.* 2009, A&A, 501, 189

Tolstoy, E., *et al.* 2009, ARA&A, 47, 371