ON BINARY CODES FROM CONICS IN PG(2, q)

ADONUS L. MADISON AND JUNHUA WU^{†,*}

ABSTRACT. Let **A** be the incidence matrix of passant lines and internal points with respect to a conic in PG(2, q), where q is an odd prime power. In this article, we study both geometric and algebraic properties of the column \mathbb{F}_2 -null space \mathcal{L} of **A**. In particular, using methods from both finite geometry and modular presentation theory, we manage to compute the dimension of \mathcal{L} , which provides a proof for the conjecture on the dimension of the binary code generated by \mathcal{L} .

1. INTRODUCTION

Let PG(2, q) be the classical projective plane of order q with underlying 3-dimensional vector space V over \mathbb{F}_q , the finite field of order q. Throughout this article, PG(2, q) is represented via homogeneous coordinates. Namely, a point is written as a non-zero vector (a_0, a_1, a_2) and a line is written as $[b_0, b_1, b_2]$ where not all b_i (i = 1, 2, 3) are zero. The set of points

$$\mathcal{O} := \{ (1, r, r^2) \mid r \in \mathbb{F}_q \} \cup \{ (0, 0, 1) \}$$
(1.1)

give rise to a geometric object called *conic* in PG(2, q) [7]. The above set also comprises the projective solutions of the nondegenarate quadratic equation

$$Q(X_0, X_1, X_2) = X_1^2 - X_0 X_2 \tag{1.2}$$

over \mathbb{F}_q . With respect to \mathcal{O} , the lines of $\mathrm{PG}(2,q)$ are partitioned into passant lines (Pa), tangent lines (T), and secant lines (Se) accordingly as the sizes of their intersections with \mathcal{O} are 0, 1, or 2. Similarly, points are partitioned into internal points (I), conic points (\mathcal{O}) , and external points (E) accordingly as the numbers of tangent lines on which they lie are 0, 1, or 2.

In [4], one low-density parity-check binary code was constructed using the column \mathbb{F}_{2} null space \mathcal{L} of the incidence matrix \mathbf{A} of passant lines and internal points with respect to \mathcal{O} . With the help of computer software Magma, the authors made a conjecture on the dimension of \mathcal{L} as follows:

Conjecture 1.1. [4, Conjecture 4.7] Let \mathcal{L} be the \mathbb{F}_2 -null space of \mathbf{A} , and let $\dim_{\mathbb{F}_2}(\mathcal{L})$ be the dimension of \mathcal{L} . Then

$$\dim_{\mathbb{F}_2}(\mathcal{L}) = \frac{(q-1)^2}{4}.$$

The goal of this article is to confirm Cojecture 1.1. Apart from the above conjecture, the dimensions of the column \mathbb{F}_2 -null spaces of the incidence matrices of external points versus secant lines, external points versus passant lines, and passant lines versus external points, were conjectured in the aforementioned paper [4], and have been established in [17] and [19], respectively. To start, we recall that the automorphism group G of \mathcal{O} is isomorphic to PGL(2, q), and that the normal subgroup H of G is isomorphic to PSL(2, q).

Key words and phrases. Block idempotent, Brauer's theory, character, conic, general linear group, incidence matrix, low-density parity-check code, module, 2-rank.

[†]Research supported in part by NSF HBCU-UP Grant Award #0929257 at Lane College.

^{*}Corresponding author.

MADISON AND WU

Let F be an algebraic closure of \mathbb{F}_2 . The idea of proving Conjecture 1.1 is to first realize \mathcal{L} as an FH-module and then decompose it into a direct sum of its certain submodules whose dimensions can be obtained easily. More concretely speaking, we view \mathbf{A} as the matrix of the following homomorphism ϕ of free F-modules:

$$\phi: F^I \to F^I$$

which first sends an internal point to the formal sum of all internal points on its polar, and then extends linearly to the whole of F^I . Additionally, it can be shown that ϕ is indeed an *FH*-module. Consequently, computing the dimension of the column \mathbb{F}_2 -null space amounts to finding the dimension of the *F*-null space of ϕ . To this end, we investigate the underlying *FH*-module structure of \mathcal{L} by applying Brauer's theory on the 2-blocks of *H* and arrive at a convenient decomposition of \mathcal{L} .

This article is organized in the following way. In Section 2, we establish that the matrix **A** satisfying the relation $\mathbf{A}^3 \equiv \mathbf{A} \pmod{2}$ under certain orderings of its rows and columns; this relation, in turn, reveals a geometric discription of $\operatorname{Ker}(\phi)$ as well as yields a set of generating elements of $\operatorname{Ker}(\phi)$ in terms of the concept of internal neighbors. In Section 3, the parity of intersection sizes of certain subsets of H with the conjugacy classes of H are computed. We will then in Section 4 review several facts about the 2-blocks of $\operatorname{PSL}(2,q)$ and the block idempotents of the 2-blocks of $\operatorname{PSL}(2,q)$; the detailed calculations of the 2-block idempotents were performed in [17]. Combining the results in Sections 3 and 4 with Brauer's theory on blocks, we are able to decompose $\operatorname{Ker}(\phi)$ into a direct sum of all non-isomorphic simple FH-modules or this sum plus a trivial module depending on q. Consequently, the dimension of \mathcal{L} follows as a lemma.

2. Geometry of Conics

First we recall several well-known results related to the geometry of conics in PG(2, q) with q odd. The books [8] and [7] are the references for what follows.

A collineation of PG(2, q) is an automorphism of PG(2, q), which is a bijection from the set of all points and all lines of PG(2, q) to itself that maps a point to a line and a line to a point, and preserves incidence. It is well known that each element of GL(3, q), the group of all 3×3 non-singular matrices over \mathbb{F}_q , induces a collineation of PG(2, q). The proof of the following lemma is straightforward.

Lemma 2.1. Let $\mathbf{P} = (a_0, a_1, a_2)$ and $\ell = [b_0, b_1, b_2]$ be a point and a line of $\mathrm{PG}(2, q)$, respectively. Suppose that θ is a collineation of $\mathrm{PG}(2, q)$ that is induced by $\mathbf{D} \in \mathrm{GL}(3, q)$. If we use \mathbf{P}^{θ} and ℓ^{θ} to denote the images of \mathbf{P} and ℓ under θ , respectively, then $\mathbf{P}^{\theta} = (a_0, a_1, a_2)^{\theta} = (a_0, a_1, a_2)\mathbf{D}$ and $\ell^{\theta} = [b_0, b_1, b_2]^{\theta} = [c_0, c_1, c_2]$, where c_0, c_1, c_2 correspond to the first, the second, and the third coordinate of the vector $\mathbf{D}^{-1}(b_0, b_1, b_2)^{\top}$, respectively.

A correlation of PG(2, q) is a bijection from the set of points to the set of lines as well as the set of lines to the set of points that reverses inclusion. A polarity of PG(2,q) is a correlation of order 2. The image of a point **P** under a correlation σ is denoted by \mathbf{P}^{σ} , and that of a line ℓ is denoted by ℓ^{σ} . It can be shown [7, p. 181] that the non-degenerate quadratic form $Q(X_0, X_1, X_2) = X_1^2 - X_0 X_2$ induces a polarity σ (or \bot) of PG(2,q), which can be represented by the matrix

$$\mathbf{M} = \begin{pmatrix} 0 & 0 & -\frac{1}{2} \\ 0 & 1 & 0 \\ -\frac{1}{2} & 0 & 0 \end{pmatrix}.$$
 (2.1)

Lemma 2.2. ([8, p. 47]) Let $\mathbf{P} = (a_0, a_1, a_2)$ and $\ell = [b_0, b_1, b_2]$) be a point and a line of PG(2, q), respetively. If σ is the polarity represented by the non-singular symmetric matrix \mathbf{M} in (2.1), then $\mathbf{P}^{\sigma} = (a_0, a_1, a_2)^{\sigma} = [c_0, c_1, c_2]$ and $\ell^{\sigma} = [b_0, b_1, b_2]^{\sigma} = (b_0, b_1, b_2)\mathbf{M}^{-1}$, where c_0, c_1, c_2 correspond to the first, the second, the third coordinate of the column vector $\mathbf{M}(a_0, a_1, a_2)^{\top}$, respectively.

For example, if $\mathbf{P} = (x, y, z)$ is a point of PG(2, q), then its image under σ is $\mathbf{P}^{\sigma} = [z, -2y, x]$.

For convenience, we will denote the set of all non-zero squares of \mathbb{F}_q by \Box_q , and the set of non-squares by \mathbb{Z}_q ; also, \mathbb{F}_q^* is the set of non-zero elements of \mathbb{F}_q .

Lemma 2.3. ([7, p. 181–182]) Assume that q is odd.

- (i) The polarity σ above defines three bijections; that is, $\sigma : I \to Pa, \sigma : E \to Se$, and $\sigma : \mathcal{O} \to T$ are all bijections.
- (ii) A line $[b_0, b_1, b_2]$ of PG(2, q) is a passant, a tangent, or a secant to \mathcal{O} if and only if $b_1^2 4b_0b_2 \in \square_q$, $b_1^2 4b_0b_2 = 0$, or $b_1^2 4b_0b_2 \in \square_q$, respectively.
- (iii) A point (a_0, a_1, a_2) of PG(2, q) is internal, absolute, or external if and only if $a_1^2 a_0 a_2 \in \not\square_q$, $a_1^2 a_0 a_2 = 0$, or $a_1^2 a_0 a_2 \in \square_q$, respectively.

The results in the following lemma can be obtained by simple counting; see [7] for more details and related results.

Lemma 2.4. ([7, p. 170]) Using the above notation, we have

$$|T| = |\mathcal{O}| = q+1, \ |Pa| = |I| = \frac{q(q-1)}{2}, \ and \ |Se| = |E| = \frac{q(q+1)}{2}.$$
 (2.2)

Also, we have the following tables:

TABLE 1. Number of points on lines of various types

Name	Absolute points	External points	Internal points
Tangent lines	1	q	0
Secant lines	2	$\frac{q-1}{2}$	$\frac{q-1}{2}$
Passant lines	0	$\frac{q\ddot{+}1}{2}$	$\frac{q+1}{2}$

TABLE 2. Number of lines through points of various types

Name	Tangent lines	Secant lines	Skew lines
Absolute points	1	q	0
External points	2	$\frac{q-1}{2}$	$\frac{q-1}{2}$
Internal points	0	$\frac{q+1}{2}$	$\frac{q+1}{2}$

2.1. The incidence matrix. Let G be the automorphism group of \mathcal{O} in PGL(3, q) (i.e. the subgroup of PGL(3, q) fixing \mathcal{O} setwise).

Lemma 2.5. [7, p. 158] $G \cong PGL(2, q)$.

We define

$$H := \left\{ \begin{pmatrix} a^2 & ab & b^2 \\ 2ac & ad + bc & 2bd \\ c^2 & cd & d^2 \end{pmatrix} \middle| a, b, c, d \in \mathbb{F}_q, ad - bc = 1 \right\}.$$
 (2.3)

In the rest of the article, we always use ξ to denote a fixed primitive element of \mathbb{F}_q . For $a, b, c \in \mathbb{F}_q$, we define

$$\mathbf{d}(a,b,c) := \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}, \ \mathbf{ad}(a,b,c) := \begin{pmatrix} 0 & 0 & a \\ 0 & b & 0 \\ c & 0 & 0 \end{pmatrix}.$$

For the convenience of discussion, we adopt the following special representatives of G from [17]:

$$H \cup \mathbf{d}(1, \xi^{-1}, \xi^{-2}) \cdot H.$$
 (2.4)

Moreover, the following holds.

Lemma 2.6. [6] The group G acts transitively on both I (respectively, Pa) and E (respectively, Se).

The following result was proved in [17] and will be used frequently.

Lemma 2.7. [17, Lemma 2.9] Let **P** be a point not on \mathcal{O} , ℓ a non-tangent line, and $\mathbf{P} \in \ell$. Using the above notation, we have the following.

- (i) If $\mathbf{P} \in I$ and $\ell \in Pa$, then $\mathbf{P}^{\perp} \cap \ell \in E$ if $q \equiv 1 \pmod{4}$, and $\mathbf{P}^{\perp} \cap \ell \in I$ if $q \equiv 3 \pmod{4}$.
- (ii) If $\mathbf{P} \in I$ and $\ell \in Se$, then $\mathbf{P}^{\perp} \cap \ell \in I$ if $q \equiv 1 \pmod{4}$, and $\mathbf{P}^{\perp} \cap \ell \in E$ if $q \equiv 3 \pmod{4}$.
- (iii) If $\mathbf{P} \in E$ and $\ell \in Pa$, then $\mathbf{P}^{\perp} \cap \ell \in I$ if $q \equiv 1 \pmod{4}$, and $\mathbf{P}^{\perp} \cap \ell \in E$ if $q \equiv 3 \pmod{4}$.
- (iv) If $\mathbf{P} \in E$ and $\ell \in Se$, then $\mathbf{P}^{\perp} \cap \ell \in E$ if $q \equiv 1 \pmod{4}$, and $\mathbf{P}^{\perp} \cap \ell \in I$ if $q \equiv 3 \pmod{4}$.

Next we define $\Box_q - 1 := \{s - 1 \mid s \in \Box_q\}$ and $\not\Box_q - 1 := \{s - 1 \mid s \in \not\Box_q\}.$

Lemma 2.8. [18] Using the above notation,

(i) if
$$q \equiv 1 \pmod{4}$$
, then $|(\Box_q - 1) \cap \Box_q| = \frac{q-5}{4}$ and
 $|(\Box_q - 1) \cap \Box_q| = |(\Box_q - 1) \cap \Box_q| = |(\Box_q - 1) \cap \Box_q| = \frac{q-1}{4};$
(ii) if $q \equiv 3 \pmod{4}$, then $|(\Box_q - 1) \cap \Box_q| = \frac{q+1}{4}$ and

(ii) if $q \equiv 3 \pmod{4}$, then $|(\not\square_q - 1) \cap \square_q| = \frac{q+1}{4}$ and

$$|(\square_q - 1) \cap \square_q| = |(\square_q - 1) \cap \not\square_q| = |(\not\square_q - 1) \cap \not\square_q| = \frac{q - 3}{4}.$$

Definition 2.9. Let \mathbf{P} be a point not on \mathcal{O} and ℓ a line. We define E_{ℓ} and I_{ℓ} to be the set of external points and the set of internal points on ℓ , respectively, $Pa_{\mathbf{P}}$ and $Se_{\mathbf{P}}$ the set of passant lines and the set of secant lines through \mathbf{P} , respectively, and $T_{\mathbf{P}}$ the set of tangent lines through \mathbf{P} . Also, $N(\mathbf{P})$ is defined to be the set of internal points on the passant lines through \mathbf{P} including or excluding \mathbf{P} accordingly as $q \equiv 3 \pmod{4}$ or $q \equiv 1 \pmod{4}$.

Remark 2.10. Using the above notation, for $\mathbf{P} \in I$, we have $|E_{\mathbf{P}^{\perp}}| = |Se_{\mathbf{P}}| = \frac{q+1}{2}$; $|I_{\mathbf{P}^{\perp}}| = |Pa_{\mathbf{P}}| = \frac{q+1}{2}$; and $|N(\mathbf{P})| = \frac{q^2-1}{4}$ or $\frac{q^2+3}{4}$ accordingly as $q \equiv 1 \pmod{4}$ or $q \equiv 3 \pmod{4}$.

Let $\mathbf{P} \in I$, $\ell \in Pa$, $g \in G$, and $W \leq G$. Using standard notations from permutation group theory, we have $I_{\ell}^g = I_{\ell^g}$, $Pa_{\mathbf{P}}^g = Pa_{\mathbf{P}^g}$; $E_{\ell}^g = E_{\ell^g}$, $Se_{\mathbf{P}}^g = Se_{\mathbf{P}^g}$; $H_{\mathbf{P}}^g = H_{\mathbf{P}^g}$; $N(\mathbf{P})^g = N(\mathbf{P}^g)$, $(W^g)_{\mathbf{P}^g} = W_{\mathbf{P}}^g$. We will use these results later without further reference. Also, the definition of G yields that $(\mathbf{P}^{\perp})^g = (\mathbf{P}^g)^{\perp}$, where \perp is the above defined polarity of $\mathrm{PG}(2,q)$. **Proposition 2.11.** Let $\mathbf{P} \in I$ and set $K := G_{\mathbf{P}}$. Then K is transitive on $I_{\mathbf{P}^{\perp}}$, $E_{\mathbf{P}^{\perp}}$, $Pa_{\mathbf{P}}$, and $Se_{\mathbf{P}}$, respectively.

Proof: Witt's theorem [9] implies that K acts transitively on isometry classes of the form Q on the points of \mathbf{P}^{\perp} . Note that $K = G_{\mathbf{P}^{\perp}}$ by the definition of G. Dually, we must have that K is transitive on both $Pa_{\mathbf{P}}$ and $Se_{\mathbf{P}}$.

When $\mathbf{P} = (1, 0, -\xi)$, using (2.2), (2.3), and (2.4), we obtain that $K := G_{\mathbf{P}} =$

$$\left\{ \left(\begin{array}{cccc} d^{2} & cd\xi & c^{2}\xi^{2} \\ 2cd & d^{2} + c^{2}\xi & 2dc\xi \\ c^{2} & dc & d^{2} \end{array} \right) \middle| d, c \in \mathbb{F}_{q}, d^{2} - c^{2}\xi = 1 \right\} \\
\bigcup \left\{ \left(\begin{array}{cccc} d^{2} & -cd\xi & c^{2}\xi^{2} \\ 2cd & -d^{2} - c^{2}\xi & 2dc\xi \\ c^{2} & -dc & d^{2} \end{array} \right) \middle| d, c \in \mathbb{F}_{q}, -d^{2} + c^{2}\xi = 1 \right\} \\
\bigcup \left\{ \left(\begin{array}{cccc} d^{2} & cd & c^{2} \\ 2cd\xi^{-1} & d^{2} + c^{2}\xi^{-1} & 2dc \\ c^{2}\xi^{-2} & dc\xi^{-1} & d^{2} \end{array} \right) \middle| d, c \in \mathbb{F}_{q}, d^{2}\xi - c^{2} = 1 \right\} \\
\bigcup \left\{ \left(\begin{array}{cccc} d^{2} & -cd & c^{2} \\ 2cd\xi^{-1} & d^{2} + c^{2}\xi^{-1} & 2dc \\ c^{2}\xi^{-2} & dc\xi^{-1} & d^{2} \end{array} \right) \middle| d, c \in \mathbb{F}_{q}, -d^{2}\xi + c^{2} = 1 \right\} .$$

$$(2.5)$$

Recall that $N(\mathbf{P})$ for $\mathbf{P} \in I$ is the set of internal points on the passant lines through \mathbf{P} , where \mathbf{P} is included or not accordingly as $q \equiv 3 \pmod{4}$ or $q \equiv 1 \pmod{4}$.

Theorem 2.12. Let $\mathbf{P} \in I$ and $\ell \in Pa$. Then $|N(\mathbf{P}) \cap I_{\ell}| \equiv 0 \pmod{2}$.

Proof: If $\mathbf{P} \in \ell$, it is clear that

$$|N(\mathbf{P}) \cap I_{\ell}| = \begin{cases} \frac{q-1}{2}, & \text{if } q \equiv 1 \pmod{4}, \\ \frac{q+1}{2}, & \text{if } q \equiv 3 \pmod{4}, \end{cases}$$

which is even. Therefore, $|N(\mathbf{P}) \cap I_{\ell}| \equiv 0 \pmod{2}$ for this case.

If $\ell = \mathbf{P}^{\perp}$, by Lemma 2.7(i), we have

$$|N(\mathbf{P}) \cap I_{\ell}| = \begin{cases} 0, & \text{if } q \equiv 1 \pmod{4}, \\ \frac{q+1}{2}, & \text{if } q \equiv 3 \pmod{4}, \end{cases}$$

which is even. Hence, $|N(\mathbf{P}) \cap I_{\ell}| \equiv 0 \pmod{2}$ for this case.

Now we assume that neither $\ell = \mathbf{P}^{\perp}$ nor $\mathbf{P} \in \ell$. As G is transitive on Pa and preserves incidence, we may take $\ell = \mathbf{P}_1^{\perp} = [1, 0, -\xi^{-1}]$, where $\mathbf{P}_1 = (1, 0, -\xi) \in I$. Since **P** is either on a passant line through \mathbf{P}_1 or on a secant line through \mathbf{P}_1 , the rest it to show that $|N(\mathbf{P}) \cap I_{\ell}|$ is even for any **P** on a line through \mathbf{P}_1 with $\mathbf{P} \notin \ell$ and $\mathbf{P} \neq \mathbf{P}_1$.

Case I. P is a point on a secant line through \mathbf{P}_1 and $\mathbf{P} \notin \ell$.

Since $K = G_{\mathbf{P}_1}$ acts transitively on $Se_{\mathbf{P}_1}$ by Proposition 2.11, it is enough to establish that $|N(\mathbf{P}) \cap I_{\ell}|$ is even for an arbitrary internal point on a *special* secant line, ℓ_1 say, through \mathbf{P}_1 . To this end, we may take $\ell_1 = [0, 1, 0]$. It is clear that

$$I_{\ell_1} = \{ (1, 0, -\xi^j) \mid 0 \le j \le q - 1, \ j \text{ odd} \}$$

and

$$I_{\ell} = \{ (1, s, \xi) \mid s \in \mathbb{F}_q, \ s^2 - \xi \in \mathcal{P}_q \}.$$

Hence, if $\mathbf{P} = (1, 0, -\xi^j) \in I_{\ell_1}$ then

$$D_j = \left\{ \left[1, -\frac{\xi^{1-j}+1}{s}, \frac{1}{\xi^j} \right] \middle| s \in \mathbb{F}_q^*, \ s^2 - \xi \in \mathcal{P}_q \right\} \cup \{ [0, 1, 0] \}$$

consists of the lines through both \mathbf{P} and the points on ℓ . Note that the number of passant lines in D_j is determined by the number of s satisfying both

$$\frac{1}{s^2}(\xi^{1-j}+1)^2 - \frac{4}{\xi^j} \in \mathcal{P}_q$$
(2.6)

and

Since, $s \neq 0$ and whenver s satisfies both (2.6) and (2.7), so does -s, we see that $|N(\mathbf{P}) \cap I_{\ell}|$ must be even in this case.

Case II. P is an internal point on a passant line through \mathbf{P}_1 and $\mathbf{P} \notin \ell$.

By Lemma 2.7, we may assume that $\mathbf{P} \in \mathbf{P}_3^{\perp}$, where $\mathbf{P}_3 = (1, x, \xi) \in I_{\ell}$ with $x \in \mathbb{F}_q^*$ and $x^2 - \xi \in \mathbb{P}_q$. Here $\mathbf{P}_3^{\perp} = [1, -\frac{2x}{\xi}, \frac{1}{\xi}]$ is a passant line through \mathbf{P}_1 . Let $K = G_{\mathbf{P}_1}$. Using (2.5), we can obtain $L := K_{\mathbf{P}_3}$ as follows: if $q \equiv 1 \pmod{4}$,

$$L = \left\{ \begin{pmatrix} \frac{\xi}{x^2 - \xi} & \frac{x\xi}{x^2 - \xi} & \frac{x^2\xi}{x^2 - \xi} \\ -\frac{2x}{x^2 - \xi} & -\frac{2x\xi}{x^2 - \xi} & -\frac{2x\xi}{x^2 - \xi} \\ \frac{x^2}{(x^2 - \xi)\xi} & \frac{x}{x^2 - \xi} & \frac{\xi}{x^2 - \xi} \end{pmatrix}, \begin{pmatrix} 0 & 0 & -1 \\ 0 & -\frac{1}{\xi} & 0 \\ -\frac{1}{\xi^2} & 0 & 0 \end{pmatrix} \right\}$$
$$\bigcup \left\{ \begin{pmatrix} -\frac{x^2}{(x^2 - \xi)\xi} & -\frac{x}{x^2 - \xi} & -\frac{\xi}{x^2 - \xi} \\ \frac{2x}{(x^2 - \xi)\xi} & \frac{x^2 + \xi}{(x^2 - \xi)\xi} & \frac{2x}{x^2 - \xi} \\ -\frac{1}{(x^2 - \xi)\xi} & -\frac{x}{(x^2 - \xi)\xi} & -\frac{x^2}{(x^2 - \xi)\xi} \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\};$$
(2.8)

if $q \equiv 3 \pmod{4}$,

$$L = \left\{ \begin{pmatrix} \frac{\xi}{x^2 - \xi} & \frac{x\xi}{x^2 - \xi} & \frac{x^2\xi}{x^2 - \xi} \\ -\frac{2x}{x^2 - \xi} & -\frac{x^2 + \xi}{x^2 - \xi} & -\frac{2x\xi}{x^2 - \xi} \\ \frac{x^2}{x^2 - \xi} & \frac{x}{x^2 - \xi} & \frac{\xi}{x^2 - \xi} \end{pmatrix}, \begin{pmatrix} 0 & 0 & -\xi \\ 0 & -1 & 0 \\ -\frac{1}{\xi} & 0 & 0 \end{pmatrix} \right\}$$
$$\cup \left\{ \begin{pmatrix} -\frac{x^2}{x^2 - \xi} & -\frac{x\xi}{x^2 - \xi} & -\frac{\xi^2}{x^2 - \xi} \\ \frac{2x}{x^2 - \xi} & \frac{x^2 + \xi}{x^2 - \xi} & \frac{2x\xi}{x^2 - \xi} \\ -\frac{1}{x^2 - \xi} & -\frac{x}{x^2 - \xi} & -\frac{x^2}{x^2 - \xi} \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\}.$$
(2.9)

Let $(1, y, \xi)$ be a point on ℓ . Using (2.8) and (2.9), we have that L fixes $(1, y, \xi)$ if and only if

$$xy^2 - (x^2 + \xi)y + x\xi = 0;$$

that is, y = x or $y = \frac{\xi}{x}$. Consequently, $\mathbf{P}_3 = (1, x, \xi)$ and $\ell \cap \mathbf{P}_3^{\perp} = (1, \frac{\xi}{x}, \xi)$ are the only points of the form (1, s, t) on ℓ fixed by L. Since $\mathbf{P} \in \mathbf{P}_3^{\perp}$, $\mathbf{P} \neq \mathbf{P}_1$ and $\mathbf{P} \neq \mathbf{P}_3^{\perp} \cap \ell$, $\mathbf{P} = (1, \frac{\xi+n}{2x}, n)$ for some $n \neq \xi$. Now if we denote by \mathbf{V} the set of passant lines through \mathbf{P} that meet ℓ in an internal point, then it is clear that $|\mathbf{V}| = |N(\mathbf{P}) \cap I_{\ell}|$. Direct computations give us that $L_{\mathbf{P}} \cong \mathbb{Z}_2$. Since \mathbf{P}_3 and \mathbf{P} are both fixed by $L_{\mathbf{P}}$, it follows that both $\ell_{\mathbf{P}_3,\mathbf{P}}$ and \mathbf{P}_3^{\perp} are fixed by $L_{\mathbf{P}}$. Note that when $q \equiv 3 \pmod{4}$, both \mathbf{P}_3^{\perp} and $\ell_{\mathbf{P}_3,\mathbf{P}}$ are in \mathbf{V} ; and when $q \equiv 1 \pmod{4}$, neither $\ell_{\mathbf{P}_3,\mathbf{P}}$ nor \mathbf{P}_3^{\perp} is in \mathbf{V} . If there were another line ℓ' through \mathbf{P} which is distinct from both \mathbf{P}_3^{\perp} and $\ell_{\mathbf{P}_3,\mathbf{P}}$ and which is also fixed by $L_{\mathbf{P}}$, then $L_{\mathbf{P}}$ would fix at least three points on $\ell = \mathbf{P}^{\perp}$, namely, $\ell' \cap \ell$, $\mathbf{P}_3^{\perp} \cap \ell$, and \mathbf{P}_3 . Since no further point of the form (1, s, t) except for \mathbf{P}_3 and $\ell \cap \mathbf{P}_3^{\perp}$ can be fixed by L due to the above discussion, we must have $\ell' \cap \ell = (0, 1, 0) \in E_{\ell}$. So $\ell' \notin \mathbf{V}$. Using the fact that $L_{\mathbf{P}}$ preserves incidence, we conclude that when $q \equiv 1 \pmod{4}$, $L_{\mathbf{P}}$ has $\frac{|\mathbf{V}|}{2}$ orbits of length 2 on \mathbf{V} ; and when $q \equiv 3 \pmod{4}$, $L_{\mathbf{P}}$ has two orbits of length 1, namely, $\{\mathbf{P}_3^{\perp}\}$ and $\{\ell_{\mathbf{P}_3,\mathbf{P}}\}$, and $\frac{|\mathbf{V}|-2}{2}$ orbits of length 2 on **V**. Either forces $|\mathbf{V}|$ to be even. Therefore, $|N(\mathbf{P}) \cap I_{\ell}|$ is even.

Recall that **A** is the incidence matrix of Pa and I whose columns are indexed by the internal points $\mathbf{P}_1, \mathbf{P}_2, ..., \mathbf{P}_N$ and whose rows are indexed by the passant lines $\mathbf{P}_1^{\perp}, \mathbf{P}_2^{\perp}, ..., \mathbf{P}_N^{\perp}$; and **A** is symmetric. For the convenience of discussion, for $\mathbf{P} \in I$, we define

$$\widehat{N(\mathbf{P})} = \begin{cases} N(\mathbf{P}) \cup \{\mathbf{P}\}, & \text{if } q \equiv 1 \pmod{4}, \\ N(\mathbf{P}) \setminus \{\mathbf{P}\}, & \text{if } q \equiv 3 \pmod{4}. \end{cases}$$

That is, $\widehat{N(\mathbf{P})}$ is the set of the internal points on the passant lines through \mathbf{P} including \mathbf{P} . It is clear that for $\mathbf{P} \notin \ell$, $|N(\mathbf{P}) \cap I_{\ell}| = |\widehat{N(\mathbf{P})} \cap I_{\ell}|$.

Lemma 2.13. Using the above notation, we have $\mathbf{A}^3 \equiv \mathbf{A} \pmod{2}$, where the congruence means entry-wise congruence.

Proof: Since the (i, j)-entry of $\mathbf{A}^2 = \mathbf{A}^\top \mathbf{A}$ is the standard dot product of the *i*-th row of \mathbf{A}^\top and *j*-th column of \mathbf{A} , we have

$$(\mathbf{A}^2)_{i,j} = (\mathbf{A}^\top \mathbf{A})_{i,j} = \begin{cases} \frac{q+1}{2}, & \text{if } i = j, \\ 1, & \text{if } \ell_{\mathbf{P}_i, \mathbf{P}_j} \in Pa, \\ 0, & \text{otherwise.} \end{cases}$$

Therefore, the *i*-th row of $\mathbf{A}^2 \pmod{2}$ indexed by \mathbf{P}_i can be viewed as the characteristic row vector of $\widehat{N(\mathbf{P}_i)}$.

If $\mathbf{P}_i \in \mathbf{P}_j^{\perp}$, then $(\mathbf{A}^3)_{i,j} = ((\mathbf{A}^2)\mathbf{A}^{\top})_{i,j} = q$ since $(\mathbf{A}^2)_{i,i} = \frac{q+1}{2}$ and there are $\frac{q-1}{2}$ internal points other than \mathbf{P}_i on \mathbf{P}_j^{\perp} that are connected with \mathbf{P}_i by the passant line \mathbf{P}_j^{\perp} . If $\mathbf{P}_i \notin \mathbf{P}_j^{\perp}$, then $(\mathbf{A}^3)_{i,j} = ((\mathbf{A}^{\top}\mathbf{A})\mathbf{A}^{\top})_{i,j} \equiv |\widehat{N(\mathbf{P}_i)} \cap I_{\mathbf{P}_j^{\perp}}| = |N(\mathbf{P}_i) \cap I_{\mathbf{P}_j^{\perp}}| \equiv 0 \pmod{2}$ by Theorem 2.12. Consequently,

$$(\mathbf{A}^3)_{i,j} \equiv \begin{cases} 1 \pmod{2}, & \text{if } \mathbf{P}_i \in \mathbf{P}_j^{\perp}, \\ 0 \pmod{2}, & \text{if } \mathbf{P}_i \notin \mathbf{P}_j^{\perp}. \end{cases}$$

The lemma follows immediately.

3. The Conjugacy Classes and Intersection Parity

In this section, we present detailed information about the conjugacy classes of H and study their intersections with some special subsets of H.

3.1. Conjugacy classes. The conjugacy classes of H can be read off in terms of the map T = tr(g) + 1, where tr(g) is the trace of g.

Lemma 3.1. [17, Lemma 3.2] The conjugacy classes of H are given as follows.

- (i) $D = \{ \mathbf{d}(1, 1, 1) \};$
- (ii) F^+ and F^- , where $F^+ \cup F^- = \{g \in H \mid T(g) = 4, g \neq \mathbf{d}(1, 1, 1)\};$
- (iii) $[\theta_i] = \{g \in H \mid T(g) = \theta_i\}, 1 \le i \le \frac{q-5}{4} \text{ if } q \equiv 1 \pmod{4}, \text{ or } 1 \le i \le \frac{q-3}{4} \text{ if } q \equiv 3 \pmod{4}, \text{ or } 1 \le i \le \frac{q-3}{4} \text{ if } q \equiv 3 \pmod{4}, \text{ where } \theta_i \in \Box_q, \theta_i \ne 4, \text{ and } \theta_i 4 \in \Box_q;$
- (iv) $[0] = \{g \in H \mid T(g) = 0\};$
- (v) $[\pi_k] = \{g \in H \mid T(g) = \pi_k\}, 1 \le k \le \frac{q-1}{4} \text{ if } q \equiv 1 \pmod{4}, \text{ or } 1 \le k \le \frac{q-3}{4} \text{ if } q \equiv 3 \pmod{4}, \text{ where } \pi_i \in \Box_q, \pi_k \neq 4, \text{ and } \pi_k 4 \in \not\Box_q.$

Remark 3.2. The set $F^+ \cup F^-$ forms one conjugacy class of G, and splits into two equalsized classes F^+ and F^- of H. For our purpose, we denote $F^+ \cup F^-$ by [4]. Also, each of D, $[\theta_i]$, [0], and $[\pi_k]$ forms a single conjugacy class of G. The class [0] consists of all the elements of order 2 in H.

In the following, for convenience, we frequently use C to denote any one of D, [0], [4], $[\theta_i]$, or $[\pi_k]$. That is,

$$C = D, [0], [4], [\theta_i], \text{ or } [\pi_k].$$
 (3.1)

3.2. Intersection properties.

Definition 3.3. Let $\mathbf{P}, \mathbf{Q} \in I$, $W \subseteq I$, and $\ell \in Pa$. We define $\mathcal{H}_{\mathbf{P},\mathbf{Q}} = \{h \in H \mid (\mathbf{P}^{\perp})^h \in Pa_{\mathbf{Q}}\}$, $\mathcal{S}_{\mathbf{P},\ell} = \{h \in H \mid (\mathbf{P}^{\perp})^h = \ell\}$, and $\mathcal{U}_{\mathbf{P},W} = \{h \in H \mid \mathbf{P}^h \in W\}$. That is, $\mathcal{H}_{\mathbf{P},\mathbf{Q}}$ consists of all the elements of H that map the passant line \mathbf{P}^{\perp} to a passant line through $\mathbf{Q}, \mathcal{S}_{\mathbf{P},\ell}$ is the set of elements of H that map \mathbf{P}^{\perp} to the passant line ℓ , and $\mathcal{U}_{\mathbf{P},W}$ is the set of elements of H that map \mathbf{P} .

Using the above notation, we have that $\mathcal{H}^{g}_{\mathbf{P},\mathbf{Q}} = \mathcal{H}_{\mathbf{P}^{g},\mathbf{Q}^{g}}, \ \mathcal{S}^{g}_{\mathbf{P},\ell} = \mathcal{S}_{\mathbf{P}^{g},\ell^{g}}, \ \text{and} \ \mathcal{U}^{g}_{\mathbf{P},W} = \mathcal{U}_{\mathbf{P}^{g},W^{g}}, \ \text{where} \ \mathcal{H}^{g}_{\mathbf{P},\mathbf{Q}} = \{g^{-1}hg \mid h \in \mathcal{H}_{\mathbf{P},\mathbf{Q}}\}, \ \mathcal{S}^{g}_{\mathbf{P},\ell} = \{g^{-1}hg \mid h \in \mathcal{S}_{\mathbf{P},\mathbf{Q}}\}, \ \text{and} \ \mathcal{U}^{g}_{\mathbf{P},W} = \{h^{g} \mid h \in \mathcal{U}_{\mathbf{P},W}\}.$ Moreover, it is true that $(C \cap \mathcal{H}_{\mathbf{P},\mathbf{Q}})^{g} = C \cap \mathcal{H}_{\mathbf{P}^{g},\mathbf{Q}^{g}}$ and $(C \cap \mathcal{U}_{\mathbf{P},W})^{g} = C \cap \mathcal{U}_{\mathbf{P}^{g},W^{g}}.$ In the following discussion, we will use these results without further reference.

Next we compute the size of the intersection of each conjugacy class of H with K which is a stabilizer of an internal point in H.

Corollary 3.4. Let $\mathbf{P} \in I$ and $K = H_{\mathbf{P}}$. Then we have

- (i) $|K \cap D| = 1;$ (ii) $|K \cap [4]| = 0;$
- (iii) $|K \cap [\pi_k]| = 2;$
- (iv) $|K \cap [\theta_i]| = 0;$
- (v) $|K \cap [0]| = \frac{q+1}{2}$ or $\frac{q-1}{2}$ accordingly as $q \equiv 1 \pmod{4}$ or $q \equiv 3 \pmod{4}$.

Proof: Let $\mathbf{Q} = (1, 0, -\xi)$ and $K_1 = H_{\mathbf{Q}}$. As H is transitive on I, $\mathbf{Q}^g = \mathbf{P}$ for some $g \in H$. Moreover,

$$K \cap C| = |K_1^g \cap C| = |(K_1 \cap C)^g|$$

Therefore, to prove the corollary, it is sufficient to assume that $\mathbf{P} = \mathbf{Q}$.

It is obvious that $|D \cap K| = 1$. Let $g \in K \cap C$. Using (2.5), we have that the quadruples (a, b, c, d) determining g satisfy the following equations

$$\begin{aligned}
bd - ac\xi &= 0 \\
b^2 - a^2\xi &= -\xi(d^2 - c^2\xi) \\
ad - bc &= 1 \\
a + d &= s,
\end{aligned}$$
(3.2)

where $s^2 = 0, 4, \pi_k, \theta_i$. The equations in (3.2) yield (1) $a = d = \frac{s}{2}, c^2 = \frac{s^2-4}{4\xi}, b^2 = \frac{(s^2-4)\xi}{4}$ and (2) $a = -d, s = 0, c^2\xi - 1 = a^2$. From Case (1), we have $|K \cap [\pi_k]| = 2$ for each $[\pi_k]$ and $|K \cap C| = 0$ for $C = [\theta_i]$, [4]; moreover, if $q \equiv 3 \pmod{4}$, we obtain one group element $\operatorname{ad}(-\xi, -1, \xi^{-1}) \in K \cap [0]$ in Case (1). Since the number of $t \in \not{\square}_q$ satisfying $t - 1 \in \square_q$ is $\frac{q-1}{4}$ or $\frac{q-3}{4}$ accordingly as $q \equiv 1 \pmod{4}$ or $q \equiv 3 \pmod{4}$ by Lemma 2.10, the number of $c \in \mathbb{F}_q^*$ satisfying $c^2\xi - 1 \in \square_q$ is $2|(\not{\square}_q - 1) \cap \square_q|$ which is $\frac{q-1}{2}$ or $\frac{q-3}{2}$ accordingly as $q \equiv 1 \pmod{4}$. When $q \equiv 1 \pmod{4}$, c = 0 also satisfies $c^2\xi - 1 \in \square_q$. Therefore, Case (1) and Case (2) give rise to $\frac{q+1}{2}$ or $\frac{q-1}{2}$ different group elements in $K \cap [0]$ depending on q. The corollary now is proved. In the following lemmas, we investigate the parity of $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ for $C \neq [0]$ and $\mathbf{P}, \mathbf{Q} \in I$. Recall that $\ell_{\mathbf{P},\mathbf{Q}}$ is the line through \mathbf{P} and \mathbf{Q} .

Lemma 3.5. Let $\mathbf{P}, \mathbf{Q} \in I$. Suppose that C = D, [4], $[\pi_k] (1 \le k \le \frac{q-1}{4})$, or $[\theta_i] (1 \le i \le \frac{q-5}{4})$.

First assume that $q \equiv 1 \pmod{4}$.

- (i) If $\ell_{\mathbf{P},\mathbf{Q}} \in Pa_{\mathbf{P}}$, then $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is always even.
- (ii) If $\ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$, $\mathbf{Q} \notin \mathbf{P}^{\perp}$, and $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is odd, then $C = [\theta_{i_1}]$ or $[\theta_{i_2}]$.
- (iii) If $\mathbf{Q} \in \ell_{\mathbf{P},\mathbf{Q}} \cap \mathbf{P}^{\perp}$ and $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is odd, then C = D.

Now assume that $q \equiv 3 \pmod{4}$.

- (iv) If $\ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$, then $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is always even.
- (v) If $\ell_{\mathbf{P},\mathbf{Q}} \in Pa_{\mathbf{P}}$, $\mathbf{Q} \notin \mathbf{P}^{\perp}$, and $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is odd, then $C = [\pi_{i_1}]$ or $[\pi_{i_2}]$.

(vi) If $\mathbf{Q} \in \ell_{\mathbf{P},\mathbf{Q}} \cap \mathbf{P}^{\perp}$ and $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is odd, then C = D.

Proof: We only provide the detailed proof for the case when $q \equiv 1 \pmod{4}$. Since G acts transitively on I and preserves incidence, without loss of generality, we may assume that $\mathbf{P} = (1, 0, -\xi)$ and let $K = G_{\mathbf{P}}$.

Since K is transitive on both $Pa_{\mathbf{P}}$ and $Se_{\mathbf{P}}$ by Proposition 2.11 and $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C| = |(\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C)^g| = |\mathcal{H}_{\mathbf{P},\mathbf{Q}^g} \cap C|$, we may assume that \mathbf{Q} is on either ℓ_1 or ℓ_2 , where $\ell_1 = [1,0,\xi^{-1}] \in Pa_{\mathbf{P}}$ and $\ell_2 = [0,1,0] \in Se_{\mathbf{P}}$.

Case I. $\mathbf{Q} \in \ell_1$.

In this case, $\mathbf{Q} = (1, x, -\xi)$ for some $x \in \mathbb{F}_q^*$ and $x^2 + \xi \in \mathbb{P}_q$, and

$$Pa_{\mathbf{Q}} = \{ [1, s, (1+sx)\xi^{-1}] \mid s \in \mathbb{F}_q, s^2 - 4(1+sx)\xi^{-1} \in \mathcal{P}_q \}$$

Using (2.5), we obtain that

$$K_{\mathbf{Q}} = \{\mathbf{d}(1,1,1), \mathbf{ad}(1,-\xi^{-1},\xi^{-2})\}$$

It is obvious that $\mathbf{d}(1,1,1)$ fixes each line in $Pa_{\mathbf{Q}}$. From

$$\mathbf{ad}(1, -\xi^{-1}, \xi^{-2})^{-1}(1, s, (1+sx)\xi^{-1})^{\top} = ((1+sx)\xi, -s\xi, 1)^{\top},$$

it follows that a line of the form $[1, s, (1 + sx)\xi^{-1}]$ is fixed by $K_{\mathbf{Q}}$ if and only if s = 0 or $s = -2x^{-1}$. Further, since $[1, -2x^{-1}, -\xi^{-1}]$ is a secant line, we obtain that $K_{\mathbf{Q}}$ on $Pa_{\mathbf{Q}}$ has one orbit of length 1, i.e. $\{\ell_1 = [1, 0, \xi^{-1}]\}$, and all other orbits, whose representatives are \mathcal{R}_1 , have length 2. From

$$|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C| = |\mathcal{S}_{\mathbf{P},\ell_1} \cap C| + 2\sum_{\ell \in \mathcal{R}_1} |\mathcal{S}_{\mathbf{P},\ell} \cap C|$$

it follows that the parity of $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is determined by that of $|\mathcal{S}_{\mathbf{P},\ell_1} \cap C|$. Here we used the fact that $|\mathcal{S}_{\mathbf{P},\ell} \cap C| = |\mathcal{S}_{\mathbf{P},\ell'} \cap C|$ if $\{\ell,\ell'\}$ is an orbit of $K_{\mathbf{P}}$ on $Pa_{\mathbf{Q}}$. Meanwhile, it is clear that $|\mathcal{S}_{\mathbf{P},\ell_1} \cap D| = 0$.

Note that the quadruples (a, b, c, d) that determine group elements in $\mathcal{S}_{\mathbf{P},\ell_1} \cap C$ are the solutions to the following equations

$$\begin{array}{rcl} -2cd + 2ab\xi^{-1} &= & 0\\ c^2 - a^2\xi^{-1} &= & (d^2 - b^2\xi^{-1})\xi^{-1}\\ a + d &= & s\\ ad - bc &= & 1, \end{array}$$
(3.3)

where $s^2 = 4$, π_k , θ_i , and that if one of b and c is zero, so is the other. If b = c = 0and $2 \in \Box_q$ then the above (3.3) give 4 group elements in [2] and 0 element is any other classes. If neither b nor c is zero, then the first two equations in (3.3) yield $b = \pm \sqrt{-1\xi c}$. Combining with the last two equations in (3.3), we obtain 0, 4 or 8 quadruples (a, b, c, d) satisfying the above equations, among which both (a, b, c, d) and (-a, -b, -c, -d) appear at the same time. Since (a, b, c, d) and (-a, -b, -c, -d) give rise to the same group element, we conclude that $|\mathcal{S}_{\mathbf{P},\ell_1} \cap C|$ is 0, 2, or 4.

Case II. $\mathbf{Q} \in \ell_2$, $\mathbf{Q} \notin \mathbf{P}^{\perp}$, and $\mathbf{Q} \neq \mathbf{P}$.

In this case, $\mathbf{Q} = (1, 0, -y)$ for some $y \in \mathbb{Z}_q$ and $y \neq \pm \xi$. Using (2.5), we obtain that

$$K_{\mathbf{Q}} = \{ \mathbf{d}(1, 1, 1), \mathbf{d}(-1, 1, -1) \}.$$

Moreover, $K_{\mathbf{Q}}$ on $Pa_{\mathbf{Q}} = \{[1, s, y^{-1}] \mid s \in \mathbb{F}_q, s^2 - 4y^{-1} \in \mathbb{P}_q\}$ has one orbit of length 1, that is, $\{\ell_4 = [1, 0, y^{-1}]\}$, and all other orbits are of length 2. Similar arguments as above show that the parity of $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is the same as that of $|\mathcal{S}_{\mathbf{P},\ell_4} \cap C|$. So the rest is to find the parity of $|\mathcal{S}_{\mathbf{P},\ell_4} \cap C|$. The group elements in $\mathcal{S}_{\mathbf{P},\ell_4} \cap C$ are determined by the quadruples (a, b, c, d) satisfying the following equations:

$$\begin{array}{rcl} -2cd + 2ab\xi^{-1} &= & 0\\ c^2 - a^2\xi^{-1} &= & (d^2 - b^2\xi^{-1})y^{-1}\\ a + d &= & s\\ ad - bc &= & 1. \end{array}$$
(3.4)

Note that if one of b and c is zero, so is the other. If neither b nor c is zero, then the first two equations in (3.4) yield $b = \pm \sqrt{-\xi y}c$ and $a = \pm \sqrt{-\xi y^{-1}}d$. Combining with the last two, the above quadruples (a, b, c, d) yield 0, 2, or 4 group elements in $[s^2]$. If b = c = 0, then ad = 1, $d^2 = \pm \sqrt{-y\xi^{-1}}$ and $a^2 = \pm \sqrt{-\xi y^{-1}}$; and so

$$s^{2} = \sqrt{-\xi y^{-1}} + \sqrt{-y\xi^{-1}} + 2 \quad \text{or} \quad s^{2} = -\sqrt{-\xi y^{-1}} - \sqrt{-y\xi^{-1}} + 2.$$

Since $(\sqrt{-\xi y^{-1}} + \sqrt{-y\xi^{-1}} + 2)(-\sqrt{-\xi y^{-1}} - \sqrt{-y\xi^{-1}} + 2) = (\sqrt{\xi y^{-1}} + \sqrt{y\xi^{-1}})^2$, the above quadruples (a, b, c, d) yield 0 or 1 group elements in two classes $[\theta_{i_1}]$ and $[\theta_{i_2}]$ where $\theta_{i_1} = \sqrt{-\xi y^{-1}} + \sqrt{-y\xi^{-1}} + 2$ and $\theta_{i_2} = -\sqrt{-\xi y^{-1}} - \sqrt{-y\xi^{-1}} + 2$. The above analysis shows that if $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is odd then $C = [\theta_{i_1}]$ or $[\theta_{i_2}]$ in this case.

Case III. $\mathbf{Q} = \ell_2 \cap \mathbf{P}^{\perp}$.

In this case, $\mathbf{Q} = (1, 0, \xi)$ and the set of passant lines through \mathbf{Q} is

$$Pa_{\mathbf{Q}} = \{ [1, u, -\xi^{-1}] \mid u \in \mathbb{F}_q, u^2 + \xi \in \mathbb{P}_q \}.$$

Using (2.5), we obtain that

$$K_{\mathbf{Q}} = \{ \mathbf{d}(1,1,1), \mathbf{d}(-1,1,-1), \mathbf{ad}(-1,-\xi^{-1},-\xi^{-2}), \mathbf{ad}(1,-\xi^{-1},\xi^{-2}) \}$$

Therefore, among the orbits of $K_{\mathbf{Q}}$ on $Pa_{\mathbf{Q}}$, $\{[1, 0, -\xi^{-1}]\}$ is the only one of length 1 and all others are of length 2. Hence, the parity of $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C|$ is the same as that of $|\mathcal{S}_{\mathbf{P},\mathbf{P}} \cap C|$ which is the same as that of $|K \cap C|$; by Corollary 3.4, it follows that $|K \cap C|$ is odd if and only if C = D.

For $\mathbf{Q} \in I$, we denote by $\overline{N(\mathbf{Q})}$ the complement of $N(\mathbf{Q})$ in I, that is, $\overline{N(\mathbf{Q})} = I \setminus N(\mathbf{Q})$.

Lemma 3.6. Let \mathbf{P} and \mathbf{Q} be two distinct internal points.

Assume that $q \equiv 1 \pmod{4}$.

- (i) If $\ell_{\mathbf{P},\mathbf{Q}} \in Pa_{\mathbf{P}}$ and $|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap C|$ is odd, then $C = [\pi_k]$ for one k or C = D.
- (ii) If $\ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$, then $|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap C|$ is even.

Assume that $q \equiv 3 \pmod{4}$.

- (iii) If $\ell_{\mathbf{P},\mathbf{Q}} \in Pa_{\mathbf{P}}$, then $|\mathcal{U}_{\mathbf{P},\overline{N(\mathbf{Q})}} \cap C|$ is even.
- (iv) If $\ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$ and $|\mathcal{U}_{\mathbf{P},\overline{N(\mathbf{Q})}} \cap C|$ is odd, then $C = [\theta_i]$ for one *i* or C = D.

Proof: Without loss of generality, we can choose $\mathbf{P} = (1, 0, -\xi)$. Since $K = G_{\mathbf{P}}$ acts transitively on both $Pa_{\mathbf{P}}$ and $Se_{\mathbf{P}}$, we may assume that $\mathbf{Q} \neq \mathbf{P}$ is on either a special passant line $\ell_1 = [1, 0, \xi^{-1}]$ or a special secant line $\ell_2 = [0, 1, 0]$ through \mathbf{Q} .

Case I. $\ell_1 = \ell_{\mathbf{P},\mathbf{Q}} \in Pa_{\mathbf{P}}$.

In this case, $\mathbf{Q} = (1, x, -\xi)$ for some $x \in \mathbb{F}_q$ with $u^2 + \xi \in \mathbb{P}_q$ and its internal neighbor is $N(\mathbf{Q}) = \{(1, u, -\xi) \mid u^2 + \xi \in \mathbb{P}_q\} \setminus \{(1, x, -\xi)\}$ by definition. As $\mathbf{P} \in N(\mathbf{Q})$, it is obvious that $|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap D|=1$. Since the action of $K_{\mathbf{Q}}$ on $Pa_{\mathbf{Q}}$ has one orbit of length 1, i.e. ℓ_1 , and all others are of length 2, whose representatives form the set \mathcal{R}_1 , we obtain that

$$|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap C| = \sum_{\ell \in Pa_{\mathbf{Q}}} \sum_{\mathbf{P}_{1} \in I_{\ell} \setminus \{\mathbf{Q}\}} |\mathcal{U}_{\mathbf{P},\mathbf{P}_{1}} \cap C|$$

$$= \sum_{\mathbf{P}_{1} \in I_{\ell_{1}} \setminus \{\mathbf{Q}\}} |\mathcal{U}_{\mathbf{P},\mathbf{P}_{1}} \cap C| + 2 \sum_{\ell \in \mathcal{R}} \sum_{\mathbf{P}_{1} \in I_{\ell} \setminus \{\mathbf{Q}\}} |\mathcal{U}_{\mathbf{P},\mathbf{P}_{1}} \cap C|.$$
(3.5)

Now let $\mathbf{P}_1 = (1, u, -\xi) \in I_{\ell_1} \setminus {\mathbf{Q}}$. Then the number of group elements that map \mathbf{P} to \mathbf{P}_1 is determined by the quadruples (a, b, c, d) which are the solutions to the following system of equations:

$$\begin{array}{rcl}
ab - cd\xi &=& u(a^2 - c^2\xi) \\
b^2 - d^2\xi &=& -\xi(a^2 - c^2\xi) \\
a + d &=& s \\
ad - bc &=& 1.
\end{array}$$
(3.6)

The first two equations in (3.6) yield $a^2 - c^2 \xi = A$ (or -A) where $A = \sqrt{\xi(u^2 + \xi^{-1})}$.

Now using $b^2 - d^2\xi = \mp \xi A$, we obtain

$$(b+c\xi)^2 = s^2\xi - (2+A)\xi$$
 (or $s^2\xi - (2-A)\xi$).

If both $s^2\xi - (2+A)\xi$ and $s^2\xi - (2-A)\xi$ are squares, we set $B_+ = \sqrt{s^2\xi - (2+A)\xi}$ and $B_- = \sqrt{s^2\xi - (2-A)\xi}$, then

$$a = \frac{1}{2s\xi} [s^2\xi - (B_{\pm} - 2B_{\pm}\xi c)] \text{ (or } \frac{1}{2s\xi} [s^2\xi - (B_{\pm} + 2B_{\pm}\xi c)])$$

and

$$d = \frac{1}{2s\xi} [s^2\xi + (B_{\pm} - 2B_{\pm}\xi c)] \text{ (or } \frac{1}{2s\xi} [s^2\xi + (B_{\pm} + 2B_{\pm}\xi c)]);$$

combining with the last two equations of (3.6), we have

$$\left(\xi - \frac{B_{\pm}^2}{s^2}\right)c^2 + \left(\frac{B_{\pm}^3}{s^2\xi} - B_{\pm}\right)c + \left(\frac{s^2}{4} - \frac{B_{\pm}^4}{4s^2\xi^2} - 1\right) = 0 \tag{3.7}$$

or

$$\left(\xi - \frac{B_{\pm}^2}{s^2}\right)c^2 - \left(\frac{B_{\pm}^3}{s^2\xi} - B_{\pm}\right)c + \left(\frac{s^2}{4} - \frac{B_{\pm}^4}{4s^2\xi^2} - 1\right) = 0.$$
(3.8)

The discriminant of (3.7) or (3.8) is

$$\Delta = (1 - \frac{B_{\pm}^2}{s^2 \xi})(B_{\pm}^2 - s^2 \xi + 4\xi) = \frac{4\xi u^2}{s^2 (u^2 + \xi)} \in \Box_q.$$

Consequently, the equations in (3.6) have 8 solutions and yield 4 different group elements.

If one of $s^2\xi - (2 + A)\xi$ and $s^2\xi - (2 - A)\xi$ is a square and the other is nonsquare, similar arguments as above give that the equations in (3.6) have 4 solutions and produce 2 different group element.

MADISON AND WU

If one of $s^2\xi - (2+A)\xi$ and $s^2\xi - (2-A)\xi$ is zero, then s^2 is one of 2+A and 2-A; and moreover it is one of π_k for $1 \le k \le \frac{q-1}{4}$ since $(2+A)(2-A) = \frac{4u^2}{u^2+\xi} \in \mathbb{Z}_q$ and $-1 \in \square_q$. Consequently, the equations in (3.6) yield either 1 or 3 group elements in $[s^2]$.

Therefore, if $|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap C|$ is odd, then C = D or $[\pi_k]$ for one k.

Case II. $\ell_2 = \ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$ and $\mathbf{Q} \notin \mathbf{P}^{\perp}$.

Then $\mathbf{Q} = (1, 0, -y)$ for $y \notin \not \square_q$ and $y \neq \pm \xi$. From the proof of Case II in Lemma 3.5, we have that $K_{\mathbf{Q}} = \{\mathbf{d}(1, 1, 1), \mathbf{ad}(-1, 1, -1)\}$, and among the orbits of $K_{\mathbf{Q}}$ on $Pa_{\mathbf{P}}, K_{\mathbf{Q}}$ has only one orbit of length 1, that is, $\ell_4 = [1, 0, y^{-1}]$; and all other orbits are of length 2 whose representatives form the set \mathcal{R} . Since $|\mathcal{U}_{\mathbf{P},I_{\ell_i}} \cap C| = |\mathcal{U}_{\mathbf{P},I_{\ell_j}} \cap C|$ where $\ell_i, \ell_j \in Pa_{\mathbf{P}}$ and $\ell_j = \ell_i^g$ for $g \in K_{\mathbf{Q}}$, we obtain that

$$|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap C| = \sum_{\ell \in Pa_{\mathbf{Q}}} \sum_{\mathbf{P}_{1} \in I_{\ell} \setminus \{\mathbf{Q}\}} |\mathcal{U}_{\mathbf{P},\mathbf{P}_{1}} \cap C|$$

$$= \sum_{\mathbf{P}_{1} \in I_{\ell_{4}} \setminus \{\mathbf{Q}\}} |\mathcal{U}_{\mathbf{P},\mathbf{P}_{1}} \cap C| + 2 \sum_{\ell \in \mathcal{R}} \sum_{\mathbf{P}_{1} \in I_{\ell} \setminus \{\mathbf{Q}\}} |\mathcal{U}_{\mathbf{P},\mathbf{P}_{1}} \cap C|.$$
(3.9)

Moreover, since the orbits of $K_{\mathbf{Q}}$ on $I_{\ell_4} \setminus \{\mathbf{Q}\}$, whose representatives form the set \mathcal{R}_1 , are of length 2 and $|\mathcal{U}_{\mathbf{P},\mathbf{P}_1} \cap C| = |\mathcal{U}_{\mathbf{P},\mathbf{P}_2} \cap C|$ for $\mathbf{P}_2 = \mathbf{P}_1^g$, the first term of the last expression in (3.9) can be rewritten as

$$2\sum_{\mathbf{P}_1\in\mathcal{R}_1}|\mathcal{U}_{\mathbf{P},\mathbf{P}_1}\cap C|.$$

So $|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap C|$ is even in this case.

Case III. $\mathbf{P} = \ell_2 \cap \mathbf{P}^{\perp}$.

In this case, we have $\mathbf{Q} = (1, 0, \xi)$. Among the orbits of $K_{\mathbf{Q}}$ on $Pa_{\mathbf{P}}$, only one has length 1, i.e. \mathbf{P}^{\perp} . Moreover, all the orbits of $K_{\mathbf{Q}}$ on $I_{\mathbf{P}^{\perp}} \setminus {\{\mathbf{Q}\}}$ are of length 2. Hence $|\mathcal{U}_{\mathbf{P},N(\mathbf{Q})} \cap C|$ is even.

The case when $q \equiv 3 \pmod{4}$ can be established in the same way and we eliminate the detail.

4. Group Algebra FH

4.1. **2-Blocks of H.** In this section we recall several results on the 2-blocks of $H \cong PSL(2,q)$. We refer the reader to [14] or [2] for a general introduction on this subject.

Let **R** be the ring of algebraic integers in the complex field \mathbb{C} . We choose a maximal ideal **M** of **R** containing 2**R**. Let $F = \mathbf{R}/\mathbf{M}$ be the residue field of characteristic 2, and let $*: \mathbf{R} \to F$ be the natural ring homomorphism. Define

$$\mathbf{S} = \{ \frac{r}{s} \mid r \in \mathbf{R}, \ s \in \mathbf{R} \setminus \mathbf{M} \}.$$

$$(4.1)$$

Then it is clear that the map $*: \mathbf{S} \to F$ defined by $(\frac{r}{s})^* = r^*(s^*)^{-1}$ is a ring homomorphism with kernel $\mathcal{P} = \{\frac{r}{s} \mid r \in \mathbf{M}, s \in \mathbf{R} \setminus \mathbf{M}\}$. In the rest of this article, F will always be the field of characteristic 2 constructed as above. Note that F is an algebraic closure of \mathbb{F}_2 .

As a convention, we use Irr(H) and IBr(H) to denote the set of irreducible ordinary characters and the set of irreducible Brauer characters of H, respectively.

In the following, the irreducible characters of H are classified according to the character tables of PSL(2, q) displayed in Appendix.

Lemma 4.1. ([10], [11], [15]) The irreducible ordinary characters of H are:

- (i) $1 = \chi_0, \gamma, \chi_1, \dots, \chi_{\frac{q-1}{4}}, \beta_1, \beta_2, \phi_1, \dots, \phi_{\frac{q-5}{4}}$ if $q \equiv 1 \pmod{4}$, where $1 = \chi_0$ is the trivial character, γ is the character of degree q, χ_s for $1 \leq s \leq \frac{q-1}{4}$ are the characters of degree q-1, ϕ_r for $1 \leq r \leq \frac{q-5}{4}$ are the characters of degree q+1, and β_i for i = 1, 2 are the characters of degree $\frac{q+1}{2}$;
- (ii) $1 = \chi_0, \chi_1, \dots, \chi_{\frac{q-3}{4}}, \beta_1, \eta_2, \eta_1, \dots, \phi_{\frac{q-3}{4}}$ if $q \equiv 3 \pmod{4}$, where $1 = \chi_0$ is the trivial character, γ is the character of degree q, χ_s for $1 \leq s \leq \frac{q-1}{3}$ are the characters of degree q-1, ϕ_r for $1 \leq r \leq \frac{q-3}{4}$ are the characters of degree q+1, and η_i for i = 1, 2 are the characters of degree $\frac{q-1}{2}$;

The following lemma illumstrates how the irreducible ordinary characters of H are partitioned into 2-blocks.

Lemma 4.2. [17, Lemma 4.1] First assume that $q \equiv 1 \pmod{4}$ and $q - 1 = m2^n$, where $2 \nmid m$.

- (i) The principal block B_0 of H contains $2^{n-2} + 3$ irreducible characters $\chi_0 = 1, \gamma$, $\beta_1, \beta_2, \phi_{i_1}, \dots, \phi_{i_{(2^n-2-1)}}, \text{ where } \chi_0 = 1 \text{ is the trivial character of } H, \gamma \text{ is the irreducible character of degree } q \text{ of } H, \beta_1 \text{ and } \beta_2 \text{ are the two irreducible characters}$ of degree $\frac{q+1}{2}$, and ϕ_{i_k} for $1 \leq k \leq 2^{n-2}-1$ are distinct irreducible characters of
- degree q + 1 of H. (ii) H has $\frac{q-1}{4}$ blocks B_s of defect 0 for $1 \leq s \leq \frac{q-1}{4}$, each of which contains an
- (ii) If max 4 could be got a go
- Now assume that $q \equiv 3 \pmod{4}$ and $q+1 = m2^n$, where $2 \nmid m$.
- (iv) The principal block B_0 of H contains $2^{n-2} + 3$ irreducible characters $\chi_0 = 1, \gamma$, $\eta_1, \eta_2, \chi_{i_1}, \dots, \chi_{i_{(2^{n-2}-1)}}, \text{ where } \chi_0 = 1 \text{ is the trivial character of } H, \gamma \text{ is the irreducible character of degree } q \text{ of } H, \eta_1 \text{ and } \eta_2 \text{ are the two irreducible characters}$ of degree $\frac{q-1}{2}$, and χ_{i_k} for $1 \leq k \leq 2^{n-2}-1$ are distinct irreducible characters of
- (v) H has ^{q-3}/₄ blocks B_r of defect 0 for 1 ≤ r ≤ ^{q-3}/₄, each of which contains an irreducible ordinary character φ_r of degree q + 1.
 (vi) If m ≥ 3, then H has ^{m-1}/₂ blocks B'_t of defect n − 1 for 1 ≤ t ≤ ^{m-1}/₂, each of which contains 2ⁿ⁻¹ irreducible ordinary characters χ_{t_i} for 1 ≤ i ≤ 2ⁿ⁻¹.

Moreover, the above blocks form all the 2-blocks of H.

Remark 4.3. Parts (i) and (iv) are from Theorem 1.3 in [13] and their proofs can be found in Chapter 7 of III in [2]. Parts (ii) and (v) are special cases of Theorem 3.18 in [14]. Parts (iii) and (vi) are proved in Sections II and VIII of [3].

4.2. Block Idempotents. Let Bl(H) be the set of 2-blocks of H. If $B \in Bl(H)$, we write

$$f_B = \sum_{\chi \in Irr(B)} e_{\chi},$$

where $e_{\chi} = \frac{\chi(1)}{|H|} \sum_{g \in H} \chi(g^{-1})g$ is a central primitive idempotent of $\mathbf{Z}(\mathbb{C}H)$ and Irr(B) = $Irr(H) \cap B$. For future use, we define $IBr(B) = IBr(H) \cap B$. Since f_B is an element of $\mathbf{Z}(\mathbb{C}H)$, we may write

$$f_B = \sum_{C \in cl(H)} f_B(\widehat{C})\widehat{C},$$

where cl(H) is the set of conjugacy classes of H, \hat{C} is the sum of elements in the class C, and

$$f_B(\widehat{C}) = \frac{1}{|H|} \sum_{\chi \in Irr(B)} \chi(1)\chi(x_C^{-1})$$
 (4.2)

with a fixed element $x_C \in C$.

Theorem 4.4. Let $B \in Bl(H)$. Then $f_B \in \mathbf{Z}(\mathbf{S}H)$. In other words, $f_B(\widehat{C}) \in \mathbf{S}$ for each block of H.

Proof: It follows from Corollary 3.8 in [14].

We extend the ring homomorphism $*: \mathbf{S} \to F$ to a ring homomorphism $*: \mathbf{S}H \to FH$ by setting $(\sum_{g \in H} s_g g)^* = \sum_{g \in H} s_g^* g$. Note that * maps $\mathbf{Z}(\mathbf{S}H)$ onto $\mathbf{Z}(FH)$ via $(\sum_{C \in cl(H)} s_C \widehat{C})^* = \sum_{C \in cl(H)} s_C^* \widehat{C}$. Now we define

$$e_B = (f_B)^* \in \mathbf{Z}(FH),$$

which is the *block idempotent* of *B*. Note that $e_B e_{B'} = \delta_{BB'} e_B$ for *B*, $B' \in Bl(H)$, where $\delta_{BB'}$ equals 1 if B = B', 0 otherwise. Also $1 = \sum_{B \in Bl(H)} e_B$.

All the block idempotents of the 2-blocks of H are given in the following lemma; see [17] for the detailed calculations.

Lemma 4.5. [17, Lemma 4.4] First assume that $q \equiv 1 \pmod{4}$ and $q - 1 = m2^n$ with $2 \nmid m$.

- 1. Let B_0 be the principal block of H. Then (a) $e_{B_0}(\widehat{D}) = 1$; (b) $e_{B_0}(\widehat{F^+}) = e_{B_0}(\widehat{F^-}) \in F$; (c) $e_{B_0}(\widehat{[\theta_i]}) \in F$, $e_{B_0}(\widehat{[0]}) = 0$; (d) $e_{B_0}(\widehat{[\pi_k]}) = 1$.
- 2. Let B_s be any block of defect 0 of H. Then (a) $e_{B_s}(\widehat{D}) = 0$; (b) $e_{B_s}(\widehat{F^+}) = e_{B_s}(\widehat{F^-}) = 1$; (c) $e_{B_s}(\widehat{[0]}) = e_{B_s}(\widehat{[\theta_i]}) = 0$; (d) $e_{B_s}(\widehat{[\pi_k]}) \in F$.
- 3. Suppose $m \ge 3$ and let B'_t be any block of defect n-1 of H. Then (a) $e_{B'_t}(\widehat{D}) = 0$; (b) $e_{B'_t}(\widehat{F^+}) = e_{B'_t}(\widehat{F^-}) = 1$; (c) $e_{B'_t}(\widehat{[\theta_i]}) \in F$, $e_{B'_t}(\widehat{[0]}) = 0$; (d) $e_{B'_t}(\widehat{[\pi_k]}) = 0$.

Now assume that $q \equiv 3 \pmod{4}$. Suppose that $q + 1 = m2^n$ with $2 \nmid m$.

- 4. Let B_0 be the principal block of H. Then (a) $e_{B_0}(\widehat{D}) = 1$; (b) $e_{B_0}(\widehat{F^+}) = e_{B_0}(\widehat{F^-}) \in F$; (c) $e_{B_0}([\widehat{\theta_i}]) = 1$; (d) $e_{B_0}([\widehat{0}]) = 0$, $e_{B_0}([\widehat{\pi_k}]) \in F$.
- 5. Let B_r be any block of defect 0 of H. Then (a) $e_{B_r}(\widehat{D}) = 0$; (b) $e_{B_r}(\widehat{F^+}) = e_{B_r}(\widehat{F^-}) = 1$; (c) $e_{B_r}(\widehat{[0]}) = e_{B_r}(\widehat{[\pi_k]}) = 0$; (d) $e_{B_r}(\widehat{[\theta_i]}) \in F$. 6. Suppose that $m \geq 3$ and let B'_t be any block of defect n-1 of H. Then (a)
- 6. Suppose that m ≥ 3 and let B'_t be any block of defect n 1 of H. Then (a) $e_{B'_t}(\widehat{D}) = 0$; (b) $e_{B'_t}(\widehat{F^+}) = e_{B'_t}(\widehat{F^-}) = 1$; (c) $e_{B'_t}(\widehat{[\theta_i]}) = 0$; (d) $e_{B'_t}(\widehat{[0]}) = 0$, $e_{B'_t}(\widehat{[\pi_k]}) \in F$.

Let M be an **S**H-module. We denote the reduction $M/\mathcal{P}M$, which is an FH-module, by \overline{M} . Then the following lemma is apparent.

Lemma 4.6. Let M be an SH-module and $B \in Bl(H)$. Using the above notation, we have

$$\overline{M \cdot f_b} = \overline{M} \cdot e_B$$

i.e. reduction commutes with projection onto a block B.

5. Linear Maps and Their Matrices

Let F be the algebraic closure of \mathbb{F}_2 defined in Section 4. Recall that for $\mathbf{P} \in I$, $N(\mathbf{P})$ is the set of external points on the passant lines through \mathbf{P} with \mathbf{P} included or excluded accordingly as $q \equiv 3 \pmod{4}$ or $q \equiv 1 \pmod{4}$. We define \mathbf{D} to be the incidence matrix of $N(\mathbf{P})$ ($\mathbf{P} \in I$) and I. Correspondingly, the rows of \mathbf{D} can be viewed as the characteristic vectors of $N(\mathbf{P})$. In the following, we always regard both \mathbf{D} and \mathbf{A} as matrices over F. Moreover, it is clear that $\mathbf{D} = \mathbf{A}^2 + \mathbf{I}$, where \mathbf{I} is the identity matrix of proper size.

Definition 5.1. For $W \subseteq \in I$, we define C_W to be the row characteristic vector of W with respect to I, i.e. C_W is a 0-1 row vector of length |I| with entries indexed by the internal points and the entry of C_W is 1 if and only if the point indexing the entry is in W. If $W = \{\mathbf{P}\}$, as a convention, we write C_W as $C_{\mathbf{P}}$.

Let k be the complex field \mathbb{C} , the algebraic closure F of \mathbb{F}_2 , or the ring **S** in (4.1). Let k^I be the free k-module with the base $\{\mathcal{C}_{\mathbf{P}} \mid \mathbf{P} \in I\}$, respectively. If we extend the action of H on the basis elements of k^I , which is defined by $\mathcal{C}_{\mathbf{Q}} \cdot h = \mathcal{C}_{\mathbf{Q}^h}$ for $\mathbf{P} \in I$ and $h \in H$, linearly to k^I , then k^I is a kH-permutation modules. Since H is transitive on I, we have

$$k^I = \operatorname{Ind}_K^H(1_k),$$

where K is the stabilizer of an internal point in H and $\operatorname{Ind}_{K}^{H}(1_{k})$ is the kH-module induced from 1_{k} .

The decomposition of $1 \uparrow_{K}^{H}$, the character of $\operatorname{Ind}_{K}^{H}(1_{k})$, into a sum of the irreducible ordinary characters of H is given as follows.

Lemma 5.2. Let K be the stabilizer of an internal point in H.

Assume that $q \equiv 1 \pmod{4}$. Let χ_s , $1 \leq s \leq \frac{q-1}{4}$, be the irreducible characters of degree q-1, ϕ_r , $1 \leq r \leq \frac{q-5}{4}$, irreducible characters of degree q+1, γ the irreducible character of degree q, and β_j , $1 \leq j \leq 2$, irreducible characters of degree $\frac{q+1}{2}$.

(i) If $q \equiv 1 \pmod{8}$, then

$$1_K \uparrow_K^H = 1_H + \sum_{s=1}^{(q-1)/4} \chi_s + \gamma + \beta_1 + \beta_2 + \sum_{j=1}^{(q-9)/4} \phi_{r_j},$$

where ϕ_{r_j} , $1 \le j \le \frac{q-9}{4}$, may not be distinct. (ii) If $q \equiv 5 \pmod{8}$, then

$$1_K \uparrow_K^H = 1_H + \sum_{s=1}^{(q-1)/4} \chi_s + \gamma + \sum_{j=1}^{(q-5)/4} \phi_{r_j},$$

where ϕ_{r_j} , $1 \leq j \leq \frac{q-5}{4}$, may not be distinct.

Next assume that $q \equiv 3 \pmod{4}$. Let χ_s , $1 \leq s \leq \frac{q-3}{4}$, be the irreducible characters of degree q - 1, ϕ_r , $1 \leq r \leq \frac{q-3}{4}$, the irreducible characters of degree q + 1, γ the irreducible character of degree q, and η_j , $1 \leq j \leq 2$, the irreducible characters of degree $\frac{q-1}{2}$.

(iii) If $q \equiv 3 \pmod{8}$, then

$$1_K \uparrow_K^H = 1_H + \sum_{r=1}^{(q-3)/4} \phi_r + \eta_1 + \eta_2 + \sum_{j=1}^{(q-3)/4} \chi_{s_j},$$

where χ_{s_j} , $1 \leq j \leq \frac{q-3}{4}$, may not be distinct.

(iv) If $q \equiv 7 \pmod{8}$, then

$$1_K \uparrow_K^H = 1_H + \sum_{r=1}^{(q-3)/4} \phi_r + \sum_{j=1}^{(q+1)/4} \chi_{s_j},$$

where χ_{s_j} , $1 \leq j \leq \frac{q+1}{4}$, may not be distinct.

Proof: We provide the proof for the case when $q \equiv 1 \pmod{4}$.

Let 1_H be the trivial character of H. By the Frobenius reciprocity [5],

$$\left\langle 1_{K}\uparrow_{K}^{H},1_{H}\right\rangle _{H}=\left\langle 1_{K},1_{H}\downarrow_{K}^{H}\right\rangle _{K}=1$$

Let χ_s be an irreducible character of degree q-1 of H, where $1 \leq s \leq \frac{q-1}{4}$. We denote the number of elements of K lying in the class $[\pi_k]$ by d_k . Then $d_k = 2$ by Lemma 3.4(iii), and so

$$\left< 1_K \uparrow_K^H, \chi_s \right>_H = \left< 1_K, \chi_s \downarrow_K^H \right>_K = \frac{1}{|K|} \sum_{g \in K} \chi_s \downarrow_K^H (g)$$

= $\frac{1}{q+1} [(1)(q-1) + 2 \sum_{k=1}^{(q-1)/4} (-\delta^{(2k)s} - \delta^{-(2k)s})]$
= 1,

where

$$\sum_{k=1}^{q-1)/4} (-\delta^{(2k)s} - \delta^{-(2k)s}) = -(1 + \delta^{2s} + (\delta^{2s})^2 + \dots + (\delta^{2s})^{(q-1)/2} - 1)$$
$$= -\frac{1 - \delta^{(q+1)s}}{1 - \delta^{2s}} + 1$$
$$= 1$$

since $\delta^{q+1} = 1$.

(

Let γ be the irreducible character of degree q of H. Then

$$\left< 1_K \uparrow_K^H, \gamma \right>_H = \left< 1_K, \gamma \downarrow_K^H \right>_K = \frac{1}{|K|} \sum_{g \in K} \gamma \downarrow_K^H (g)$$

= $\frac{1}{q+1} [(1)(q) + (2)(-1)(\frac{q-1}{4}) + (1)(\frac{q+1}{2})]$
= 1.

Let β_j be any irreducible character of degree $\frac{q+1}{2}$ of H. Then

$$\left\langle 1_K \uparrow_K^H, \beta_j \right\rangle_H = \frac{1}{|K|} \sum_{g \in K} \beta_j \downarrow_K^H (g)$$

= $\frac{1}{q+1} [(1)(\frac{q+1}{2}) + (2)(\frac{q-1}{4})(0) + (\frac{q+1}{2})(-1)^{(q-1)/4}].$ (5.1)

Consequently, if $q \equiv 1 \pmod{8}$, then $(-1)^{\frac{q-1}{4}} = 1$, and so $\langle 1_K \uparrow_K^H, \beta_j \rangle_H = 1$; otherwise, $(-1)^{\frac{q-1}{4}} = -1$, and so $\langle 1_K \uparrow_K^H, \beta_j \rangle_H = 0$.

Since the sum of the degrees of $\overline{1}$, χ_s , γ , and β_j is less than the degree of $1 \uparrow_K^H$ and only the irreducible characters of degree q + 1 of H have not been taken into account yet, we see that all the irreducible constituents of

$$1_{K}\uparrow_{K}^{H}-1_{H}-\sum_{s=1}^{(q-1)/4}\chi_{s}-\gamma-\beta_{1}-\beta_{2} \text{ or } 1_{K}\uparrow_{K}^{H}-1_{H}-\sum_{s=1}^{(q-1)/4}\chi_{s}-\gamma$$

must have degree q + 1.

Corollary 5.3. Using the above notation,

- (i) if $q \equiv 1 \pmod{4}$, then the character of $\operatorname{Ind}_{K}^{H}(1_{\mathbb{C}}) \cdot f_{B_{s}}$ is χ_{s} for each block B_{s} of defect 0;
- (ii) if $q \equiv 3 \pmod{4}$, then the character of $\operatorname{Ind}_{K}^{H}(1_{\mathbb{C}}) \cdot f_{B_{r}}$ is ϕ_{r} for each block B_{r} of defect 0.

Proof: The corollary follows from Lemma 4.2 and Lemma 5.2.

Since H preserves incidence, it is obvious that, for $\mathbf{P} \in I$ and $h \in H$,

$$\mathcal{C}_{N(\mathbf{P})} \cdot h = \mathcal{C}_{N(\mathbf{P}^h)}$$

In the rest of the article, we always view $C_{\mathbf{P}}$ as a vector over F. Consider the maps ϕ and μ from F^I to F^I defined by extending

$$\mathcal{C}_{\mathbf{P}} \mapsto \mathcal{C}_{\mathbf{P}^{\perp}}, \mathcal{C}_{\mathbf{P}} \mapsto \mathcal{C}_{N(\mathbf{P})}$$

linearly to F^{I} , respectively. Then it is clear that as F-linear maps, the marices of ϕ and μ , are **A** and **D**, respectively, and for $\mathbf{x} \in F^{I}$, $\phi(\mathbf{x}) = \mathbf{x}\mathbf{A}$ and $\mu(\mathbf{x}) = \mathbf{x}\mathbf{D}$. Moreover, we have the following result since H is transitive on I and preserves incidence:

Lemma 5.4. The maps ϕ and μ are both FH-module homomorphisms from F^I to F^I .

We will always use **0** and **0** to denote the all-zero row vector of length |I| and the all-zero matrix of size $|I| \times |I|$, respectively; and we denote by $\hat{\mathbf{J}}$ and \mathbf{J} the all-one row vector of length |I| and the all-one matrix of size $|I| \times |I|$.

Proposition 5.5. As FH-modules, $F^{I} = \text{Im}(\phi) \oplus \text{Ker}(\phi)$, where $\text{Im}(\phi)$ and $\text{Ker}(\phi)$ are the image and kernel of ϕ , respectively.

Proof: It is clear that $\operatorname{Ker}(\phi) \subseteq \operatorname{Ker}(\phi^2)$. If $\mathbf{x} \in \operatorname{Ker}(\phi^2)$, then $\mathbf{x} \in \operatorname{Ker}(\phi)$ since

$$\phi(\mathbf{x}) = \phi^3(\mathbf{x}) = \phi(\phi^2(\mathbf{x})) = \mathbf{0}.$$

Therefore, $\operatorname{Ker}(\phi^2) = \operatorname{Ker}(\phi)$. Furthermore, since $\operatorname{Ker}(\phi) \subseteq \operatorname{Ker}(\phi^2) \subseteq \operatorname{Ker}(\phi^3) \subseteq \cdots$, we have $\operatorname{Ker}(\phi^i) = \operatorname{Ker}(\phi)$ for $i \geq 2$. Applying the Fitting decomposition theorem [12, p. 285] to the operator ϕ , we can find an i such that $F^E = \operatorname{Ker}(\phi^i) \oplus \operatorname{Im}(\phi^i)$. From the above discussions, we must have $F^E = \operatorname{Ker}(\phi) \oplus \operatorname{Im}(\phi)$. \Box

Corollary 5.6. As FH-modules, $\operatorname{Ind}_{K}^{H}(1_{F}) \cong \operatorname{Ker}(\phi) \oplus \operatorname{Im}(\phi)$.

Proof: The conclusion follows immediately from Proposition 5.5 and the fact that $\operatorname{Ind}_{K}^{H}(1_{F}) \cong F^{E}$.

Using the above notation, we set $\mathbf{C} = \mathbf{D} + \mathbf{J}$, where \mathbf{J} is the all-one matrix of proper size. Then the matrix \mathbf{C} can be viewed as the incidence matrix of $\overline{N(\mathbf{P})}$ ($\mathbf{P} \in I$) and I, and so $\mathcal{C}_{\mathbf{P}}\mathbf{C} = \mathcal{C}_{\overline{N(\mathbf{P})}}$.

Let μ_2 be the FH-homomorphism from F^I to F^I whose matrix with respect to the natural basis is **C**. The following proposition is clear.

Proposition 5.7. Using the above notation, we have $\text{Ker}(\phi) = \text{Im}(\mu)$.

Furthermore, we have the following decomposition of $\text{Ker}(\phi)$.

Lemma 5.8. Assume that $q \equiv 3 \pmod{4}$. Then as FH-modules, we have $\operatorname{Ker}(\phi) = \langle \hat{\mathbf{J}} \rangle \oplus \operatorname{Im}(\mu_2)$, where $\langle \hat{\mathbf{J}} \rangle$ is the trivial FH-module generated by $\hat{\mathbf{J}}$.

 \square

Proof: Let $\mathbf{y} \in \langle \hat{\mathbf{J}} \rangle \cap \operatorname{Im}(\mu_2)$. Then $\mathbf{y} = \mu_2(\mathbf{x}) = \lambda \hat{\mathbf{J}}$ for some $\lambda \in F$ and $\mathbf{x} \in F^I$. Or equivalently, we have $\mu_2(\mathbf{x}) = \mathbf{x}\mathbf{C} = \mathbf{x}(\mathbf{A}^2 + \mathbf{I} + \mathbf{J}) = \lambda \hat{\mathbf{J}}$. Note that $\mathbf{J}^2 = \mathbf{J}$ and $\hat{\mathbf{J}}\mathbf{J} = \hat{\mathbf{J}}$ since $2 \nmid |I|$ when $q \equiv 3 \pmod{4}$. Moreover, $\mathbf{A}^2\mathbf{J} = \hat{\mathbf{0}}$ as each row of \mathbf{A}^2 , viewed as the characteristic vector of $\widehat{N(\mathbf{P})}$, has an even number of 1s. Consequently,

$$\lambda \hat{\mathbf{J}} = \lambda \hat{\mathbf{J}} \mathbf{J} = \mathbf{x} (\mathbf{A}^2 + \mathbf{I} + \mathbf{J}) \mathbf{J} = \mathbf{x} (\mathbf{A}^2 \mathbf{J} + \mathbf{I} \mathbf{J} + \mathbf{J}^2) = \mathbf{x} (\hat{\mathbf{0}} + \mathbf{J} + \mathbf{J}) = \mathbf{0}.$$

It follows that $\lambda = 0$. Therefore, we must have $\langle \hat{\mathbf{J}} \rangle \cap \operatorname{Im}(\mu_2) = \mathbf{0}$.

It is obvious that $\langle \hat{\mathbf{J}} \rangle + \operatorname{Im}(\mu_2) \subseteq \operatorname{Ker}(\phi)$. Let $\mathbf{x} \in \operatorname{Ker}(\phi)$. Then $\mathbf{x} = \mathbf{y}(\mathbf{A}^2 + \mathbf{I})$ for some $\mathbf{y} \in F^I$. Since $\mathbf{y}\mathbf{J} = \langle \mathbf{y}, \hat{\mathbf{J}} \rangle \hat{\mathbf{J}}$, we obtain that $\mathbf{x} = \mathbf{y}(\mathbf{A}^2 + \mathbf{I} + \mathbf{J}) + \langle \mathbf{y}, \hat{\mathbf{J}} \rangle \hat{\mathbf{J}}$, where $\langle \mathbf{y}, \hat{\mathbf{J}} \rangle$ is the standard inner product of the vectors \mathbf{y} and $\hat{\mathbf{J}}$. Hence $\mathbf{x} \in \langle \hat{\mathbf{J}} \rangle + \operatorname{Im}(\mu_2)$ and so $\operatorname{Ker}(\phi) = \langle \hat{\mathbf{J}} \rangle \oplus \operatorname{Im}(\mu_2)$.

6. STATEMENT AND PROOF OF MAIN THEOREM

The main theorem is stated as follows.

Theorem 6.1. Let $Ker(\phi)$ be defined as above. As FH-modules,

(i) if $q \equiv 1 \pmod{4}$, then

$$\operatorname{Ker}(\phi) = \bigoplus_{s=1}^{(q-1)/4} M_s,$$

where M_s for $1 \leq s \leq \frac{q-1}{4}$ are pairwise non-isomorphic simple FH-modules of dimension q-1;

(ii) if $q \equiv 3 \pmod{4}$, then

$$\operatorname{Ker}(\phi) = \langle \hat{\mathbf{J}} \rangle \oplus (\bigoplus_{r=1}^{(q-3)/4} M_r),$$

where M_r for $1 \leq s \leq \frac{q-3}{4}$ are pairwise non-isomorphic simple FH-modules of dimension q+1 and $\langle \hat{\mathbf{J}} \rangle$ is the trivial FH-module generated by the all-one column vector of length |I|.

To prove the main theorem, we need refer to the following lemma.

Lemma 6.2. Let $q - 1 = 2^n m$ or $q + 1 = 2^n m$ with $2 \nmid m$ accordingly as $q \equiv 1 \pmod{4}$ or $q \equiv 3 \pmod{4}$. Using the above notation,

- (i) if $q \equiv 1 \pmod{4}$, then $\operatorname{Ker}(\phi) \cdot e_{B_0} = \mathbf{0}$, $\operatorname{Im}(\phi) \cdot e_{B_s} = \mathbf{0}$ for $1 \leq s \leq \frac{q-1}{4}$, and $\operatorname{Ker}(\phi) \cdot e_{B'_*} = \mathbf{0}$ for $m \geq 3$ and $1 \leq t \leq \frac{m-1}{2}$;
- (ii) if $q \equiv 3 \pmod{4}$, then $\operatorname{Im}(\mu_2) \cdot e_{B_0} = \mathbf{0}$, $\operatorname{Im}(\phi) \cdot e_{B_r} = \mathbf{0}$ for $1 \leq r \leq \frac{q-3}{4}$, and $\operatorname{Im}(\mu_2) \cdot e_{B'_t} = \mathbf{0}$ for $m \geq 3$ and $1 \leq t \leq \frac{m-1}{2}$.

Proof: It is clear that $Im(\phi)$, $Ker(\phi)$, and $Im(\mu_2)$ are generated by

$$\{\mathcal{C}_{\mathbf{P}^{\perp}} \mid \mathbf{P} \in I\}, \ \{\mathcal{C}_{N(\mathbf{P})} \mid \mathbf{P} \in I\}, \ \text{and} \ \{\mathcal{C}_{\overline{N(\mathbf{P})}} \mid \mathbf{P} \in I\}$$

over F, respectively. Now let $B \in Bl(H)$. Since

$$\begin{aligned} \mathcal{C}_{\mathbf{P}^{\perp}} \cdot e_B &= \sum_{C \in cl(H)} e_B(\widehat{C}) \sum_{h \in C} \mathcal{C}_{\mathbf{P}^{\perp}} \cdot h \\ &= \sum_{C \in cl(H)} e_B(\widehat{C}) \sum_{h \in C} \mathcal{C}_{(\mathbf{P}^{\perp})^h}, \\ &= \sum_{C \in cl(H)} e_B(\widehat{C}) \sum_{h \in C} \sum_{\mathbf{Q} \in (\mathbf{P}^{\perp})^h \cap I} \mathcal{C}_{\mathbf{Q}}, \end{aligned}$$

we have

$$\mathcal{C}_{\mathbf{P}^{\perp}} \cdot e_B = \sum_{\mathbf{Q} \in I} \mathcal{S}_1(B, \mathbf{P}, \mathbf{Q}) \mathcal{C}_{\mathbf{Q}}$$

where

$$\mathcal{S}_1(B, \mathbf{P}, \mathbf{Q}) := \sum_{C \in cl(H)} |\mathcal{H}_{\mathbf{P}, \mathbf{Q}} \cap C| e_B(\widehat{C}).$$

Similarly $\mathcal{C}_{N(\mathbf{P})} \cdot e_B = \sum_{\mathbf{Q} \in I} \mathcal{S}_2(B, \mathbf{P}, \mathbf{Q}) \mathcal{C}_{\mathbf{Q}}$ and $\mathcal{C}_{\overline{N(\mathbf{P})}} \cdot e_B = \sum_{\mathbf{Q} \in I} \mathcal{S}_3(B, \mathbf{P}, \mathbf{Q}) \mathcal{C}_{\mathbf{Q}}$, where

$$\mathcal{S}_2(B, \mathbf{P}, \mathbf{Q}) = \sum_{C \in Cl(H)} |\mathcal{U}_{\mathbf{P}, N(\mathbf{Q})} \cap C| e_B(\widehat{C})$$

and

$$\mathcal{S}_3(B, \mathbf{P}, \mathbf{Q}) = \sum_{C \in Cl(H)} |\mathcal{U}_{\mathbf{P}, \overline{N(\mathbf{Q})}} \cap C| e_B(\widehat{C}).$$

Assume first that $q \equiv 1 \pmod{4}$. If $\ell_{\mathbf{P},\mathbf{Q}} \in Pa_{\mathbf{P}}$, then $S_1(B_s, \mathbf{P}, \mathbf{Q}) = 0$ for each s since $|\mathcal{H}_{\mathbf{P},\mathbf{Q}} \cap C| = 0$ in F for each $C \neq [0]$ by Lemma 3.6(i) and $e_{B_s}(\widehat{[0]}) = 0$ by Lemmas 4.5 2(c); and by Lemma 3.6(i) and Lemma 4.5 1(a), 1(c), 1(d), 3(a), 3(c), 3(d), we obtain

$$S_2(B_0, \mathbf{P}, \mathbf{Q}) = e_{B_0}(\widehat{[0]}) + e_{B_0}(\widehat{[\pi_k]}) + e_{B_0}(\widehat{D}) = 0 + 1 + 1 = 0$$

and

$$S_2(B'_t, \mathbf{P}, \mathbf{Q}) = e_{B'_t}(\widehat{[0]}) + e_{B'_t}(\widehat{[\pi_k]}) + e_{B'_t}(\widehat{D}) = 0 + 0 + 0 = 0.$$

If $\ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$ and $\mathbf{Q} \notin \mathbf{P}^{\perp}$, then by Lemma 3.5(ii) and Lemma 4.5 2(c) we obtain

$$S_1(B_s, \mathbf{P}, \mathbf{Q}) = e_{B_s}(\widehat{[0]}) + e_{B_s}(\widehat{[\theta_{i_1}]}) + e_{B_s}(\widehat{[\theta_{i_1}]}) = 0 + 0 + 0 = 0;$$

and by Lemma 4.5 1(c), 3(c), and Lemma 3.6(ii), $S_2(B_0, \mathbf{P}, \mathbf{Q}) = e_{B_0}(\widehat{[0]}) = 0$ and $S_2(B'_t, \mathbf{P}, \mathbf{Q}) = e_{B'_t}(\widehat{[0]}) = 0.$

If $\ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$ and $\mathbf{Q} \in \mathbf{P}^{\perp}$, then by Lemma 3.5(iii) and Lemma 4.5 2(a) and 2(c) we obtain $S_1(B_s, \mathbf{P}, \mathbf{Q}) = e_{B_s}(\widehat{[0]}) + e_{B_s}(\widehat{D}) = 0 + 0 = 0$; and from Lemma 3.6(ii) and Lemma 4.5 1(c) and 3(c), it follows that $S_2(B_0, \mathbf{P}, \mathbf{Q}) = e_{B_0}(\widehat{[0]}) = 0$ and $S_2(B'_t, \mathbf{P}, \mathbf{Q}) = e_{B'_t}(\widehat{[0]}) = 0$.

Next we assume that $q \equiv 3 \pmod{4}$. If $\ell_{\mathbf{P},\mathbf{Q}} \in Pa_{\mathbf{P}}$ and $\mathbf{Q} \notin \mathbf{P}^{\perp}$, then by Lemma 3.5(v) and Lemma 4.5 5(c), we have

$$S_1(B_r, \mathbf{P}, \mathbf{Q}) = e_{B_r}(\widehat{[0]}) + e_{B_r}(\widehat{[\pi_{k_1}]}) + e_{B_r}(\widehat{[\pi_{k_2}]}) = 0 + 0 + 0 = 0;$$

and by Lemma 3.6(iii) and Lemma 4.5 4(d) and 6(d), we obtain $S_3(B_0, \mathbf{P}, \mathbf{Q}) = e_{B_0}(\widehat{[0]}) = 0$ and $S_3(B'_t, \mathbf{P}, \mathbf{Q}) = e_{B'_t}(\widehat{[0]}) = 0.$

If $\mathbf{Q} = \ell_{\mathbf{P},\mathbf{Q}} \cap \mathbf{P}^{\perp}$, then by Lemmas 3.6(iii) and 3.5(iii), and 4(d), 5(a), 5(c), 6(d) of Lemma 4.5, $S_3(B_0, \mathbf{P}, \mathbf{Q}) = e_{B_0}(\widehat{[0]}) = 0$, $S_1(B_r, \mathbf{P}, \mathbf{Q}) = e_{B_r}(\widehat{[0]}) + e_{B_r}(\widehat{D}) = 0 + 0 = 0$, and $S_3(B'_t, \mathbf{P}, \mathbf{Q}) = e_{B'_t}(\widehat{[0]}) = 0$.

If $\ell_{\mathbf{P},\mathbf{Q}} \in Se_{\mathbf{P}}$, then by Lemmas 3.6(iv) and 3.5(iv), and 4(a), 4(c), 4(d), 5(c), 6(a), 6(c), 6(d) of Lemma 4.5,

$$S_3(B_0, \mathbf{P}, \mathbf{Q}) = e_{B_0}(\widehat{[0]}) + e_{B_0}(\widehat{D}) + e_{B_0}(\widehat{[\theta_i]}) = 0 + 1 + 1 = 0,$$

 $S_1(B_r, \mathbf{P}, \mathbf{Q}) = e_{B_r}(\widehat{[0]}) = 0, \text{ and}$ $S_3(B'_t, \mathbf{P}, \mathbf{Q}) = e_{B'_t}(\widehat{[0]}) + e_{B'_t}(\widehat{D}) + e_{B'_t}(\widehat{[\theta_i]}) = 0 + 0 + 0 = 0.$

Proof of Theorem 6.1: Let B be a 2-block of defect 0 of H. Then by Lemma 4.6, we have

$$F^I \cdot e_B = \overline{\mathbf{S}^I \cdot f_B}.$$

Therefore, by Corollary 5.3, $F^I \cdot e_B = N$, where N is the simple FH-module of dimension q-1 or q+1 lying in B accordingly as $q \equiv 1 \pmod{4}$ or $q \equiv 3 \pmod{4}$.

Assume that $q \equiv 1 \pmod{4}$ and $q - 1 = m2^n$ with $2 \nmid m$. Since

$$1 = e_{B_0} + \sum_{s=1}^{(q-1)/4} e_{B_s} + \sum_{t=1}^{(m-1)/2} e_{B'_t}$$

 $\operatorname{Ker}(\phi) \cdot e_{B_0} = \mathbf{0}$ and $\operatorname{Ker}(\phi) \cdot e_{B'_{\star}} = \mathbf{0}$, then

$$\operatorname{Ker}(\phi) = \bigoplus_{B \in Bl(H)} \operatorname{Ker}(\phi) \cdot e_B = \bigoplus_{s=1}^{(q-1)/4} \operatorname{Ker}(\phi) \cdot e_{B_s} = \bigoplus_{s=1}^{(q-1)/4} N_s,$$

where N_s is the simple module of dimension q-1 lying in B_s for each s by the discussion in the first paragraph.

Now assume that $q \equiv 3 \pmod{4}$. Lemma 5.8 yields $\operatorname{Ker}(\phi) = \langle \mathbf{J} \rangle \oplus \operatorname{Im}(\mu_2)$. Since $\operatorname{Im}(\mu_2) \cdot e_{B_0} = \mathbf{0}$ and $\operatorname{Im}(\mu_2) \cdot e_{B'_{\star}} = \mathbf{0}$, applying the same argument as above, we have

$$\operatorname{Im}(\mu_2) = \bigoplus_{r=1}^{(q-3)/4} M_r,$$

where each M_r is a simple FH-module of dimension q + 1. Consequently,

$$\operatorname{Ker}(\phi) = \langle \hat{\mathbf{J}} \rangle \oplus \big(\bigoplus_{r=1}^{(q-3)/4} M_r \big)$$

Now Conjecture 1.1 follows as a corollary.

Corollary 6.3. Let \mathcal{L} be the \mathbb{F}_2 -null space of \mathbf{A} . Then

$$\dim_{\mathbb{F}_2}(\mathcal{L}) = \frac{(q-1)^2}{4}.$$

Proof: By Theorem 6.1 and the fact that $\dim_{\mathbb{F}_2}(\mathcal{L}) = \dim_{\mathbb{F}_2}(\operatorname{Ker}(\phi))$, when $q \equiv 1 \pmod{4}$, we have

$$\dim_{\mathbb{F}_2}(\mathcal{L}) = \sum_{i=1}^{(q-1)/4} (q-1),$$

and when $q \equiv 3 \pmod{4}$, we have

$$\dim_{\mathbb{F}_2}(\mathcal{L}) = 1 + \sum_{i=1}^{(q-3)/4} (q+1),$$

both of which are equal to $\frac{(q-1)^2}{4}$.

MADISON AND WU

APPENDIX

The character tables of PSL(2, q) were obtained by Jordan [10] and Schur [15] independently, from which we can deduce the character tables of H as follows. Let $\epsilon \in \mathbb{C}$ be a primitive (q - 1)-th root of unity and $\delta \in \mathbb{C}$ a primitive (q + 1)-th root of unity.

Number	1	2	$\frac{q-5}{4}$	1	$\frac{q-1}{4}$
Size	1	$\frac{q^2-1}{2}$	q(q+1)	$\frac{q(q+1)}{2}$	q(q-1)
Representative	D	F^{\pm}	$[heta_i]$	[0]	$[\pi_k]$
ϕ_r	q+1	1	$\epsilon^{(2i)r} + \epsilon^{-(2i)r}$	$2(-1)^{r}$	0
γ	q	0	1	1	-1
1	1	1	1	1	1
χ_s	q-1	-1	0	0	$-\delta^{(2k)s} - \delta^{-(2k)s}$
β_1	$\frac{q+1}{2}$	$\frac{1}{2}(1\pm\sqrt{q})$	$\zeta(heta_i)$	$(-1)^{(q-1)/4}$ $(-1)^{(q-1)/4}$	0
eta_2	$\frac{\overline{q+1}}{2}$	$\frac{1}{2}(1\mp\sqrt{q})$	$\zeta(heta_i)$	$(-1)^{(q-1)/4}$	0

TABLE 3. Character table of H when $q \equiv 1 \pmod{4}$

Here $s = 1, 2, ..., \frac{q-1}{4}, r = 1, 2, ..., \frac{q-5}{4}, k = 1, 2, ..., \frac{q-1}{4}, i = 1, 2, ..., \frac{q-5}{4}$, and $\zeta(\theta_i) = 1$ or -1.

TABLE 4. Character table of H when $q \equiv 3 \pmod{4}$

Number	1	2	$\frac{q-3}{4}$	1	$\frac{q-3}{4}$
Size	1	$\frac{q^2-1}{2}$	q(q+1)	$\frac{q(q-1)}{2}$	q(q-1)
Representative	D	F^{\pm}	$[heta_i]$	[0]	$[\pi_k]$
ϕ_r	q+1	1	$\epsilon^{(2i)r} + \epsilon^{-(2i)r}$	0	0
γ	q	0	1	-1	-1
1	1	1	1	1	1
χ_s	q-1	-1	0	$-2(-1)^{s}$	$-\delta^{(2k)s}-\delta^{-(2k)s}$
η_1	$\frac{q-1}{q-1}$	$\frac{1}{2}(-1 \pm \sqrt{-q})$	0	$(-1)^{(q+5)/4}$	$-\zeta(\pi_k)$
η_2	$\frac{q-1}{2}$	$\frac{1}{2}(-1 \mp \sqrt{-q})$	0	$(-1)^{(q+5)/4}$	$-\zeta(\pi_k)$

Here $s = 1, 2, ..., \frac{q-3}{4}, r = 1, 2, ..., \frac{q-3}{4}, k = 1, 2, ..., \frac{q-3}{4}, i = 1, 2, ..., \frac{q-3}{4}$, and $\zeta(\pi_k) = 1$ or -1.

References

- E. F. Assmus Jr. and D. Key, *Designs and Their Codes*, Cambridge University Press, New York, 1992.
- [2] R. Brauer, Some applications of the theory of blocks of characters of finite groups. I, II, III, and IV, J. Algebra 1 (1964), 152–167, 304–334; 3 (1966), 225–255; 17 (1971), 489–521.
- [3] R. Burkhardt, Die zerlegungsmatrizen der gruppen PSL(2, p^f), J. Algebra 40 (1976), 75–96.
- [4] S. Droms, K. E. Mellinger, and C. Meyer, LDPC codes generated by conics in the classical projective plane, *Des. Codes Cryptogr.* 40 (2006), 343–356.
- [5] G. Frobenius, Uber relationen zwischen den charakteren einer gruppe and denen ihrer untergruppen, S'ber. Akad. Wiss. Berlin (1898), 501–515; Ges. Abh. III, 104–118.
- [6] R. H. Dye, Hexagons, Conics, A₅ and PSL₂(K), J. Lond. Math. Soc. (2) 44 (1991), 270–286.
- [7] J. W. P. Hirschfeld, Projective Geometries over Finite Fields, Second Edition, Oxford University Press, Oxford, 1998.

- [8] D. R. Hughes and F. C. Piper, *Projective Planes*, Graduate Texts in Mathematics 6, Springer-Verlag, New York Inc.
- [9] B. Huppert, Endliche Gruppen I, Springer, Berlin, 1976.
- [10] H. E. Jordan, Group-characters of various types of linear groups, Amer. J. Math. 29 (1907), 387–405.
- [11] G. Karpilovsky, Group Representations, Volume 3, North-Holland, Amsterdam, 1993.
- [12] T.-Y. Lam, A First Course in Noncommutative Rings, Graduate Texts in Mathematics 131, Springer-Verlag, New York Inc.
- [13] P. Landrock, The principal block of finite groups with dihedral Sylow 2-subgroups J. Algebra 39 (1976), 410–428.
- [14] G. Navarro, Characters and Blocks of Finite Groups, London Mathematical Society Lecture Note Series 250, Cambridge University Press, Cambridge, 1998.
- [15] I. Schur, Untersuchungen über die darstellung der endlichen gruppen durch gebrochene lineare substitutionen, J. Reine Angew. Math. 132 (1907), 85–137.
- [16] B. Segre, Ovals in a finite projective plane, Canad. J. Math. 7 (1955), 414–416.
- [17] P. Sin, J. Wu, and Q. Xiang, Dimensions of some binary codes arising from a conic in PG(2,q), J. Combin. Theory (A) **118** (2011), 853–878.
- [18] T. Store, Cyclotomy and Difference Sets, Markham, Chicago, 1967.
- [19] J. Wu, Proofs of two conjectures on the dimensions of binary codes, submitted.

DEPARTMENT OF MATHEMATICS, LANE COLLEGE, JACKSON, TN, USA *E-mail address*: jwu@lanecollege.edu

LANE COLLEGE, JACKSON, TN, USA *E-mail address*: adonus_madison@lanecollege.edu