

Average Sequence Dissimilarity under the multi-species coalescent with constant population size

Joseph Heled¹

¹*Department of Computer Science, University of Auckland, Auckland, NZ*

April 5, 2011

1 Average Sequence dissimilarity

The sequence dissimilarity between two aligned sequences is defined as the fraction of positions with non-equal characters. We will show how to analytically derive the average sequence dissimilarity (ASD) within and between species under a simplified multi-species coalescent setup. To generate the sequences, we first draw a random pure birth (Yule, 1925) species tree S with n taxa and birth rate λ . The population size for each branch is constant and equal to N_e . A gene tree G compatible with S is randomly drawn using the multi-species coalescent, and finally sequences for the tips of G are simulated using the Jukes-Cantor (Jukes and Cantor, 1969) model with a mutation rate of μ .

ASD within species

Under those condition computing the ASD between two individuals within the same species is fairly straightforward. First, since the two lineages are always in a population of size N_e , regardless of species, the density of the time to coalescence is simply the exponential distribution with rate $1/N_e$ (Kingman, 1982)

$$f_{CW}(t) = \frac{1}{N_e} e^{-t/N_e}. \quad (1)$$

The evolutionary distance between two individuals whose common ancestor is t time units in the past equals $2t\mu$, and so the dissimilarity under the JC model is

$$JC_\mu(t) = \frac{3}{4} \left(1 - e^{-\frac{4}{3}2\mu t}\right). \quad (2)$$

The ASD is given by integrating the product of density and value over time,

$$ASD_W(N_e, \mu, \lambda) = \int_0^\infty f_{CW}(t)JC_\mu(t)dt = \frac{6\mu N_e}{3 + 8\mu N_e}. \quad (3)$$

ASD between species

The calculation of the ASD between species requires more work. While the dissimilarity as a function of time is given by equation (2) as before, the density of the time of coalescence depends on the Yule process. If the two populations diverged x time units ago, the coalescence density of two lineages from those species would be $f_{CW}(t)$ shifted by x . Now, assuming the density of the divergence time of the two species is $f_Y(x)$, The density of two lineages from different species coalescing at time t is obtained by integrating over all possible divergence times before the coalescence –

$$f_{CB}(t) = \int_0^t f_Y(x)f_{CW}(t-x)dx. \quad (4)$$

The ASD formula is similar to the within species case (Equation 3):

$$ASD_B(N_e, \mu, \lambda) = \int_0^\infty f_{CB}(t)JC_\mu(t)dt \quad (5)$$

We still need to figure out $f_Y(x)$. Since all species pairs are identical for our purposed, arbitrarily pick a pair $\{u, v\}$. First, partition all ranked species topologies with n taxa into $n - 1$ groups $\psi_1, \psi_2, \dots, \psi_{n-1}$, where ψ_i contains all topologies with i ($1 \leq i \leq n - 1$) surviving lineages at the time of the common ancestor of u and v .

Let c_i be the size pf ψ_i . To compute c_i , note that there are $\frac{(k-2)(k-3)}{2} + 2(k-2) = \frac{(k-2)(k+1)}{2}$ ways to coalesce k lineages into $k-1$ without coalescing u and v , giving,

$$\begin{aligned} c_i &= \prod_{k=2}^i \frac{k(k-1)}{2} \prod_{k=i+2}^n \frac{(k-2)(k+1)}{2} \\ &= \frac{(n-2)!(n+1)!}{2^{n-2}(i+1)(i+2)} \\ &= \frac{n(n-1)\mathcal{R}_{n-1}}{(i+1)(i+2)}. \end{aligned} \quad (6)$$

Where \mathcal{R}_n is the total number of ranked trees with n taxa. The density of a tree with a topology from ψ_i and with height h of the common ancestor of $\{u, v\}$ is (Joseph and Drummond (2011), Appendix C)

$$g_i(h) = \frac{1}{\mathcal{R}_n} (n-i) \binom{n}{i} \lambda e^{-\lambda(i+1)h} (1 - e^{-\lambda h})^{n-i-1}. \quad (7)$$

Putting equations (6) and (7) together, we have the required density -

$$f_Y(x) = \sum_{i=1}^{n-1} c_i g_i(x). \quad (8)$$

After some very tedious expansions and simplifications, we can obtain a closed form formula

$$\begin{aligned} ASD_B(N_e, \mu, \lambda) &= \frac{6\mu N_e}{8\mu N_e + 3} + \frac{3(n+1)}{4(n-1)(w-1)(w\lambda N_e + 1)} \\ &\quad \left((w+1) \binom{w+n}{n}^{-1} + w \frac{n-1}{n+1} - 1 \right), \end{aligned} \quad (9)$$

where $w = \frac{8\mu}{3\lambda}$. Note that we use the extended binomial since w is real.

Unlike the simple formula for within species, the ASD between species is not independent of n , and the number of terms increase with n when the formula is expanded. For example, The ASD between species for $n = 3$ is

$$\frac{6\mu (32\mu^2 N_e + 60\lambda\mu N_e + 27\lambda^2 N_e + 12\mu + 18\lambda)}{(4\mu + 3\lambda) (8\mu + 9\lambda) (8\mu N_e + 3)} \quad (10)$$

References

- H. Joseph and A. Drummond. Calibrated Tree Priors for Relaxed Phylogenetics and Divergence Time Estimation. *ArXiv e-prints*, March 2011.
- T.H. Jukes and CR Cantor. Evolution of protein molecules. *Mammalian protein metabolism*, 3:21–132, 1969.
- J. F. C. Kingman. The coalescent. *Stoch. Proc. Appl*, 13:235–248, 1982.
- G.U. Yule. A mathematical theory of evolution, based on the conclusions of Dr. JC Willis, FRS. *Philosophical Transactions of the Royal Society of London. Series B, Containing Papers of a Biological Character*, pages 21–87, 1925.