# Some further q-series identities related to divisor functions

Victor J. W. Guo<sup>1</sup> and Cai Zhang<sup>2</sup>

Department of Mathematics, East China Normal University, Shanghai 200062, People's Republic of China <sup>1</sup>jwguo@math.ecnu.edu.cn, http://math.ecnu.edu.cn/~jwguo, <sup>2</sup>51060601083@alumni.ecnu.cn

**Abstract.** We give new generalizations of some q-series identities of Dilcher and Prodinger related to divisor functions. Some interesting special cases are also deduced, including an identity related to overpartitions studied by Corteel and Lovejoy.

Keywords. Prodinger's identity; Dilcher's identity; Lagrange interpolation; l'Hôpital's rule.

MR Subject Classifications: Primary 11B65; Secondary 30E05

#### 1 Introduction

In the paper [15], Uchimura proved the following identity

$$\sum_{k=1}^{\infty} (-1)^{k-1} \frac{q^{\binom{k+1}{2}}}{(q;q)_k (1-q^k)} = \sum_{k=1}^{\infty} \frac{q^k}{1-q^k},$$
(1.1)

where  $(x;q)_N = (1-x)(1-xq)\cdots(1-xq^{N-1})$  for  $N \ge 0$ . It was pointed out by Dilcher [3] that (1.1) was known much earlier; see [12]. The identity (1.1) has caught the interests of several authors. Van Hamme [17] (see also [1,10]) gave the following finite form:

$$\sum_{k=1}^{n} (-1)^{k-1} {n \brack k} \frac{q^{\binom{k+1}{2}}}{1-q^k} = \sum_{k=1}^{n} \frac{q^k}{1-q^k},$$
(1.2)

where the q-binomial coefficient is defined as

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{cases} \frac{(q;q)_n}{(q;q)_k (q;q)_{n-k}}, & \text{if } 0 \le k \le n, \\ 0, & \text{otherwise.} \end{cases}$$

Uchimura [16] obtained a generalization of (1.2) as follows:

$$\sum_{k=1}^{n} (-1)^{k-1} {n \brack k} \frac{q^{\binom{k+1}{2}}}{1 - q^{k+m}} = \sum_{k=1}^{n} \frac{q^k}{1 - q^k} {k+m \brack m}^{-1}, \quad m \ge 0.$$
 (1.3)

Dilcher [3] established the following multiple series generalization of (1.2):

$$\sum_{k=1}^{n} (-1)^{k-1} {n \brack k} \frac{q^{\binom{k}{2} + km}}{(1-q^k)^m} = \sum_{1 \le k_1 \le \dots \le k_m \le n} \frac{q^{k_1 + \dots + k_m}}{(1-q^{k_1}) \cdots (1-q^{k_m})}.$$
 (1.4)

Prodinger [13] proved that

$$\sum_{\substack{k=0\\k\neq m}}^{n} (-1)^{k-1} {n \brack k} \frac{q^{\binom{k+1}{2}}}{1-q^{k-m}} = (-1)^m q^{\binom{m+1}{2}} {n \brack m} \sum_{\substack{k=0\\k\neq m}}^{n} \frac{q^k}{1-q^{k-m}}, \quad 0 \le m \le n.$$
 (1.5)

Fu and Lascoux [4,5] gave some further generalizations of (1.3)–(1.5). Prodinger [14] and Zeng [18] gave different proofs of Fu and Lascoux's identities.

In this paper, we shall give some other generalizations of Prodinger's identity (1.5) and Dilcher's identity (1.4) as well as a symmetric generalization of (1.2).

**Theorem 1.1.** For  $n \ge 0$  and  $0 \le l, m \le n$ , there holds

$$\sum_{\substack{k=0\\k\neq m}}^{n} \begin{bmatrix} n\\k \end{bmatrix} \frac{(q/z;q)_k (zq^{-l};q)_{n-k}}{1-q^{k-m}} z^k$$

$$= (-1)^m q^{\binom{m+1}{2}} {n \brack m} (zq^{-l};q)_l (zq^{-m};q)_{n-l} \left( \sum_{k=0}^{n-l-1} \frac{zq^{k-m}}{1 - zq^{k-m}} - \sum_{\substack{k=0\\k\neq m}}^n \frac{q^{k-m}}{1 - q^{k-m}} \right). \quad (1.6)$$

Noticing that

$$\lim_{z \to 0} (q/z; q)_k z^k = (-1)^k q^{\binom{k+1}{2}},$$

when l = 0 and z tends to 0, the identity (1.6) reduces to (1.5).

**Theorem 1.2.** For  $m, n \ge 1$ , there holds

$$\sum_{k=1}^{n} {n \brack k} \frac{(q^{m}/z; q)_{k}(z; q)_{n-k}}{(zq^{-m}; q)_{m+n}(1 - q^{k})^{m}} z^{k}$$

$$= -\sum_{k_{1}=1}^{n} \frac{q^{k_{1}}}{(1 - zq^{k_{1}-1})(1 - q^{k_{1}})} \sum_{k_{2}=1}^{k_{1}} \frac{q^{k_{2}}}{(1 - zq^{k_{2}-2})(1 - q^{k_{2}})} \cdots \sum_{k_{m}=1}^{k_{m}-1} \frac{q^{k_{m}}}{(1 - zq^{k_{m}-m})(1 - q^{k_{m}})}.$$
(1.7)

Similarly, when z tends to 0, the identity (1.7) reduces to (1.4). It should be mentioned here that (1.7) is not a consequence of the following key identity appearing in [18, (2)]:

$$\sum_{1 \le i_1 \le i_2 \le \dots \le i_m \le N} a_{i_1} a_{i_2} \cdots a_{i_m} = \sum_{k=1}^N \left( \prod_{j=1, j \ne k}^N (1 - a_j / a_k)^{-1} \right) a_k^m,$$

for the right-hand side of (1.7) cannot be written as a complete symmetric function of m indeterminates  $a_1, a_2, \ldots, a_m$ .

Our third theorem is the following symmetric generalization of the l=m=0 case of Theorem 1.1.

**Theorem 1.3.** For  $m, n \geq 0$ , there holds

$$\sum_{k=1}^{n} \frac{(q/z;q)_{k}(vq^{m};q)_{k}(z;q)_{n-k}(z;q)_{m}}{(q;q)_{k}(v;q)_{k}(q;q)_{n-k}(q^{k};q)_{m+1}} z^{k} - \sum_{k=1}^{m} \frac{(q/z;q)_{k}(vq^{n};q)_{k}(z;q)_{m-k}(z;q)_{n}}{(q;q)_{k}(v;q)_{k}(q;q)_{m-k}(q^{k};q)_{n+1}} z^{k}$$

$$= \frac{(1-zq^{-1})(z;q)_{m}(z;q)_{n}}{(q;q)_{m}(q;q)_{n}} \left( \sum_{k=1}^{m} \frac{q^{k}}{(1-zq^{k-1})(1-q^{k})} - \sum_{k=1}^{n} \frac{q^{k}}{(1-zq^{k-1})(1-q^{k})} \right). (1.8)$$

The rest of the paper is organized as follows. We shall prove Theorem 1.1 in Section 2 and give some special cases in Section 3. Theorem 1.2 will be proved in Section 4, and a class of binomial sums will be evaluated in Section 5. Finally, the proof of Theorem 1.3 and some interesting conclusions will be given in Section 6.

# 2 Proof of Theorem 1.1

Recall that the Lagrange interpolation formula states that a polynomial of degree  $\leq n$  that passes through the n+1 points  $(x_0, f(x_0)), (x_1, f(x_1)), \ldots, (x_n, f(x_n))$  is given by

$$f(x) = \sum_{k=0}^{n} f(x_k) \prod_{\substack{j=0 \ j \neq k}}^{n} \frac{x - x_j}{x_k - x_j}$$

(see, for example, [11]).

For  $0 \le l \le n$ , consider the Lagrange interpolation formula for  $(xz;q)_{n-l}$  at the values  $q^{-i}$   $(0 \le i \le n)$  of x, we immediately obtain

**Lemma 2.1.** For  $n \ge 0$  and  $0 \le l \le n$ , there holds

$$\sum_{k=0}^{n} {n \brack k} \frac{(q/z;q)_k (zq^{-l};q)_{n-k}}{1 - xq^k} z^k = \frac{(q;q)_n (xz;q)_{n-l} (zq^{-l};q)_l}{(x;q)_{n+1}}.$$
 (2.1)

Proof of Theorem 1.1. From (2.1) it follows that

$$\begin{split} &\sum_{\substack{k=0\\k\neq m}}^{n} {n\brack k} \frac{(q/z;q)_k(zq^{-l};q)_{n-k}}{1-xq^k} z^k \\ &= \frac{(q;q)_n(xz;q)_{n-l}(zq^{-l};q)_l}{(x;q)_{n+1}} - {n\brack m} \frac{(q/z;q)_m(zq^{-l};q)_{n-m}}{1-xq^m} z^m \\ &= \frac{1}{(x;q)_m(xq^{m+1};q)_{n-m}} \\ &\times \left(\frac{(q;q)_n(xz;q)_{n-l}(zq^{-l};q)_l}{1-xq^m} - {n\brack m} \frac{(q/z;q)_m(zq^{-l};q)_{n-m}(x;q)_m(xq^{m+1};q)_{n-m}}{1-xq^m} z^m\right). \end{split}$$

It is easy to see that

$$(q;q)_n(xz;q)_{n-l}(zq^{-l};q)_l - \begin{bmatrix} n \\ m \end{bmatrix} (q/z;q)_m(zq^{-l};q)_{n-m}(x;q)_m(xq^{m+1};q)_{n-m}z^m = 0$$

if  $x = q^{-m}$ . By l'Hôpital's rule, we have

$$\begin{split} &\lim_{x\to q^{-m}} \left(\frac{(q;q)_n(xz;q)_{n-l}(zq^{-l};q)_l}{1-xq^m} - \begin{bmatrix} n\\m \end{bmatrix} \frac{(q/z;q)_m(zq^{-l};q)_{n-m}(x;q)_m(xq^{m+1};q)_{n-m}}{1-xq^m} z^m \right) \\ &= \lim_{x\to q^{-m}} (q;q)_n(zq^{-l};q)_l \sum_{k=0}^{n-l-1} \frac{zq^{k-m}(xz;q)_{n-l}}{1-xzq^k} \\ &- \lim_{x\to q^{-m}} \begin{bmatrix} n\\m \end{bmatrix} (q/z;q)_m(zq^{-l};q)_{n-m} z^m \sum_{\substack{k=0\\k\neq m}}^n \frac{q^{k-m}(x;q)_m(xq^{m+1};q)_{n-m}}{1-xq^k} \\ &= (q;q)_n(zq^{-m};q)_{n-l}(zq^{-l};q)_l \left( \sum_{k=0}^{n-l-1} \frac{zq^{k-m}}{1-zq^{k-m}} - \sum_{\substack{k=0\\k\neq m}}^n \frac{q^{k-m}}{1-q^{k-m}} \right). \end{split}$$

Therefore, letting  $x \to q^{-m}$  in (2.2) and applying the relation

$$\frac{(q;q)_n}{(q^{-m};q)_m(q;q)_{n-m}} = (-1)^m q^{\binom{m+1}{2}} \begin{bmatrix} n \\ m \end{bmatrix},$$

we complete the proof.

## 3 Consequences of Theorem 1.1

Letting l = 0 in Theorem 1.1, we get

Corollary 3.1. For  $n \ge 0$  and  $0 \le m \le n$ , there holds

$$\sum_{\substack{k=0\\k\neq m}}^{n} {n \brack k} \frac{(q/z;q)_k(z;q)_{n-k}}{1-q^{k-m}} z^k = (-1)^m q^{\binom{m+1}{2}} {n \brack m} (zq^{-m};q)_n \left( \sum_{k=0}^{n-1} \frac{zq^{k-m}}{1-zq^{k-m}} - \sum_{\substack{k=0\\k\neq m}}^{n} \frac{q^{k-m}}{1-q^{k-m}} \right). \tag{3.1}$$

Letting  $n \to \infty$  in (3.1), we obtain

$$\sum_{\substack{k=0\\k\neq m}}^{\infty} \frac{(q/z;q)_k z^k}{(q;q)_k (1-q^{k-m})} = (-1)^m q^{\binom{m+1}{2}} \frac{(zq^{-m};q)_m}{(q;q)_m} \left( \sum_{k=0}^{\infty} \frac{zq^{k-m}}{1-zq^{k-m}} - \sum_{\substack{k=0\\k\neq m}}^{\infty} \frac{q^{k-m}}{1-q^{k-m}} \right). \tag{3.2}$$

For m = 0, the above identity (3.2) reduces to

$$\sum_{k=1}^{\infty} \frac{(q/z;q)_k z^k}{(q;q)_k (1-q^k)} = \sum_{k=0}^{\infty} \frac{zq^k}{1-zq^k} - \sum_{k=1}^{\infty} \frac{q^k}{1-q^k}.$$
 (3.3)

Now, letting z = -q in (3.3), we are led to

$$\sum_{k=1}^{\infty} \frac{(-1;q)_k (-q)^k}{(q;q)_k (1-q^k)} = -\sum_{k=1}^{\infty} \frac{2q^k}{1-q^{2k}},$$

of which a combinatorial interpretation was given by Corteel and Lovejoy [2, Theorem 4.4].

Letting m = 0 in Theorem 1.1, we get

Corollary 3.2. For  $n \ge 0$  and  $0 \le l \le n$ , there holds

$$\sum_{k=1}^{n} {n \brack k} \frac{(q/z;q)_k (zq^{-l};q)_{n-k}}{1-q^k} z^k = (zq^{-l};q)_n \left( \sum_{k=1}^{n-l} \frac{zq^{k-1}}{1-zq^{k-1}} - \sum_{k=1}^{n} \frac{q^k}{1-q^k} \right). \tag{3.4}$$

Furthermore, letting l = 0 in (3.4), and observing that

$$\frac{zq^{k-1}}{1-zq^{k-1}} - \frac{q^k}{1-q^k} = -\frac{q^k(1-zq^{-1})}{(1-zq^{k-1})(1-q^k)},$$
(3.5)

we obtain

Corollary 3.3. For  $n \ge 0$ , there holds

$$\sum_{k=1}^{n} {n \brack k} \frac{(q/z;q)_k(z;q)_{n-k}}{1-q^k} z^k = (zq^{-1};q)_{n+1} \sum_{k=1}^{n} \frac{q^k}{(1-zq^{k-1})(1-q^k)}.$$
 (3.6)

### 4 Proof of Theorem 1.2

*Proof.* We proceed by induction on m and on n. For m = 1, the identity (1.7) reduces to (3.6). Assume that (1.7) holds for some  $m \ge 1$ . We need to show that it also holds for m + 1, namely,

$$\sum_{k=1}^{n} {n \brack k} \frac{(q^{m+1}/z;q)_k(z;q)_{n-k}}{(zq^{-m-1};q)_{m+n+1}(1-q^k)^{m+1}} z^k$$

$$= -\sum_{k_0=1}^{n} \frac{q^{k_0}}{(1-zq^{k_0-1})(1-q^{k_0})} \sum_{k_1=1}^{k_0} \frac{q^{k_1}}{(1-zq^{k_1-2})(1-q^{k_1})} \cdots \sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m}}{(1-zq^{k_m-m-1})(1-q^{k_m})}.$$
(4.1)

We shall prove this induction step (4.1) by induction on n, following the proofs in [10] and [3, Theorem 4].

For n = 1, both sides of (4.1) are equal to

$$-\frac{q^{m+1}}{(zq^{-m};q)_{m+1}(1-q)^{m+1}},$$

and it is true. We assume that (4.1) holds for n-1. In order to show that it also holds for n, we have to check that the difference between (4.1) for n and (4.1) for n-1 is a true identity. This difference is

$$\sum_{k=1}^{n} \frac{(q^{m+1}/z;q)_{k}}{(1-q^{k})^{m+1}} z^{k} \left( \begin{bmatrix} n \\ k \end{bmatrix} \frac{(z;q)_{n-k}}{(zq^{-m-1};q)_{m+n+1}} - \begin{bmatrix} n-1 \\ k \end{bmatrix} \frac{(z;q)_{n-k-1}}{(zq^{-m-1};q)_{m+n}} \right) \\
= -\frac{q^{n}}{(1-zq^{n-1})(1-q^{n})} \sum_{k_{1}=1}^{n} \frac{q^{k_{1}}}{(1-zq^{k_{1}-2})(1-q^{k_{1}})} \cdots \sum_{k_{m}=1}^{k_{m}-1} \frac{q^{k_{m}}}{(1-zq^{k_{m}-m-1})(1-q^{k_{m}})}.$$
(4.2)

Now using the relation

$$\begin{bmatrix} n \\ k \end{bmatrix} \frac{(z;q)_{n-k}}{(zq^{-m-1};q)_{m+n+1}} - \begin{bmatrix} n-1 \\ k \end{bmatrix} \frac{(z;q)_{n-k-1}}{(zq^{-m-1};q)_{m+n}} = \begin{bmatrix} n \\ k \end{bmatrix} \frac{(zq^{-1};q)_{n-k}(1-q^k)q^{n-k}}{(zq^{-m-1};q)_{m+n}(1-q^n)(1-zq^{n-1})},$$

we see that (4.2) is equivalent to (1.7) with z being replaced by  $zq^{-1}$ . This proves that (4.1) holds for all  $n \ge 1$ , and consequently (1.7) holds for all  $m \ge 1$  and all  $n \ge 1$ .

Similarly to the proof of Theorem 1.2, we can prove that

**Theorem 4.1.** For  $m, n \ge 1$ , there holds

$$\sum_{k=1}^{n} {n \brack k} \frac{(q^{m}/z;q)_{k}(z/q;q)_{n-k}}{(zq^{-m};q)_{n+m-1}(1-q^{k})^{m}} z^{k}$$

$$= \sum_{k_{1}=1}^{n} \frac{q^{k_{1}}}{(1-zq^{k_{1}-2})(1-q^{k_{1}})} \sum_{k_{2}=1}^{k_{1}} \frac{q^{k_{2}}}{(1-zq^{k_{2}-3})(1-q^{k_{2}})} \cdots \left(\sum_{k_{m}=1}^{k_{m-1}-1} \frac{zq^{k_{m}-m}}{1-zq^{k_{m}-m}} - \sum_{k_{m}=1}^{k_{m-1}} \frac{q^{k_{m}}}{1-q^{k_{m}}}\right).$$

$$(4.3)$$

By taking the limit as  $n \to \infty$  in Theorems 1.2 and 4.1, we obtain the following two results.

Corollary 4.2. For  $m \geq 1$ , there holds

$$\begin{split} &\sum_{k=1}^{\infty} \frac{(q^m/z;q)_k}{(zq^{-m};q)_m(q;q)_k(1-q^k)^m} z^k \\ &= -\sum_{k=1}^{\infty} \frac{q^{k_1}}{(1-zq^{k_1-1})(1-q^{k_1})} \sum_{k=1}^{k_1} \frac{q^{k_2}}{(1-zq^{k_2-2})(1-q^{k_2})} \cdots \sum_{k=1}^{k_{m-1}} \frac{q^{k_m}}{(1-zq^{k_m-m})(1-q^{k_m})}. \end{split}$$

Corollary 4.3. For  $m \geq 1$ , there holds

$$\begin{split} &\sum_{k=1}^{\infty} \frac{(q^m/z;q)_k}{(zq^{-m};q)_{m-1}(q;q)_k(1-q^k)^m} z^k \\ &= \sum_{k_1=1}^{\infty} \frac{q^{k_1}}{(1-zq^{k_1-2})(1-q^{k_1})} \sum_{k_2=1}^{k_1} \frac{q^{k_2}}{(1-zq^{k_2-3})(1-q^{k_2})} \cdots \left(\sum_{k_m=1}^{k_{m-1}-1} \frac{zq^{k_m-m}}{1-zq^{k_m-m}} - \sum_{k_m=1}^{k_{m-1}} \frac{q^{k_m}}{1-q^{k_m}}\right). \end{split}$$

### 5 Some binomial identities

As was pointed out by Andrews and Uchimura [1], the partial sum

$$H_n(q) := \sum_{k=1}^n \frac{q^k}{1 - q^k}$$

may be considered as a q-analogue of the harmonic number  $H_n := \sum_{k=1}^n 1/k$ . Thus, the identity (1.2) is a q-analogue of the celebrated identity

$$\sum_{k=1}^{n} (-1)^{k-1} \binom{n}{k} \frac{1}{k} = \sum_{k=1}^{n} \frac{1}{k}.$$
 (5.1)

(see, for example, [7, (1.46)]). Dilcher [3] has obtained a multiple generalization of (5.1) by taking the limit as  $q \to 1$  in (1.4) as follows:

$$\sum_{k=1}^{n} (-1)^{k-1} \binom{n}{k} \frac{1}{k^m} = \sum_{1 \le k_m \le k_{m-1} \le \dots \le k_1 \le n} \frac{1}{k_1 k_2 \cdots k_m}.$$
 (5.2)

To obtain a further generalization of (5.2), we multiply both sides of (1.7) by  $(1-q^k)^{2m}$ , replace z by  $q^x$ , and let q tend to 1. Then we get the following result.

Corollary 5.1. For  $m, n \ge 1$ , there holds

$$\sum_{k=1}^{n} \binom{n}{k} \frac{(m-x)_k(x)_{n-k}}{(x-m)_{m+n}k^m} = -\sum \frac{1}{k_1 k_2 \cdots k_m (x+k_1-1)(x+k_2-2) \cdots (x+k_m-m)},$$
(5.3)

where  $(x)_N = x(x+1)\cdots(x+N-1)$  and the sum ranges over all integers  $1 \le k_m \le k_{m-1} \le \cdots \le k_1 \le n$ .

It is clear that when x tends to  $\infty$ , the identity (5.3) (multiplying both sides by  $-x^m$ ) reduces to Dilcher's identity (5.2). On the other hand, taking the limit as  $x \to m$  in (5.3), we are led to

$$\sum_{k=1}^{n} \binom{n}{k} \frac{(k-1)!(m+n-k-1)!}{k^m}$$

$$= \sum_{1 \le k_m \le k_{m-1} \le \dots \le k_1 \le n} \frac{(m-1)!(m+n-1)!}{k_1 k_2 \cdots k_m (k_1+m-1)(k_2+m-2) \cdots k_m}.$$

Similarly, we can derive the following result from (4.3).

Corollary 5.2. For  $m, n \ge 1$ , there holds

$$\sum_{k=1}^{n} {n \choose k} \frac{(m-x)_k (x-1)_{n-k}}{(x-m)_{m+n-1} k^m}$$

$$= \sum \frac{\sum_{j=1}^{k_{m-1}-1} \frac{1}{x+j-m} - \sum_{j=1}^{k_{m-1}} \frac{1}{j}}{k_1 k_2 \cdots k_{m-1} (x+k_1-2)(x+k_2-3) \cdots (x+k_{m-1}-m)},$$

where the sum is over all integers  $1 \le k_{m-1} \le \cdots \le k_1 \le n$ .

#### 6 Proof of Theorem 1.3

We start with the following identity appearing in Guo and Zeng [9]:

$$\frac{(xz, yz; q)_m}{(q, xyz; q)_m} \sum_{k=0}^n \frac{(x, y, vq^m; q)_k(z; q)_{n-k}}{(q, v, xyzq^m; q)_k(q; q)_{n-k}} z^k = \frac{(xz, yz; q)_n}{(q, xyz; q)_n} \sum_{k=0}^m \frac{(x, y, vq^n; q)_k(z; q)_{m-k}}{(q, v, xyzq^n; q)_k(q; q)_{m-k}} z^k,$$
(6.1)

which can be obtained applying Sears's transformation [6, p. 360, (III.15)]. Letting y = q/z in (6.1), we get

$$\sum_{k=0}^{n} \frac{(q/z;q)_{k}(vq^{m};q)_{k}(z;q)_{n-k}(xz;q)_{m}}{(q;q)_{k}(v;q)_{k}(q;q)_{n-k}(xq^{k};q)_{m+1}} z^{k} = \sum_{k=0}^{m} \frac{(q/z;q)_{k}(vq^{n};q)_{k}(z;q)_{m-k}(xz;q)_{n}}{(q;q)_{k}(v;q)_{k}(q;q)_{m-k}(xq^{k};q)_{n+1}} z^{k},$$

which can be rewritten as

$$\sum_{k=1}^{n} \frac{(q/z;q)_{k}(vq^{m};q)_{k}(z;q)_{n-k}(xz;q)_{m}}{(q;q)_{k}(v;q)_{k}(q;q)_{n-k}(xq^{k};q)_{m+1}} z^{k} - \sum_{k=1}^{m} \frac{(q/z;q)_{k}(vq^{n};q)_{k}(z;q)_{m-k}(xz;q)_{n}}{(q;q)_{k}(v;q)_{k}(q;q)_{m-k}(xq^{k};q)_{n+1}} z^{k}$$

$$= \frac{(z;q)_{m}(q;q)_{n}(xz;q)_{n}(xq;q)_{m} - (z;q)_{n}(q;q)_{m}(xz;q)_{m}(xq;q)_{n}}{(q;q)_{m}(q;q)_{n}(xq;q)_{m}(xq;q)_{n}(1-x)}.$$
(6.2)

It is obvious that

$$(z;q)_m(q;q)_n(xz;q)_n(xq;q)_m - (z;q)_n(q;q)_m(xz;q)_m(xq;q)_n = 0$$
 if  $x = 1$ .

Let  $x \to 1$  in (6.2). Then applying l'Hôpital's rule and the relation (3.5), we complete the proof of Theorem 1.3.

Letting  $v \to 0$  or  $v \to \infty$  in (1.8), we obtain the following two corollaries.

Corollary 6.1. For  $m, n \geq 0$ , there holds

$$\sum_{k=1}^{n} \frac{(q/z;q)_{k}(z;q)_{n-k}(z;q)_{m}}{(q;q)_{k}(q;q)_{n-k}(q^{k};q)_{m+1}} z^{k} - \sum_{k=1}^{m} \frac{(q/z;q)_{k}(z;q)_{m-k}(z;q)_{n}}{(q;q)_{k}(q;q)_{m-k}(q^{k};q)_{n+1}} z^{k}$$

$$= \frac{(1-zq^{-1})(z;q)_{m}(z;q)_{n}}{(q;q)_{m}(q;q)_{n}} \left(\sum_{k=1}^{m} \frac{q^{k}}{(1-zq^{k-1})(1-q^{k})} - \sum_{k=1}^{n} \frac{q^{k}}{(1-zq^{k-1})(1-q^{k})}\right).$$

Corollary 6.2. For  $m, n \ge 0$ , there holds

$$\sum_{k=1}^{n} \frac{(q/z;q)_{k}(z;q)_{n-k}(z;q)_{m}}{(q;q)_{k}(q;q)_{n-k}(q^{k};q)_{m+1}} q^{mk}z^{k} - \sum_{k=1}^{m} \frac{(q/z;q)_{k}(z;q)_{m-k}(z;q)_{n}}{(q;q)_{k}(q;q)_{m-k}(q^{k};q)_{n+1}} q^{nk}z^{k}$$

$$= \frac{(1-zq^{-1})(z;q)_{m}(z;q)_{n}}{(q;q)_{m}(q;q)_{n}} \left(\sum_{k=1}^{m} \frac{q^{k}}{(1-zq^{k-1})(1-q^{k})} - \sum_{k=1}^{n} \frac{q^{k}}{(1-zq^{k-1})(1-q^{k})}\right).$$

Taking the limit as  $z \to 0$ , the above two corollaries reduce to

Corollary 6.3. For  $m, n \geq 0$ , there holds

$$\sum_{k=1}^{n} \frac{(-1)^{k} q^{\binom{k+1}{2}}}{(q;q)_{k}(q;q)_{n-k}(q^{k};q)_{m+1}} - \sum_{k=1}^{m} \frac{(-1)^{k} q^{\binom{k+1}{2}}}{(q;q)_{k}(q;q)_{m-k}(q^{k};q)_{n+1}} \\
= \frac{1}{(q;q)_{m}(q;q)_{n}} \left( \sum_{k=1}^{m} \frac{q^{k}}{1-q^{k}} - \sum_{k=1}^{n} \frac{q^{k}}{1-q^{k}} \right).$$
(6.3)

Corollary 6.4. For  $m, n \ge 0$ , there holds

$$\sum_{k=1}^{n} \frac{(-1)^{k} q^{mk + \binom{k+1}{2}}}{(q;q)_{k}(q;q)_{n-k}(q^{k};q)_{m+1}} - \sum_{k=1}^{m} \frac{(-1)^{k} q^{nk + \binom{k+1}{2}}}{(q;q)_{k}(q;q)_{m-k}(q^{k};q)_{n+1}} \\
= \frac{1}{(q;q)_{m}(q;q)_{n}} \left( \sum_{k=1}^{m} \frac{q^{k}}{1-q^{k}} - \sum_{k=1}^{n} \frac{q^{k}}{1-q^{k}} \right).$$
(6.4)

It is clear that both (6.3) and (6.4) reduce to (1.2) if m = 0. Moreover, if we perform the substitution  $q \to q^{-1}$ , the identities (6.3) and (6.4) can respectively be written as

$$\begin{split} &\sum_{k=1}^{n} \frac{(-1)^{k} q^{\binom{k+1}{2} + (m-n)k}}{(q;q)_{k}(q;q)_{n-k}(q^{k};q)_{m+1}} - \sum_{k=1}^{m} \frac{(-1)^{k} q^{\binom{k+1}{2} + (n-m)k}}{(q;q)_{k}(q;q)_{m-k}(q^{k};q)_{n+1}} \\ &= \frac{1}{(q;q)_{m}(q;q)_{n}} \left( \sum_{k=1}^{m} \frac{1}{1-q^{k}} - \sum_{k=1}^{n} \frac{1}{1-q^{k}} \right), \\ &\sum_{k=1}^{n} \frac{(-1)^{k} q^{\binom{k+1}{2} - nk}}{(q;q)_{k}(q;q)_{n-k}(q^{k};q)_{m+1}} - \sum_{k=1}^{m} \frac{(-1)^{k} q^{\binom{k+1}{2} - mk}}{(q;q)_{k}(q;q)_{m-k}(q^{k};q)_{n+1}} \\ &= \frac{1}{(q;q)_{m}(q;q)_{n}} \left( \sum_{k=1}^{m} \frac{1}{1-q^{k}} - \sum_{k=1}^{n} \frac{1}{1-q^{k}} \right). \end{split}$$

Finally, letting  $n \to \infty$  in (6.3) and (6.4), we obtain two different generalizations of (1.1):

$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1} q^{\binom{k+1}{2}}}{(q;q)_k (q^k;q)_{m+1}} - \sum_{k=1}^{m} \frac{(-1)^{k-1} q^{\binom{k+1}{2}}}{(q;q)_{m-k} (1-q^k)} = \frac{1}{(q;q)_m} \sum_{k=m+1}^{\infty} \frac{q^k}{1-q^k},$$

$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1} q^{mk+\binom{k+1}{2}}}{(q;q)_k (q^k;q)_{m+1}} = \frac{1}{(q;q)_m} \sum_{k=m+1}^{\infty} \frac{q^k}{1-q^k}.$$

Acknowledgements. This work was partially supported by the Fundamental Research Funds for the Central Universities, Shanghai Rising-Star Program (#09QA1401700), Shanghai Leading Academic Discipline Project (#B407), and the National Science Foundation of China (#10801054).

#### References

- [1] G.E. Andrews, and K. Uchimura, Identities in combinatorics, IV: Differentiation and harmonic numbers, Utilitas Math. 28 (1985), 265–269.
- [2] S. Corteel and J. Lovejoy, Overpartitions, Trans. Amer. Math. Soc 356 (2004), 1623–1635.
- [3] K. Dilcher, Some q-series identities related to divisor functions, Discrete Math. 145 (1995), 83–93.
- [4] A.M. Fu and A. Lascoux, q-Identities from Lagrange and Newton interpolation, Adv. Appl. Math. 31 (2003), 527–531.
- [5] A.M. Fu and A. Lascoux, q-Identities related to overpartitions and divisor functions, Electron. J. Combin. 12 (2005), #R38.
- [6] G. Gasper and M. Rahman, Basic hypergeometric series, Encyclopedia of Mathematics and Its Applications, Vol. 96, Second Edition, Cambridge University Press, Cambridge, 2004.
- [7] H.W. Gould, Combinatorial Identities, A standardized set of tables listing 500 binomial coefficient summations, Morgantown, W. Va., 1972.
- [8] V.J.W. Guo, Elementary proofs of some q-identities of Jackson and Andrews-Jain, Discrete Math. 295 (2005), 63–74.
- [9] V.J.W. Guo and J. Zeng, A combinatorial proof of a symmetric q-Pfaff-Saalschütz identity, Electron. J. Combin. 12 (2005), #N2.
- [10] M. Hoffman, Solution to problem 6407 (proposed by L. Van Hamme), Amer. Math. Monthly 91 (1984), 315–316.
- [11] H. Jeffreys and B.S. Jeffreys, Lagrange's Interpolation Formula, §9.011 in Methods of Mathematical Physics, 3rd Ed., Cambridge, England, Cambridge University Press, 1988, p. 260.
- [12] J.C. Kluyver, Vraagstuk XXXVII. (Solution by S.C. van Veen), Wiskundige Opgaven (1919), 92–93.
- [13] H. Prodinger, Some applications of the q-Rice formula, Random Structures Algorithms 19 (2001), 552-557.
- [14] H. Prodinger, q-Identities of Fu and Lascoux proved by the q-Rice formula, Quaest. Math. 27 (2004), 391–395.
- [15] K. Uchimura, An identity for the divisor generating function arising from sorting theory, J. Combin. Theory Ser. A 31 (1981), 131–135.
- [16] K. Uchimura, A generalization of identities for the divisor generating function, Utilitas Mathematica 25 (1984), 377–379.
- [17] L. Van Hamme, Advanced problem 6407, Amer. Math. Monthly 40 (1982), 703–704.
- [18] J. Zeng, On some q-identities related to divisor functions, Adv. Appl. Math. 34 (2005), 313–315.