arXiv:1104.1142v1 [math.AG] 6 Apr 2011

INVARIANT HYPERSURFACES OF ENDOMORPHISMS OF THE PROJECTIVE 3-SPACE

DE-QI ZHANG

ABSTRACT. We consider surjective endomorphisms f of degree > 1 on the projective n-space \mathbb{P}^n with n = 3, and f^{-1} -stable hypersurfaces V. We show that V is a hyperplane (i.e., deg(V) = 1) but with four possible exceptions; it is conjectured that deg(V) = 1 for any $n \ge 2$; cf. [7], [3].

Dedicated to Prof. Miyanishi on the occasion of his 70th birthday

1. INTRODUCTION

We work over the field \mathbb{C} of complex numbers. In this paper, we study properties of f^{-1} -stable prime divisors of X for endomorphisms $f: \mathbb{P}^3 \to \mathbb{P}^3$. Below is our main result.

Theorem 1.1. Let $f : \mathbb{P}^3 \to \mathbb{P}^3$ be an endomorphism of degree > 1 and V an irreducible hypersurface such that $f^{-1}(V) = V$. Then either $\deg(V) = 1$, i.e., V is a hyperplane, or V equals one of the four cubic hypersurfaces $V_i = \{S_i = 0\}$, where S_i 's are as follows, with suitable projective coordinates:

- (1) $S_1 = X_3^3 + X_0 X_1 X_2;$
- (2) $S_2 = X_0^2 X_3 + X_0 X_1^2 + X_2^3;$
- (3) $S_3 = X_0^2 X_2 + X_1^2 X_3;$
- (4) $S_4 = X_0 X_1 X_2 + X_0^2 X_3 + X_1^3$.

We are unable to rule out the four cases in Theorem 1.1 and do not know whether there is any endomorphism $f_{V_i}: V_i \to V_i$ of deg $(f_{V_i}) > 1$ for i = 2, 3 or 4, but see Examples 1.5 (for V_1) and 1.2 below.

Example 1.2. There are many endomorphisms $f_{V'}: V' \to V'$ of deg $(f_{V'}) > 1$ for the normalization V' of $V = V_i$ (i = 3, 4), where $V' \simeq \mathbb{F}_1$ in either case (cf. Remark 1.3 below). Conjecture 1.4 below asserts that $f_{V'}$ is not lifted from any endomorphism $f : \mathbb{P}^3 \to \mathbb{P}^3$ restricted to the non-normal cubic surface V. Indeed, consider the endomorphism $f_{\mathbb{P}^2}: \mathbb{P}^2 \to \mathbb{P}^2$ ($[X_0, X_1, X_2] \to [X_0^q, X_1^q, X_2^q]$) with $q \ge 2$. It lifts to an endomorphism

²⁰⁰⁰ Mathematics Subject Classification. 37F10, 32H50, 14E20, 14J45.

Key words and phrases. endomorphism, iteration, projective 3-space.

The author is supported by an ARF of NUS.

 $f_{\mathbb{F}_1}: \mathbb{F}_1 \to \mathbb{F}_1$ of deg $(f_{\mathbb{F}_1}) = q^2$, where $\mathbb{F}_1 \to \mathbb{P}^2$ is the blowup of the point [0, 0, 1] fixed by $f_{\mathbb{P}^2}^{-1}$.

Remark 1.3. Below are some remarks about Theorem 1.1.

- (1) The non-normal locus of V_i (i = 3, 4) is a single line C and stabilized by f^{-1} . Let $\sigma : V'_i \to V_i$ (i = 3, 4) be the normalization. Then V'_i is the (smooth) Hirzebruch surface \mathbb{F}_1 (i.e., the one-point blowup of \mathbb{P}^2 ; see [1, Theorem 1.5], [16]) with the conductor $\sigma^{-1}(C) \subset V'_i$ a smooth section at infinity (for V_3), and the union of the negative section and a fibre (for V_4), respectively. $f|V_i|$ lifts to a (polarized) endomorphism $f_{V'_i}: V'_i \to V'_i$.
- (2) V_1 (resp. V_2) is unique as a normal cubic (or degree three del Pezzo) surface of Picard number one and with the singular locus Sing $V_1 = 3A_2$ (resp. Sing $V_2 = E_6$); see [17, Theorem 1.2], and [9, Theorem 4.4] for the anti-canonical embedding of V_i in \mathbb{P}^3 . V_1 contains exactly three lines of triangle-shaped whose three vertices form the singular locus of V_1 . And V_2 contains a single line on which lies its unique singular point. f^{-3} fixes the singular point(s) of V_i (i = 1, 2).
- (3) f^{-1} (or its positive power) does not stabilize the only line L on V_2 by using [14, Theorem 4.3.1] since the pair (V_2, L) is not log canonical at the singular point of V_2 . For V_1 , we do not know whether f^{-1} (or its power) stabilizes the three lines.

1.4. A motivating conjecture. Here are some motivations for our paper. It is conjectured that every hypersurface $V \subset \mathbb{P}^n$ stabilized by the inverse f^{-1} of an endomorphism $f : \mathbb{P}^n \to \mathbb{P}^n$ of deg(f) > 1, is linear. This conjecture is still open when $n \geq 3$ and V is singular, since the proof of [3] is incomplete as we were informed by an author. The smooth hypersurface case was settled in the affirmative in any dimension by Cerveau - Lins Neto [4] and independently by Beauville [2]. See also [15, Theorem 1.5 in arXiv version], [18] and [19] for related results.

By Theorem 1.1, this conjecture is true when n = 3 but with four exceptional cubic surfaces V_i which we could not rule out.

From the dynamics point of view, as seen in Dinh-Sibony [5, Theorem 1.3, Corollary 1.4], $f : \mathbb{P}^n \to \mathbb{P}^n$ behaves nicely *exactly* outside those f^{-1} -stabilized subvarieties. We refer to Fornaess-Sibony [7], and [5] for further references.

A smooth hypersurface X in \mathbb{P}^{n+1} with deg $(X) \geq 3$ and $n \geq 2$, has no endomorphism $f_X : X \to X$ of degree > 1 (cf. [4], [2, Theorem]). However, singular X may have plenty of endomorphisms f_X of arbitrary degrees as shown in Example 1.5 below. Conjecture 1.4 asserts that such f_X can not be extended to an endomorphism of \mathbb{P}^{n+1} .

3

Example 1.5. We now construct many polarized endomorphisms for some degree n + 1 hypersurface $X \subset \mathbb{P}^{n+1}$, with X isomorphic to the V_1 in Theorem 1.1 when n = 2. Let

$$f = (F_0, \ldots, F_n) : \mathbb{P}^n \to \mathbb{P}^n$$

 $(n \ge 2)$, with $F_i = F_i(X_0, \ldots, X_n)$ homogeneous, be any endomorphism of degree $q^n > 1$, such that $f^{-1}(S) = S$ for a reduced degree n + 1 hypersurface $S = \{S(X_0, \ldots, X_n) = 0\}$. So S must be normal crossing and linear: $S = \sum_{i=0}^n S_i$ (cf. [15, Theorem 1.5 in arXiv version]). Thus we may assume that $f = (X_0^q, \ldots, X_n^q)$ and $S_i = \{X_i = 0\}$. The relation $S \sim (n+1)H$ with $H \subset \mathbb{P}^n$ a hyperplane, defines

$$\pi: X = Spec \oplus_{i=0}^{n} \mathcal{O}(-iH) \to \mathbb{P}^{n}$$

which is a Galois $\mathbb{Z}/(n+1)$ -cover branched over S so that $\pi^*S_i = (n+1)T_i$ with the restriction $\pi|T_i: T_i \to S_i$ an isomorphism.

This X is identifiable with the degree n + 1 hypersurface

$$\{Z^{n+1} = S(X_0, \dots, X_n)\} \subset \mathbb{P}^{n+1}$$

and has singularity of type $z^{n+1} = xy$ over the intersection points of S locally defined as xy = 0. Thus, when n = 2, we have $\text{Sing } X = 3A_2$ and X is isomorphic to the V_1 in Theorem 1.1 (cf. Remark 1.3). We may assume that

$$f^*S(X_0,\ldots,X_n) = S(X_0,\ldots,X_n)^q$$

after replacing $S(X_0, \ldots, X_n)$ by a scalar multiple, so f lifts to an endomorphism

$$g = (Z^q, F_0, \dots, F_n)$$

of \mathbb{P}^{n+1} (with homogeneous coordinates $[Z, X_0, \ldots, X_n]$), stabilizing X, so that

$$g_X := g|X : X \to X$$

is a polarized endomorphism of deg $(g_X) = q^n$ (cf. [15, Lemma 2.1]). Note that $g^{-1}(X)$ is the union of q distinct hypersurfaces

$$\{Z = \zeta^i S(X_0, \dots, X_n)\} \subset \mathbb{P}^{n+1}$$

(all isomorphic to X), where $\zeta := \exp(2\pi\sqrt{-1}/q)$.

This X has only Kawamata log terminal singularities and $\operatorname{Pic} X = (\operatorname{Pic} \mathbb{P}^{n+1}) | X (n \ge 2)$ is of rank one, using Lefschetz type theorem [12, Example 3.1.25] when $n \ge 3$. We have $f^{-1}(S_i) = S_i$ and $g_X^{-1}(T_i) = T_i$, where $0 \le i \le n$.

When n = 2, the relation $(n + 1)(T_1 - T_0) \sim 0$ gives rise to an étale-in-codimension-one $\mathbb{Z}/(n+1)$ -cover

$$\tau: \mathbb{P}^n \simeq \widetilde{X} \to X$$

so that $\sum_{i=0}^{n} \tau^* T_i$ is a union of n+1 normal crossing hyperplanes; indeed, τ restricted over $X \setminus \text{Sing } X$, is its universal cover (cf. [13, Lemma 6]), so that g_X lifts up to \widetilde{X} . A similar result *seems* to be true for $n \geq 3$, by considering the 'composite' of the $\mathbb{Z}/(n+1)$ -covers given by $(n+1)(T_i - T_0) \sim 0$ $(1 \leq i < n)$.

2. Proofs of Theorem 1.1 and Remark 1.3

We use the standard notation in Hartshorne's book and [11].

2.1. We now prove Theorem 1.1 and Remark 1.3. By [15, Theorem 1.5 in arXiv version], we may assume that $V \subset \mathbb{P}^3$ is an irreducible rational *singular* cubic hypersurface.

We first consider the case where V is non-normal. Such V is classified in [6, Theorem 9.2.1] to the effect that either $V = V_i$ (i = 3, 4) or V is a cone over a nodal or cuspidal rational planar cubic curve B. The description in Remark 1.3 on V_3, V_4 and their normalizations, is given in [16, Theorem 1.1], [1, Theorem 1.5, Case (C), (E1)]; the f^{-1} -invariance of the non-normal locus C is proved in [15, Proposition 5.4 in arXiv version].

We consider and will rule out the case where V is a cone over B. Since V is normal crossing in codimension 1 (cf. [15, Theorem 1.5 or Proposition 5.4 in arXiv version]), the base B of the cone V is nodal. Let P be the vertex of the cone V, and $L \subset V$ the generating line lying over the node of B. Then $f_V := f|V$ satisfies the assertion that $f_V^{-1}(P) = P$. Indeed, the normalization V' of V is a cone over a smooth rational (twisted) cubic curve (in \mathbb{P}^3), i.e., the contraction of the (-3)-curve on the Hirzebruch surface \mathbb{F}_3 of degree 3; f_V lifts to an endomorphism $f_{V'}$ of V' so that the conductor $C' \subset V'$ is preserved by $f_{V'}^{-1}$ (cf. [15, Proposition 5.4 in arXiv version]) and consists of two distinct generating lines L_i (lying over L). Thus $f_{V'}^{-1}$ fixes the vertex $L_1 \cap L_2$ (lying over P). Hence $f_V^{-1}(P) = P$ as asserted.

By [15, Lemma 5.9 in arXiv version], $f : \mathbb{P}^3 \to \mathbb{P}^3$ (with deg $(f) = q^3 > 1$ say) descends, via the projection $\mathbb{P}^3 \dots \to \mathbb{P}^2$ from the point P, to an endomorphism $h : \mathbb{P}^2 \to \mathbb{P}^2$ with deg $(h) = q^2 > 1$ so that $h^{-1}(B) = B$. This and deg(B) = 3 > 1 contradict the linearity property of h^{-1} -stable curves in \mathbb{P}^2 (see e.g. Theorem 1.5 and the references in [15, arXiv version]).

2.2. Next we consider the case where $V \subset \mathbb{P}^3$ is a normal rational *singular* cubic hypersurface. By the adjunction formula,

$$-K_V = -(K_{\mathbb{P}^3} + V)|V \sim H|V$$

which is ample, where $H \subset \mathbb{P}^3$ is a hyperplane. Since K_V is a Cartier divisor, V has only Du Val (or rational double, or ADE) singularities. Let

$$\sigma: V' \to V$$

be the minimal resolution. Then

$$K_{V'} = \sigma^* K_V \sim \sigma^* (-H|V)$$

For $f: \mathbb{P}^3 \to \mathbb{P}^3$, we can apply the result below to $f_V := f|V$.

Lemma 2.3. Let $V \subset \mathbb{P}^3$ be a normal cubic surface, and $f_V : V \to V$ an endomorphism such that $f_V^*(H|V) \sim qH|V$ for some q > 1 and the hyperplane $H \subset \mathbb{P}^3$. Let

$$S(V) = \{G \subset V \mid G : irreducible, G^2 < 0\}$$

be the set of negative curves on V, and set

$$E_V := \sum_{E \in S(V)} E.$$

Replacing f_V by its positive power, we have:

(1) If $f_V^*G \equiv aG$ for some Weil divisor $G \neq 0$, then a = q. We have

$$f_V^*(L|V) \sim q(L|V)$$

for every divisor L on \mathbb{P}^3 . Especially, $\deg(f_V) = q^2$; $K_V \sim -H|V$ satisfies $f_V^*K_V \sim qK_V$.

- (2) S(V) is a finite set. $f_V^* E = qE$ for every $E \in S(V)$. So $f_V^* E_V = qE_V$.
- (3) A curve $E \subset V$ is a line in \mathbb{P}^3 if and only if E is equal to $\sigma(E')$ for some (-1)-curve $E' \subset V'$.
- (4) Every curve $E \in S(V)$ is a line in \mathbb{P}^3 .
- (5) We have

$$K_V + E_V = f_V^*(K_V + E_V) + \Delta$$

for some effective divisor Δ containing no line in S(V), so that the ramification divisor

$$R_{f_V} = (q-1)E_V + \Delta$$

In particular, the cardinality $\#S(V) \leq 3$, and the equality holds exactly when $K_V + E_V \sim_{\mathbb{Q}} 0$; in this case, f_V is étale outside the three lines of S(V) and $f_V^{-1}(\operatorname{Sing} V)$.

Proof. For (1) and (2), we refer to [15, Lemma 2.1] and [14, Proposition 3.6.8] and note that $L \sim bH$ for some integer b.

(3) We may assume that $E' \simeq \mathbb{P}^1$, where $E' := \sigma'(E)$ is the proper transform of E. (3) is true because E is a line if and only if

$$1 = E.H|V(=E'.\sigma^*(H|V) = E'.(-K_{V'})),$$

and by the genus formula $-2 = 2g(E') - 2 = (E')^2 + E' K_{V'}$.

(4) $E' := \sigma'(E)$ satisfies $E'.K_{V'} = E.K_V < 0$ and $(E')^2 \le E'.\sigma^*E = E^2 < 0$. Hence E' is a (-1)-curve by the genus formula. Thus (4) follows from (3).

(5) The first part is true because, by (2), the ramification divisor $R_{f_V} = (q-1)E_V +$ (other effective divisors). Also, by (1) and (2), $\Delta \sim (1-q)(K_V + E_V)$. Since $K_V \cdot E = -1$ for every $E \in S(V)$ (by (4)), we have

$$0 \le -K_V \cdot \Delta = -K_V \cdot (1-q)(K_V + E_V) = (q-1)(3 - \#S(V)).$$

Now the second part of (5) follows from this and the fact that $\Delta = 0$ if and only if $-K_V \Delta = 0$ since $-K_V$ is ample. The last part of (5) follows from the purity of branch loci and the description of R_{f_V} in (5).

2.4. We now prove Theorem 1.1 and Remark 1.3 for the normal cubic surface V. We use the notation in Lemma 2.3. Suppose that the Picard number

$$\rho := \rho(V) \ge 3.$$

Since K_V is not nef and by the minimal model program for klt surfaces, there is a composite

$$V = V_{\rho} \stackrel{\tau_{\rho}}{\to} V_{\rho-1} \cdots \stackrel{\tau_3}{\to} V_2$$

of birational extremal contractions such that

$$\rho(V_i) = i.$$

Let

 $E_i \subset V_i$

be the exceptional (irreducible) divisor of $\tau_i : V_i \to V_{i-1}$. Since V is Du Val, either E_i is contained in the smooth locus $V_i \setminus \text{Sing}(V_i)$ and is a (-1)-curve, or E_i contains exactly one singular point

$$P_i \in \operatorname{Sing} V_i$$

of type A_{n_i} so that $\tau_i(E_i) \in V_{i-1}$ is a smooth point. In particular, every V_i is still Du Val. Let

$$V'_i \to V_i$$

be the minimal resolution. Since $-K_{V_i}$ is the pushforward of the ample divisor $-K_V$, it is ample. So V_i is still a Gorenstein del Pezzo surface. Noting that $K_{V'_i}$ is the pullback of K_{V_i} , we have

$$(K_{V_{i-1}}^2) = K_{V_{i-1}}^2 = K_{V_i}^2 + (n_i + 1) \ge 3 + (0+1) = 4$$

for all $3 \leq i \leq \rho$.

Note that the proper transform

 $E_i(V) \subset V$

of $E_i \subset V_i$ is a negative curve. Since f_V^{-1} stabilizes every negative curve in S(V) and especially $E_i(V)$ (when f is replaced by its positive power, as seen in Lemma 2.3), f_V descends to

$$f_i: V_i \to V_i.$$

The V'_2 and $S(V'_2)$, the set of negative curves on V'_2 , are classified in [17, Figure 6]. Since $K^2_{V_2} \ge 4$, $(V'_2, S(V'_2))$ is as described in one of the last 10 cases in [ibid.]. For example, we write

$$V_2 = V_2(2A_2 + A_1)$$

if Sing V_2 consist of two points of type A_2 and one point of type A_1 .

Except the four cases

$$V_2(D_4), V_2(4A_1), V_2(A_3), V_2(2A_1)$$

in [ibid.], exactly two (-1)-curves in $S(V'_2)$ map to intersecting negative curves

 $M_i \in S(V_2)$

so that

 $S(V) = \{E_{\rho}, M_1(V), M_2(V)\}$

with

 $M_i(V) \subset V$

the proper transform of M_i , so

$$K_V + E_\rho + M_1(V) + M_2(V) \sim_{\mathbb{Q}} 0$$

(cf. Lemma 2.3) and hence $K_{V_2} + M_1 + M_2 \sim_{\mathbb{Q}} 0$, which is impossible by a simple calculation and blowing down V'_2 to its relative minimal model.

For each of the above four exceptional cases, we may assume that f_2^{-1} stabilizes both extremal rays $\mathbb{R}_+[M_i]$ of the closed cone $\overline{\text{NE}}(V_2)$ of effective 1-cycles, with

 $M_i \subset V_2$

the image of some (-1)-curve on V'_2 , where both extremal rays are of fibre type in the cases $V_2(D_4)$ and $V_2(4A_1)$, where the first (resp. second) is of fibre type (resp. birational type) in the cases $V_2(A_3)$ and $V_2(2A_1)$. Let

$$F_i (\sim 2M_i)$$

with i = 1, 2, or with i = 1 only, be the fibre of the extremal fibration

$$\varphi_i = \Phi_{|2M_i|} : V \to B_i \simeq \mathbb{P}^1$$

passing through the point $(\tau_3 \circ \cdots \circ \tau_{\rho})(E_{\rho})$. Then the proper transform

$$F_i(V) \subset V$$

of F_i is a negative curve so that

$$E_V = F_1(V) + F_2(V) + E_{\mu}$$

in the cases $V_2(D_4)$ and $V_2(4A_1)$, and

$$E_V = F_1(V) + M_2(V) + E_{\rho}$$

in the cases $V_2(A_3)$ and $V_2(2A_1)$ (cf. Lemma 2.3). Then $K_V + E_V \sim_{\mathbb{Q}} 0$, and hence $K_{V_2} + F_1 + F_2 \sim_{\mathbb{Q}} 0$ or $K_{V_2} + F_1 + M_2 \sim_{\mathbb{Q}} 0$ where the latter is impossible by a simple calculation as in the early paragraph. Thus

$$K_{V_2} + F_1 + F_2 \sim_{\mathbb{Q}} 0.$$

By making use of Lemma 2.3 (1) or (2), $f_2^*F_i = qF_i$, and f_2 descends to an endomorphism

$$f_{B_1}: B_1 \to B_1$$

of degree q. Thus the ramification divisor of f_{B_1} is of degree 2(q-1) by the Hurwitz formula, and is hence equal to

$$(q-1)P + \sum (b_i - 1)P_i$$

with

$$\sum (b_i - 1) = q - 1$$

where $P \in B_1$ so that F_1 lies over P. But then

$$R_{f_2} \ge (q-1)(F_1+F_2) + \sum (b_i-1)F'_i$$

where F'_i are fibres of φ_1 lying over P_i , so that

$$K_{V_2} + F_1 + F_2 \ge f_2^*(K_{V_2} + F_1 + F_2) + \sum (b_i - 1)F_i'$$

which is impossible since $K_{V_2} + F_1 + F_2 \sim_{\mathbb{Q}} 0$.

2.5. Consider the case $\rho(V) = 2$. Then the minimal resolution

$$V' \to V$$

and its negative curves are described in one of the first five cases in [17, Figure 6].

For the case $V = V(A_5)$, two (-1)-curves on V' map to two negative curves

 M_1, M_2

on V. Note that $f_V^*(M_i) = q^*M_i$ (see Lemma 2.3). There is a contraction

 $V \to \mathbb{P}^2$

of M_1 so that the image of M_2 is a plane conic preserved by f_P^{-1} where

$$f_P: \mathbb{P}^2 \to \mathbb{P}^2$$

is the descent of f_V (of degree $q^2 > 1$), contradicting [15, Theorem 1.5(4) in arXiv version].

For the case $V(2A_2 + A_1)$, there are exactly five (-1)-curves

 $M'_i \subset V'$

with $M_i \subset V$ their images. Moreover, $M'_1 \cdot M'_2 = 1$ and both M_i (i = 1, 2) are negative curves on V; each M'_j (j = 3, 4) meets the isolated (-2)-curve; M_1 and M_3 (reap. M_2 and M_4) meet the same component of one (resp. another) (-2)-chain of type A_2 . We have

$$M_1 + M_2 \sim 2L$$

for some integral Weil divisor L, by considering a relative minimal model of V'. In fact, $M_1 + 3M_2 \sim 4M_3$. Since $f_V^*(M_1 + M_2) = q(M_1 + M_2)$ (see Lemma 2.3), f_V lifts to some

 $g: U \to U.$

Here the double cover (given by the relation $M_1 + M_2 \sim 2L$)

$$\pi: U = Spec \oplus_{i=0}^{1} \mathcal{O}(-iL) \to V$$

is branched along $M_1 + M_2$. Indeed, when 2 /q, the normalization

 \hat{U}

of the fibre product of $\pi: U \to V$ and $f_V: V \to V$ is isomorphic to U and we take g to be the first projection $\hat{U} \to U$; when $2 \mid q$, we have $\hat{U} = V \coprod V$ and let g be the composite of $\pi: U \to V$, the inclusion $V \cup \emptyset \to V \coprod V$ and the first projection $\hat{U} \to U$. Now Sing Uconsists of a type A_1 singularity lying over $M_1 \cap M_2$ and four points in $\pi^{-1}(\text{Sing } V)$ of type $A_1, A_1, \frac{1}{3}(1, 1)$ and $\frac{1}{3}(1, 1)$, and every M_j (j = 3, 4) splits into two negative curves on U which are hence preserved by g^{-1} (as in Lemma 2.3 after f_V is replaced by its positive power). Thus $f_V^{-1}(M_i) = M_i$ $(1 \le i \le 4)$. As in the proof of Lemma 2.3,

 $E'_V := M_1 + M_2 + M_3$ satisfies $K_V + E'_V \sim_{\mathbb{Q}} 0$ and f_V is étale over $V \setminus (E'_V \cup \operatorname{Sing} V)$. The latter contradicts the fact that $f^*_V M_4 = q M_4$ and hence f_V has ramification index q along M_4 .

For $V = V(A_4 + A_1)$, there are exactly four (-1)-curves

$$M'_i \subset V'$$

with $M_i \subset V$ their images. We may so label that the five (-2)-curves and M'_j (j = 1, 2, 3) form a simple loop:

$$M'_1 - (-2) - (-2) - (-2) - (-2) - M'_2 - M'_3 - (-2) - M'_1;$$

both M_j (j = 2, 3) are negative curves on V. We can verify that $M_3 + 2M_2 \sim 3(M_5 - M_2)$, and $M_1 + M_2 \sim M_5$, where M_5 is the image of a curve $M'_5 \simeq \mathbb{P}^1$ and M'_5 is the smooth fibre (passing through the intersection point of M'_3 and the isolated (-2)-curve) of the \mathbb{P}^1 -fibration on V' with a singular fibre consisting of M'_1 , M'_2 and the (-2)-chain of type A_4 sitting in between them. As in the case $V(2A_2 + A_1)$, f_V lifts to

$$g: U \to U$$

on the triple cover U defined by the relation $M_3 + 2M_2 \sim 3(M_5 - M_2)$, so that each M_i (i = 4, 5) splits into three negative curves on U preserved by g^{-1} . Hence $f_V^{-1}(M_i) = M_i$ (i = 2, ..., 5). But then for $E'_V := M_2 + M_3 + M_4$ we have $K_V + E'_V \sim_{\mathbb{Q}} 0$ as in the proof of Lemma 2.3 so that f_V is étale over $V \setminus (E'_V \cup \text{Sing } V)$, contradicting the fact that f_V has ramification index q along M_5 .

For $V = V(D_5)$, the lonely (-1)-curve M'_1 and the intersecting (-1)-curves $M'_2 \cup M'_3$ on V' satisfy $2M_1 \sim M_2 + M_3$ where

$$M_i \subset V$$

denotes the image of M'_i (indeed, the three M'_i together with the five (-2)-curves form the support of two singular fibres and a section in a \mathbb{P}^1 -fibration). As in the case $V(2A_2+A_1)$, f_V lifts to

 $g: U \to U$

on the double cover U defined by the relation $2M_1 \sim M_2 + M_3$, so that M_1 splits into two negative curves on U preserved by g^{-1} . Thus $f_V^{-1}(M_1) = M_1$. Hence (V, M_1) is log canonical (cf. [14, Theorem 4.3.1]). But (V, M_1) is not log canonical because M'_1 meets the (-2)-tree of type- D_5 in a manner different from the classification of [10, Theorem 9.6]. We reach a contradiction.

For $V = V(A_3 + 2A_1)$, let $M'_1 \subset V'$ be the (-1)-curve meeting the middle component of the (-2)-chain of type A_3 , let M'_3 be the (-1)-curve meeting two isolated (-2)-curves, and let M'_2 be the (-1)-curve meeting both M'_1 and M'_3 . Then the images

 $M_i \subset V$

of M'_i satisfy $2M_1 \sim 2M_3$ (indeed, M'_1 , M'_3 and the five (-2)-curves form the support of two singular fibres in some \mathbb{P}^1 -fibration). The relation $2(M_1 - M_3) \sim 0$ defines a double cover

$$\pi: U = Spec \oplus_{i=0}^{1} \mathcal{O}(-i(M_1 - M_3)) \to V$$

étale over $V \setminus \text{Sing } V$. In fact, π restricted over $V \setminus \text{Sing } V$, is the universal cover over it, so U is again a Gorenstein del Pezzo surface and hence the irregularity q(U) = 0. Thus f_V lifts to

$$q: U \to U.$$

Now $\pi^{-1}(\operatorname{Sing} V)$ consists of two smooth points and the unique singular point of U (of type A_1), and each $\pi^* M_i$ (i = 1, 2) splits into two negative curves $M_i(1)$, $M_i(2)$ on U preserved by g^{-1} ; thus $f_V^* M_i = q M_i$ and $g^* M_i(j) = q M_i(j)$ (as in Lemma 2.3 (1)).

We assert that f_V^{-1} permutes members of the pencil

$$\Lambda := |M_1 + M_2|.$$

It suffices to show that g^{-1} permutes members of the irreducible pencil

 Λ_U

(parametrized by \mathbb{P}^1 for q(U) = 0) which is the pullback of Λ . Now $\pi^*(M_1 + M_2)$ splits into two members

$$M_1(j) + M_2(j) = \operatorname{div}(\xi_j)$$

(in local equation; j = 1, 2) which are preserved by g^{-1} and span Λ_U . We may assume that $g^*\xi_j = \xi_j^q$ after replacing the equation by a scalar multiple. Then the g^* -pullback of every member div $(a\xi_1 + b\xi_2)$ in Λ_U is equal to div $(a\xi_1^q + b\xi_2^q)$ and hence is the union of members in Λ_U because we can factorize $a\xi_1^q + b\xi_2^q$ as a product of linear forms in ξ_1, ξ_2 . This proves the assertion.

By the assertion and since $f_V^*(M_1+M_2) = q(M_1+M_2)$, f_V descends to an endomorphism

$$f_B: B \to B$$

of degree q on the curve $B \simeq \mathbb{P}^1$ parametrizing the pencil Λ . We have $f_B^* P_0 = q P_0$ for the point parametrizing the member $M_1 + M_2$ of Λ . Write $K_B = f_B^* K_B + R_{f_B}$, where the ramification divisor

$$R_{f_B} = (q-1)P_0 + \Delta_B$$

with $\Delta_B = \sum (b_i - 1)Q_j$ of degree q - 1 for some $b_i \ge 2$. Thus the ramification divisor

$$R_{f_V} = (q-1)(M_1 + M_2) + \Delta_V$$

with $\Delta_V = \sum (b_i - 1)F_i +$ (other effective divisor), where

 $F_i \in \Lambda$

is parametrized by Q_i . On the other hand, one can verify that $-K_V \sim 2M_1 + M_2$, by blowing down to a relative minimal model of V'; indeed, M_2 is a double section of the \mathbb{P}^1 -fibration

 $\varphi := \Phi_{|2M_1|} : V \to \mathbb{P}^1.$

So $-M_1 \sim K_V + M_1 + M_2 = f_V^*(K_V + M_1 + M_2) + \Delta_V$ and hence, by Lemma 2.3 (1),

$$(b_1 - 1)F_1 \le \Delta_V \sim (1 - q)(K_V + M_1 + M_2) \sim (q - 1)M_1.$$

This is impossible because F_1 is horizontal to the half fibre M_1 of φ . Indeed, $F_1.M_1 = (M_1 + M_2).M_1 = M_2.M_1 = 1$.

2.6. Consider the last case $\rho(V) = 1$. Since $K_V^2 = 3$, we have

 $V = V(3A_2), V(E_6), \text{ or } V(A_1 + A_5),$

and the minimal resolution

 $V' \to V$

and the negative curves on V' are described in [17, Figure 5]. For the first two cases, V is isomorphic to V_i (i = 1, or 2) in Theorem 1.1 by the uniqueness result in [17, Theorem 1.2] and by [9, Theorem 4.4].

For $V = V(A_1 + A_5)$, the images

 $M_i \subset V$

of the two (-1)-curves $M'_i \subset V'$ satisfy $2M_1 \sim 2M_2$; indeed, M'_i together with the six (-2)-curves form the support of two singular fibres and a section in some \mathbb{P}^1 -fibration. Let

 $\pi: U \to V$

be the double cover given by the relation $2(M_1 - M_2) \sim 0$. In fact, π restricted over $V \setminus \text{Sing } V$, is the universal cover over it. So f_V lifts to

$$g: U \to U.$$

As in the case $V(A_3 + 2A_1)$, if we let M'_1 be the one meeting the second component of the (-2)-chain of type A_5 , then π^*M_1 splits into two negative curves on U preserved by g^{-1} . Thus $f_V^{-1}(M_1) = M_1$, and, as in the case $V(D_5)$ above, contradicts [14, Theorem 4.3.1] and [10, Theorem 9.6].

This completes the proof of Theorem 1.1 for normal cubic surfaces and hence the whole of Theorem 1.1. To determine the equations of V_i (i = 1, 2), we can check that the equations in Theorem 1.1 possess the right combination of singularities and then use

the very ampleness of $-K_{V_i}$ to embed V_i in \mathbb{P}^3 as in [9] and the uniqueness of V_i up to isomorphism, and hence up to projective transformation by [9] (cf. [17, Theorem 1.2]).

2.7. Now we prove Remark 1.3. From Lemma 2.3 till now, we did not assume the hypothesis (*) that f_V is the restriction of some $f : \mathbb{P}^3 \to \mathbb{P}^3$ whose inverse stabilizes V. From now on till the end of the paper, we assume this hypothesis (*).

For $V = V(E_6)$ or $V(3A_2)$, the relation $V \sim 3H$ defines a triple cover

$$\pi: X = Spec \oplus_{i=0}^{2} \mathcal{O}(-iH) \to V$$

branched along V. Then

$$X = \{Z^3 = V(X_0, \dots, X_3)\} \subset \mathbb{P}^4$$

is a cubic hypersurface, where we let $V(X_0, \ldots, V_3)$ be the cubic form defining $V \subset \mathbb{P}^3$. Our π^{-1} restricts to a bijection π^{-1} : Sing $V \to \text{Sing } X$. As in Example 1.5, f lifts to $(Z^q, f) : \mathbb{P}^4 \to \mathbb{P}^4$ stabilizing X, so that the restriction

$$g = (Z^q, f) | X : X \to X$$

is also a lifting of f. By the Lefschetz type theorem [12, Example 3.1.25], $\operatorname{Pic}(X) = (\operatorname{Pic}(\mathbb{P}^4))|X$.

For $V = V(E_6)$, V' contains only one (-1)-curve M' and hence V contains only one line M (the image of M') by Lemma 2.3. Note that

$$\{Q\} := \operatorname{Sing} V \subset M.$$

Let

 $\Pi_{MM} \subset \mathbb{P}^3$

(say given by $X_3 = 0$) be the unique plane such that

$$\Pi_{MM}|V=3M.$$

Indeed, 3M belongs to the complete linear system |H|V|, and the exact sequence

$$0 \to \mathcal{O}(-2H) \to \mathcal{O}(H) \to \mathcal{O}_V(H) \to 0$$

and the vanishing of $H^1(\mathbb{P}^3, -2H)$ (e.g. by the Kodaira vanishing) imply

$$H^0(\mathbb{P}^3, \mathcal{O}(H)) \simeq H^0(V, \mathcal{O}_V(H)).$$

Our $\pi^* \Pi_{MM}$ is a union of three 2-planes

$$L_i \subset \mathbb{P}^4$$

because the restriction of π over Π_{MM} is given by the equation

$$Z^{3} = V(X_{0}, \dots, X_{3}) | \Pi_{MM} = M(X_{0}, \dots, X_{2})^{3}$$

where $M(X_0, \ldots, X_2)$ is a linear equation of $M \subset \Pi_{MM}$. This and the fact that $\pi^* \Pi_{MM}$ is a generator of $\operatorname{Pic}(X) = (\operatorname{Pic}(\mathbb{P}^3)) | X$, imply that the Weil divisor L_1 is not a Cartier divisor on X. Since Sing X consists of a single point P lying over $\{Q\} = \operatorname{Sing} V, L_1$ is not Cartier at P and hence X is not factorial at P. Thus X is not Q-factorial at P because the local π_1 of P is trivial by a result of Milnor (cf. the proof of [10, Lemma 5.1]). Hence $g^{-1}(P)$ contains no smooth point (cf. [11, Lemma 5.16]) and must be equal to Sing $X = \{P\}$. Thus $f^{-1}(Q) = Q$ because $\pi^{-1}(Q) = P$.

2.8. Before we treat the case $V(3A_2)$, we make some remarks. Up to isomorphism, there is only one $V(3A_2)$ (cf. [17, Theorem 1.2]). Set $V := V(3A_2)$. There is a Gorenstein del Pezzo surface W such that $\rho(W) = 1$, Sing W consists of four points β_i of Du Val type A_2 ,

$$\pi_1(W \setminus \operatorname{Sing} W) = (\mathbb{Z}/(3))^{\oplus 2}$$

and there is a Galois triple cover $V \to W$ étale over $W \setminus \{\beta_1, \beta_2, \beta_3\}$ so that a generator

$$h \in \operatorname{Gal}(V/W)$$

permutes the three singular points of V lying over β_4 (cf. [13, Figure 1, Lemma 6]). Since the embedding $V \subset \mathbb{P}^3$ is given by the complete linear system $|-K_V|$ (cf. [9]) and $h^*(-K_V) \sim -K_V$, our h extends to a projective transformation of \mathbb{P}^3 , also denoted as h. Since h(V) = V,

$$h^*V(X_0, \dots, X_3) = cV(X_0, \dots, X_3)$$

for some nonzero constant c. This h lifts to a projective transformation of \mathbb{P}^4 , also denoted as h, stabilizing the above triple cover $X \subset \mathbb{P}^4$ of \mathbb{P}^3 by defining $h^*Z = \sqrt[3]{cZ}$. Then this h permutes the three singular points of X lying over Sing V.

2.9. For $V = V(3A_2)$, V' has exactly three (-1)-curves M'_i and their images M_i are therefore the only lines on V (cf. Lemma 2.3). The graph $\sum M_i$ is triangle-shaped whose vertices (the intersection $M_i \cap M_j$) are the three points in Sing V. The sum of the three (-1)-curves M'_i and three (-2)-chains of type A_2 is linearly equivalent to $-K_{V'}$ and hence $\sum M_i \sim -K_V$; also $2M_a \sim M_b + M_c$ so long $\{a, b, c\} = \{1, 2, 3\}$; indeed, the three M'_i and the six (-2)-curves form the support of two singular fibres and two cross-sections of some \mathbb{P}^1 -fibration. Thus $3M_i \sim -K_V \sim H|V$. As argued in the case $V(E_6)$, there is a unique hyperplane Π_i such that

$$\Pi_i | V = 3M_i;$$

our $\pi^*\Pi_i$ is a union of three 2-planes L_{ij} in \mathbb{P}^4 (sharing a line lying over $M_i \subset \mathbb{P}^3$), L_{i1} is not a Cartier divisor on X, the X is not \mathbb{Q} -factorial at least at one of the two points (and hence at both points, since the above h permutes Sing X) in $L_{i1} \cap \text{Sing } X$ (lying over $M_i \cap \operatorname{Sing} V$), and $g^{-1}(\operatorname{Sing} X) = \operatorname{Sing} X$. Thus $f^{-1}(\operatorname{Sing} V) = \operatorname{Sing} V$. Hence f^{-3} fixes each point in $\operatorname{Sing} V$.

This completes the proof of Remark 1.3 for normal cubic surfaces and hence the whole of Remark 1.3.

Remark 2.10. The proof of Theorem 1.1 actually shows: if $f_V : V \to V$ is an endomorphism (not necessarily the restriction of some $f : \mathbb{P}^3 \to \mathbb{P}^3$) of deg $(f_V) > 1$ of a Gorenstein normal del Pezzo surface with $K_V^2 = 3$ (i.e., a normal cubic surface), then V is equal to V_1 or V_2 in Theorem 1.1 in suitable projective coordinates.

References

- [1] M. Abe and M. Furushima, On non-normal del Pezzo surfaces, Math. Nachr. 260 (2003), 3–13.
- [2] A. Beauville, Endomorphisms of hypersurfaces and other manifolds, Internat. Math. Res. Notices 2001, no. 1, 53–58.
- [3] J.-V. Briend, S. Cantat and M. Shishikura, Linearity of the exceptional set for maps of $\mathbb{P}_k(\mathbb{C})$, Math. Ann. **330** (2004), 39–43.
- [4] D. Cerveau and A. Lins Neto, Hypersurfaces exceptionnelles des endomorphismes de CP(n), Bol. Soc. Brasil. Mat. (N.S.) 31 (2000), no. 2, 155–161.
- [5] T. -C. Dinh and N. Sibony, Equidistribution speed for endomorphisms of projective spaces, Math. Ann. 347 (2010), no. 3, 613–626.
- [6] I. V. Dolgachev, Topics in Classical Algebraic Geometry. Part I, January 19, 2009, at: http://www.math.lsa.umich.edu/~idolga/topics1.pdf
- [7] J. E. Fornaess and N. Sibony, Complex dynamics in higher dimension. I, Complex analytic methods in dynamical systems (Rio de Janeiro, 1992), pp. 201–231, Astérisque 222, Soc. Math. France, 1994.
- [8] C. D. Hacon and J. McKernan, On Shokurov's rational connectedness conjecture, Duke Math. J. 138 (2007), no. 1, 119–136.
- F. Hidaka and K. Watanabe, Normal Gorenstein surfaces with ample anti-canonical divisor, Tokyo J. Math. 4 (1981), no. 2, 319–330.
- [10] Y. Kawamata, Crepant blowing-ups of 3-dimensional canonical singularities and its application to degeneration of surfaces, Ann. of Math. 127 (1988), 93–163.
- [11] J. Kollár and S. Mori, Birational geometry of algebraic varieties, Cambridge Tracts in Math. 134, Cambridge Univ. Press, 1998.
- [12] R. Lazarsfeld, Positivity in algebraic geometry. I. Classical setting: line bundles and linear series. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, Vol. 48. Springer-Verlag, Berlin, 2004.
- [13] M. Miyanishi and D. -Q. Zhang, Gorenstein log del Pezzo surfaces of rank one, J. Algebra, 118 (1988), 63-84.
- [14] N. Nakayama, On complex normal projective surfaces admitting non-isomorphic surjective endomorphisms, Preprint 2 September 2008.
- [15] N. Nakayama and D. -Q. Zhang, Polarized endomorphisms of complex normal varieties, Math. Ann. 346 (2010), no. 4, 991–1018; also: arXiv:0908.1688v1.
- [16] M. Reid, Nonnormal del Pezzo surfaces, Publ. Res. Inst. Math. Sci. Kyoto Univ. 30 (1994), 695–727.

- [17] Q. Ye, On Gorenstein log del Pezzo surfaces, Japan. J. Math. 28 (2002), no. 1, 87–136.
- [18] D. -Q. Zhang, Polarized endomorphisms of uniruled varieties (with an Appendix by Y. Fujimoto and N. Nakayama), Compos. Math. 146 (2010) 145 - 168.
- [19] D. -Q. Zhang, Invariant hypersurfaces of endomorphisms of projective varieties, Preprint 2009.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE 10 LOWER KENT RIDGE ROAD, SINGAPORE 119076 *E-mail address:* matzdq@nus.edu.sg