Nonlinear Convergence Sets of Divergent Power Series

Buma L. Fridman, Daowei Ma and Tejinder S. Neelon

To Professor Józef Siciak on his 80th birthday

ABSTRACT. A nonlinear generalization of convergence sets of formal power series, in the sense of Abhyankar-Moh $[\mathbf{AM}]$, is introduced. Given a family $y = \varphi_s(t, x) = sb_1(x)t + b_2(x)t^2 + ...$ of analytic curves in $\mathbb{C} \times \mathbb{C}^n$ passing through the origin, $\operatorname{Conv}_{\varphi}(f)$ of a formal power series $f(y, t, x) \in \mathbb{C}[[y, t, x]]$ is defined to be the set of all $s \in \mathbb{C}$ for which the power series $f(\varphi_s(t, x), t, x)$ converges as a series in (t, x). We prove that for a subset $E \subset \mathbb{C}$ there exists a divergent formal power series $f(y, t, x) \in \mathbb{C}[[y, t, x]]$ such that $E = \operatorname{Conv}_{\varphi}(f)$ if and only if E is a F_{σ} set of zero capacity. This generalizes the results of P. Lelong and A. Sathaye for the linear case $\varphi_s(t, x) = st$.

We say that formal power series $f(z) = \sum_{|\alpha|=0}^{\infty} a_{\alpha} z^{\alpha}$ is convergent if there exists a constant C such that $|a_{\alpha}| \leq C^{|\alpha|}$, for all $\alpha \in \mathbb{Z}_{+}^{n}$. (Here we have used multiindex notation: \mathbb{Z}^n_+ denote the set of all *n*-tuples $\alpha := (\alpha_1, \alpha_2, ..., \alpha_n)$ of integers $\alpha_i \ge 0$, if $z = (z_1, z_2, ..., z_n)$ and $\alpha \in \mathbb{Z}_+^n$, then $z^{\alpha} = z_1^{\alpha_1} z_2^{\alpha_2} ... z_n^{\alpha_n}$, and $|\alpha| := \alpha_1 + \alpha_2 + ... + \alpha_n$ denotes the length of $\alpha \in \mathbb{Z}_{+}^{n}$.) A series f is called divergent if it is not convergent. A divergent power series may still converge when restricted to a certain set of lines or planes through the origin. For example, Abhyankar and Moh $[\mathbf{AM}]$ considered the convergence set $\operatorname{Conv}(f)$ of a series f defined to be the set of all $s \in \mathbb{C}$ for which $f(sz_2, z_2, ..., z_n)$ converges as a series in $(z_2, z_3, ..., z_n)$. The convergence set of divergent series can be empty or an arbitrary countable set (see examples below). The Abhyankar-Moh paper proved that the one dimensional Hausdorff measure of the convergence set of a divergent series is zero. In the case when n = 2, Pierre Lelong had earlier proved that if $\operatorname{Conv}(f)$ is not contained in a F_{σ} set of zero capacity then the series f is necessarily convergent, and conversely, given any set Econtained in a F_{σ} set of zero capacity a divergent power series f can be constructed so that $E \subset \operatorname{Conv}(f)$. This result has been rediscovered, independently, by several authors (see e.g. [LM], [Ne], [Sa], and see also [FM], [FM1], [Ne1], [Ri], for other related results). The optimal result was obtained by Sathaye [Sa] who strengthened the results of Abhyankar-Moh and Lelong by proving that a necessary and sufficient condition for a set $S \subset C$ to be equal to the convergence set of a divergent power series f(z) is that S is F_{σ} -set of transfinite diameter zero i.e. $S = \bigcup_{j=1}^{\infty} E_j$ where

²⁰⁰⁰ Mathematics Subject Classification. Primary: 26E05, 32A05, 30C85, 40A05.

Key words and phrases. convergence sets, formal power series, analytic functions, capacity, polar sets.

each E_j is closed set of transfinite diameter zero. These results can be viewed as an optimal and formal analogs of Hartogs' Theorem on separate analyticity in several complex variables.

In this article, we consider 'nonlinear' convergence sets of formal power series $f(y,t,x) \in \mathbb{C}[[y,t,x]], x = (x_1, x_2, ..., x_n)$ by restricting f(y,t,x) along a oneparameter family of 'tangential' perturbations of a fixed analytic curve $y = \varphi(t,x)$ through the origin. Throughout this paper, $\varphi(t,x) := \sum_{j=1}^{\infty} b_j(x)t^j$ will denote a fixed convergent power series where $b_j(x) := \sum_{i=0}^{\infty} b_{ji}x^i, j = 1, 2, 3, ...,$ are convergent power series in x with complex coefficients. We assume that $b_{10} = 1$.

For $s \in \mathbb{C}$, we put $\varphi_s(t,x) = \varphi(s,t,x) = sb_1(x)t + \sum_{j=2}^{\infty} b_j(x)t^j$. Define the φ -convergence set of a series $f(y,t,x) \in \mathbb{C}[[y,t,x]]$, as follows.

 $\operatorname{Conv}_{\varphi}(f) := \{ s \in \mathbb{C} : f(\varphi(s, t, x), t, x) \text{ converges as a series in } (t, x) \}.$

If K is an infinite compact set in \mathbb{C} , then for each positive integer n there is a unique monic polynomial $P_{n,K}(z)$ (called the Chebychev polynomial) such that

$$\rho_n(K) := \max_{z \in K} |P_{n,K}(z)| = \min_{P_n} (\max_{z \in K} |P_n(z)|)$$

where minimum is taken over the set of all monic polynomials $P_n(z) = z^n + a_1 z^{n-1} + \cdots + a_n$. The limit $\rho(K) := \lim_n (\rho_n(K))^{1/n}$ exists and is called the Chebychev constant of K. For a compact subset K of \mathbb{C} , $\rho(K)$ coincides with the logarithmic capacity $\mathfrak{c}(K)$ and the transfinite diameter $d_{\infty}(K)$ of K (see [Ah], Chapter 2). If $E = \bigcup_{n=1}^{\infty} K_n$, where K_n are compact sets of zero capacity, then $\mathfrak{c}(E) = 0$. A subset E of \mathbb{C} is zero capacity if and only if it is polar i.e. $E \subset \{z : u(z) = -\infty\}$ for some nonconstant subharmonic function $u : \mathbb{C} \to [-\infty, \infty)$. A F_{σ} set E in \mathbb{C} is said to have zero capacity if $\mathfrak{c}(E \cap \{|z| \le r\}) = 0$ for every $r \ge 0$.

Theorem. Let $\varphi(t, x)$ be as above, and let E be a subset of \mathbb{C} . There exists a divergent formal power series $f(y, t, x) \in \mathbb{C}[[y, t, x]]$ such that $E = \operatorname{Conv}_{\varphi}(f)$ if and only if E is an F_{σ} set of zero capacity.

PROOF. Suppose that E is an F_{σ} set with $\mathfrak{c}(E) > 0$. By replacing E with a compact subset $K \subset E$ of positive capacity we can assume that E is compact. Let $f(y,t,x) := \sum_{i,j,k} a_{ijk} y^i t^j x^k \in \mathbb{C}[[y,t,x]]$ be such that $g(s;t,x) := f(\varphi(s,t,x),t,x)$ converges for each $s \in E$. We need to show that f is convergent. Rewrite f as $f(y,t,x) = \sum_{i,j} a_{ij}(x) y^i t^j$, where $a_{ij}(x) := \sum_{k=0}^{\infty} a_{ijk} x^k \in \mathbb{C}[[x]]$, and

$$g(s;t,x) := \sum_{q \ge 0, k \ge 0} \lambda_{qk}(s) t^q x^k := \sum_{p \ge 0, q \ge 0} d_{pq}(x) s^p t^q.$$

It is clear that $\lambda_{qk}(s)$ is a polynomial of degree at most q, and thus

 $d_{pq}(x) = 0$, for p > q.

Let $d_{pq}(x) := \sum_{k \ge 0} d_{pqk} x^k$, and write $\lambda_{qk}(s) := \sum_{p=0}^q d_{pqk} s^p$. We have

$$d_{pq}(x) = \sum' a_{ij}(x) \frac{i!}{p!m_2!m_3!\cdots} b_1(x)^p b_2(x)^{m_2} b_3(x)^{m_3}\cdots$$

where the summation \sum' is taken over all nonnegative integers i, j, m_2, m_3, \ldots satisfying

$$j + p + 2m_2 + 3m_3 + \dots = q$$
 and $p + m_2 + m_3 + \dots = i$.

Since

$$\begin{array}{rcl} & d_{qq}(x) & = & a_{q,0}(x)b_1(x)^q, \\ (1) & d_{q-1,q}(x) & = & a_{q-1,1}(x)b_1(x)^{q-1}, \\ & d_{q-2,q}(x) & = & a_{q-2,2}(x)b_1(x)^{q-2} + a_{q-1,0}(x)(q-1)b_1(x)^{q-2}b_2(x), \\ & d_{q-k,q}(x) & = & a_{q-k,k}(x)b_1(x)^{q-k} + \text{terms involving } a_{ij}(x) \text{ with } i+j < q, \end{array}$$

it follows that $a_{ij}(x)$ can be solved uniquely in terms of $d_{pq}(x)$. In particular, if $d_{pq}(x) = 0$ for all p, q then all $a_{ij}(x) = 0$ for all i, j.

For each $s \in E$, there is a constant C_s such that $|\lambda_{qk}(s)| \leq C_s^{q+k}$ for all $q+k \geq 1$, since the power series $\sum_{a,k} \lambda_{qk}(s) t^q x^k$ converges. For each positive integer n, set

$$E_n = \{ s \in E : |\lambda_{qk}(s)| \le n^{q+k} \quad \forall q+k \ge 1 \}.$$

The sets E_n are closed and $E = \bigcup_{n=1}^{\infty} E_n$. There is a positive integer N such that $E' := \bigcup_{n=1}^{N} E_n$ has positive capacity. It follows that $|\lambda_{qk}(s)| \leq N^{q+k}$ for $q+k \geq 1$ and for $s \in E'$. By the Bernstein-Walsh inequality (see [**FM**], Lemma 1.4), there is a constant $C_{E'} \ge 1$ such that $|d_{pqk}| \le C_{E'}^q N^{q+k} \le (C_{E'}N)^{q+k}$. For some $\tau > 0$, g(s; t, x) represents a holomorphic function in $\Delta_{\tau} \times \Delta_{\tau} \times \Delta_{\tau}$,

where $\Delta_{\tau} = \{z \in \mathbb{C} : |z| < \tau\}$. Shrinking τ , if necessary, we may assume that

$$\min\{|b_1(x)| : x \in \mathbb{C}, |x| \le \tau\} \ge 1/2$$

and

(2)
$$\sum_{q,k} |b_{qk}| \tau^{q+k} < \infty, \quad \sum_{p,q,k} |d_{pqk}| \tau^{p+k} 2^q (\tau + \sum |b_{ij}| \tau^{i+j-1})^q < \infty.$$

The map $\psi : \mathbb{C}^{n+2} \to \mathbb{C}^{n+2}$ defined by $\psi(s,t,x) := (\varphi(s,t,x),t,x)$ is holomorphic near the origin and is injective on $Q = \{(s, t, x) \in \Delta_{\tau} \times \Delta_{\tau} \times \Delta_{\tau} : t \neq 0\}$. It follows that there is a holomorphic function G(u, v, w) defined on $\psi(Q)$ such that $g = G \circ \psi$ on Q.

We now prove that G extends holomorphically to a neighborhood of the origin. Choose a δ , $0 < \delta < \tau/2$, sufficiently small so that the set

$$\Gamma := \{ (u, v, w) \in \mathbb{C}^3 : |u| \le \delta^2, |v| = \delta, |w| \le \delta \}$$

is contained in $\psi(Q)$. The function G extends holomorphically to a neighborhood of the origin, if for $|u_0| < \delta^2$, $|w_0| < \delta$,

$$I_k(u_0, w_0) := \frac{1}{2\pi\sqrt{-1}} \int_{|v|=\delta} v^k G(u_0, v, w_0) \, dv = 0, \forall k = 0, 1, 2, \dots$$

For fixed w_0 and u_0 , write

$$\left(\frac{u_0 - (\varphi(t, w_0) - tb_1(w_0))}{tb_1(w_0)}\right)^p := \sum_{j=-p}^{\infty} c_{pj}(u_0, w_0)t^j.$$

By making use of (2), and substituting the above series expansion into the integrand in $I_k(u_0, w_0)$, we obtain

$$I_k(u_0, w_0) = \frac{1}{2\pi\sqrt{-1}} \int_{|t|=\delta} t^k g\left(\frac{u_0 - (\varphi(t, w_0) - tb_1(w_0))}{tb_1(w_0)}, t, w_0\right) dt$$

= $\sum d_{pq}(w_0) c_{pj}(u_0, w_0),$

where the sum is over all p, q, j with q+j = -k-1 and $j \ge -p$. Since q+j = -k-1and $j \ge -p$ imply $q = -j - k - 1 < -j \le p$, and since $d_{pq} = 0$ for q < p, we see that $I_k(u_0, w_0) = 0$ for k = 0, 1, 2, ... and for all (u_0, w_0) with $|u_0| < \delta^2$, $|w_0| < \delta$. Therefore, G extends holomorphically to a neighborhood of the origin.

Now $g = G \circ \psi$ on $\Delta_{\tau} \times \Delta_{\tau} \times \Delta_{\tau}$. Hence $g = G \circ \psi$ as a formal power series. Since $g = f \circ \psi$, we see that $\hat{f} \circ \psi := \sum \hat{d}_{pq}(x)s^p t^q = 0$, where $\hat{f} := \sum_{i,j} \hat{a}_{ij}(x)y^i t^j = f - G$. It follows that all $\hat{d}_{pq}(x)$, and hence $\hat{a}_{ij}(x)$ are all 0. This proves f is convergent as $\hat{f} = 0$ and $f \equiv G$.

Conversely, suppose E is an F_{σ} set with $\mathfrak{c}(E) = 0$. We construct a divergent power series f(y, t, x) such that $\operatorname{Conv}_{\varphi}(f) = E$.

By Theorem 6.1 of [**Ri**], there exists an increasing sequence $\{q_j\}$ of positive integers and a sequence of polynomials $\{P_j(s)\}$ with $\deg(P_j) \leq q_j$, for all j = 1, 2, 3, ..., such that the series $\psi_s(t) = \sum_j P_j(s)t^{q_j}$ converges for each $s \in E$, and diverges for each $s \notin E$. Set $g(s; t, x) := \psi_s(t) := \sum_{p,q} d_{pq}(x)s^pt^q$. We solve (1) for $a_{ij}(x)$ in terms of $d_{pq}(x)$, and set $f(y, t, x) = \sum a_{ij}(x)y^it^j$. Then, $f(\varphi(s, t, x), t, x) = g(s; t, x)$. Therefore, f(y, t, x) diverges, and $\operatorname{Conv}_{\varphi}(f) = E$. \Box

Corollary 1. For any $f(y, t, x) \in \mathbb{C}[[y, t, x]]$, either $\mathfrak{c}(\operatorname{Conv}_{\varphi}(f)) = 0$ or $\operatorname{Conv}_{\varphi}(f) = \mathbb{C}$.

Finally, we list some properties of φ -convergence sets that follow directly from the corresponding properties of polar sets.

Corollary 2. If $f \in \mathbb{C}[[y, t, x]]$ is divergent, then for any F_{σ} subset $E \subseteq \operatorname{Conv}_{\varphi}(f)$, there exists a divergent series $h(y, t, x) \in \mathbb{C}[[y, t, x]]$ such that $E = \operatorname{Conv}_{\varphi}(h)$.

Corollary 3. For any finite or infinite sequence $f_1, f_2, ..., f_n, ...$ in $\mathbb{C}[[y, t, x]]$, then there exists an $h \in \mathbb{C}[[y, t, x]]$ such that $\operatorname{Conv}_{\varphi}(h) = \bigcup_{i=1}^{\infty} \operatorname{Conv}_{\varphi}(f_i)$.

Corollary 4. Let $f \in \mathbb{C}[[y, t, x]]$, $\lambda \in \mathbb{C}$ and $P : \mathbb{C} \to \mathbb{C}$ be a monic polynomial function. There are series g and h in $\mathbb{C}[[y, t, x]]$ such that $\operatorname{Conv}_{\varphi}(g) = P^{-1}(\operatorname{Conv}_{\varphi}(f))$ and $\operatorname{Conv}_{\varphi}(h) = \lambda \cdot \operatorname{Conv}_{\varphi}(f)$.

PROOF. The corollary follows since

$$\mathfrak{c} \left(P^{-1} \left(\operatorname{Conv}_{\varphi}(f) \right) \right) = (\mathfrak{c} (\operatorname{Conv}_{\varphi}(f)))^{1/\deg P}$$
 and
 $\mathfrak{c} \left(\lambda \cdot \operatorname{Conv}_{\varphi}(f) \right) = |\lambda| \cdot \mathfrak{c} (\operatorname{Conv}_{\varphi}(f)).$

Since a locally polar set is polar by Josefson's theorem (see [Kl]), we have the following.

Corollary 5. An F_{σ} subset S of \mathbb{C} is locally a φ -convergence set of divergent power series if and only if it is φ -convergence set of a divergent power series. That is, if for every $s \in S$ there exists a neighborhood U of s in \mathbb{C} and a divergent power series f_U such that $S \cap U = \operatorname{Conv}_{\varphi}(f_U)$ then there exists a divergent power series f such that $S = \operatorname{Conv}_{\varphi}(f)$.

Example 1. (cf. **[AM]**) Let $\varphi_k(t, x)$ be a sequence of convergent power series with $\varphi_k(0, 0) = 0$. Then

(3)
$$f(y,t,x) := \sum_{n=1}^{\infty} n^n \prod_{j=1}^n \left[y - \varphi_j(t,x) \right] \text{ and }$$

(4)
$$g(y,t,x) := \sum_{i=1}^{\infty} \left[i!t^i + i!^2 (y - \varphi_j(t,x))^i \right]$$

are divergent power series but for each k = 1, 2, ..., the series $f(\varphi_k(t, x), t, x)$ converges while $g(\varphi_k(t, x), t, x)$ diverges. The empty set and finite or countable sets being sets of capacity zero are φ -convergence sets of divergent series. If $\{s_j\}_{j=1}^{\infty}$ is a sequence of complex numbers by taking $\varphi_j(t, x) = s_j t + \varphi(t, x)$ for all j in g(y, t, x) in (4), and by setting $\varphi_j(t, x) = \varphi(t, x)$ for all j in f(y, t, x) we obtain concrete examples of series with $\text{Conv}_{\varphi}(g) = \{s_i\}$ and $\text{Conv}_{\varphi}(f) = \phi$.

Example 2. The Cantor set C in \mathbb{R} has Hausdorff dimension $\ln 2/\ln 3$, hence of positive capacity, any series f(y, t, x) for which $f(st + \varphi(t, x), t, x)$ converges for all $s \in C$ is necessarily convergent. However, there exist modifications of Cantor sets that have capacity zero, and therefore are φ -convergence sets of divergent power series.

Remark 1. The closure of a φ -convergence set is not necessarily a φ -convergence set. For example, the countable set \mathbb{Q} is a φ -convergence set of divergent f in (3) but its closure \mathbb{R} being nonpolar can not be a φ -convergence set of a divergent series.

Remark 2. The situation is quite different, as one would expect, when restrictions of functions are considered. For example, for any positive integer k, it is elementary to construct a function $f : \mathbb{R}^n \to \mathbb{R}$ that is exactly k-times differentiable but whose restriction to every line in \mathbb{R}^n is real-analytic. The function

$$f(x,y) := \begin{cases} (x^2 + y^2) \exp\left(-\frac{y^2}{x^2 + y^4}\right) & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0) \end{cases}$$

is in $C^{\infty}(\mathbb{R}^2)$ and for all $m \neq 0$ the single-variable function f(mt, t) is real-analytic. However, f is not a real-analytic function as it fails to be real-analytic along the y-axis. Is there a C^{∞} function $f : \mathbb{R}^n \to \mathbb{R}$ which is not real-analytic but whose restriction to *every* line is real-analytic? The answer is negative as it was shown by J. Bochnak [**Bo**] and J. Siciak [**Si**] that if a C^{∞} function $f : \mathbb{R}^n \to \mathbb{R}$ is real-analytic on every line segment through a point x_0 , then f is real-analytic in a neighborhood of x_0 . (See [**Ne**] and [**Ne1**] for C^{∞} -analogs of the Bochnak-Siciak theorem.)

References

- [Ah] L.V. Ahlfors, Conformal invariants: topics in geometric function theory, McGraw-Hill, N.Y., 1973.
- [AM] S.S. Abhyankar, T.T. Moh, A reduction theorem for divergent power series, J. Reine Angew. Math., 241 (1970), 27–33.
- [Bo] Bochnak, J., Analytic Functions in Banach Spaces, Studia Mathematica. T, XXXV (1970) 273-292.
- [FM] B.L. Fridman, D. Ma, Osgood-Hartogs type properties of power series and smooth functions, *Pacific J. Math.*, 251 (2011), 67–79.
- [FM1] B.L. Fridman, D. Ma, Testing holomorphy on curves, Israel J. Math., to appear.
- [Kl] M. Klimek, *Pluripotential Theory*, Clarendon Press, New York, 1991.
- [Le] P. Lelong, On a problem of M. A. Zorn, Proc. Amer. Math. Soc., 2 (1951), 11–19.
- [LM] N. Levenberg, and R.E. Molzon, Convergence sets of a formal power series, Math. Z., 197 (1988), 411–420.
- [Ne] T.S. Neelon, A Bernstein-Walsh type inequality and applications, Canadian Math. Bull. 49 (2) 2006 pp. 256-264.
- [Ne1] T.S. Neelon, Restrictions of power series and functions to algebraic surfaces, Analysis (Munich), 29 (2009), no. 1, 1–15.
- [Ri] J. Ribon, Holomorphic extensions of formal objects, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 3 (2004), no. 4, 657–680.

- [Re] R. Ree, On a problem of Max A. Zorn, Bull. Amer. Math. Soc., 55 (1949), 575–576.
- [Sa] A. Sathaye, Convergence sets of divergent power series, J. Reine Angew. Math., 283 (1976), 86–98.
- [Si] Siciak, J., A Characterization of Analytic Functions of n Real Variables, Studia Mathematica. T, XXXV (1970) 293-297.

fridman@math.wichita.edu, Department of Mathematics, Wichita State University, Wichita, KS 67260-0033, USA

 $\rm DMA@MATH.wichita.edu,$ Department of Mathematics, Wichita State University, Wichita, KS 67260-0033, USA

Neelon@csusm.edu, Department of Mathematics, California State University San Marcos, San Marcos CA 92096-0001, USA

6