

On Sisterhood in the Gale-Shapley Matching Algorithm

Yannai A. Gonczarowski^{*} and Ehud Friedgut[†]

Abstract

Lying in order to manipulate the Gale-Shapley matching algorithm has been studied in [2] and [3] and was shown to be generally more appealing to the proposed-to side (denoted as the women in [1]) than to the proposing side (denoted as men there). It can also be shown that in the case of lying women, for every woman who is better-off due to lying, there exists a man who is worse-off.

In this paper, we show that an even stronger dichotomy between the goals of the sexes holds, namely, if no woman is worse-off then no man is better-off, while a form of sisterhood between the lying and the “innocent” women also holds, namely, if none of the former are worse-off, then neither is any of the latter.

This paper is based upon an undergraduate (“Amirim”) thesis of the first author.

1 Background

1.1 The Gale-Shapley Algorithm

In order to standardize the notation used throughout this article, and for sake of self-containment, let us quickly recap the scenario introduced in [1] and the Gale-Shapley Algorithm introduced there:

^{*}Einstein Institute of Mathematics, Hebrew University, Jerusalem, Israel. Email: yannai@gonch.name

[†]Einstein Institute of Mathematics, Hebrew University, Jerusalem, Israel. Email: ehud.friedgut@gmail.com

Let W and M be equally-sized finite sets of women and men, respectively. Let each member of these sets have a strict order of preference regarding the members of the set this member does not belong to.

Definition 1.1. *A one-to-one map between W and M is called a matching.*

Definition 1.2. *A matching is called unstable under the given orders of preference if there exist two matched couples (w, m) and (\tilde{w}, \tilde{m}) such that w prefers \tilde{m} over m and \tilde{m} prefers w over \tilde{w} . A matching which is not unstable is called stable.*

Theorem 1.3 ([1]). *The following algorithm stops and yields a stable matching between W and M (in particular, such a matching exists), and no stable matching is better for any man $m \in M$: The algorithm is divided into steps, to which we shall refer as “nights”. On each night, each man serenades under the window of the woman he prefers most among all women who have not (yet) rejected him, and then each woman, under whose window more than one man serenades, rejects every man who serenades under her window, except for the man she prefers most among these men. The algorithm stops on a night on which no man is rejected by any woman, and then each woman is matched with whoever has serenaded under her window on this night.*

Generalizations to this algorithm which include different sizes for W and M , preference lists which do not include all the members of the opposite sex (“blacklisting”), and one-to-many matchings, are also presented in [1] and will be discussed in section 3.

1.2 Previous Results

As quoted above, the stable matching given by the Gale-Shapley algorithm is optimal (out of all stable matchings) for each man. A somewhat reverse claim holds regarding the women:

Theorem 1.4 ([6], derived from a more general theorem by Knuth). *No stable matching is worse for any woman $w \in W$ than the matching given by the Gale-Shapley algorithm.*

The benefits of the Gale-Shapley matching algorithm for the men have been demonstrated even further in [2], where it was shown that no man can get a better match by lying about his preferences (i.e., manipulate the algorithm by declaring a false order of preference, assuming the algorithm

is run according to the “true” preferences of all women and all other men) and that no subset of the men can all get better matches by lying in a coordinated fashion.

These observations lead to the analysis of the profitability of lying by women in [3], where it is proven that if more than one stable matching exists, then at least one woman can get a better match by lying about her preferences.

It is easy to show that any woman who is better off as a result of someone’s lie (whoever that liar or those liars may be and whatever their lie may be) is matched (due to the lie) to someone who is now worse off. (Indeed, if they are both better off, then the original match can not be stable.) In other words, for every woman who is better-off, some man is worse-off.

As mentioned above, in the next section we will show that an even stronger dichotomy between the goals of the sexes holds, while a form of sisterhood between the lying and the “innocent” women also holds.

2 Monogamous Matchings

2.1 The Theorem

Let W and M be equally-sized finite sets of women and men, respectively. Let each member of these sets be endowed with a strict order of preference regarding the members of the other set. These will be referred to as “the true preferences”, the application of the Gale-Shapley algorithm according to these preferences will be referred to as OA (original algorithm), and resulting matching will be referred to as “the original matching”.

Assume that a subset of the women, denoted L (for liars) declare false orders of preference for themselves. Call the application of the Gale-Shapley algorithm according to these false preferences for the members of L , and according to the real preferences of other members of W (referred to, henceforth, as innocent) and of all the members of M , NA , and the resulting matching — “the new matching”.

Definition 2.1. *A person $p \in W \cup M$ is said to be “better-off” (resp. “worse-off”) if p prefers, according to their true order of preference, their match according to the new matching (resp. the original matching) over their match according to the original matching (resp. the new matching).*

Let us now phrase our main result for the above conditions.

Theorem 2.2. *Under the above conditions, if no lying woman is worse-off, then:*

- (a) *No woman is worse-off.*
- (b) *No man is better-off.*

2.2 Proofs for a Special Case

Definition 2.3. *A woman $l \in L$ is said to be lying in a personally-optimal way if, all other orders of preference being the same, there is no other false order of preference she can declare which will result in her being even better-off, i.e. her being matched with a man she (truly) prefers over the man matched to her by the new matching.*

Theorem 2.2 is easily provable in the special case that all women in L lie in a personally-optimal way, as it is possible to show that in this case, the new matching is stable under the true preferences, and thus, no woman is worse-off, and no man is better-off, than under the original matching. Nonetheless, it will be shown below that in some cases, when liars may coordinate their lies, it is logical to lie in a non-personally-optimal way, and that in these cases the resulting matching might be unstable under the true preferences.

It can be noted that if we examine a “lying game” between the member of L , where a player’s strategy is a declaration of a specific order of preference for themselves, and the utility for each player is determined by the position of her new match on their true preference list, then in this game, the conditions of the special case described above are met if and only if the set of lies constitutes a Nash equilibrium.

Roth (private communication, Dec. 2007) suggested the following sketch of a proof to part a of Theorem 2.2:

- If women can do better than to state their true preferences, they can do so by truncating their preferences.
- Truncating preferences is the opposite of extending preferences (as discussed in [6] in the context of adding new players)
- When any woman extends her preferences, it harms the other women.

(As already noted, truncating preference lists is discussed in section 3 below.) While this also proves Theorem 2.2 in the special case discussed above (in which all women lie in a personally-optimal way), it appears not to prove Theorem 2.2 itself (i.e., with no additional assumptions), as while it is true that if a woman can do better than to state her true preferences then she can do better by truncating them (and even more so, there always exists a truncation of her true preferences which constitutes a personally-optimal lie for her), it turns out that there may exist some man whom she can secure for herself by submitting false preferences, but not by truncating her true preferences. This will be illustrated by an example below.

2.3 When a Lie Needs Not be Optimal

As was shown in section 2.2, Theorem 2.2 is easily provable if each lying woman lies in a personally-optimal way, or if the new matching is stable under everyone's true preferences.

Before continuing to the main proof, let us give an example of a scenario with the following properties:

1. No woman can do better by lying alone (while all others tell the truth).
2. In every conspiracy by more than one woman to lie so that none of them are worse-off and at least one is better-off, there exists a woman who does not lie in a personally-optimal way. In other words, it is logical to lie in a non-optimal way.
3. The resulting matching is not stable under the true preferences and can not be achieved by simply truncating women's true preference lists (even if that is allowed).

This example includes four women (w_1, \dots, w_4) and four men (m_1, \dots, m_4). The orders of preference for the women fulfill:

- w_1 : First choice: m_3 , second choice: m_1 .
- w_2 : First choice: m_3 , second choice: m_1 .
- w_3 : Prefers m_2 over m_1 and prefers m_1 over m_3 .

- w_4 : Any order of preference.

The orders of preference for the men fulfill:

Man	1st choice	2nd choice	3rd choice	4th choice
m_1	w_1	w_3	w_2	w_4
m_2	w_2	w_3	any	any
m_3	w_3	w_2	w_1	w_4
m_4	w_1	w_4	any	any

Let us examine OA :

Night	Window	w_1	w_2	w_3	w_4
	1		m_4, m_1	m_2	m_3
2		m_1	m_2	m_3	m_4

It is clear from this examination that any conspiring subset of the women wishing to make a difference (in the scenario which does not allow truncating preference lists) has to include w_1 . Also, it can be verified that w_1 can not lie alone (while all others tell the truth) and become better-off. Indeed, in order to become better-off, w_1 has to be matched, under the new matching, with m_3 , and since he prefers w_2 over her, this entails his rejection by w_2 , but since he is the first choice of w_2 , this means w_2 has to lie and declare that she prefers another man over him. Let us assume, then, that $L = \{w_1, w_2\}$. (It can be verified that the result would not change even if we admit more women to L .)

It can easily be verified that, given the true orders of preference for everyone except w_1 and w_2 , there exists exactly one combination of false orders of preference for these two women (up to changes in their preferences regarding men who will not reach their windows as long as everyone else tells the truth) which will cause them both to be better-off:

- w_1 must declare she prefers m_3 over m_4 and m_4 over m_1 .
- w_2 must declare she prefers m_1 over m_3 and m_3 over m_2 .

Let us examine NA under these false orders of preference:

Night	Window			
	w_1	w_2	w_3	w_4
1	m_4, m_1	m_2	m_3	
2	m_4	m_2	m_1, m_3	
3	m_4	m_2, m_3	m_1	
4	m_4	m_3	m_1, m_2	
5	m_4	m_1, m_3	m_2	
6	m_3, m_4	m_1	m_2	
7	m_3	m_1	m_2	m_4

Indeed, w_1 and w_2 are both better-off (and, as Theorem 2.2 states, no other woman is worse-off, and the even innocent w_3 is better-off as well) however this lie is clearly not personally-optimal for w_2 , since she can declare any order of preference under which her first choice is m_3 , and become even better-off by being matched with him, but if she tries to do that (either by telling the truth, or by lying), then w_1 becomes worse-off and in this case it is better for w_1 to tell the truth which, as stated earlier, will cause the new matching to be identical to the old matching, and hence will cause w_2 to be matched with m_2 , ending up in a worse situation than had she lied the above-described non-personally-optimal lie.

In the above-described lying game between w_1 and w_2 , the utility for each of the players given these strategies is higher than her utility in any Nash equilibrium, and this is the only pair of strategies (up to the degrees of freedom discussed above) with this property.

It should be noted that the resulting new matching is neither stable under the original preferences (as w_2 and m_3 truly prefer each other over their respective matches) nor can it be achieved by simply truncating the preference lists of w_1 and w_2 (as any such truncation will result in w_2 either having a blank list or a list with m_3 as her first choice), giving, as promised, an example where the above proofs do not hold.

2.4 A General Proof

Before proving the general case of Theorem 2.2, let us first introduce a few notations: For each person $p \in W \cup M$, let us denote the person from the opposite sex matched with p under the original matching (resp. new matching) with $O(p)$ (resp. $N(p)$).

Definition 2.4. A woman $w \in W$ is said to be a rejecter if she rejected $N(w)$ during OA . In this case, let the man who serenaded under her window on the night of OA on which she rejected $N(w)$, but whom she did not reject on that night (and whom, therefore, she prefers over $N(w)$), be denoted $B(w)$.

Lemma 2.5. If a man $m \in M$ never serenaded under the window of a woman $w \in W$ during NA (resp. OA), then m prefers $N(m)$ (resp. $O(m)$) over w .

Proof. Since m approaches women according to his own order of preference, and since he never reached w 's window during NA (resp. OA), then he must have ended up being matched, under the new (resp. old) matching, to a woman he prefers over w , and by definition, that woman is $N(m)$ (resp. $O(m)$). \square

By the following Lemma, we need only prove part b of Theorem 2.2.

Lemma 2.6. If a woman $w \in W$ is worse-off, then $O(w)$ is better-off.

Proof. Since w is worse-off, w prefers $O(w)$ over $N(w)$ according to her true order of preference. Since it is given that no liar is worse-off, then w is not a liar, and therefore she declared that she prefers $O(w)$ over $N(w)$ also during NA . Therefore, since w prefers $N(w)$ over any other man who serenaded under her window during NA , it follows that $O(w)$ could not have serenaded under her window on any night during NA . Thus, by Lemma 2.5, $O(w)$ prefers $N(O(w))$ over $w = O(O(w))$, making $O(w)$ better-off. \square

Lemma 2.7. If a man $m \in M$ is better-off, then $N(m)$ is a rejecter.

Proof. Since m is better-off, m prefers $O(m)$ over $N(m)$. Therefore, since m approaches women according to his own order of preference, m would not serenade under $O(m)$'s window before first having been rejected by $N(m)$. Now, since m serenades under $O(m)$'s window during the last night of OA , it follows that he must have been rejected by $N(m)$ on some earlier night during OA . Since $N(m)$ rejected $m = N(N(m))$ during OA , then she is a rejecter. \square

Lemma 2.8. If a woman $w \in W$ is a rejecter, then she is worse-off.

Proof. By induction and transitivity, w prefers $O(w)$ over any man she rejected during OA and therefore, being a rejecter, over $N(w)$, and is thus worse-off. \square

Lemma 2.9. *If a woman $w \in W$ is a rejecter, then $B(w)$ prefers $N(B(w))$ over w .*

Proof. By Lemma 2.8, w is not a liar, and as such, her order of preference during NA is her true order of preference. Since w is a rejecter, w prefers $B(w)$ over $N(w)$. Therefore, as w truly prefers $N(w)$ over any other man who serenaded under her window during NA , it follows that $B(w)$ could not have serenaded under w 's window during NA . Thus, by Lemma 2.5, $B(w)$ prefers $N(B(w))$ over w . \square

Lemma 2.10. *If a woman $w \in W$ is a rejecter, then*

1. $N(B(w))$ is a rejecter.
2. During OA , w rejected $N(w)$ on a strictly later night than the night on which $N(B(w))$ rejected $B(w)$.

Proof. Throughout this proof, for the sake of conciseness, wherever we discuss whether a rejection or serenading occurred, or when it occurred, we always refer to OA . Since w is a rejecter, then according to Lemma 2.9, $B(w)$ prefers $N(B(w))$ over w . Also note that by definition of $B(w)$, $B(w)$ serenaded under w 's window on some night. By these two observations, and since $B(w)$ approaches women according to his own order of preference, $N(B(w))$ rejected $B(w)$, and did so on a night strictly earlier than any night on which $B(w)$ serenaded under w 's window. Since $N(B(w))$ rejected $B(w) = N(N(B(w)))$, then she is a rejecter. Moreover, since, by definition of $B(w)$, w rejected $N(w)$ on a night on which $B(w)$ serenaded under her window as well, then this night is a strictly later night than the night on which $N(B(w))$ rejected $B(w)$. \square

To complete the proof of Theorem 2.2, assume for contradiction that there exists a better-off man $m \in M$. Let us denote $w_1 = N(m)$, and by Lemma 2.7, she is a rejecter. Now, for each $i \in \mathbb{N}$, assume by induction that w_i is a rejecter and set $w_{i+1} = N(B(w_i))$. By Lemma 2.10, w_{i+1} is a rejecter. Moreover, by that lemma, during OA , w_i rejected $N(w_i)$ on a night strictly later than the night on which w_{i+1} rejected $B(w_i) = N(N(B(w_i))) = N(w_{i+1})$.

From finiteness of W , there must exist $i < j$ such that $w_i = w_j$, but by induction, since $i < j$, then during OA , w_i rejected $N(w_i)$ on a night strictly later than the night on which w_j rejected $N(w_j)$ — a contradiction. \square

Corollary 2.11. *Under the conditions of Theorem 2.2, every better-off woman, is matched to a worse-off man.*

3 Generalizations

3.1 Polygamous Matchings

In [1], a one-to-many version of the algorithm was specified and proposed as a way to assign students to colleges (on each algorithm-step, the students apply to their favorite not-yet-rejected-by college, and then each college rejects all applicants except for the most preferred ones, according to the quota of this college) and proved that the resulting matching is stable and that it is optimal (within all stable matches) for each applicant. It can also be shown that it is the worst (within all stable matches) for each college.

In [4], it was made public that the assignment of medical interns to hospitals in the USA has been done using a similar algorithm since 1951, however, in this algorithm the roles were switched and the hospitals were the ‘proposers’ and the resulting matching is optimal (within all stable matches) for each hospital and worst for each intern.

In this subsection, we will prove a generalization of Theorem 2.2 for these scenarios. In order to ease the transition from the previous section, we will maintain the notation of women and men, and refer to the above scenarios as “the polygamous scenarios”. Let us redefine our notation for these scenarios:

Let W and M be finite lists of women and men, respectively, and let each person be endowed, as before, with a strict order of preference with regards to the members of the opposite sex. For each person $p \in W \cup M$, define n_p to be the “quota” of this person, i.e., the amount of spouses from the opposite sex this person seeks. As our goal is the generalization of both algorithms described above, the reader may assume, for ease of readability, that either all women are monogamous ($\forall w \in W : n_w = 1$) or all men are monogamous ($\forall m \in M : n_m = 1$), however the rest of this paper holds verbatim even if this is not the case. (If both the women and men are monogamous we are reduced to the scenario given in the previous section.) We will also assume for now that $\sum_{w \in W} n_w = \sum_{m \in M} n_m$. (This guarantees that when the algorithm stops, each person p is matched with

exactly n_p people of the opposite sex — this reduces in the monogamous case to W and M being of identical size.)

Definition 3.1. *A map between W and M , mapping each $p \in W \cup M$ to exactly n_p members of the set p does not belong to, is called a matching.*

The definition of instability of a matching requires another detail which was inferred from the other requirements in the monogamous scenario:

Definition 3.2. *A matching is called unstable under the given orders of preference if there exist two matched couples (w, m) and (\tilde{w}, \tilde{m}) such that:*

1. w is not matched with \tilde{m} .
2. w prefers \tilde{m} over m .
3. \tilde{m} prefers w over \tilde{w} .

As before, a matching which is not unstable is called stable.

On each night of the polygamous algorithm, each man $m \in M$ serenades under the windows of n_m women that he prefers the most out of all women who have not (yet) rejected him, and then each woman $w \in W$, under whose window more than n_w men serenade, rejects every man who serenaded under her window, except for the n_w men she prefers most among them.

Now, as before, assume that a subset of the women, denoted L , declare false orders of preference for themselves.

For a person $p \in P$, Let us denote the set of people of the opposite sex matched with this person under the original matching (resp. the new matching) with $O(p) = \{o_1^p, \dots, o_{n_p}^p\}$ (resp. $N(p) = \{n_1^p, \dots, n_{n_p}^p\}$) such that p prefers o_i^p over o_{i+1}^p (resp. prefers n_i^p over n_{i+1}^p).

Before we phrase the polygamous version of Theorem 2.2, we have to redefine the circumstances under which a person is said to be better-, or worse-off. It should be noted that while in the monogamous scenario each person's order of preference yields a full order on the set of possible matches for that person (i.e., people of the opposite sex), in the polygamous case each person's order of preference yields a partial order on the set of possible matches for that person (i.e., n_p -tuples of people of the opposite sex). This introduces an asymmetry between the following two definitions, which did not exist in the monogamous scenario.

Definition 3.3. A woman $w \in W$ is said to be weakly better-off if for each $1 \leq i \leq n_w$, w does not prefer o_i^w over n_i^w .

Definition 3.4. A man $m \in M$ is said to have gained only worse matches if he prefers each member of $O(w)$ over each member of $N(w) \setminus O(w)$. (Note that this condition is met in the special case in which $N(w) = O(w)$.)

Theorem 3.5. Under the above conditions, if all lying women are weakly better-off, then:

- (a) All women are weakly better-off.
- (b) All men have gained only worse matches.

The special case of Theorem 3.5 in which all women are polygamous and men are monogamous can be easily proven by reduction to the conditions of Theorem 2.2 by “replicating” each polygamous woman $w \in W$ into n_w monogamous women $\{(w, i)\}_{i=1}^{n_w}$, each having the same order of preference as w . For each man $m \in M$, replace w on his list of preferences with these women, in such a way that he prefers (w, i) over $(w, i + 1)$ for all i . A reduction along these lines was used in [2] to generalize certain properties of the monogamous scenario to the polygamous-women scenario and it was shown there that the men matched with $(w, 1), \dots, (w, n_w)$ by the monogamous algorithm are exactly those matched with w by the polygamous algorithm. Furthermore, in the notations of this paper, it can be shown that the man matched with (w, i) by *OA* is o_i^w and by *NA* is n_i^w , thus completing the reduction since this yields that w is weakly better-off if and only if none of the monogamous women (w, i) is worse-off, and since it is clear that as all men are monogamous, if a man is not better-off under the reduction, then he has only gained worse matches before the reduction.

Unfortunately, “replicating” each man in a similar same way will not yield a proof for even the monogamous women - polygamous men scenario as easily, for it is possible for a woman w to be weakly better-off before the reduction by maintaining the same match m , but to be worse-off under the reduction because her match is e.g. $(m, 2)$ instead of $(m, 1)$. Also, every replicated monogamous man not being better-off under the reduction does not necessarily imply that every polygamous man has gained only worse matches before the reduction. To make things even worse, in

the general case where both men and women may be polygamous, running the monogamous algorithm after such a “replication” does not even produce the same matching as would be produced by the polygamous algorithm, as it may lead to situations such as a couple matched to each other with multiplicity greater than 1.

Indeed, in order to prove Theorem 3.5 in its general form we have to retrace our steps and revisit the inner workings of the proof of Theorem 2.2, rewriting it to accommodate for the generalizations we introduced. We will now redefine some of the definitions used in that proof, and then give its generalized form.

Definition 3.6. *A woman $w \in W$ is said to be a rejecter if she rejected any of the members of $N(w)$ during OA . Let us denote the set of all such rejected members $R(w)$.*

Definition 3.7. *A man $m \in M$ is said to be a rejectee if there exists a rejecter $w \in N(m)$ such that $m \in R(w)$. Similarly to the proof of Theorem 2.2, a key role will be played by a man $B(w, m)$ for which, in a sense, m was rejected. The exact definition of this man will appear in Lemma 3.11.*

As before, we will begin with a Lemma by which we need only prove part b of Theorem 3.5.

Lemma 3.8. *If a woman $w \in W$ is not weakly better-off, then there exists $m \in O(w)$ who has not gained only worse matches.*

Proof. Since it is given that all liars are weakly better-off, it follows that w is not a liar, and therefore her order of preference during NA is her true order of preference. Since w is not weakly better-off, there exists $1 \leq i \leq n_w$ such that w prefers o_i^w over n_i^w and thus prefers o_1^w, \dots, o_i^w over n_i^w . By induction and by the definition of the polygamous algorithm, there are exactly $i - 1$ men serenaded under w 's window during NA and that she (truly) prefers over n_i^w (these are n_1^w, \dots, n_{i-1}^w), so by the pigeonhole principle, there exists $1 \leq j \leq i$ such that o_j^w did not serenade under w 's window during NA . Therefore, since o_j^w approaches women according to his own order of preference, o_j^w prefers all women in $N(o_j^w)$ over w . Since $|N(o_j^w)| = n_{o_j^w} = |O(o_j^w)|$, and since $w \in O(o_j^w) \setminus N(o_j^w)$, it follows that there exists $\tilde{w} \in N(o_j^w) \setminus O(o_j^w)$ and as stated, o_j^w prefers her over w and hence (as $w \in O(o_j^w)$) o_j^w has not gained only worse matches. \square

Lemma 3.9. *If a man $m \in M$ has not gained only worse matches, then he is a rejectee.*

Proof. Since m has not gained only worse matches, there exist $w \in O(m)$ and $\tilde{w} \in N(m) \setminus O(m)$ such that m prefers \tilde{w} over w . Since $w \in O(m)$, it follows that m serenaded under w 's window during OA . Since m approaches women according to his own order of preference, he would not have serenaded under w 's window during OA without having serenaded on the same night, or on a previous night, under \tilde{w} 's window. However, since m is not matched with \tilde{w} at the end of OA , then \tilde{w} must have rejected him during OA and therefore, by definition, $m \in R(\tilde{w})$, and thus m is a rejectee. \square

Lemma 3.10. *If a woman $w \in W$ is a rejecter, then she is not weakly better-off.*

Proof. Since w is a rejecter, there exists $r \in R(w)$ and by the definition of $R(w)$ there exists $1 \leq i \leq n_w$ such that $n_i^w = r$. Since, by definition of $R(w)$, w rejected r during OA , and since, by induction and transitivity, w prefers each member of $O(w)$ over any man she rejected during OA , it follows that she prefers o_i^w over $r = n_i^w$, and is, thus, not weakly better-off. \square

Lemma 3.11. *If a woman $w \in W$ is a rejecter, then for each $r \in R(w)$ there exists a man $B(w, r)$ such that*

1. $B(w, r)$ serenaded under w 's window during OA on the night on which she rejected r , but $B(w, r)$ was not rejected by her on that night.
2. $B(w, r)$ prefers each member of $N(B(w, r))$ over w .

(If more than one such man exists, define $B(w, r)$ to be one of these men, arbitrarily.)

Proof. By Lemma 3.10, w is not a liar, and as such, her order of preference during NA is her true order of preference. Let B be the set of all men who serenaded under her window on the night during OA on which she rejected r , but who were not rejected by her on that night. By definition of the algorithm, w prefers each member of B over r , and $|B| = n_w$. Since w did not reject r during NA (since $r \in N(w)$), and since the order of preference of w during NA is her true order of preference, then not all member of B serenaded under her window during NA (for she is matched under NA to the set of the n_w men that she prefers most out of all the men

who serenaded under her window during NA). Define, therefore, $B(w, r)$ to be a member of B who did not serenade under w 's window during NA . Since $B(w, r)$ did not serenade under w 's window during NA , and since he approaches women according to his own order of preference, it follows that he prefers each member of $N(B(w, r))$ over w . \square

Lemma 3.12. *If a woman $w \in W$ is a rejecter, then for each $r \in R(w)$ there exists a woman $\tilde{w} \in N(B(w, r))$ such that*

1. \tilde{w} is a rejecter.
2. $B(w, r) \in R(\tilde{w})$
3. During OA , w rejected r on a strictly later night than the night on which \tilde{w} rejected $B(w, r)$.

Proof. Once again, throughout this proof, for the sake of conciseness, wherever we discuss whether a rejection or serenading occurred, or when it occurred, we always refer to OA . Since w is a rejecter, then by Lemma 3.11, $B(w, r)$ serenaded under w 's window (on the night on which she rejected r) and $B(w, r)$ prefers each member of $N(B(w, r))$ over w . Hence, as $B(w, r)$ approaches $n_{B(w, r)}$ women each night according to his own order of preference, and as $|N(B(w, r))| = n_{B(w, r)}$, it follows that $B(w, r)$ serenaded on earlier nights under each of the windows of $N(B(w, r))$ and was rejected by at least one of them on a night strictly earlier than any night on which he serenaded under w 's window — let us denote such a woman \tilde{w} . Since \tilde{w} rejected $B(w, r)$ (to whom she is matched under the new matching), it follows that she is a rejecter and that $B(w, r) \in R(\tilde{w})$. Moreover, since, by definition of $B(w, r)$, w rejected r on a night on which $B(w, r)$ serenaded under her window as well, then this night is a strictly later night than the night on which \tilde{w} rejected $B(w, r)$. \square

To complete the proof of Theorem 3.5, assume for contradiction that there exists a man $m_1 \in M$ who has not gained only worse matches. By Lemma 3.9, m_1 is a rejectee, therefore there exists a rejecter $w_1 \in N(m_1)$ such that $m_1 \in R(w_1)$. Now, for each $i \in \mathbb{N}$, assume by induction that w_i is a rejectee and that $m_i \in R(w_i)$ and set, by Lemma 3.11, $m_{i+1} = B(w_i, m_i)$. By Lemma 3.12, there exists a rejecter $w_{i+1} \in N(m_{i+1})$. Moreover, by that lemma, during OA , w_i rejected m_i on a night strictly later than the night on which w_{i+1} rejected m_{i+1} .

From finiteness of $W \times M$, there must exist $i < j$ such that $w_i = w_j$ and $m_i = m_j$, but by induction, since $i < j$, then during OA , w_i rejected m_i on a night strictly later than the night on which w_j rejected m_j — a contradiction. \square

3.2 Blacklists and Mismatched Quotas

As mentioned before, in [1], it is not required that the sum of the quotas of all colleges be the same as the number of applicants, resulting in some colleges not fulfilling their quotas or some applicants not being accepted to any college. Moreover, it is allowed for a college to remove some of the students from its preference list, indicating that the college is unwilling to accept these candidates even if it means that its quota will not be met. Similarly, it is allowed for an applicant to remove some of the colleges from their preference lists, indicating their unwillingness to attend these colleges even at the risk of not being accepted to any college.

The modified algorithm-step for this scenario (in the notations used throughout this document) is that if, by a certain night, the number of women who have not blacklisted or (yet) rejected a man $m \in M$, and are not blacklisted by him, is less than n_m , then on that night he will serenade under the windows of all these women. (Otherwise, if there are at least n_m such women, he will, as before, serenade under the windows of the n_m women that he prefers most out of them.)

Let us adjust our notations for this scenario, and then prove that the result of this paper still holds under it:

Definition 3.13. *A map between W and M , mapping each $p \in W \cup M$ to at most n_p members of the set p does not belong to, is called a matching.*

Definition 3.14. *A matching is called unstable under the given orders of preference if either the conditions of Definition 3.2 are met, or if there exists a matched couple (w, m) and a woman (resp. man) p such that:*

1. p is matched with less than n_p spouses.
2. p has not blacklisted m (resp. w).
3. m (resp. w) prefers p over w (resp. m).

Once again, a matching which is not unstable is called stable.

For a person $p \in P$, let us denote $O(p)$ and $N(p)$ as before, and for each $1 \leq i \leq |O(p)|$ (resp. $1 \leq i \leq |N(p)|$), let us denote o_i^p (res. n_i^p) as before as well.

Definition 3.15. *A woman $w \in W$ is said to be weakly better-off if the following conditions hold:*

1. $N(w)$ contains none of the men blacklisted by w .
2. $|O(w)| \leq |N(w)|$
3. For each $1 \leq i \leq O(w)$, w does not prefer o_i^w over n_i^w .

If, in addition, either for some $1 \leq i \leq O(w)$, w prefers n_i^w over o_i^w (“ w has improved her matches”), or $|O(w)| < |N(w)|$ (“ w has gained matches”), then w is said to be better-off.

The definition of a man having gained only worse matches remains unchanged. Specifically, it should be emphasized that this definition does not require that $|N(m)| \leq |O(m)|$, but we will show in Corollary 3.17 that under the above conditions the inverse is an impossibility. Similarly, the same corollary will show that under the above conditions, a woman may not be better-off due to gaining matches, but only due to improving them. It should also be noted that, by the definition of the algorithm, it is not possible for any person declaring their true preferences to be matched, under the new matching, to any person blacklisted by them.

Corollary 3.16. *Theorem 3.5 holds under the above conditions.*

Proof. Corollary 3.16 can be proven by reducing to the conditions of Theorem 3.5 by introducing, for each person p , n_p new monogamous people of the opposite sex $\emptyset_1^p, \dots, \emptyset_{n_p}^p$, each of whom prefers p the most (the rest of their orders of preference can be arbitrary). The original (resp. new) order of preference for p will now be: first, all of the original people of the opposite sex who are not originally (resp. newly) blacklisted by p , ordered according to her or his given original (resp. new) order of preference, then $\emptyset_1^p, \dots, \emptyset_{n_p}^p$ in this order, and then, in an arbitrary order, the original (resp. new) blacklist of p and the people newly-introduced for the other people of the same sex as p . It is straight-forward to verify that the quotas of all women (original and newly-introduced) match the quotas of all men (original and newly-introduced). It is left for the reader to verify that the

old (resp. new) matching before the reduction is identical to the old (resp. new) matching under the reduction, after the newly-introduced people have been removed from it, that each of the original women is weakly better-off before the reduction iff she is weakly better-off under the reduction, and that if each of the original men has gained only worse matches under the reduction, then he has also gained only worse matches before the reduction. \square

Corollary 3.17. *Under the conditions of Corollary 3.16, for each person $p \in W \cup M$, it holds that $|N(p)| = |O(p)|$.*

Proof. Observe that for each person $p \in W \cup M$, their old (resp. new) matches under the reduction are their old (resp. new) matches before the reduction, “padded” by as many people as needed from the top of $\emptyset_1^p, \dots, \emptyset_{n_p}^p$ to fulfill their quota of n_p matches. Assume, for contradiction, that there exists a man $m \in M$ such that (before the reduction) $|N(m)| > |O(m)|$. Let $o = |O(m)|$ (before the reduction), then under the reduction, m is matched with $\emptyset_{n_p-o}^p$ under the original matching, but not under the new matching. Recall that $\emptyset_{n_p-o}^p$ prefers p the most, so she is not weakly better-off — a contradiction. We have thus shown that for each man $m \in M$, $|N(m)| \leq |O(m)|$, and since all women are weakly better-off, then for each woman $w \in W$, $|N(w)| \geq |O(w)|$, thus we have

$$\sum_{w \in W} N(w) \geq \sum_{w \in W} O(w) = \sum_{m \in M} O(m) \geq \sum_{m \in M} N(m) = \sum_{w \in W} N(w)$$

which yields that all members of this inequality are actually equal. Since for every woman $w \in W$, $|N(w)| \geq |O(w)|$, (res. for every man $m \in M$, $|N(m)| \leq |O(m)|$), and since summing over all women (resp. all men) we get an equality, it follows that the equality holds for each woman (resp. each man) separately. \square

It should be noted that in [5], Corollary 3.17 is proved using an entirely different approach for the special case of the monogamous scenario where the new match is stable, and in [6], it is proved also for the polygamous scenario where the new match is stable. Corollary 3.17 generalizes these results, as it does not require stability of the new match.

Corollary 3.18. *Under the conditions of Theorem 2.2, if $|L| = 1$ and the lying woman is better-off, then so is some innocent woman too.*

Proof. Since the liar is better-off, then she is newly matched with some man to whom she was not originally matched, and since the number of matches for this man remained the same, then he is no longer matched with some woman to whom he was originally matched, and hence the set of matches for that woman has changed, and thus, since she is weakly better-off, she is better-off. \square

4 Summary and Open Problems

We have shown throughout this paper that even in very general scenarios, a form of sisterhood exists in the Gale-Shapley matching algorithm, both in terms of not harming each other, and in terms of not helping any man.

If one were to define the concepts of “weakly worse-off” and “having gained only better matches” along the lines of definitions 3.3 and 3.4, then one would find that gaining only worse matches is stronger than (i.e. a special case of) being weakly worse-off and gaining only better matches is stronger case of being better-off. Thus, it follows that in the scenarios discussed in this paper, in a sense, the minimum possible damage to a man is greater than the minimum possible gain for a woman.

It is interesting to check whether, under various conditions, the overall damage to men is, in any sense, usually greater than the overall gain for women (and thus resulting in damage to the entire population as a whole, which contrasts the sisterhood which exists amongst the women).

The example given in this paper (and other examples given in the first author’s undergraduate thesis, upon which this paper is based) had to be crafted very delicately. It is interesting to check whether, when the orders of preference are determined by a random model, lies can be very beneficial (either for the lying women, and for their innocent colleagues), or whether the utility gained from such a lie is usually relatively small. Furthermore, in many of these examples, and in many examples specified in the literature, the preferences of every person differ greatly from those of each of their colleagues. In the real-world scenario of colleges and applicants, however, it is reasonable to expect the preferences of most applicants to be similar, and the same applies to the preferences of most colleges. There will always be differences, but it seems that they are likely to be local, such as a permutation on colleges which follow each other in the order of preference. It seems unlikely, for example, for a certain col-

lege to be highly rated by half of the applicants and poorly rated by the rest. (All this applies to the preferences of colleges as well, naturally.) It is interesting to try and build a probabilistic model along these lines, that matches the observed behaviors by medical interns and colleges throughout the years, and to check the questions raised above under such a model. In addition, since in the real world the knowledge of each player is not full, it can be interesting to check whether, given only knowledge of a statistical model for the preferences of the rest of the players (and possibly full knowledge of the preferences of just a few players), there exist strategies (for individual players, or for small sets of players) which are expected to be better for these players than telling the truth. It can also be interesting to check what happens if such strategies are simultaneously applied by more than one conspiring set of players.

Acknowledgments: We would like to thank Sergiu Hart for useful conversations and for providing us with references to the literature.

References

- [1] D. Gale and L. S. Shapley, College admissions and the stability of marriage. *American Mathematical Monthly* 69 (1962), 9–15.
- [2] L. E. Dubins and D. Freedman, Machiavelli and the Gale-Shapley algorithm. *American Mathematical Monthly* 88 (1981) 485–494.
- [3] D. Gale and M. Sotomayor, Ms. Machiavelli and the stable matching problem. *American Mathematical Monthly* 92 (1985) 261–268.
- [4] A.E. Roth, The evolution of the labor market for medical interns and residents: a case study in game theory, *Journal of Political Economy* 92 (1984) 991–1016.
- [5] D. Gale and M. Sotomayor, Some remarks on the stable matching problem. *Discrete Applied Mathematics* 11 (1985) 223–232.
- [6] A. E. Roth and M. A. O. Sotomayor, *Two-Sided Matchings* (Cambridge University Press, 1990).