STANLEY DEPTH AND COMPLETE *k*-PARTITE HYPERGRAPHS

MUHAMMAD ISHAQ AND MUHAMMAD IMRAN QURESHI

ABSTRACT. We give an upper bound for the Stanley depth of the edge ideal of a complete k-partite hypergraph and as an application we give an upper bound for the Stanley depth of a monomial ideal in a polynomial ring S. We also give a lower and an upper bound for the cyclic module S/I associated to the complete k-partite hypergraph.

Key words: Monomial ideals, Stanley decomposition, Stanley depth. 2000 Mathematics Subject Classification: Primary 13C15, Secondary 13P10, 13F20, 05E45, 05C65.

1. INTRODUCTION

Let K be a field, $S = K[x_1, \ldots, x_n]$ be the polynomial ring in n variables over a field K and M be a finitely generated \mathbb{Z}^n -graded S-module. Let $u \in M$ be a homogeneous element in M and Z a subset of the set of variables $Z \subset \{x_1, \ldots, x_n\}$. We denote by uK[Z] the K-subspace of M generated by all elements uv where v is a monomial in K[Z]. If uK[Z] is a free K[Z]-module, the \mathbb{Z}^n -graded K-space $uK[Z] \subset M$ is called a Stanley space of dimension |Z|. A Stanley decomposition of M is a presentation of the \mathbb{Z}^n -graded K-vector space M as a finite direct sum of Stanley spaces

$$\mathcal{D}: M = \bigoplus_{i=1}^{s} u_i K[Z_i].$$

The number

sdepth
$$\mathcal{D} = \min\{|Z_i| : i = 1, \dots, s\}$$

is called the Stanley depth of decomposition \mathcal{D} and the number

sdepth $M := \max\{ \operatorname{sdepth} \mathcal{D} : \mathcal{D} \text{ is a Stanley decomposition of } M \}$

is called the Stanley depth of M. This is a combinatorial invariant and does not depend on the characteristic of K. The following open conjecture is due to Stanley [13]:

 $\operatorname{depth} M \leq \operatorname{sdepth} M,$

for all finitely generated \mathbb{Z}^n -graded S-modules M.

Let $\mathbf{H} = (V, E)$ denote a hypergraph with vertex set V and hyperedge set E. A

The authors would like to express their gratitude to ASSMS of GC University Lahore for creating a very appropriate atmosphere for research work. This research is partially supported by HEC Pakistan.

hyperedge $e \in E$ is a subset of the vertices. That is, $e \subset V$ for each $e \in E$. A hypergraph is called complete k-partite if the vertices are partitioned into k disjoint subsets V_i , i = 1, ..., k and E consists of all hyperedges containing exactly one vertex from each of the k subsets.

In this paper we try to answer the following question asked by B. Nill and K. Vorwerk in [8].

Question 1.1 ([8]). Let I be the edge ideal of a complete k-partite hypergraph \mathbf{H}_d^k . Here, \mathbf{H}_d^k has kd vertices divided into k independent sets $V^{(i)}$ (for i = 1, ..., k) each with d vertices $v_1^{(i)}, \ldots, v_d^{(i)}$, and \mathbf{H}_d^k has d^k hyperedges consisting of exactly k vertices. Then I is squarefree monomial ideal in the polynomial ring $K[v_j^{(i)} : i \in \{1, ..., k\}, j \in \{1, ..., d\}]$:

$$I = (v_1^{(1)}, \dots, v_d^{(1)}) \cdots (v_1^{(k)}, \dots, v_d^{(k)}).$$

What is sdepth(S/I) in this case?

We consider this question even in more general frame. We consider the case where each vertex set $V^{(i)}$ is not necessarily of the same cardinality. Let I be the edge ideal of a complete k-partite hypergraph \mathbf{H}^k , where \mathbf{H}^k has n vertices divided into k independent sets $V^{(i)}$ (for i = 1, ..., k) each with d_i vertices $v_1^{(i)}, ..., v_{d_i}^{(i)}$, and \mathbf{H}^k has $d_1 d_2 \cdots d_k$ hyperedges consisting of exactly k vertices. To each vertex set $V^{(i)}$ we associate a set of variables $\{x_{i_1}, ..., x_{i_{d_i}}\}$ and set $S = K[(x_{i_j})]$. Now let $V^{(i)}$ and $V^{(j)}$ be two vertex sets, $\{x_{i_1}, ..., x_{i_{d_i}}\}$ and $\{x_{j_1}, ..., x_{j_{d_j}}\}$ be the sets of variables associated to $V^{(i)}$ and $V^{(j)}$ respectively. Since $V^{(i)}$ and $V^{(j)}$ are independent we have $\{x_{i_1}, ..., x_{i_{d_i}}\} \cap \{x_{j_1}, ..., x_{j_{d_j}}\} = \emptyset$. Then I is the squarefree monomial ideal in the polynomial ring S:

$$I = P_1 P_2 \cdots P_k = P_1 \cap P_2 \cap \cdots \cap P_k,$$

where $P_i = (x_{i_1}, \ldots, x_{i_{d_i}})$ and $\sum_{i=1}^k P_i = \mathfrak{m} = (x_1, \ldots, x_n)$. We give a tight upper bound to sdepth(I) see our Theorem 2.6 and as an application we give an upper

bound to sdepth(I) see our Theorem 2.0 and as an application we give an upper bound for the Stanley depth of a monomial ideal (see Theorem 2.14). In some cases we are able to give the exact values of sdepth(I) (see our corollaries 2.9, 2.10). We also give a tight lower bound to sdepth(S/I) see our Theorem 3.1. Our Proposition 3.8 gives an upper bound for Stanley depth of S/I but is big in general. We owe thanks to the Referee who inspired us Theorem 2.2.

2. Edge ideals of a complete k-partite hypergraph

We recall the method of Herzog et al. [3] for computing the Stanley depth of a monomial ideal I using posets. For $c \in \mathbb{N}^n$, let x^c denote the monomial $x_1^{c(1)}x_2^{c(2)}\cdots x_n^{c(n)}$ and let $I = (x^{a_1}, x^{a_2}, \ldots, x^{a_m})$ be a monomial ideal of S. Let $g \in \mathbb{N}^n$ be the componentwise maximum of a_i . Then characteristic poset of I with respect to g (see [3]), denoted by \mathcal{P}_I^g is in fact the set

$$\mathcal{P}_I^g = \{ c \in \mathbb{N}^n \mid c \le g, \text{ there is } i \text{ such that } c \ge a_i \},\$$

where \leq denotes the partial order in \mathbb{N}^n which is given by componentwise comparison. For every $a, b \in \mathcal{P}_I^g$ with $a \leq b$, define the interval [a, b] to be $\{c \in \mathcal{P}_I^g : a \leq c \leq b\}$. Let $\mathcal{P}: \mathcal{P}_I^g = \bigcup_{i=1}^r [c_i, d_i]$ be a partition of \mathcal{P}_I^g , define $\rho(d_i) := |\{j : d_i(j) = g(j)\}|$. Define the Stanley depth of a partition \mathcal{P} to be

$$\operatorname{sdepth}(\mathcal{P}) = \min_{[c_i, d_i] \in \mathcal{P}} \rho(d_i)$$

and the Stanley depth of the poset \mathcal{P}_{I}^{g} to be $\operatorname{sdepth}(\mathcal{P}_{I}^{g}) := \max_{\mathcal{P}} \operatorname{sdepth}(\mathcal{P})$, where maximum is taken over all the partitions \mathcal{P} of \mathcal{P}_{I}^{g} into intervals. Herzog et al., showed in [3] that $\operatorname{sdepth}(I) = \operatorname{sdepth}(\mathcal{P}_{I}^{g})$. Next lemma is a small extension of [2, Lemma 1.1], its proof is given for the sake of our completeness.

Lemma 2.1. Let r, m and a be positive integers with r < m and $v_1, \ldots, v_m \in K[x_2, \ldots, x_n]$ be some monomials of S. Let $I = (x_1^a v_1, \ldots, x_1^a v_r, v_{r+1}, \ldots, v_m)$ and $I' = (x_1^{a+1}v_1, \ldots, x_1^{a+1}v_r, v_{r+1}, \ldots, v_m)$ be monomial ideals of S. Then

$$sdepth(I) = sdepth(I').$$

Proof. Let $\mathcal{P} : \mathcal{P}_I^g = \bigcup_{i=1}^r [c_i, d_i]$ be a partition of \mathcal{P}_I^g such that $\mathrm{sdepth}(\mathcal{P}) = \mathrm{sdepth}(I)$, Define

$$c'_{i} = \begin{cases} c_{i}, & \text{if } c_{i}(1) < a \\ c_{i} + e_{1}, & \text{if } c_{i}(1) = a \end{cases} \quad and \quad d'_{i} = \begin{cases} d_{i}, & \text{if } d_{i}(1) < a - 1 \\ d_{i} + e_{1}, & \text{if } d_{i}(1) \ge a - 1 \end{cases}.$$

Let $g' := (g(1) + 1, g(2), \ldots, g(n)) \in \mathbb{N}^n$. Let $\mathcal{P}_{I'}^{g'}$ be the characteristic poset of I'with respect to g'. We claim that there exists a partition $\mathcal{P}' : \mathcal{P}_{I'}^{g'} = \bigcup_{i=1}^r [c'_i, d'_i]$ of $\mathcal{P}_{I'}^{g'}$. Note that $\rho(d_i) = \rho(d'_i)$ for all $i \in [r]$. Indeed, $d'_i(1) = g'(1) = a + 1$ if and only if $d_i(1) = g(1) = a$ and $d'_i(j) = g(j)$ if and only if $d_i(j) = g(j)$, for all $j \geq 2$. Therefore sdepth $(I) \leq$ sdepth(I'). Now we have to prove our claim. First we show that $\mathcal{P}_{I'}^{g'} = \bigcup_i^r [c'_i, d'_i]$. Let $\alpha \in \mathcal{P}_{I'}^{g'}$. If $\alpha(1) \leq a$ then $\alpha \in \mathcal{P}_I^g$ that is $\alpha \in [c_i, d_i]$ for some i, because if $\alpha \notin \mathcal{P}_I^g$ then we have $x_1^{a+1}v_k|x^{\alpha}$ for some $k \leq r$ and therefore $\alpha(1) = a + 1$, a contradiction.

If $\alpha(1) < a$, it follows that $c_i(1) < a$ and therefore $c'_i = c_i$. We get $c'_i = c_i \le d_i \le d'_i$, thus $\alpha \in [c'_i, d'_i]$. If $\alpha(1) = a$, it follows that $d_i(1) = a$ and $d'_i(1) = a + 1$. Suppose that $\alpha \notin [c'_i, d'_i]$. Then $\alpha - e_1 \in \mathcal{P}_I^g$ because $\alpha \in \mathcal{P}_{I'}^{g'}$, $\alpha(1) \neq a + 1$ and thus $x_1^{a+1}v_k$ does not divide x^{α} for all $k \le r$. It follows that $\alpha - e_1 \in [c_j, d_j]$ for some $j \ne i$. But $(\alpha - e_1)(1) = a - 1$ which implies $d_j(1) \ge a - 1$. It follows that $\alpha \le d_j + e_1 = d'_j$ and therefore $\alpha \in [c'_j, d'_j]$, because $c'_j = c_j$.

If $\alpha(1) = a + 1$, it follows that $\alpha - e_1 \in \mathcal{P}_I^g$ and therefore $\alpha - e_1 \in [c_i, d_i]$ for some i. Indeed, if $x_1^{a+1}v_k|x^{\alpha}$ for some k then $x_1^a v_k|x^{\alpha-e_1}$, else if $v_k|x^{\alpha}$ for some $k \ge r+1$ then $v_k|x^{\alpha-e_1}$. But $(\alpha - e_1)(1) = a$ and therefore $d_i(1) = a$. We get $\alpha \le d_i + e_1 = d'_i$ and thus $\alpha \in [c'_i, d'_i]$.

Now, we must prove that for any $i \neq j$, we have $[c'_i, d'_i] \cap [c'_j, d'_j] = \emptyset$. Assume that there exists some $\alpha \in [c'_i, d'_i] \cap [c'_j, d'_j]$. If $\alpha(1) < a$ then $\alpha \leq d_k$ and $\alpha \geq c'_k = c_k$ for k = i, j. Suppose that $\alpha(1) \geq a$. Then $d'_j(1) \geq a$. If $c_i(1) < a$ then $\alpha - e_1 \geq c_i = c'_i$. If $c_i(1) = a_1$ then $\alpha - e_1 \geq c'_i - e_1 = c_i$. Hence $\alpha - e_1 \in [c_i, d_i]$ and similarly, $\alpha - e_1 \in [c_j, d_j]$, which gives again a contradiction.

Now since $x_1 \notin I'$ and $I': x_1 = I$ then by [11, Proposition 2] we have sdepth $(I) \ge$ sdepth(I'). This completes the proof.

Theorem 2.2. Let $I = \bigcap_{i=1}^{k} Q_i \subset S$ be a monomial ideal such that each Q_i is irreducible and $G(\sqrt{Q_i}) \cap G(\sqrt{Q_j}) = \emptyset$ for all $i \neq j$, then $\operatorname{sdepth}(I) = \operatorname{sdepth}(\sqrt{I})$. Proof. We may suppose that $\operatorname{ht}(Q_i) \geq 2$ for all i because if for example $\operatorname{ht}(Q_k) = 1$, then we may remove Q_k since $I \cong \bigcap_{i=1}^{k-1}Q_i$. Let $\{v_1, v_2, \ldots, v_m\}$ be the set of minimal monomial generators of I. If I is squarefree then $I = \sqrt{I}$. In the case that I is not squarefree, we may assume that x_1^2 divides some v_i . We may further assume that $x_1|v_i$ for $i = 1, \ldots, r$ and x_1 does not divide v_i for i > r. Then, since $G(\sqrt{Q_i}) \cap G(\sqrt{Q_j}) = \emptyset$ for all $i \neq j$, it follows that there exists an integer $a \geq 1$ such that $x_1^{a+1}|v_i$ for $i = 1, \ldots, r$. Now we apply Lemma 2.1 and conclude that sdepth $(I) = \operatorname{sdepth}(v_1/x_1, \ldots, v_r/x_1, v_{r+1}, \ldots, v_m)$. Thus an obvious induction argument completes the proof.

Remark 2.3. In the setting of Theorem 2.2, if Q_i are not irreducible for all i then the result is false. For example if n = 4, $I = (x_1^2, x_1x_2, x_2^2) \cap (x_3^2, x_3x_4, x_4^2)$ and \mathcal{P} is a partition of \mathcal{P}_I^g , g = (2, 2, 2, 2, 2) then we must have 9 intervals [a, b] in \mathcal{P} starting with the generators a of I but only 8 monomials b are in \mathcal{P}_I^g with $\rho(b) = 3$, the biggest one $x_1^2 x_2^2 x_3^2 x_4^2$ cannot be taken. Thus sdepth I < 3. But clearly sdepth(\sqrt{I}) = 3.

Lemma 2.4. Let I be a squarefree monomial ideal of S generated by monomials of degree d. Let A be the number of monomials of degree d and B be the number of monomials of degree d + 1 in I. Then

$$d \leq \operatorname{sdepth}(I) \leq d + \lfloor \frac{B}{A} \rfloor,$$

where $|a|, a \in \mathbb{Q}$, denotes the largest integer which is not greater than a.

Proof. We follow the proof of [4, Theorem 2.8](see also [6] and [7]). Let k := sdepth(I). The poset \mathcal{P}_I^g has a partition $\mathcal{P} : \mathcal{P}_I^g = \bigcup_{i=1}^s [c_i, d_i]$, satisfying sdepth(\mathcal{P}) = k. For each interval $[c_i, d_i]$ in \mathcal{P} with $|c_i| = d$ we have $|d_i| \ge k$. Also there are $|d_i| - |c_i|$ subsets of cardinality d+1 in this interval. Since these intervals are disjoint, counting the number of subsets of cardinality d and d+1 we have $(k-d)A \le B$ that is $k \le d + \frac{B}{A}$. Hence sdepth(I) $\le d + \lfloor \frac{B}{A} \rfloor$. The other inequality follows by [7, Lemma 2.1].

Corollary 2.5. Let I be a squarefree monomial ideal of S generated by monomials of degree d. If $\binom{n}{d+1} < |G(I)|$ then sdepth(I) = d.

Theorem 2.6. Let $I = \bigcap_{i=1}^{k} P_i$ be a monomial ideal in S where each P_i is a monomial prime ideal and $\sum_{i=1}^{k} P_i = \mathfrak{m}$. Suppose that $G(P_i) \cap G(P_j) = \emptyset$ for all $i \neq j$. Then

$$\operatorname{sdepth}(I) \le \frac{n+k}{2}$$

Proof. Note that I is generated by monomials of degree k. Let $|G(P_i)| = d_i$ then clearly the number of monomials of degree k in I is $d_1d_2\cdots d_k$. Now let $v \in I$ be a monomial of degree k+1, then v is divisible by at least one variable from $G(P_i)$ for all $1 \leq i \leq k$ and one of these sets contains two variables from $\sup(v) := \{x_j : x_j | v\}$. We may suppose that all $d_i \geq 2$ because if for example $d_k = 1$, then we may remove P_k because $I \cong \bigcap_{i=1}^{k-1} P_i$. Fix first $G(P_1)$ such that $\sup(v)$ contains two variables from $G(P_1)$ then the number of such monomials is $\binom{d_1}{2}d_2d_3\cdots d_k$. Now let $\sup(v)$ contains two variables from $G(P_2)$ then the number of such monomials is $\binom{d_2}{2}d_1d_3\cdots d_k$. Continuing in the same way for all $G(P_i)$, $3 \leq i \leq k$. We get the total number of monomials of degree k + 1 is

$$\sum_{i=1}^{k} \binom{d_i}{2} d_1 \cdots d_{i-1} d_{i+1} \cdots d_k = \frac{d_1 d_2 \cdots d_k}{2} (\sum_{i=1}^{k} d_i - k) = \frac{d_1 d_2 \cdots d_k}{2} (n-k).$$

Now by Lemma 2.4 we have

$$sdepth(I) \le k + \frac{1}{d_1 d_2 \cdots d_k} \cdot \frac{d_1 d_2 \cdots d_k}{2} (n-k) = k + \frac{n-k}{2} = \frac{n+k}{2}.$$

Let $I = \bigcap_{i=1}^{k} Q_i$ be a monomial ideal such that each Q_i is irreducible and $G(\sqrt{Q_i}) \cap G(\sqrt{Q_j}) = \emptyset$ for all $i \neq j$, $\operatorname{ht}(Q_i) = d_i$ and $\sum_{i=1}^{k} \sqrt{Q_i} = \mathfrak{m}$. We define a set $A := \{Q_i : \operatorname{ht}(Q_i) \text{ is odd }\}.$

Corollary 2.7. Let $I = \bigcap_{i=1}^{k} Q_i$ be a monomial ideal such that each Q_i is irreducible and $G(\sqrt{Q_i}) \cap G(\sqrt{Q_j}) = \emptyset$ for all $i \neq j$, $\operatorname{ht}(Q_i) = d_i$ and $\sum_{i=1}^{k} \sqrt{Q_i} = \mathfrak{m}$. Then $\frac{n+|A|}{2} \leq \operatorname{sdepth}(I) \leq \lfloor \frac{n+k}{2} \rfloor$.

Proof. By [9, Lemma 1.2] we have $\lceil \frac{d_1}{2} \rceil + \lceil \frac{d_2}{2} \rceil + \dots + \lceil \frac{d_k}{2} \rceil \leq \text{sdepth}(I)$. Now by Theorem 2.2 we have $\text{sdepth}(I) = \text{sdepth}(\sqrt{I})$ and by Theorem 2.6 the required result follows.

Remark 2.8. With the hypothesis from the above Corollary, n is odd if and only if |A| is odd, thus n + |A| is always even.

Corollary 2.9. Let $I = \bigcap_{i=1}^{k} Q_i$ be a monomial ideal such that each Q_i is irreducible and $G(\sqrt{Q_i}) \cap G(\sqrt{Q_j}) = \emptyset$ for all $i \neq j$ and $\sum_{i=1}^{k} \sqrt{Q_i} = \mathfrak{m}$. Suppose that |A| = k, then

$$\operatorname{sdepth}(I) = \frac{n+k}{2}.$$

Proof. The proof follows by Corollary 2.7 and Remark 2.8.

Corollary 2.10. Let $I = \bigcap_{i=1}^{k} Q_i$ be a monomial ideal such that each Q_i is irreducible and $G(\sqrt{Q_i}) \cap G(\sqrt{Q_j}) = \emptyset$ for all $i \neq j$ and $\sum_{i=1}^{k} \sqrt{Q_i} = \mathfrak{m}$. Suppose that k is odd and |A| = k - 1 then,

$$sdepth(I) = \frac{n+k-1}{2}$$

Proof. Since k is odd then k-1 is even and since |A| = k-1 is even thus n is even and so n+k-1 is even. The result follows by Corollary 2.7.

Remark 2.11. The bounds found above for sdepth(I) are certainly sufficient to show Stanley's Conjecture in this case because sdepth(I) $\geq k = \text{depth}(I)$, as it is done in [9, Theorem 1.4] and [5, Theorem 3.2] in a more general frame. It is a difficult combinatorial question to find the precise value of sdepth(I). It is hard to find sdepth(I) for example in the case k = 2 and n even(see [4, Corollary 2.10]). We believe that in this case sdepth(I) = (n/2) + 1 but we can prove it only for small n. The trouble is that we must construct a partition \mathcal{P} on \mathcal{P}_I^g such that sdepth(\mathcal{P}) = (n/2) + 1. For example if n = 6 and $I = (x_1, x_2) \cap (x_3, \ldots, x_6)$ then we have the partition $\mathcal{P}_I^g = [13, 1345] \cup [14, 1456] \cup [15, 1256] \cup [16, 1236] \cup [23, 1234] \cup$ $[24, 1245] \cup [25, 2356] \cup [26, 2346] \cup (\cup_C [C, C])$, where $C \subset [6]$ with |C| = 5, 6, or $C \in \{1235, 1246, 1346, 1356, 2345, 2456\}$. Thus sdepth(I) = 4 = (n/2) + 1.

Corollary 2.12. Let I be the edge ideal of a complete k-partite hypergraph \mathbf{H}_d^k . Then

$$\operatorname{sdepth}(I) = \frac{n+k}{2}, \quad \text{if } d \text{ is } odd;$$

 $\frac{n}{2} \leq \operatorname{sdepth}(I) \leq \frac{n+k}{2}, \quad \text{if } d \text{ is } even.$

In the next theorem we give an upper bound for the Stanley depth of any monomial ideal of S. We will need the following lemma.

Lemma 2.13. ([5, Lemma 4.4]) Let $I \subset \hat{S} = S[x_{n+1}]$ be a monomial ideal, x_{n+1} being a new variable. If $I \cap S \neq (0)$, then sdepth_S $(I \cap S) \geq \text{sdepth}_{\hat{S}} I' - 1$.

Theorem 2.14. Let I be a monomial ideal and let $Min(S/I) = \{P_1, \ldots, P_s\}$ with $\sum_{i=1}^{s} P_i = \mathfrak{m}$. Let $d_i := |G(P_i) \setminus G(\sum_{i \neq j}^{s} P_j)|$, and $r := |\{d_i : d_i \neq 0\}|$. Suppose that $r \ge 1$. Then

$$sdepth(I) \le (2n + r - \sum_{i=1}^{s} d_i)/2.$$

Proof. We may assume that P'_i 's be prime ideals such that $G(P'_i) = G(P_i) \setminus G(\sum_{i \neq j}^s P_j)$ and $\operatorname{ht}(P'_i) \neq 0$ for all $1 \leq i \leq r \leq s$. We have $G(P'_i) \cap G(P'_j) = \emptyset$ for all $i \neq j$. Let $A := \bigcup_{i=1}^{r} G(P'_i)$, S' = K[A] and $B := \{x_1, \ldots, x_n\} \setminus \bigcup_{i=1}^{r} G(P'_i)$, then $|B| = n - \sum_{i=1}^{r} d_i$. Now applying Lemma 2.13 by recurrence on the set B we have

$$\operatorname{sdepth}_{S}(\sqrt{I}) \leq \operatorname{sdepth}_{S'}\left(\bigcap_{i=1}^{r} P'_{i}\right) + |B|.$$

By Theorem 2.6 we have sdepth_{S'} $\left(\bigcap_{i=1}^{r} P'_{i}\right) \leq \frac{|A|+r}{2}$. But |A| = n - |B| thus we have

$$sdepth(\sqrt{I}) \le \frac{n - |B| + r}{2} + |B| = \frac{2n + r - \sum_{i=1}^{s} d_i}{2}$$

By [4, Corollary 2.2] sdepth(I) \leq sdepth(\sqrt{I}) and we get sdepth(I) \leq $(2n + r - \sum_{i=1}^{s} d_i)/2$.

Example 2.15. Let $I = (x_1 \dots, x_9) \cap (x_9, \dots, x_{18}) \cap (x_{18}, \dots, x_{27}) \cap (x_{27}, \dots, x_{36}) \subset K[x_1, \dots, x_{36}]$. We have $d_1 = 8$, $d_2 = 8$, $d_3 = 8$, $d_4 = 9$ and s = 4, then by Theorem 2.14 we have sdepth $(I) \leq 23$.

Remark 2.16. In the above example we have that $sdepth(I) \leq 23$ but by [5, Theorem 2.1] we have only $sdepth(I) \leq 31$.

3. Cyclic modules associated to a complete k-partite hypergraph

Theorem 3.1. Let $S = K[x_1, \ldots, x_n]$ be a polynomial ring and Q_1, Q_2, \ldots, Q_k monomial irreducible ideals of S such that $G(\sqrt{Q_i}) \cap G(\sqrt{Q_j}) = \emptyset$ for all $i \neq j$. Let $r_i := \operatorname{ht}(Q_i), \sum_{i=1}^k r_i = n$. If $I = Q_1 \cap Q_2 \cap \ldots \cap Q_k$, then

$$\operatorname{sdepth}(S/I) \geq \min\left\{n - r_1, \min_{2 \leq i \leq k} \left\{ \lceil \frac{r_1}{2} \rceil + \ldots + \lceil \frac{r_{i-1}}{2} \rceil + r_{i+1} + \ldots + r_k \right\} \right\},$$

where $[a], a \in \mathbb{Q}$, denotes the smallest integer which is not less than a.

Proof. As a K-linear space S/I is isomorphic to the direct sum of some multigraded modules as,

$$S/I \cong S/Q_1 \oplus (Q_1/Q_1 \cap Q_2) \oplus (Q_1 \cap Q_2/Q_1 \cap Q_2 \cap Q_3) \oplus \dots$$
$$\oplus (Q_1 \cap \dots \cap Q_{k-1}/Q_1 \cap \dots \cap Q_k),$$

and we have $sdepth(S/I) \ge$

min{sdepth S/Q_1 , sdepth $Q_1/(Q_1 \cap Q_2), \ldots$, sdepth $(Q_1 \cap \ldots \cap Q_{k-1})/(Q_1 \cap \ldots \cap Q_k)$ }. By [10, Lemma 1.1] sdepth $(S/Q_1) = n - r_1$. Now since

$$(Q_1 \cap \ldots \cap Q_{i-1})/(Q_1 \cap \ldots \cap Q_i) \cong (Q_1 \cap \ldots \cap Q_{i-1} + Q_i)/Q_i$$

and

$$(Q_1 \cap \ldots \cap Q_{i-1} + Q_i)/Q_i \cong Q_1 \cap \ldots \cap Q_{i-1} \cap K[x_j \mid x_j \notin G(\sqrt{Q_i})]$$

Now by using [9, Lemma 1.2] and [3, Lemma 3.6] we have

sdepth
$$((Q_1 \cap \ldots \cap Q_{i-1})/(Q_1 \cap \ldots \cap Q_i)) \ge \lceil \frac{r_1}{2} \rceil + \ldots + \lceil \frac{r_{i-1}}{2} \rceil + r_{i+1} + \ldots + r_k.$$

Thus

$$\min\{\operatorname{sdepth} S/Q_1, \operatorname{sdepth} Q_1/(Q_1 \cap Q_2), \dots, \operatorname{sdepth}(Q_1 \cap \dots \cap Q_{k-1})/(Q_1 \cap \dots \cap Q_k)\} \\ \geq \min\left\{n - r_1, \min_{2 \le i \le k} \left\{ \lceil \frac{r_1}{2} \rceil + \dots + \lceil \frac{r_{i-1}}{2} \rceil + r_{i+1} + \dots + r_k \right\} \right\}.$$

Remark 3.2. The theorem says in particular that Stanley's conjecture holds for S/I in the above settings, because $\operatorname{sdepth}(I) \ge k-1 = \operatorname{depth}(S/I)$ as it was noticed in [5, Theorem 3.2]. The found lower bound depends on the numbering of (r_i) as shows the following example.

Example 3.3. Let $I = P_1 \cap P_2 \cap P_3 \cap P_4 \subset K[x_1, \ldots, x_{14}]$ where $G(P_i) \cap G(P_j) \neq \emptyset$ for all $i \neq j$, $P_1 + P_2 + P_3 + P_4 = (x_1, \ldots, x_{14})$ and $\operatorname{ht}(P_1) = 5$, $\operatorname{ht}(P_2) = 4$, $\operatorname{ht}(P_3) = 3$, $\operatorname{ht}(P_4) = 2$. Then by above theorem we have $\operatorname{sdepth}(S/I) \geq \min\{9, \min\{8, 7, 7\}\} =$ 7. Now let us reorder these primes or equivalently consider that $\operatorname{ht}(P_1) = 4$, $\operatorname{ht}(P_2) =$ 5, $\operatorname{ht}(P_3) = 2$, $\operatorname{ht}(P_4) = 3$. Then again by the above theorem we have $\operatorname{sdepth}(S/I) \geq \min\{10, \min\{7, 8, 6\}\} = 6$.

Corollary 3.4. Let $r_1 \ge r_2 = \ldots = r_k$, then

$$\operatorname{sdepth}(S/I) \ge \lceil \frac{r_1}{2} \rceil + \ldots + \lceil \frac{r_{k-1}}{2} \rceil.$$

Corollary 3.5. Let I be the edge ideal of a complete k-partite hypergraph \mathbf{H}_{d}^{k} . Then

$$\operatorname{sdepth}(S/I) \ge \lceil \frac{d}{2} \rceil (k-1).$$

Remark 3.6. In Corollary 3.5 if we take k = 2 then we have $\operatorname{sdepth}(S/I) \ge \lceil \frac{d}{2} \rceil$ but by [10] and [8] we have $\operatorname{sdepth}(S/I) = \lceil \frac{d}{2} \rceil$. This shows that the bound is equal to the actual value in this case. If d is odd in Corollary 3.5 then $\operatorname{sdepth}(S/I) \ge \frac{n+k}{2} - \frac{d+1}{2}$. Now assume that $\operatorname{sdepth}(I) \ge \operatorname{sdepth}(S/I) + 1$, as A. Rauf asks in [12], (for k = 2this inequality is true [10]). Then by Theorem 2.6 we have $\operatorname{sdepth}(S/I) \le \frac{n+k}{2} - 1$. If this is the case then our lower bound by Corollary 3.5 could be a reasonable one, as it is clear that for d = 3 we have $\frac{n+k}{2} - 2 \le \operatorname{sdepth}(S/I) \le \frac{n+k}{2} - 1$.

Next we give an upper bound for the Stanley depth of S/I with $I = P_1 \cap P_2 \cap \cdots \cap P_k$, (P_i) being monomial prime ideals such that $G(P_i) \cap G(P_j) = \emptyset$. Let $\operatorname{ht}(P_i) = r_i$. By [3, Lemma 3.6] it is enough to consider that $P_1 + P_2 + \cdots + P_k = \mathfrak{m}$. Let $\mathcal{D} : S/I = \bigoplus_{i=1}^p u_i K[Z_i]$ be a Stanley decomposition. Then Z_i cannot have in the same time variables from all $G(P_i)$, otherwise $u_i K[Z_i]$ will not be a free $K[Z_i]$ -module. Suppose $u_1 = 1$ and $Z_1 \subset \{x_{r_1+1}, \ldots, x_n\}$.

Lemma 3.7. Let $r_2 \ge r_3 \ge \cdots \ge r_k \ge 1$. Then sdepth(\mathcal{D}) $< \lceil \frac{r_1}{r_1} \rceil + r_2 + r_3 + r_3$

$$\operatorname{depth}(\mathcal{D}) \leq \lceil \frac{r_1}{2} \rceil + r_2 + r_3 + \dots + r_{k-1}$$

Proof. We can assume that $P_1 = (x_1, \ldots, x_{r_1})$ and

$$\psi: P_1 \cap K[x_1, \dots, x_{r_1}] \hookrightarrow S/I$$

be the inclusion given by

 $K[x_1,\ldots,x_{r_1}] \hookrightarrow S/I.$

Then $P_1 \cap K[x_1, \ldots, x_{r_1}] = \bigoplus_i \psi^{-1}(u_i K[Z_i])$. If $\psi^{-1}(u_i K[Z_i]) \neq 0$ implies there exists $u_i f \in u_i K[Z_i]$ with $u_i f \in P_1 \cap K[x_1, \ldots, x_{r_1}]$, but then $u_i \neq 1$ and so $u_i \in P_1 \cap K[x_1, \ldots, x_{r_1}]$. Let $Z'_i = Z_i \cap \{x_1, \ldots, x_{r_1}\}$. Then $\psi^{-1}(u_i K[Z_i]) = u_i K[Z'_i]$ and we get a Stanley decomposition of $J := P_1 \cap K[x_1, \ldots, x_{r_1}]$. Since sdepth $(J) = \lceil \frac{r_1}{2} \rceil$ by [1] it follows that

$$|Z'_i| \le \operatorname{sdepth}(J) = \lceil \frac{r_1}{2} \rceil.$$

But Z_i cannot have variables from all $G(P_j)$ and we get $Z_i \subset \bigcup_{1=e\neq j}^k G(P_e)$ for some $j \neq 1$. Therefore

$$Z_i \subset \{Z'_i \cup G(P_2) \cup G(P_3) \cup \dots \cup G(P_k)\} \setminus G(P_j) \text{ for some } j \neq 1.$$

Thus we have

sdepth(
$$\mathcal{D}$$
) $\leq \lceil \frac{r_1}{2} \rceil + r_2 + r_3 + \dots + r_{j-1} + r_{j+1} + \dots + r_k \leq \lceil \frac{r_1}{2} \rceil + r_2 + r_3 + \dots + r_{k-1}.$

Proposition 3.8. Suppose that $r_1 \ge r_2 \ge \cdots \ge r_k$, $k \ge 3$. Then sdepth $(S/I) \le \lceil \frac{r_{k-1}}{2} \rceil + r_1 + r_2 + \cdots + r_{k-2}$.

Proof. Let \mathcal{D} be a Stanley decomposition of S/I, $S/I = \bigoplus_{i=1}^{p} u_i K[Z_i]$ such that sdepth(\mathcal{D}) = sdepth(S/I). Then there exists $1 \le m \le k$ such that $Z_1 \subset \bigcup_{1=e \ne m}^{k} G(P_e)$. By the above lemma sdepth(\mathcal{D}) $\le \sum_{1=i \ne m}^{k-1} r_i + \lceil \frac{r_m}{2} \rceil$, which is enough. \Box

Corollary 3.9. Let I be the edge ideal of a complete k-partite hypergraph \mathbf{H}_{d}^{k} . Then

$$(k-1)\lceil \frac{d}{2}\rceil \le \operatorname{sdepth}(S/I) \le (k-2)d + \lceil \frac{d}{2}\rceil.$$

References

- C. Biro, D.M. Howard, M.T. Keller, W.T. Trotter, S.J. Young, *Interval partitions and Stanley depth*, J. Combin. Theory Ser. A (2009), in press, doi:10.1016/j.jcta.2009.07.008.
- M. Cimpoeaş, Stanley depth of complete intersection monomial ideals, Bull. Math. Soc. Sci. Math. Roumanie, 51(99) no 3, 2008, 205211.
- [3] J. Herzog, M. Vladoiu, X. Zheng, How to compute the Stanley depth of a monomial ideal, J. Algebra, 322(9)(2009), 3151-3169.
- [4] M. Ishaq, Upper bounds for the Stanley depth, to appear in Comm. Algebra, arXiv:1003.3471v1.
- [5] M. Ishaq, Values and bounds of the Stanley depth, to appear in Carpathian J. Math., arXiv:1010.4692v1.

- [6] M. Ishaq, M.I. Qureshi, *Stanley depth of edge ideals*, to appear in Studia Sci. Math. Hungarica, arXiv:1104.1018v1.
- [7] M.T. Keller, Y. Shen, N. Streib, S.J. Young, On the Stanley Depth of Squarefree Veronese Ideals arXiv:0910.4645v1.
- [8] B. Nill, K. Vorwerk, Stanley's conjecture, cover depth and extremal simplicial complexes, LE MATEMATICHE LXIII(2008)-Fasc. II, pp.213-228.
- [9] A. Popescu, Special Stanley decomposition, Bull. Math. Soc. Sc. Math. Roumanie, 52(101), no 4 (2010), 363-372.
- [10] D. Popescu, M.I. Qureshi, Computing the Stanley depth, J. Algebra 323 (10) (2010), 2943-2959.
- [11] D. Popescu, Bounds of Stanley depth, Analele Univ. Ovidius Constanta, 19(2011), 187-194.
- [12] A. Rauf, Depth and Stanley depth of multigraded modules, Comm. Algebra, 38 (2010), 773-784.
- [13] R.P. Stanley, Linear Diophantine equations and local cohomology, Invent. Math. 68 (1982) 175-193.

MUHAMMAD ISHAQ, ABDUS SALAM SCHOOL OF MATHEMATICAL SCIENCES, GC UNIVERSITY, LAHORE, 68-B NEW MUSLIM TOWN LAHORE, PAKISTAN.

E-mail address: ishaq_maths@yahoo.com

Muhammad Imran Qureshi, Comsats Institute of Information Technology, Vehari campus, Multan Road Vehari, Pakistan.

E-mail address: imranqureshi18@gmail.com