

Extremal sizes of subspace partitions*

O. Heden, J. Lehmann, E. Năstase, and P. Sissokho

September 8, 2017

Abstract

A *subspace partition* Π of $V = V(n, q)$ is a collection of subspaces of V such that each 1-dimensional subspace of V is in exactly one subspace of Π . The *size* of Π is the number of its subspaces. Let $\sigma_q(n, t)$ denote the *minimum size* of a subspace partition of V in which the largest subspace has dimension t , and let $\rho_q(n, t)$ denote the *maximum size* of a subspace partition of V in which the smallest subspace has dimension t . In this paper, we determine the values of $\sigma_q(n, t)$ and $\rho_q(n, t)$ for all positive integers n and t . Furthermore, we prove that if $n \geq 2t$, then the minimum size of a maximal partial t -spread in $V(n + t - 1, q)$ is $\sigma_q(n, t)$.

Keywords. Subspace partition; Vector space partitions; Partial t -spreads.

1 Introduction

Let $V = V(n, q)$ denote a vector space of dimension n over a finite field with q elements. A *subspace partition* Π of V is a collection of subspaces of V such that each 1-dimensional subspace of V is in exactly one subspace of Π . A subspace partition Π is also called a *vector space partition* (or simply a *partition*) of V . There is a rich literature about vector space partitions, see e.g. [1, 3, 5, 14, 23] and the references therein.

The *size* of Π is the number of its subspaces. Let $\sigma_q(n, t)$ denote the *minimum size* of a subspace partition of V in which the largest subspace has dimension t , and let $\rho_q(n, t)$ denote the *maximum size* of a subspace partition of V in which the smallest subspace has dimension t . The purpose of this study is to find these numbers. Since $\sigma_q(n, n) = \rho_q(n, n) = 1$, and $\sigma_q(n, 1) = \rho_q(n, 1) = (q^n - 1)/(q - 1)$, we will focus on the case $1 < t < n$. Moreover, if t divides n , then $\sigma_q(n, t) = \rho_q(n, t)$ is the size of a t -spread in V , i.e., a subspace partition of V in which all the subspaces have dimension t .

We will prove the following theorem:

*This research project was supported by grant KAW 2005.0098 from the Knut and Alice Wallenberg Foundation.

Theorem 1 *Let n, k, t , and r be integers such that $0 \leq r < t$, $k \geq 2$, and $n = kt + r$. Then*

$$\rho_q(n, t) = q^{t+r} \sum_{i=0}^{k-2} q^{it} + 1,$$

and if furthermore $1 \leq r < t$, then

$$\sigma_q(n, t) = q^{t+r} \sum_{i=0}^{k-2} q^{it} + q^{\lceil \frac{t+r}{2} \rceil} + 1.$$

This theorem improves a result of Beutelspacher [2] who in 1980 proved that

$$\sigma_q(n, t) \geq q^{\lceil \frac{n}{2} \rceil} + 1.$$

We must also remark that the last two authors of this paper recently found the value of $\sigma_q(2t + 1, t)$, see [21]. They used some equations for subspace partitions derived by the first two authors in [17]. Furthermore, our derivation of the value of $\sigma_q(n, t)$ uses arguments quite similar to those used in [21].

After some preliminary results in Section 2, we will prove our theorem in Section 3 and Section 4. Finally, in Section 5, we combine our result on $\sigma_q(n, t)$ with a construction of P. Govaerts [13] to show that the minimum size of a maximal partial t -spread in $V(n+t-1, q)$ is $\sigma_q(n, t)$ for any integer $n \geq 2t$.

2 Preliminary Results

Let Π be a subspace partition of $V = V(n, q)$, $n \geq 2$, with m_i subspaces of dimension i , $1 \leq i \leq n - 1$. Let H be any hyperplane, i.e., any $(n - 1)$ -dimensional subspace of V , and let $b_i \leq m_i$ be the number of subspaces of Π that are contained in H . We say that (m_{n-1}, \dots, m_1) is the *type* of Π and $b = (b_{n-1}, \dots, b_1)$ is the *type of the hyperplane* H (with respect to Π). Let s_b denote the number of hyperplanes in V of type b and define the set

$$B = \{b : s_b > 0\}.$$

For $1 \leq i \leq n$, let

$$\theta_i = \frac{q^i - 1}{q - 1}$$

denote the number of 1-dimensional subspaces in an i -space; then

$$h_q(n, i) = \max\{0, \theta_{n-i}\}$$

denotes the number of hyperplanes containing a given i -dimensional subspace. The following two lemmas were derived in [17].

Lemma 1 Let Π be a subspace partition of $V = V(n, q)$ of type (m_{n-1}, \dots, m_1) and let $b = (b_{n-1}, \dots, b_1)$ be the type of the hyperplane H with respect to Π . Let s_b denote the number of hyperplanes in V with type b . Assume furthermore that Π contains a subspace of dimension d and a subspace of dimension d' , with $1 \leq d, d' \leq n - 2$. Then

- (i) $\sum_{b \in B} s_b = \frac{q^n - 1}{q - 1} = h_q(n, 0)$,
- (ii) $\sum_{b \in B} b_d s_b = m_d h_q(n, d)$,
- (iii) $\sum_{b \in B} \binom{b_d}{2} s_b = \binom{m_d}{2} h_q(n, 2d)$,
- (iv) $\sum_{b \in B} b_d b_{d'} s_b = m_d m_{d'} h_q(n, d + d')$.

Lemma 2 Let Π be a subspace partition of $V = V(n, q)$ and let (b_{n-1}, \dots, b_1) be the type of the hyperplane H with respect to Π . Then the number of subspaces in Π is

$$|\Pi| = 1 + \sum_{i=1}^{n-1} b_i q^i.$$

We will also use the following lemma due to Herzog and Schönheim [18] and independently Beutelspacher [1] and Bu [5].

Lemma 3 Let n and d be integers such that $1 \leq d \leq n/2$. Then $V = V(n, q)$ admits a partition with one subspace of dimension $n - d$ and q^{n-d} subspaces of dimension d .

For $n = kt + r$, $0 \leq r < t$, and $k \geq 2$, let

$$\ell = q^r \sum_{i=0}^{k-2} q^{it}. \quad (1)$$

The following proposition is an immediate consequence of Lemma 3.

Proposition 1 Let n, k, t , and r be integers such that $0 \leq r < t$, $k \geq 2$, and $n = kt + r$. Then $V = V(n, q)$ admits a partition Π_m of size

$$|\Pi_m| = \ell \cdot q^t + 1,$$

consisting of ℓq^t subspaces of dimension t and one subspace of dimension $t + r$. If furthermore, $1 \leq r < t$, then V admits a partition Π_M of size

$$|\Pi_M| = \ell \cdot q^t + q^{\lceil \frac{t+r}{2} \rceil} + 1,$$

consisting of ℓq^t subspaces of dimension t , $q^{\lceil (t+r)/2 \rceil}$ subspaces of dimension $\lfloor (t+r)/2 \rfloor$ and one subspace of dimension $\lceil (t+r)/2 \rceil$.

We close this section by giving three relations that will be frequently used. They follow easily from the definitions of ℓ and the function θ_i ; the third is an immediate consequence of the first two:

$$\theta_{n-t} - \theta_r = \ell \theta_t, \quad (2)$$

$$\theta_{a+b} - \theta_b = q^b \theta_a, \quad (3)$$

$$\theta_n - \ell q^t \theta_t = \theta_{t+r}. \quad (4)$$

3 The minimum size

In this section we will find $\sigma_q(n, t)$, as indicated in Theorem 1. We will need the following lemma, which may be of independent interest.

Lemma 4 *Let n, k, t , and r be integers such that $k \geq 2, 1 \leq r < t$, and $n = kt + r$. Let Π be a subspace partition of $V = V(n, q)$ with no subspace of dimension higher than t . Assume furthermore that Π contains a subspace of dimension t and a subspace of dimension d , with $0 \leq d < t$. Then*

$$|\Pi| \geq q^{t+r} \sum_{i=0}^{k-2} q^{it} + q^d + 1.$$

Proof. Let Π be a subspace partition of V containing subspaces of dimension t and d with $t > d$. Since there exist subspaces of dimensions t and d in Π , we have $m_t > 0$ and $m_d > 0$. So it follows from Lemma 1(iv) that

$$\sum_{b \in B} b_t b_d s_b = m_t m_d \theta_{n-t-d} \neq 0. \quad (5)$$

Additionally,

$$\sum_{b \in B} b_t b_d s_b = \sum_{\substack{b \in B \\ 0 \leq b_t \leq \ell-1}} b_t b_d s_b + \sum_{\substack{b \in B \\ b_t \geq \ell}} b_t b_d s_b.$$

If

$$\sum_{b \in B, b_t \geq \ell} b_t b_d s_b \neq 0,$$

then there exists $b \in B$ such that $b_t \geq \ell, b_d \geq 1$, and $s_b \geq 1$. In this case, Lemma 2 yields

$$|\Pi| = \sum_{i=1}^{n-1} b_i q^i + 1 \geq b_t q^t + b_d q^d + 1 \geq \ell q^t + q^d + 1,$$

and the lemma follows. So we may assume that $\sum_{b \in B, b_t \geq \ell} b_t b_d s_b = 0$. This assumption, combined with (5) and Lemma 1(iv), yields

$$\begin{aligned} (\ell - 1)m_d \theta_{n-d} &= \sum_{b \in B} (\ell - 1) \cdot b_d s_b \\ &= \sum_{\substack{b \in B \\ 0 \leq b_t \leq \ell-1}} (\ell - 1) \cdot b_d s_b + \sum_{\substack{b \in B \\ b_t \geq \ell}} (\ell - 1) \cdot b_d s_b \\ &\geq \sum_{\substack{b \in B \\ 0 \leq b_t \leq \ell-1}} b_t \cdot b_d s_b + 0 \\ &= \sum_{\substack{b \in B \\ 0 \leq b_t \leq \ell-1}} b_t \cdot b_d s_b + \sum_{\substack{b \in B \\ b_t \geq \ell}} b_t \cdot b_d s_b \end{aligned}$$

$$\begin{aligned}
&= \sum_{b \in B} b_t b_d s_b \\
&= m_t m_d \theta_{n-t-d}
\end{aligned} \tag{6}$$

Since $m_d > 0$, dividing both sides of (6) by m_d yields

$$m_t \leq \frac{(\ell - 1) \theta_{n-d}}{\theta_{n-t-d}}.$$

We now show that this implies that

$$m_t \leq (\ell - 1)q^t + q^d. \tag{7}$$

From (3) we obtain that $\theta_{n-d} = \theta_t + q^t \theta_{n-d-t}$, and hence it remains to prove that

$$\frac{(\ell - 1)\theta_t}{\theta_{n-d-t}} \leq q^d.$$

This fact follows from Equations (2), (3) and (4):

$$q^d \theta_{n-d-t} - \ell \theta_t + \theta_t = \theta_{n-t} - \theta_d - \theta_{n-t} + \theta_r + \theta_t = \theta_t + \theta_r - \theta_d,$$

as $\theta_t > \theta_d$.

Note that Π is the disjoint union of $\mathcal{A} = \{W \in \Pi : \dim(W) = t\}$ and $\mathcal{B} = \{W \in \Pi : \dim(W) \leq t - 1\}$. By counting the 1-dimensional subspaces not taken up by \mathcal{A} , we can bound the size of \mathcal{B} by

$$|\mathcal{B}| \geq \frac{\theta_n - |\mathcal{A}| \cdot \theta_t}{\theta_{t-1}}.$$

Since $|\mathcal{A}| = m_t$, we obtain from (7) that

$$|\Pi| = |\mathcal{A}| + |\mathcal{B}| \geq m_t + \frac{\theta_n - m_t \cdot \theta_t}{\theta_{t-1}} \geq \frac{\theta_n - (\ell q^t - q^t + q^d)(\theta_t - \theta_{t-1})}{\theta_{t-1}}. \tag{8}$$

By using Equation (4), the above inequality can be further simplified

$$|\Pi| \geq \ell q^t + q^d + \frac{\theta_{t+r} + q^t(\theta_t - \theta_{t-1}) - q^d \theta_t}{\theta_{t-1}} > \ell q^t + q^d + \frac{q^t(\theta_t - \theta_{t-1}) - q^d \theta_t}{\theta_{t-1}}.$$

As furthermore,

$$q^t(\theta_t - \theta_{t-1}) = q^{2t-1} > q^d \theta_t,$$

we finally obtain

$$|\Pi| \geq \ell q^t + q^d + 1.$$

This concludes the proof of the lemma.

We now prove that under the assumptions of Theorem 1, $\sigma_q(n, t) = \ell q^t + q^{\lceil \frac{t+r}{2} \rceil} + 1$.

Proof. Let Π be a subspace partition of $V = V(n, q)$ in which the largest subspace has dimension t . Let $\beta = \lceil (t+r)/2 \rceil$. If there is a subspace of dimension d in Π with $\beta \leq d < t$, then by Lemma 4

$$|\Pi| \geq \ell q^t + q^d + 1 \geq \ell q^t + q^\beta + 1. \quad (9)$$

It remains to consider the case where every subspace in Π has either dimension t or a dimension less than or equal to $\beta - 1$.

If there exists a hyperplane H of type b with $b_t \geq \ell + 1$, then by Lemma 2

$$|\Pi| = \sum_{i=1}^{n-1} b_i q^i + 1 \geq (\ell + 1)q^t + 1 \geq \ell q^t + q^\beta + 1, \quad (10)$$

where the last inequality holds since $\beta \leq t$.

So now assume that if $s_b \neq 0$ then $b_t \leq \ell$. Then Lemma 1(ii) yields

$$m_t \theta_{n-t} = \sum_{b \in B} b_t s_b \leq \ell \cdot \sum_{b \in B} s_b = \ell \cdot \theta_n. \quad (11)$$

From (2), we derive $\ell \theta_t < \theta_{n-t}$. By combining this inequality with (3), we obtain

$$\ell \theta_n = \ell(q^t \theta_{n-t} + \theta_t) = \ell q^t \theta_{n-t} + \ell \theta_t < \ell q^t \theta_{n-t} + \theta_{n-t}.$$

Consequently, (11) yields

$$m_t < \ell q^t + 1. \quad (12)$$

Note that Π is the disjoint union of $\mathcal{A} = \{W \in \Pi : \dim(W) = t\}$ and $\mathcal{B} = \{W \in \Pi : \dim(W) \leq \beta - 1\}$. By Equation (12) and since m_t is an integer, we may assume that $m_t \leq \ell q^t$. So by using Equation (3), we obtain that

$$|\Pi| = |\mathcal{A}| + |\mathcal{B}| \geq m_t + \frac{\theta_n - m_t \cdot \theta_t}{\theta_{\beta-1}} = \frac{\theta_n - m_t(\theta_t - \theta_{\beta-1})}{\theta_{\beta-1}} \geq \frac{\theta_n - \ell q^t(\theta_t - \theta_{\beta-1})}{\theta_{\beta-1}},$$

and hence from (4), and the fact that $\theta_{\beta-1}(q^\beta + 1) \leq \theta_{t+r}$, we conclude that

$$|\Pi| \geq \ell q^t + \frac{\theta_n - \ell q^t \theta_t}{\theta_{\beta-1}} = \ell q^t + \frac{\theta_{t+r}}{\theta_{\beta-1}} \geq \ell q^t + q^\beta + 1. \quad (13)$$

Summarizing the distinct cases we have considered, we thus obtain

$$\sigma_q(n, t) \geq \ell q^t + q^\beta + 1. \quad (14)$$

Finally, by using Proposition 1 we may conclude that

$$\sigma_q(n, t) = \ell q^t + q^\beta + 1.$$

4 The maximum size

In this section we now prove that under the assumptions of Theorem 1, $\rho_q(n, t) = \ell q^t + 1$.

Proof. Let Π be a subspace partition of $V = V(n, q)$ in which the smallest subspace has dimension t . Suppose $|\Pi| > \ell q^t + 1$. The type of Π is $(m_{n-1}, \dots, m_t, 0, \dots, 0)$. Let H be any hyperplane of V , and let $(b_{n-1}, \dots, b_t, 0, \dots, 0)$ be the type of H with respect to Π . Then by Lemma 2, we have

$$|\Pi| = 1 + \sum_{i=t}^{n-1} b_i q^i = 1 + q^t \sum_{i=t}^{n-1} b_i q^{i-t}.$$

Thus, $|\Pi| \equiv 1 \pmod{q^t}$, and by our above assumption on $|\Pi|$, we have $|\Pi| \geq \ell q^t + q^t + 1$. As the dimension of each member of Π is at least t , we may use relation (4) and the fact that $(q^t + 1)\theta_t = \theta_{2t}$ to conclude that

$$\theta_n \geq |\Pi| \theta_t \geq (\ell q^t + q^t + 1)\theta_t = \theta_n - \theta_{t+r} + \theta_{2t},$$

which is a contradiction as $\theta_{2t} > \theta_{t+r}$. Thus $|\Pi| \leq \ell q^t + 1$. Since Π is an arbitrary partition, we obtain

$$\rho_q(n, t) \leq \ell q^t + 1. \tag{15}$$

Hence, from Proposition 1 now follows that

$$\rho_q(n, t) = \ell q^t + 1.$$

5 Application to maximal partial t -spreads

A *partial t -spread* of $V = V(n, q)$ is a collection $\mathcal{S} = \{W_1, \dots, W_k\}$ of t -dimensional subspaces of V such that $W_i \cap W_j = \{0\}$ for $i \neq j$. The *size* of \mathcal{S} is its cardinality $|\mathcal{S}|$. If $V = \bigcup_{W \in \mathcal{S}} W$, then \mathcal{S} is called a *t -spread*. A partial t -spread is called *maximal* if it cannot be extended to a larger one. Maximal partial t -spreads have been extensively studied, see e.g. [4, 9, 11, 13, 15, 19, 20]. They can be used to construct error-correcting codes [6, 8], orthogonal arrays [7, 10], and recently factorial designs [22].

We let $\tau_q(n, t)$ denote the *minimum number* of subspaces in any maximal partial t -spread of $V(n, q)$. A maximal partial t -spread \mathcal{S} of $V(n, q)$ such that $|\mathcal{S}| = \tau_q(n, t)$, is called a *minimum size* maximal partial t -spread. Let n and t be fixed integers and let k and r be the unique integers defined by $n = kt + r$ and $0 \leq r < t$. Beutelspacher [1] showed that if $r = 0$ and $k \geq 2$, then

$$\tau_q(n + t - 1, t) = \sigma_q(n, t) = \frac{q^{kt} - 1}{q^t - 1}.$$

For $r > 0$, P. Govaerts [13] proved several results related to the number $\tau_q(n + t - 1, t)$. In particular, he provided the following upper bound for $\tau_q(n + t - 1, t)$.

Lemma 5 (Govaerts [13]) *Let n and $t > 1$ be integers such that $n \geq 2t$. Then there exist (see page 610 in [13] for a construction) maximal partial t -spreads of $V(n+t-1, q)$ of size $\sigma_q(n, t)$. Consequently, $\tau_q(n+t-1, t) \leq \sigma_q(n, t)$.*

We will prove the following theorem.

Theorem 2 *Let n and $t > 1$ be integers such that $n \geq 2t$. Then $\tau_q(n+t-1, t) = \sigma_q(n, t)$.*

The method employed to prove Theorem 2 will be the same as was used in [21] to prove $\tau_q(3t, t) = \sigma_q(2t+1, t)$. In particular, we will use Theorem 1 in Section 1. We first introduce the relevant definitions and a useful Lemma due to Govaerts [13]. A set of points B , i.e., 1-spaces of V , is called a *blocking set* with respect to the t -spaces of V if $W \cap B \neq \{0\}$ for any t -space W in V . Note that any $(n-t+1)$ -dimensional subspace of V is a blocking set with respect to the t -spaces of V . Such blocking sets are called *trivial*. The following lemma follows from the results of Govaerts (see Case 2, page 612 in [13]).

Lemma 6 (Govaerts [13]) *Let n and $t > 1$ be integers such that $n \geq 2t$. If \mathcal{S} is a minimum size maximal partial t -spread of $V(n, q)$, then $\bigcup_{W \in \mathcal{S}} W$ contains a trivial blocking set.*

In the proof of Theorem 2 we will also use the following proposition.

Proposition 2 *Let d, d' , and n be integers such that $0 < d' < d \leq n/2$. Then*

$$\sigma_q(n, d) < \sigma_q(n, d').$$

Proof. We will prove that $\sigma_q(n, t) < \sigma_q(n, t-1)$ holds, for $1 < t \leq n/2$.

If t divides n , then $\sigma_q(n, t) = \theta_n/\theta_t$. Consequently, by Theorem 1 and with the use of Equation (4), we note that it is always true that

$$\frac{\theta_n}{\theta_t} \leq \sigma_q(n, t) < \frac{\theta_n}{\theta_t} + q^\beta,$$

where $0 \leq r = n - kt < t$ and $\beta = \lceil (t+r)/2 \rceil$. As $\theta_t > q\theta_{t-1}$ and $q^\beta < \theta_n/\theta_t$, we thus get

$$\sigma_q(n, t) < 2\frac{\theta_n}{\theta_t} \leq q\frac{\theta_n}{\theta_t} < \frac{\theta_n}{\theta_{t-1}} \leq \sigma_q(n, t-1).$$

Proof. [Theorem 2] By Lemma 5, we have $\tau_q(n+t-1, t) \leq \sigma_q(n, t)$. So, it remains to show that

$$\tau_q(n+t-1, t) \geq \sigma_q(n, t). \tag{16}$$

Let \mathcal{S} be a minimum size maximal partial t -spread in $V(n+t-1, q)$. Then by Lemma 6, $A = \bigcup_{W \in \mathcal{S}} W$ contains a trivial blocking set. In other words, there exists an n -dimensional subspace $B \subseteq A$. Let

$$\Pi_{\mathcal{S}} = \{W \cap B : W \in \mathcal{S}\}.$$

Since B is a blocking set with respect to t -spaces, we have $W \cap B \neq \{0\}$ for any $W \in \mathcal{S}$. Thus, $\Pi_{\mathcal{S}}$ is a subspace partition of $B \cong V(n, q)$ containing subspaces of dimensions at most t . If $\Pi_{\mathcal{S}}$ contains a t -subspace, then it follows from Theorem 1 and the minimality of \mathcal{S} that

$$\tau_q(n + t - 1, t) = |\mathcal{S}| = |\Pi_{\mathcal{S}}| \geq \sigma_q(n, t).$$

If $\Pi_{\mathcal{S}}$ does not contain any t -subspace, then each subspace in $\Pi_{\mathcal{S}}$ has dimension at most $t - 1$ (and contains at most θ_{t-1} 1-dimensional subspaces). So the theorem now follows from the fact that the function $\sigma_q(n, t)$ is antimotone in t by Proposition 2.

6 Some remarks

Let Π be a subspace partition of $V = V(n, q)$ consisting of n_i subspaces of dimension d_i , for $1 \leq i \leq k$. Let us assume that $d_1 < d_2 < \dots < d_k$ (and $n_1 n_2 \dots n_k \neq 0$). In [16] a lower bound on n_1 was given as a function of q , d_1 and d_2 , and, as easily verified from that result, it is always true that $n_1 \geq \sigma_q(d_2, d_1)$. Working on the results of this paper has given us many indications that the following conjecture holds.

Conjecture 1 *Let Π be a subspace partition of $V(n, q)$ with $n_i > 0$ subspaces of dimension d_i , $1 \leq i \leq k$, and where $d_1 < \dots < d_k$. Then, for any integer j , $1 \leq j < k$, we have*

$$n_1 + \dots + n_j \geq \sigma_q(d_{j+1}, d_j).$$

Let us also remark that for $n \leq 2t - 1$, the problem of determining the minimum size $\tau_q(n + t - 1, t)$ of a maximal partial t -spread in $V(n + t - 1, q)$ is still open. For $t = 2$ and $n = 3$, the following lower bound was achieved by Glynn [12]:

$$\tau_q(4, 2) \geq 2q,$$

while the following two upper bounds are due to Gács and Szönyi [11]:

$$\tau_q(4, 2) \leq (2 \log_2 q + 1)q + 1, \quad \text{if } q \text{ odd,}$$

and

$$\tau_q(4, 2) \leq (6.1 \ln q + 1)q + 1, \quad \text{if } q > q_0 \text{ even.}$$

Acknowledgment. The last three authors of this paper wish to thank their host Olof Heden and the Department of Mathematics at KTH for their warm hospitality while working on this paper during their visit there.

References

- [1] A. Beutelspacher, Partial spreads in finite projective spaces and partial designs, *Math. Zeit.* 145 (1975), 211–229.
- [2] A. Beutelspacher, Blocking sets and partial spreads in finite projective spaces, *Geom. Dedicata* 9 (1980), 425–449.
- [3] A. Blinco, S. El-Zanati, G. Seelinger, P. Sissokho, L. Spence, and C. Vanden Eynden, On Vector space partitions and uniformly resolvable designs, *Des. Codes Cryptogr.* 15 (2008), 69–77.
- [4] A. Blokhuis, A. Brouwer, and H. Wilbrink, Blocking sets in $\text{PG}(2, p)$ for small p , and partial spreads in $\text{PG}(3, 7)$, *Advanc. Geom.* 3 (2003), 245–243.
- [5] T. Bu, Partitions of a vector space, *Disc. Math.* 31 (1980), 79–83.
- [6] W. Clark and L. Dunning, Partial partitions of vector spaces arising from the construction of byte error control codes, *Ars Combin.* 33 (1992), 161–177.
- [7] C. Colbourn and M. Greig, Orthogonal arrays of index more than one, in: The CRC handbook of combinatorial designs. Edited by C. Colbourn and J. Dinitz. *CRC Press Series on Discrete Mathematics and its Applications*, CRC Press, Boca Raton, FL (2007), 219–228.
- [8] D. Drake and J. Freeman, Partial t -spreads and group constructible (s, r, μ) -nets, *J. Geom.* 13 (1979), 211–216.
- [9] J. Eisfeld, L. Storme, and P. Sziklai, On the spectrum of the sizes of maximal partial line spreads in $\text{PG}(2n, q)$, $n \geq 3$, *Des. Codes Cryptogr.* 36 (2005), 101–110.
- [10] S. El-Zanati, H. Jordon, G. Seelinger, P. Sissokho, and L. Spence, The maximum size of a partial 3-spread in a finite vector space over $\text{GF}(2)$, *Des. Codes Cryptogr.* 54 (2010), 101–107.
- [11] A. Gács and T. Szönyi, On maximal partial spreads in $\text{PG}(n, q)$, *Des. Codes Cryptogr.* 29 (2003), 123–129.
- [12] D. G. Glynn, A lower bound for maximal partial spreads in $\text{PG}(3, q)$, *Ars Combin.* 13 (1982), 39–40.
- [13] P. Govaerts, Small maximal partial t -spreads, *Bull. Belg. Math. Soc. Simon Stevin* 12 (2005), 607–615.
- [14] O. Heden, Partitions of finite abelian groups, *Europ. J. Combin.* 7 (1986), 11–25.
- [15] O. Heden, A Maximal Partial Spread of Size 45 in $\text{PG}(3, 7)$, *Des. Codes Cryptogr.* 22 (2001), 331–334.

- [16] O. Heden, On the length of the tail of a vector space partition, *Disc. Math.* 309 (2009), 6169–6180.
- [17] O. Heden and J. Lehmann, Some necessary conditions for vector space partitions, *submitted*, available at arXiv:1101.3745v1.
- [18] M. Herzog and J. Schönheim, Group partition, factorization and the vector covering problem, *Canad. Math. Bull.* 15(2) (1972), 207–214.
- [19] S. Hong and A. Patel, A general class of maximal codes for computer applications, *IEEE Trans. Comput.* C-21 (1972), 1322–1331.
- [20] D. Jungnickel and L. Storme, A note on maximal partial spreads with deficiency $q + 1$, q even, *J. Combin. Theory Ser. A*, 102 (2003), 443–446.
- [21] E. Năstase and P. Sissokho, The minimum size of a finite subspace partition, *to appear in Linear Algebr. and its Appl.*
- [22] Pritam Ranjan, Derek R. Bingham, and Angela M. Dean, Existence and construction of randomization defining contrast subspaces for regular factorial designs, *Ann. Statist.*, 37 (2009), 3580–3599.
- [23] G. Spera, On Partitions of Finite Vector Spaces, arXiv:0902.3075v1 [math.CO] (2009) 14 pp.

O. Heden (olohed@math.kth.se), Department of Mathematics, KTH, S-100 44 Stockholm, Sweden.

J. Lehmann (jlehmann@math.uni-bremen.de), Department of Mathematics, Bremen University, Bibliothekstrasse 1 - MZH, 28359 Bremen, Germany.

E. Năstase (nastasee@xavier.edu): Department of Mathematics and Computer Science, Xavier University, 3800 Victory Parkway, Cincinnati, Ohio 45207.

P. Sissokho (psissok@ilstu.edu): Mathematics Department, Illinois State University, Normal, Illinois 61790.