CONGRUENCES INVOLVING $\binom{2k}{k}^2 \binom{3k}{k} m^{-k}$

ZHI-Hong Sun

School of the Mathematical Sciences, Huaiyin Normal University, Huaian, Jiangsu 223001, PR China Email: zhihongsun@yahoo.com

Homepage: http://www.hytc.edu.cn/xsjl/szh

ABSTRACT. Let p > 3 be a prime, and let m be an integer with $p \nmid m$. In the paper, based on the work of Brillhart and Morton, by using the work of Ishii and Deuring's theorem for elliptic curves with complex multiplication we solve some conjectures of Zhi-Wei Sun concerning $\sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} m^{-k} \pmod{p^2}$.

MSC: Primary 11A07, Secondary 33C45, 11E25, 11G07, 11L10, 05A10, 05A19 Keywords: Congruence; Legendre polynomial; character sum; elliptic curve; binary quadratic form

1. Introduction.

For positive integers a, b and n, if $n = ax^2 + by^2$ for some integers x and y, we briefly say that $n = ax^2 + by^2$. Let p > 3 be a prime. In 2003, Rodriguez-Villegas[RV] posed some conjectures on supercongruences modulo p^2 . One of his conjectures is equivalent to

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{108^k} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

This conjecture has been solved by Mortenson[Mo] and Zhi-Wei Sun[Su2].

Let \mathbb{Z} be the set of integers, and for a prime p let \mathbb{Z}_p be the set of rational numbers whose denominator is coprime to p. Recently the author's brother Zhi-Wei Sun[Su1] posed many conjectures for $\sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} m^{-k} \pmod{p^2}$, where p > 3 is a prime and $m \in \mathbb{Z}$ with $p \nmid m$. For example, he conjectured that (see [Su1, Conjecture A13])

$$(1.1) \qquad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k} \equiv \begin{cases} 0 \pmod{p^2} & \text{if } p \equiv 7, 11, 13, 14 \pmod{15}, \\ 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 15y^2 \equiv 1, 4 \pmod{15}, \\ 20x^2 - 2p \pmod{p^2} & \text{if } p = 5x^2 + 3y^2 \equiv 2, 8 \pmod{15}. \end{cases}$$

Let $\{P_n(x)\}$ be the Legendre polynomials given by (see [MOS, pp. 228-232], [G, (3.132)-(3.133)])

(1.2)
$$P_n(x) = \frac{1}{2^n} \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{k} (-1)^k \binom{2n-2k}{n} x^{n-2k} = \frac{1}{2^n \cdot n!} \cdot \frac{d^n}{dx^n} (x^2 - 1)^n,$$

The author is supported by the Natural Sciences Foundation of China (grant No. 10971078).

where [a] is the greatest integer not exceeding a. From (1.2) we see that

$$(1.3) P_n(-x) = (-1)^n P_n(x).$$

Let $(\frac{a}{m})$ be the Jacobi symbol. For a prime p > 3, in [S2] the author showed that

$$(1.4) P_{\left[\frac{p}{3}\right]}(t) \equiv \sum_{k=0}^{[p/3]} {2k \choose k} {3k \choose k} \left(\frac{1-t}{54}\right)^k \equiv \sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} \left(\frac{1-t}{54}\right)^k \pmod{p}.$$

We note that $p \mid {2k \choose k} {3k \choose k}$ for $\frac{p}{3} < k < p$. In the paper, using the work of Brillhart and Morton[BM] we prove that

$$(1.5) P_{\left[\frac{p}{3}\right]}(t) \equiv -\left(\frac{p}{3}\right) \sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p}\right) \pmod{p}.$$

Based on (1.5) and the work of Ishii[I], we determine

$$P_{\left[\frac{p}{3}\right]}(t) \pmod{p} \quad \text{for} \quad t = \frac{5}{4}, \frac{5}{\sqrt{-2}}, \frac{\sqrt{-11}}{4}, \frac{1}{\sqrt{2}}, \sqrt{5}, \frac{9}{20}\sqrt{5}, \frac{\sqrt{17}}{4}, \frac{5}{32}\sqrt{41}, \frac{53}{500}\sqrt{89}.$$

For instance, if $p \equiv 1, 4 \pmod{5}$ is a prime, then

$$P_{[\frac{p}{3}]}(\sqrt{5}) \equiv \left\{ \begin{array}{ll} 2x(\frac{x}{3}) \pmod{p} & \text{if } p = x^2 + 15y^2 \equiv 1,4 \pmod{15}, \\ 0 \pmod{p} & \text{if } p \equiv 11,14 \pmod{15}. \end{array} \right.$$

Let p > 3 be a prime, $m \in \mathbb{Z}_p$, $m \not\equiv 0 \pmod{p}$ and $t = \sqrt{1 - 108/m}$. In the paper we show that

(1.6)
$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k} \equiv P_{\left[\frac{p}{3}\right]}(t)^2 \pmod{p}$$

and that

(1.7)
$$P_{\left[\frac{p}{3}\right]}(t) \equiv 0 \pmod{p} \quad \text{implies} \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k} \equiv 0 \pmod{p^2}.$$

On the basis of (1.6) and (1.7), we prove some congruences for $\sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} m^{-k}$ in the cases $m=8,64,216,-27,-192,-8640,-12^3,-48^3,-300^3$. Thus we partially solve some conjectures posed by Zhi-Wei Sun in [Su1]. As two examples, for odd primes $p \neq 11$ we have

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k} \equiv \begin{cases} x^2 \pmod{p} & \text{if } (\frac{p}{11}) = 1 \text{ and so } 4p = x^2 + 11y^2, \\ 0 \pmod{p^2} & \text{if } (\frac{p}{11}) = -1, \end{cases}$$

for odd primes p with $(\frac{17}{p}) = 1$ we have

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-12)^{3k}} \equiv \left\{ \begin{array}{l} x^2 \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 51y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{array} \right.$$

2. A general congruence modulo p^2 .

Lemma 2.1. Let m be a nonnegative integer. Then

$$\sum_{k=0}^{m} \binom{2k}{k}^2 \binom{3k}{k} \binom{k}{m-k} (-27)^{m-k} = \sum_{k=0}^{m} \binom{2k}{k} \binom{3k}{k} \binom{2(m-k)}{m-k} \binom{3(m-k)}{m-k}.$$

We prove the lemma by using WZ method and Mathematica. Clearly the result is true for m = 0, 1. Since both sides satisfy the same recurrence relation

$$81(m+1)(3m+2)(3m+4)S(m) - 3(2m+3)(9m^2 + 27m + 22)S(m+1) + (m+2)^3S(m+2) = 0,$$

we see that the lemma is true. The proof certificate for the left hand side is

$$-\frac{729k^2(m+2)(m-2k)(m-2k+1)}{(m-k+1)(m-k+2)},$$

and the proof certificate for the right hand side is

$$\frac{9k^2(3m-3k+1)(3m-3k+2)(9m^2-9mk+30m-14k+24)}{(m-k+1)^2(m-k+2)^2}$$

Theorem 2.1. Let p be an odd prime and let x be a variable. Then

$$\sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} (x(1-27x))^k \equiv \left(\sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} x^k\right)^2 \pmod{p^2}.$$

Proof. It is clear that

$$\sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} (x(1-27x))^k$$

$$= \sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} x^k \sum_{r=0}^k {k \choose r} (-27x)^r$$

$$= \sum_{m=0}^{2(p-1)} x^m \sum_{k=0}^{\min\{m,p-1\}} {2k \choose k}^2 {3k \choose k} {k \choose m-k} (-27)^{m-k}.$$

Suppose $p \le m \le 2p-2$ and $0 \le k \le p-1$. If $k > \frac{p}{2}$, then $p \mid \binom{2k}{k}$ and so $p^2 \mid \binom{2k}{k}^2$. If $k < \frac{p}{2}$, then $m-k \ge p-k > k$ and so $\binom{k}{m-k} = 0$. Thus, from the above and Lemma 2.1

we deduce that

$$\sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} (x(1-27x))^k$$

$$\equiv \sum_{m=0}^{p-1} x^m \sum_{k=0}^m {2k \choose k}^2 {3k \choose k} {k \choose m-k} (-27)^{m-k}$$

$$= \sum_{m=0}^{p-1} x^m \sum_{k=0}^m {2k \choose k} {3k \choose k} {2(m-k) \choose m-k} {3(m-k) \choose m-k}$$

$$= \sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} x^k \sum_{m=k}^{p-1} {2(m-k) \choose m-k} {3(m-k) \choose m-k} x^{m-k}$$

$$= \sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} x^k \sum_{r=0}^{p-1-k} {2r \choose r} {3r \choose r} x^r$$

$$= \sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} x^k (\sum_{r=0}^{p-1} {2r \choose r} {3r \choose r} x^r - \sum_{r=p-k}^{p-1} {2r \choose r} {3r \choose r} x^r)$$

$$= (\sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} x^k)^2 - \sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} x^k \sum_{r=p-k}^{p-1} {2r \choose r} {3r \choose r} x^r \pmod{p^2}.$$

If $\frac{2p}{3} \le k \le p-1$, then $\binom{2k}{k}\binom{3k}{k} = \frac{(3k)!}{k!^3} \equiv 0 \pmod{p^2}$. If $0 \le k \le \frac{p}{3}$ and $p-k \le r \le p-1$, then $\frac{2p}{3} \le r \le p-1$ and so $\binom{2r}{r}\binom{3r}{r} = \frac{(3r)!}{r!^3} \equiv 0 \pmod{p^2}$. If $\frac{p}{3} < k < \frac{2p}{3}$ and $p-k \le r \le p-1$, then $r \ge p-k > \frac{p}{3}$, $\binom{2k}{k}\binom{3k}{k} = \frac{(3k)!}{k!^3} \equiv 0 \pmod{p}$ and $\binom{2r}{r}\binom{3r}{r} = \frac{(3r)!}{r!^3} \equiv 0 \pmod{p}$. Hence, for $0 \le k \le p-1$ and $p-k \le r \le p-1$ we have $p^2 \mid \binom{2k}{k}\binom{3k}{k}\binom{2r}{r}\binom{3r}{r}$ and so

$$\sum_{k=0}^{p-1} {2k \choose k} {3k \choose k} x^k \sum_{r=p-k}^{p-1} {2r \choose r} {3r \choose r} x^r \equiv 0 \pmod{p^2}.$$

Therefore the result follows.

Corollary 2.1. Let p > 3 be a prime and $m \in \mathbb{Z}_p$ with $m \not\equiv 0 \pmod{p}$. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k} \equiv \left(\sum_{k=0}^{p-1} \binom{2k}{k} \binom{3k}{k} \left(\frac{1-\sqrt{1-108/m}}{54}\right)^k\right)^2 \pmod{p^2}.$$

Proof. Taking $x = \frac{1 - \sqrt{1 - 108/m}}{54}$ in Theorem 2.1 we deduce the result.

3. Congruences for $P_{[p/3]}(t) \pmod{p}$ **.** Let $W_n(x)$ be the Deuring polynomial given by

(3.1)
$$W_n(x) = \sum_{k=0}^n \binom{n}{k}^2 x^k.$$

It is known that ([G,(3.134)],[BM])

(3.2)
$$W_n(x) = (1-x)^n P_n\left(\frac{1+x}{1-x}\right).$$

Let p > 3 be a prime, $m, n \in \mathbb{Z}_p$ and $4m^3 + 27n^2 \not\equiv 0 \pmod{p}$. From [Mor, Theorem 3.3] we have

$$(3.3) \sum_{x=0}^{p-1} \left(\frac{x^3 + mx + n}{p} \right)$$

$$\equiv -(-48m)^{\frac{1-(\frac{p}{3})}{2}} (864n)^{\frac{1-(\frac{-1}{p})}{2}} (-16(4m^3 + 27n^2))^{\left[\frac{p}{12}\right]} J_p \left(\frac{2^8 \cdot 3^3 m^3}{4m^3 + 27n^2} \right) \pmod{p},$$

where $J_p(t)$ is a certain Jacobi polynomial given by

(3.4)
$$J_p(t) = 1728^{\left[\frac{p}{12}\right]} P_{\left[\frac{p}{12}\right]}^{\left(-\frac{1}{3}\left(\frac{p}{3}\right), -\frac{1}{2}\left(\frac{-1}{p}\right)\right)} \left(1 - \frac{t}{864}\right)$$

and

$$P_k^{(\alpha,\beta)}(x) = \frac{1}{2^k} \sum_{r=0}^k \binom{k+\alpha}{r} \binom{k+\beta}{k-r} (x-1)^{k-r} (x+1)^r.$$

Theorem 3.1. Let p > 3 be a prime and $t \in \mathbb{Z}_p$. Then

$$P_{\left[\frac{p}{3}\right]}(t) \equiv -\left(\frac{p}{3}\right) \sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p}\right) \pmod{p}.$$

Proof. It is well known that $P_n(1) = 1$. Since $P_{\left[\frac{p}{3}\right]}(1) = 1$ and

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 3x - 2}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{(x+1)^2 (x-2)}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{x-2}{p} \right) - \left(\frac{-1-2}{p} \right) = -\left(\frac{p}{3} \right),$$

we see that the result is true for $t \equiv 1 \pmod{p}$. Since $P_{[\frac{p}{3}]}(-1) = (-1)^{[\frac{p}{3}]} P_{[\frac{p}{3}]}(1) = (\frac{p}{3})$ and

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 27x + 54}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{(-3x)^3 - 27(-3x) + 54}{p} \right) = \left(\frac{-3}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 3x - 2}{p} \right) = -1,$$

we see that the result is also true for $t \equiv -1 \pmod{p}$.

Now we assume $t \not\equiv \pm 1 \pmod{p}$. Set $W_n(x) = \sum_{k=0}^n \binom{n}{k}^2 x^k$. From [BM, Theorem 6] we know that

$$W_{\left[\frac{p}{3}\right]}\left(1 - \frac{x}{27}\right) \equiv u_p(x)(x - 27)^{\left[\frac{p}{12}\right]} J_p\left(\frac{x(x - 24)^3}{x - 27}\right) \pmod{p},$$

where $J_p(x)$ is a certain Jacobi polynomial given by (3.4) and

$$u_p(x) = \begin{cases} 1 & \text{if } p \equiv 1 \pmod{12}, \\ -3(x-24) & \text{if } p \equiv 5 \pmod{12}, \\ x^2 - 36x + 216 & \text{if } p \equiv 7 \pmod{12}, \\ -3(x-24)(x^2 - 36x + 216) & \text{if } p \equiv 11 \pmod{12}. \end{cases}$$

Set x = 54/(t+1). We then have

$$W_{[\frac{p}{2}]}((t-1)/(t+1))$$

$$\equiv \begin{cases} \left(\frac{27(1-t)}{1+t}\right)^{\left[\frac{p}{12}\right]} J_p\left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right) \pmod{p} & \text{if } p \equiv 1 \pmod{12}, \\ \frac{18(4t-5)}{t+1} \left(\frac{27(1-t)}{1+t}\right)^{\left[\frac{p}{12}\right]} J_p\left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right) \pmod{p} & \text{if } p \equiv 5 \pmod{12}, \\ \frac{108(2t^2-14t+11)}{(t+1)^2} \left(\frac{27(1-t)}{1+t}\right)^{\left[\frac{p}{12}\right]} J_p\left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right) \pmod{p} & \text{if } p \equiv 7 \pmod{12}, \\ \frac{1944(4t-5)(2t^2-14t+11)}{(t+1)^3} \left(\frac{27(1-t)}{1+t}\right)^{\left[\frac{p}{12}\right]} J_p\left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right) \pmod{p} & \text{if } p \equiv 1 \pmod{12}, \end{cases}$$

By (3.2) we have

$$(3.6) W_{\left[\frac{p}{3}\right]}\left(\frac{t-1}{t+1}\right) = \left(1 - \frac{t-1}{t+1}\right)^{\left[\frac{p}{3}\right]} P_{\left[\frac{p}{3}\right]}\left(\frac{1 + (t-1)/(t+1)}{1 - (t-1)/(t+1)}\right) = \left(\frac{2}{t+1}\right)^{\left[\frac{p}{3}\right]} P_{\left[\frac{p}{3}\right]}(t).$$

If $p \equiv 2 \pmod{3}$ and $t \equiv \frac{5}{4} \pmod{p}$, from the above we get

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5}{4}\right) = \left(\frac{\frac{5}{4}+1}{2}\right)^{\left[\frac{p}{3}\right]} W_{\left[\frac{p}{3}\right]}\left(\frac{\frac{5}{4}-1}{\frac{5}{4}+1}\right) \equiv 0 \pmod{p}.$$

On the other hand,

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{x^3 - 27/4}{p} \right) = \sum_{y=0}^{p-1} \left(\frac{y - 27/4}{p} \right) = 0.$$

Thus the result is true when $p \equiv 2 \pmod 3$ and $t \equiv \frac{5}{4} \pmod p$. If $p \equiv 3 \pmod 4$ and $2t^2 - 14t + 11 \equiv 0 \pmod p$, from (3.5) and (3.6) we deduce that

$$P_{\left[\frac{p}{3}\right]}(t) = \left(\frac{t+1}{2}\right)^{\left[\frac{p}{3}\right]} W_{\left[\frac{p}{3}\right]} \left(\frac{t-1}{t+1}\right) \equiv 0 \pmod{p}.$$

As

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{(-x)^3 + 3(4t - 5)(-x)}{p} \right) = -\sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x}{p} \right),$$

we see that

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x}{p} \right) = 0.$$

Thus the result is true when $p \equiv 3 \pmod 4$ and $2t^2 - 14t + 11 \equiv 0 \pmod p$. Set m = 3(4t - 5) and $n = 2(2t^2 - 14t + 11)$. Then

$$4m^3 + 27n^2 = -432(1-t)(1+t)^3$$
 and so $\frac{2^8 \cdot 3^3 m^3}{4m^3 + 27n^2} = \frac{432(5-4t)^3}{(1-t)(1+t)^3}$.

By the above we need only to assume $m \not\equiv 0 \pmod{p}$ for $p \equiv 2 \pmod{3}$ and $n \not\equiv 0 \pmod{p}$ for $p \equiv 3 \pmod{4}$. From (3.3) we see that

$$J_p\left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right) = J_p\left(\frac{2^8 \cdot 3^3 m^3}{4m^3 + 27n^2}\right)$$

$$\equiv -(-48m)^{\frac{(\frac{2}{3})-1}{2}} (864n)^{\frac{(-\frac{1}{p})-1}{2}} (-16(4m^3 + 27n^2))^{-\left[\frac{p}{12}\right]} \sum_{x=0}^{p-1} \left(\frac{x^3 + mx + n}{p}\right) \pmod{p}.$$

If $p \equiv 1 \pmod{12}$, from all the above we deduce that

$$\begin{split} &P_{\left[\frac{p}{3}\right]}(t) \\ &= \left(\frac{t+1}{2}\right)^{\left[\frac{p}{3}\right]} W_{\left[\frac{p}{3}\right]} \left(\frac{t-1}{t+1}\right) \equiv \left(\frac{t+1}{2}\right)^{\frac{p-1}{3}} \left(\frac{27(1-t)}{1+t}\right)^{\frac{p-1}{12}} J_p \left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right) \\ &\equiv -2^{-\frac{p-1}{3}} (3(t+1))^{\frac{p-1}{4}} (1-t)^{\frac{p-1}{12}} (16 \cdot 432(1-t)(1+t)^3)^{-\frac{p-1}{12}} \sum_{x=0}^{p-1} \left(\frac{x^3+mx+n}{p}\right) \\ &\equiv -\sum_{x=0}^{p-1} \left(\frac{x^3+3(4t-5)x+2(2t^2-14t+11)}{p}\right) \pmod{p}. \end{split}$$

If $p \equiv 5 \pmod{12}$, from all the above we deduce that

$$P_{\left[\frac{p}{3}\right]}(t)$$

$$= \left(\frac{t+1}{2}\right)^{\left[\frac{p}{3}\right]} W_{\left[\frac{p}{3}\right]} \left(\frac{t-1}{t+1}\right) \equiv \left(\frac{t+1}{2}\right)^{\frac{p-2}{3}} \frac{18(4t-5)}{t+1} \left(\frac{27(1-t)}{1+t}\right)^{\frac{p-5}{12}} J_p \left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right)$$

$$\equiv 2^{-\frac{p-5}{3}} 3^{\frac{p+3}{4}} (4t-5)(1+t)^{\frac{p-5}{4}} (1-t)^{\frac{p-5}{12}} (144(4t-5))^{-1}$$

$$\times (16 \cdot 432(1-t)(1+t)^3)^{-\frac{p-5}{12}} \sum_{x=0}^{p-1} \left(\frac{x^3+mx+n}{p}\right)$$

$$\equiv \sum_{x=0}^{p-1} \left(\frac{x^3+3(4t-5)x+2(2t^2-14t+11)}{p}\right) \pmod{p}.$$

If $p \equiv 7 \pmod{12}$, from all the above we deduce that

$$P_{\left[\frac{p}{3}\right]}(t) = \left(\frac{t+1}{2}\right)^{\left[\frac{p}{3}\right]} W_{\left[\frac{p}{3}\right]}\left(\frac{t-1}{t+1}\right)$$

$$\equiv \left(\frac{t+1}{2}\right)^{\frac{p-1}{3}} \frac{108(2t^2 - 14t + 11)}{(t+1)^2} \left(\frac{27(1-t)}{1+t}\right)^{\frac{p-7}{12}} J_p\left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right)$$

$$\equiv -2^{-\frac{p-7}{3}} 3^{\frac{p+5}{4}} (2t^2 - 14t + 11)(1+t)^{\frac{p-7}{4}} (1-t)^{\frac{p-7}{12}} (1728(2t^2 - 14t + 11))^{-1}$$

$$\times (16 \cdot 432(1-t)(1+t)^3)^{-\frac{p-7}{12}} \sum_{x=0}^{p-1} \left(\frac{x^3 + mx + n}{p}\right)$$

$$\equiv -\sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p}\right) \pmod{p}.$$

If $p \equiv 11 \pmod{12}$, from all the above we deduce that

$$P_{\left[\frac{p}{3}\right]}(t) = \left(\frac{t+1}{2}\right)^{\left[\frac{p}{3}\right]} W_{\left[\frac{p}{3}\right]}\left(\frac{t-1}{t+1}\right)$$

$$\equiv \left(\frac{t+1}{2}\right)^{\frac{p-2}{3}} \frac{1944(4t-5)(2t^2-14t+11)}{(t+1)^3} \left(\frac{27(1-t)}{1+t}\right)^{\frac{p-11}{12}} J_p\left(\frac{432(5-4t)^3}{(1-t)(1+t)^3}\right)$$

$$\equiv 2^{-\frac{p-11}{3}} 3^{\frac{p-11}{4}+5} (4t-5)(2t^2-14t+11)(1+t)^{\frac{p-11}{4}} (1-t)^{\frac{p-11}{12}} (48m)^{-1} (864n)^{-1}$$

$$\times \left(16 \cdot 432(1-t)(1+t)^3\right)^{-\frac{p-11}{12}} \sum_{x=0}^{p-1} \left(\frac{x^3+mx+n}{p}\right)$$

$$\equiv \sum_{x=0}^{p-1} \left(\frac{x^3+3(4t-5)x+2(2t^2-14t+11)}{p}\right) \pmod{p}.$$

This proves the theorem.

Corollary 3.1. Let p > 3 be a prime and let t be a variable. Then

$$\begin{split} &\sum_{k=0}^{[p/3]} \binom{2k}{k} \binom{3k}{k} \left(\frac{1-t}{54}\right)^k \\ &\equiv P_{\left[\frac{p}{3}\right]}(t) \equiv -\left(\frac{p}{3}\right) \sum_{x=0}^{p-1} (x^3 + 3(4t-5)x + 2(2t^2 - 14t + 11))^{\frac{p-1}{2}} \pmod{p}. \end{split}$$

Proof. From [S2, Lemma 2.3] we have $P_{\left[\frac{p}{3}\right]}(t) \equiv \sum_{k=0}^{\left[p/3\right]} {2k \choose k} {3k \choose k} (\frac{1-t}{54})^k \pmod{p}$. By Theorem 3.1 and Euler's criterion, the result is true for $t=0,1,\ldots,p-1$. Since both sides are polynomials of t with degree at most p-1. Using Lagrange's theorem we obtain the result.

Corollary 3.2. Let $p \geq 17$ be a prime and $t \in \mathbb{Z}_p$. Then

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p} \right)$$
$$= \left(\frac{p}{3} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 3(4t + 5)x + 2(2t^2 + 14t + 11)}{p} \right).$$

Proof. Since $P_{[\frac{p}{3}]}(-t) = (-1)^{[\frac{p}{3}]} P_{[\frac{p}{3}]}(t) = (\frac{p}{3}) P_{[\frac{p}{3}]}(t)$, by Theorem 3.1 we have

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p} \right)$$

$$\equiv \left(\frac{p}{3} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 3(4t + 5)x + 2(2t^2 + 14t + 11)}{p} \right) \pmod{p}.$$

By Weil's estimate ([BEW, p.183]) we have

$$\left| \sum_{x=0}^{p-1} \left(\frac{x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11)}{p} \right) \right| \le 2\sqrt{p},$$

$$\left| \sum_{x=0}^{p-1} \left(\frac{x^3 - 3(4t + 5)x + 2(2t^2 + 14t + 11)}{p} \right) \right| \le 2\sqrt{p}.$$

Since $4\sqrt{p} < p$ for $p \ge 17$, from the above we deduce the result.

Corollary 3.3. Let p > 3 be a prime. Then

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 120x + 506}{p} \right) = \left\{ \begin{array}{ll} (\frac{2}{p})L & \text{if } 3 \mid p-1, \ 4p = L^2 + 27M^2 \ and \ 3 \mid L-1, \\ 0 & \text{if } p \equiv 2 \ (\text{mod } 3). \end{array} \right.$$

Proof. It is easy to check the result for p=5,7,11,13. Now we assume $p\geq 17$. Taking $t=\frac{5}{4}$ in Corollary 3.2 we find that

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - \frac{27}{4}}{p} \right) = \left(\frac{p}{3} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 30x + \frac{253}{4}}{p} \right) = \left(\frac{p}{3} \right) \sum_{x=0}^{p-1} \left(\frac{\left(\frac{x}{2} \right)^3 - 30 \cdot \frac{x}{2} + \frac{253}{4}}{p} \right)$$
$$= \left(\frac{p}{3} \right) \left(\frac{2}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 120x + 506}{p} \right).$$

For $p \equiv 2 \pmod{3}$ it is clear that

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - \frac{27}{4}}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{x - \frac{27}{4}}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{x}{p} \right) = 0.$$

Thus the result is true when $p \equiv 2 \pmod{3}$.

Now assume $p \equiv 1 \pmod{3}$, $p = A^2 + 3B^2$, $4p = L^2 + 27M^2$ and $A \equiv L \equiv 1 \pmod{3}$. It is known that $2^{\frac{p-1}{3}} \equiv 1 \pmod{p}$ if and only if $3 \mid B$. When $3 \nmid B$ we choose the sign of B so that $B \equiv 1 \pmod{3}$. By [S1, (2.12)] we have $2^{(p-1)/3} \equiv \frac{1}{2}(-1 - \frac{A}{B}) \pmod{p}$. From [S1, (2.9)-(2.11)] we deduce that

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 27/4}{p} \right) = 1 + \sum_{x=1}^{p-1} \left(\frac{x^3 - 27/4}{p} \right)$$

$$= \begin{cases} -2A = L \pmod{p} & \text{if } 2^{\frac{p-1}{3}} \equiv 1 \pmod{p}, \\ A + 3B = L \pmod{p} & \text{if } 2^{\frac{p-1}{3}} \not\equiv 1 \pmod{p} \text{ and } B \equiv 1 \pmod{3}. \end{cases}$$

Thus

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 120x + 506}{p} \right) = \left(\frac{2}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 27/4}{p} \right) = \left(\frac{2}{p} \right) L.$$

This completes the proof.

Theorem 3.2. Let p > 3 be a prime. Then

(i) If $p \equiv 2 \pmod{3}$, then

$$\sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{(-216)^k} \equiv \sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{24^k} \equiv P_{\left[\frac{p}{3}\right]} \left(\frac{5}{4}\right) \equiv 0 \pmod{p}.$$

(ii) If $p \equiv 1 \pmod{3}$ and so $4p = L^2 + 27M^2$ with $L, M \in \mathbb{Z}$ and $L \equiv 1 \pmod{3}$, then

$$\sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{(-216)^k} \equiv \sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{24^k} \equiv P_{\left[\frac{p}{3}\right]} \left(\frac{5}{4}\right) \equiv -L \equiv \left(\frac{-2}{p}\right) \left(\frac{\frac{2(p-1)}{3}}{\left[\frac{p}{12}\right]}\right) \; (\text{mod } p).$$

Proof. Putting $t = \pm \frac{5}{4}$ in Corollary 3.1 we get

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5}{4}\right) \equiv \sum_{k=0}^{[p/3]} \frac{\binom{2k}{k}\binom{3k}{k}}{(-216)^k} \pmod{p} \quad \text{and} \quad P_{\left[\frac{p}{3}\right]}\left(-\frac{5}{4}\right) \equiv \sum_{k=0}^{[p/3]} \frac{\binom{2k}{k}\binom{3k}{k}}{24^k} \pmod{p}.$$

This together with (1.3) yields

$$\sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{(-216)^k} \equiv \left(\frac{p}{3}\right) \sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{24^k} \pmod{p}.$$

If $p \equiv 2 \pmod{3}$, by Theorem 3.1 we have

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5}{4}\right) \equiv \sum_{x=0}^{p-1} \left(\frac{x^3 - \frac{27}{4}}{p}\right) = \sum_{x=0}^{p-1} \left(\frac{x - \frac{27}{4}}{p}\right) = \sum_{x=0}^{p-1} \left(\frac{x}{p}\right) = 0 \pmod{p}.$$

Thus (i) is true.

Now assume $p \equiv 1 \pmod{3}$, $4p = L^2 + 27M^2$ and $L \equiv 1 \pmod{3}$. By Theorem 3.1 and the proof of Corollary 3.3 we have

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5}{4}\right) \equiv -\sum_{x=0}^{p-1} \left(\frac{x^3 - \frac{27}{4}}{p}\right) = -L \pmod{p}.$$

On the other hand, by the proof of Theorem 3.1,

$$\begin{split} P_{\left[\frac{p}{3}\right]}\left(\frac{5}{4}\right) &= \left(\frac{\frac{5}{4}+1}{2}\right)^{\left[\frac{p}{3}\right]} W_{\left[\frac{p}{3}\right]}\left(\frac{\frac{5}{4}-1}{\frac{5}{4}+1}\right) \\ &= \left\{ \begin{array}{l} \left(\frac{9}{8}\right)^{\frac{p-1}{3}} \left(\frac{27(1-\frac{5}{4})}{1+\frac{5}{4}}\right)^{\frac{p-1}{12}} J_p(0) \equiv (-1)^{\frac{p-1}{12}} 3^{-\frac{p-1}{4}} J_p(0) \pmod{p} & \text{if } p \equiv 1 \pmod{12}, \\ \left(\frac{9}{8}\right)^{\frac{p-1}{3}} \frac{108(2(\frac{5}{4})^2 - 14 \cdot \frac{5}{4} + 11)}{(\frac{5}{4} + 1)^2} \left(\frac{27(1-\frac{5}{4})}{1+\frac{5}{4}}\right)^{\frac{p-7}{12}} J_p(0) \\ &\equiv -8(-1)^{\frac{p-7}{12}} 3^{-\frac{p-7}{4}} J_p(0) \pmod{p} & \text{if } p \equiv 7 \pmod{12}. \end{split}$$

By the definition of $J_p(x)$, we have

$$J_{p}(0) = 1728^{\left[\frac{p}{12}\right]} \cdot 2^{-\left[\frac{p}{12}\right]} \sum_{r=0}^{\left[\frac{p}{12}\right]} {\left[\frac{p}{12}\right] - \frac{1}{3} {\left(\frac{p}{3}\right) \choose 3}} {\left[\frac{p}{12}\right] - \frac{1}{2} {\left(\frac{-1}{p}\right) \choose p}} 0^{\left[\frac{p}{12}\right] - r} 2^{r}$$
$$= 1728^{\left[\frac{p}{12}\right]} {\left[\frac{p}{12}\right] - \frac{1}{3} {\left(\frac{p}{3}\right) \choose 3}} = (-1728)^{\left[\frac{p}{12}\right]} {\left(\frac{1}{3} {\left(\frac{p}{3}\right) - 1 \choose p}\right)}.$$

Hence

$$J_p(0) \equiv (-1728)^{\left[\frac{p}{12}\right]} {\left(\frac{2(p-1)}{3}\right) \pmod{p}}$$

and therefore

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5}{4}\right) \equiv \left\{ \begin{array}{l} (-1)^{\frac{p-1}{12}} 3^{-\frac{p-1}{4}} (-1728)^{\frac{p-1}{12}} {2(p-1) \choose \frac{p-1}{12}} \right) \equiv \left(\frac{2}{p}\right) {2(p-1) \choose \frac{p-1}{12}} \pmod{p} & \text{if } 12 \mid p-1, \\ -8(-1)^{\frac{p-7}{12}} 3^{-\frac{p-7}{4}} (-1728)^{\frac{p-7}{12}} {2(p-1) \choose \frac{p-7}{12}} \right) \equiv -\left(\frac{2}{p}\right) {2(p-1) \choose \frac{p-7}{12}} \pmod{p} & \text{if } 12 \mid p-7. \end{array} \right.$$

Now putting all the above together we obtain the result.

Remark 3.1 For any prime p > 3, Zhi-Wei Sun conjectured that ([Su1, Conjecture A46])

$$\sum_{k=0}^{p-1} \frac{(3k)!}{24^k \cdot k!^3} \equiv \left(\frac{p}{3}\right) \sum_{k=0}^{p-1} \frac{(3k)!}{(-216)^k \cdot k!^3} \equiv \begin{cases} \binom{2(p-1)/3}{(p-1)/3} \pmod{p^2} & \text{if } p \equiv 1 \pmod{3}, \\ 0 \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

4. Congruences for $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k}$. Let p > 3 be a prime and $m \in \mathbb{Z}$ with $p \nmid m$. In the section we partially solve Z.W. Sun's conjectures on $\sum_{k=0}^{p-1} {2k \choose k}^2 {3k \choose k} m^{-k} \pmod{p^2}$.

Theorem 4.1. Let p > 3 be a prime, $m \in \mathbb{Z}_p$, $m \not\equiv 0 \pmod{p}$ and $t = \sqrt{1 - 108/m}$. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k} \equiv P_{\left[\frac{p}{3}\right]}(t)^2 \equiv \left(\sum_{x=0}^{p-1} (x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11))^{\frac{p-1}{2}}\right)^2 \pmod{p}.$$

Moreover, if $P_{\left[\frac{p}{3}\right]}(t) \equiv 0 \pmod{p}$ or $\sum_{x=0}^{p-1} (x^3 + 3(4t - 5)x + 2(2t^2 - 14t + 11))^{\frac{p-1}{2}} \equiv 0 \pmod{p}$, then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k} \equiv 0 \pmod{p^2}.$$

Proof. Since $\frac{1-t}{54}(1-27\cdot\frac{1-t}{54})=\frac{1}{m}$, by Theorem 2.1 we have

(4.1)
$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{m^k} \equiv \left(\sum_{k=0}^{p-1} \binom{2k}{k} \binom{3k}{k} \left(\frac{1-t}{54}\right)^k\right)^2 \pmod{p^2}.$$

Observe that $p \mid {2k \choose k} {3k \choose k}$ for $[\frac{p}{3}] < k < p$. From the above and Corollary 3.1 we see that

$$\begin{split} &\sum_{k=0}^{p-1} \binom{2k}{k} \binom{3k}{k} \left(\frac{1-t}{54}\right)^k \\ &\equiv P_{\left[\frac{p}{3}\right]}(t) \equiv -\left(\frac{p}{3}\right) \sum_{x=0}^{p-1} (x^3 + 3(4t-5)x + 2(2t^2 - 14t + 11))^{\frac{p-1}{2}} \pmod{p}. \end{split}$$

This together with (4.1) yields the result.

Theorem 4.2 ([Su1, Conjecture A8]). Let p > 3 be a prime. Then

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-192)^k} \equiv \left\{ \begin{array}{ll} L^2 \pmod{p} & \textit{if } p \equiv 1 \pmod{3} \ \textit{and so } 4p = L^2 + 27M^2, \\ 0 \pmod{p^2} & \textit{if } p \equiv 2 \pmod{3}. \end{array} \right.$$

Proof. Putting m=-192 and $t=\frac{5}{4}$ in Theorem 4.1 and then applying Theorem 3.2 we obtain the result.

Lemma 4.1. Let p be an odd prime and let a, m, n be p-adic integers. Then

$$\sum_{x=0}^{p-1} (x^3 + a^2 mx + a^3 n)^{\frac{p-1}{2}} \equiv a^{\frac{p-1}{2}} \sum_{x=0}^{p-1} (x^3 + mx + n)^{\frac{p-1}{2}} \pmod{p}.$$

Moreover, if a, m, n are congruent to some integers, then

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + a^2 m x + a^3 n}{p} \right) = \left(\frac{a}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 + m x + n}{p} \right).$$

Proof. For any positive integer k it is well known that (see [IR])

$$\sum_{k=0}^{p-1} x^k \equiv \begin{cases} p-1 \pmod{p} & \text{if } p-1 \mid k, \\ 0 \pmod{p} & \text{if } p-1 \nmid k. \end{cases}$$

Since

$$\begin{split} &\sum_{x=0}^{p-1} (x^3 + a^2 m x + a^3 n)^{\frac{p-1}{2}} \\ &= \sum_{x=0}^{p-1} \sum_{k=0}^{(p-1)/2} \binom{(p-1)/2}{k} (x^3 + a^2 m x)^k (a^3 n)^{\frac{p-1}{2} - k} \\ &= \sum_{x=0}^{p-1} \sum_{k=0}^{(p-1)/2} \binom{(p-1)/2}{k} \sum_{r=0}^k \binom{k}{r} x^{3r} (a^2 m x)^{k-r} (a^3 n)^{\frac{p-1}{2} - k} \\ &= \sum_{r=0}^{(p-1)/2} \sum_{k=r}^{(p-1)/2} \binom{(p-1)/2}{k} \binom{k}{r} (a^2 m)^{k-r} (a^3 n)^{\frac{p-1}{2} - k} \sum_{x=0}^{p-1} x^{k+2r} \\ &\equiv (p-1) \sum_{r=0}^{(p-1)/2} \binom{(p-1)/2}{p-1-2r} \binom{p-1-2r}{r} (a^2 m)^{p-1-3r} (a^3 n)^{2r-\frac{p-1}{2}} \\ &= a^{\frac{p-1}{2}} (p-1) \sum_{\frac{p-1}{4} \le r \le \frac{p-1}{3}} \binom{(p-1)/2}{p-1-2r} \binom{p-1-2r}{r} m^{p-1-3r} n^{2r-\frac{p-1}{2}} \pmod{p}, \end{split}$$

we see that the congruence in Lemma 4.1 is true.

Now suppose that a, m, n are congruent to some integers. If $a \equiv 0 \pmod{p}$, clearly

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + a^2 m x + a^3 n}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{x^3}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{x}{p} \right) = 0 = \left(\frac{a}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 + m x + n}{p} \right).$$

If $a \not\equiv 0 \pmod{p}$, then clearly

$$\sum_{x=0}^{p-1} \left(\frac{x^3 + a^2 mx + a^3 n}{p} \right) = \sum_{x=0}^{p-1} \left(\frac{(ax)^3 + a^2 m(ax) + a^3 n}{p} \right) = \left(\frac{a}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 + mx + n}{p} \right).$$

Thus the lemma is proved.

Lemma 4.2. Let p be an odd prime. Then

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 30x - 56}{p} \right)$$

$$= \begin{cases} (-1)^{\left[\frac{p}{8}\right] + 1} \left(\frac{3}{p}\right) 2c & \text{if } p \equiv 1, 3 \pmod{8}, \ p = c^2 + 2d^2 \text{ and } 4 \mid c - 1, \\ 0 & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Proof. From [BE, Theorems 5.12 and 5.17] we know that

$$\sum_{k=0}^{p-1} \left(\frac{x^3 - 4x^2 + 2x}{p} \right) = \begin{cases} (-1)^{\left[\frac{p}{8}\right] + 1} 2c & \text{if } p = c^2 + 2d^2 \equiv 1, 3 \pmod{8} \text{ with } 4 \mid c - 1, \\ 0 & \text{otherwise.} \end{cases}$$

As
$$27(x^3 - 4x^2 + 2x) = (3x - 4)^3 - 30(3x - 4) - 56$$
, we see that

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 4x^2 + 2x}{p} \right)$$

$$= \left(\frac{3}{p} \right) \sum_{x=0}^{p-1} \left(\frac{(3x - 4)^3 - 30(3x - 4) - 56}{p} \right) = \left(\frac{3}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 30x - 56}{p} \right).$$

Thus the result follows.

Lemma 4.3. Let p be an odd prime. Then

$$\begin{split} &\sum_{n=0}^{p-1} (n^3 - (15 + 30\sqrt{-2})n - 28 + 70\sqrt{-2}))^{\frac{p-1}{2}} \\ &\equiv \begin{cases} (\frac{2+\sqrt{-2}}{p})(-1)^{\left[\frac{p}{8}\right]+1}(\frac{3}{p})2c \pmod{p} & \textit{if } p = c^2 + 2d^2 \equiv 1, 3 \pmod{8} \ \textit{and } 4 \mid c - 1, \\ 0 \pmod{p} & \textit{if } p \equiv 5, 7 \pmod{8}. \end{cases} \end{split}$$

Proof. It is easily seen that

$$-15(1+2\sqrt{-2}) = -30\left(\frac{1-\sqrt{-2}}{\sqrt{-2}}\right)^2 \quad \text{and} \quad -28+70\sqrt{-2} = -56\left(\frac{1-\sqrt{-2}}{\sqrt{-2}}\right)^3.$$

Thus, by Lemmas 4.1 and 4.2 we have

$$\sum_{n=0}^{p-1} \left(n^3 - (15 + 30\sqrt{-2})n - 28 + 70\sqrt{-2} \right)^{\frac{p-1}{2}}$$

$$\equiv \left(\frac{1 - \sqrt{-2}}{\sqrt{-2}} \right)^{\frac{p-1}{2}} \sum_{n=0}^{p-1} (n^3 - 30n - 56)^{\frac{p-1}{2}}$$

$$\equiv \left(-\frac{2 + \sqrt{-2}}{2} \right)^{\frac{p-1}{2}} \sum_{n=0}^{p-1} \left(\frac{n^3 - 30n - 56}{p} \right)$$

$$\equiv \begin{cases} \left(\frac{2 + \sqrt{-2}}{p} \right) (-1)^{\left[\frac{p}{8}\right] + 1} \left(\frac{3}{p} \right) 2c \pmod{p} & \text{if } p = c^2 + 2d^2 \equiv 1, 3 \pmod{8} \text{ and } 4 \mid c - 1, \\ 0 \pmod{p} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

This proves the lemma.

Theorem 4.3. Let p be an odd prime. Then

$$P_{\left[\frac{p}{3}\right]}(5/\sqrt{-2})$$

$$\equiv \begin{cases} (-1)^{\left[\frac{p}{8}\right]}(\frac{-2-\sqrt{-2}}{p})2c \pmod{p} & \text{if } p = c^2 + 2d^2 \equiv 1, 3 \pmod{8} \text{ and } 4 \mid c - 1, \\ 0 \pmod{p} & \text{if } p \equiv 5, 7 \pmod{8} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{8^k} \equiv \begin{cases} 4c^2 \pmod{p} & \text{if } p = c^2 + 2d^2 \equiv 1, 3 \pmod{8}, \\ 0 \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Proof. From Corollary 3.1 and Lemma 4.3 we deduce that

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5}{\sqrt{-2}}\right) \equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} (n^3 + 3(-10\sqrt{-2} - 5)n - 28 + 70\sqrt{-2})^{\frac{p-1}{2}}$$

$$\equiv \begin{cases} \left(\frac{p}{3}\right) \left(\frac{2+\sqrt{-2}}{p}\right) (-1)^{\left[\frac{p}{8}\right]} \left(\frac{3}{p}\right) 2c = \left(\frac{-2-\sqrt{-2}}{p}\right) (-1)^{\left[\frac{p}{8}\right]} 2c \pmod{p} \\ \text{if } p = c^2 + 2d^2 \equiv 1, 3 \pmod{8} \text{ and } 4 \mid c - 1, \\ 0 \pmod{p} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Now taking m=8 and $t=5/\sqrt{-2}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Remark 4.1 Let p be an odd prime. Zhi-Wei Sun conjectured that ([Su1, Conjecture A5])

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{8^k} \equiv \begin{cases} 4c^2 - 2p \pmod{p^2} & \text{if } p = c^2 + 2d^2 \equiv 1, 3 \pmod{8}, \\ 0 \pmod{p^2} & \text{if } p \equiv 5, 7 \pmod{8}. \end{cases}$$

Corollary 4.1. Let p be a prime such that $p \equiv 1, 3 \pmod{8}$ and $p = c^2 + 2d^2$ with $c \equiv 1 \pmod{4}$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5d}{c}\right) \equiv \left\{ \begin{array}{ll} (\frac{c-d}{3})2c \pmod{p} & \text{if } p \equiv 1 \pmod{8}, \\ (-1)^{\frac{d-1}{2}}(\frac{c-d}{3})2c \pmod{p} & \text{if } p \equiv 3 \pmod{8}. \end{array} \right.$$

Proof. It is clear that $(\frac{d}{p}) = (\frac{p}{d}) = (\frac{c^2}{d}) = 1$. Thus

$$\begin{split} \left(\frac{2+c/d}{p}\right) &= \left(\frac{d}{p}\right) \left(\frac{c+2d}{p}\right) = \left(\frac{c+2d}{p}\right) = (-1)^{\frac{p-1}{2} \cdot \frac{2d+c-1}{2}} \left(\frac{p}{c+2d}\right) \\ &= (-1)^{\frac{p-1}{2}d} \left(\frac{(c+2d)(c-2d)+6d^2}{c+2d}\right) = (-1)^{\frac{p-1}{2}d} \left(\frac{6}{c+2d}\right) \\ &= (-1)^{\frac{p-1}{2}d+\frac{(c+2d)^2-1}{8}} \left(\frac{3}{c+2d}\right) = (-1)^{\frac{p-1+4d+2d^2}{8}} \cdot (-1)^{\frac{p-1}{2}d+\frac{c+2d-1}{2}} \left(\frac{c+2d}{3}\right) \\ &= (-1)^{\frac{p-1+4d+2d^2}{8}} \left(\frac{c-d}{3}\right) = \begin{cases} (-1)^{\frac{p-1}{8}} \left(\frac{c-d}{3}\right) & \text{if } p \equiv 1 \pmod{8}, \\ (-1)^{\frac{p-3}{8}+\frac{d+1}{2}} \left(\frac{c-d}{3}\right) & \text{if } p \equiv 3 \pmod{8}. \end{cases} \end{split}$$

By Theorem 4.3 we have

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5}{c/d}\right) \equiv (-1)^{\left[\frac{p}{8}\right]} \left(\frac{-1}{p}\right) \left(\frac{2+c/d}{p}\right) 2c \pmod{p}.$$

Now combining all the above we obtain the result.

Lemma 4.4. Let p be an odd prime and $p \neq 11$. Then

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 24 \cdot 11x + 14 \cdot 11^2}{p} \right) = \begin{cases} \left(\frac{u}{11} \right) u & \text{if } \left(\frac{p}{11} \right) = 1 \text{ and so } 4p = u^2 + 11v^2, \\ 0 & \text{if } \left(\frac{p}{11} \right) = -1. \end{cases}$$

Proof. It is known that (see [PR] and [JM])

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 96 \cdot 11x + 112 \cdot 11^2}{p} \right) = \begin{cases} \left(\frac{2}{p} \right) \left(\frac{u}{11} \right) u & \text{if } \left(\frac{p}{11} \right) = 1 \text{ and } 4p = u^2 + 11v^2, \\ 0 & \text{if } \left(\frac{p}{11} \right) = -1. \end{cases}$$

Since

$$\sum_{x=0}^{p-1} \left(\frac{x^3 - 96 \cdot 11x + 112 \cdot 11^2}{p} \right)$$

$$= \sum_{x=0}^{p-1} \left(\frac{(2x)^3 - 96 \cdot 11 \cdot 2x + 112 \cdot 11^2}{p} \right) = \left(\frac{2}{p} \right) \sum_{x=0}^{p-1} \left(\frac{x^3 - 24 \cdot 11x + 14 \cdot 11^2}{p} \right),$$

we deduce the result.

Lemma 4.5. Let $p \neq 11$ be an odd prime. Then

$$\sum_{n=0}^{p-1} (n^3 + 12(-5 + \sqrt{-11})n + 14(11 - 4\sqrt{-11}))^{\frac{p-1}{2}}$$

$$\equiv \begin{cases} (\frac{-22 + 2\sqrt{-11}}{p})(\frac{u}{11})u \pmod{p} & \text{if } (\frac{p}{11}) = 1 \text{ and so } 4p = u^2 + 11v^2, \\ 0 \pmod{p} & \text{if } (\frac{p}{11}) = -1. \end{cases}$$

Proof. It is easily seen that

$$12(-5+\sqrt{-11}) = -24 \cdot 11 \left(\frac{\sqrt{-11}+1}{2\sqrt{-11}}\right)^2 \text{ and } 14(11-4\sqrt{-11}) = 14 \cdot 11^2 \left(\frac{\sqrt{-11}+1}{2\sqrt{-11}}\right)^3.$$

Thus, by Lemma 4.1 we have

$$\sum_{n=0}^{p-1} \left(n^3 + 12(-5 + \sqrt{-11})n + 14(11 - 4\sqrt{-11}) \right)^{\frac{p-1}{2}}$$

$$\equiv \left(\frac{\sqrt{-11} + 1}{2\sqrt{-11}} \right)^{\frac{p-1}{2}} \sum_{x=0}^{p-1} \left(x^3 - 24 \cdot 11x + 14 \cdot 11^2 \right)^{\frac{p-1}{2}}$$

$$\equiv \left(\frac{-22 + 2\sqrt{-11}}{-11 \cdot 4} \right)^{\frac{p-1}{2}} \sum_{x=0}^{p-1} \left(\frac{x^3 - 24 \cdot 11x + 14 \cdot 11^2}{p} \right) \pmod{p}.$$

Now applying Lemma 4.4 we deduce the result.

Theorem 4.4. Let $p \neq 11$ be an odd prime. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{\sqrt{-11}}{4}\right) \equiv \begin{cases} -(\frac{p}{3})(\frac{-11+\sqrt{-11}}{p})(\frac{u}{11})u \pmod{p} & \text{if } (\frac{p}{11}) = 1 \text{ and so } 4p = u^2 + 11v^2, \\ 0 \pmod{p} & \text{if } (\frac{p}{11}) = -1 \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k} \equiv \begin{cases} u^2 \pmod{p} & \text{if } (\frac{p}{11}) = 1 \text{ and so } 4p = u^2 + 11v^2, \\ 0 \pmod{p^2} & \text{if } (\frac{p}{11}) = -1. \end{cases}$$

Proof. From Corollary 3.1 and Lemma 4.5 we deduce that

$$\begin{split} P_{\left[\frac{p}{3}\right]}\left(\frac{\sqrt{-11}}{4}\right) &\equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(n^3 + 3(\sqrt{-11} - 5)n + \frac{-11}{4} + 22 - 7\sqrt{-11}\right)^{\frac{p-1}{2}} \\ &\equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\left(\frac{n}{2}\right)^3 + 3(\sqrt{-11} - 5)\frac{n}{2} + \frac{77 - 28\sqrt{-11}}{4}\right)^{\frac{p-1}{2}} \\ &\equiv -\left(\frac{p}{3}\right) \left(\frac{2}{p}\right) \sum_{n=0}^{p-1} \left(n^3 + 12(-5 + \sqrt{-11})n + 14(11 - 4\sqrt{-11})\right)^{\frac{p-1}{2}} \\ &\equiv \begin{cases} -\left(\frac{p}{3}\right)\left(\frac{-11 + \sqrt{-11}}{p}\right)\left(\frac{u}{11}\right)u \pmod{p} & \text{if } \left(\frac{p}{11}\right) = 1 \text{ and so } 4p = u^2 + 11v^2, \\ 0 \pmod{p} & \text{if } \left(\frac{p}{11}\right) = -1. \end{cases} \end{split}$$

Now taking m = 64 and $t = \frac{\sqrt{-11}}{4}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Remark 4.2 Let p be an odd prime such that $p \neq 11$. Zhi-Wei Sun conjectured that ([Su1, Conjecture A4])

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{64^k} \equiv \begin{cases} u^2 - 2p \pmod{p^2} & \text{if } (\frac{p}{11}) = 1 \text{ and so } 4p = u^2 + 11v^2, \\ 0 \pmod{p^2} & \text{if } (\frac{p}{11}) = -1. \end{cases}$$

Corollary 4.2. Let p be an odd prime such that $(\frac{p}{11}) = 1$ and so $4p = u^2 + 11v^2$ with $u, v \in \mathbb{Z}$. Let $v = 2^{\alpha}v_0$ and $u - 11v = 2^{\beta}w$ with $2 \nmid v_0w$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{u}{4v}\right) \equiv (-1)^{\frac{p-1}{2} \cdot \frac{v_0 - w}{2}} \left(\frac{2}{p}\right)^{\alpha + \beta} \left(\frac{v - u}{3}\right) u \pmod{p}.$$

Proof. It is clear that

Thus, by Theorem 4.4 and the above we get

$$P_{\left[\frac{p}{3}\right]}\left(\frac{u}{4v}\right) \equiv -\left(\frac{p}{3}\right) \left(\frac{-11 + u/v}{p}\right) \left(\frac{u}{11}\right) u \equiv -\left(\frac{u^2 + 11v^2}{3}\right) \left(\frac{2}{p}\right)^{\alpha + \beta} (-1)^{\frac{v_0 - w}{2} \cdot \frac{p - 1}{2}} \left(\frac{u + v}{3}\right) u$$

$$= \left(\frac{2}{p}\right)^{\alpha + \beta} (-1)^{\frac{v_0 - w}{2} \cdot \frac{p - 1}{2}} \left(\frac{v - u}{3}\right) u \pmod{p}.$$

This proves the corollary.

Let p > 3 be a prime and let \mathbb{F}_p be the field of p elements. For $m, n \in \mathbb{F}_p$ let $\#E_p(x^3 + mx + n)$ be the number of points on the curve $E: y^2 = x^3 + mx + n$ over the field \mathbb{F}_p . It is well known that (see for example [S1, pp.221-222])

(4.2)
$$#E_p(x^3 + mx + n) = p + 1 + \sum_{x=0}^{p-1} \left(\frac{x^3 + mx + n}{p} \right).$$

Let $K = \mathbb{Q}(\sqrt{-d})$ be an imaginary quadratic field and the curve $y^2 = x^3 + mx + n$ has complex multiplication by K. By Deuring's theorem ([C, Theorem 14.16],[PV],[I]), we have

(4.3)
$$#E_p(x^3 + mx + n) = \begin{cases} p+1 & \text{if } p \text{ is inert in } K, \\ p+1-\pi-\bar{\pi} & \text{if } p = \pi\bar{\pi} \text{ in } K, \end{cases}$$

where π is in an order in K and $\bar{\pi}$ is the conjugate number of π . If $4p = u^2 + dv^2$ with $u, v \in \mathbb{Z}$, we may take $\pi = \frac{1}{2}(u + v\sqrt{-d})$. Thus,

(4.4)
$$\sum_{x=0}^{p-1} \left(\frac{x^3 + mx + n}{p} \right) = \begin{cases} \pm u & \text{if } 4p = u^2 + dv^2 \text{ with } u, v \in \mathbb{Z}, \\ 0 & \text{otherwise.} \end{cases}$$

In [Gr], [JM] and [PV] the sign of u in (4.4) was determined for those imaginary quadratic fields K with class number 1. In [LM] and [I] the sign of u in (4.4) was determined for imaginary quadratic fields K with class number 2. For general results on the sign of u in (4.4), see [M], [St], [RS] and the survey [Ri].

Lemma 4.6. Let p be a prime with $p \equiv \pm 1 \pmod{8}$. Then

$$\sum_{n=0}^{p-1} \left(\frac{n^3 + (-15 + 6\sqrt{2})n + 24 - 14\sqrt{2}}{p} \right)$$

$$= \begin{cases} 2x(\frac{2x}{3}) & \text{if } p \equiv 1,7 \pmod{24} \text{ and so } p = x^2 + 6y^2, \\ 0 & \text{if } p \equiv 17,23 \pmod{24}. \end{cases}$$

Proof. It is easy to check the result for p=7. Now assume $p \ge 17$. From [I, p.133] we know that the elliptic curve defined by the equation $y^2 = x^3 + (-21 + 12\sqrt{2})x - 28 + 22\sqrt{2}$ has complex multiplication by the order of discriminant -24. Since $4p = u^2 + 24v^2$ implies $2 \mid u$ and $p = (\frac{u}{2})^2 + 6v^2$, by (4.4) and [I, Theorem 3.1] we have

$$\sum_{n=0}^{p-1} \left(\frac{n^3 + (-21 + 12\sqrt{2})n - 28 + 22\sqrt{2}}{p} \right)$$

$$= \begin{cases} 2x(\frac{2x}{3})(\frac{1+\sqrt{2}}{p}) & \text{if } p \equiv 1,7 \pmod{24} \text{ and so } p = x^2 + 6y^2, \\ 0 & \text{if } p \equiv 17,23 \pmod{24}. \end{cases}$$

Observe that

$$\frac{-15 - 6\sqrt{2}}{-21 + 12\sqrt{2}} = (1 + \sqrt{2})^2 \quad \text{and} \quad \frac{24 + 14\sqrt{2}}{-28 + 22\sqrt{2}} = (1 + \sqrt{2})^3.$$

Using Corollary 3.2 and Lemma 4.1 we see that

$$\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (-15 + 6\sqrt{2})n + 24 - 14\sqrt{2}}{p}\right)$$

$$= \sum_{n=0}^{p-1} \left(\frac{n^3 - (15 + 6\sqrt{2})n + 24 + 14\sqrt{2}}{p}\right)$$

$$= \left(\frac{1 + \sqrt{2}}{p}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (-21 + 12\sqrt{2})n - 28 + 22\sqrt{2}}{p}\right).$$

Now putting all the above together we obtain the result.

Theorem 4.5. Let p be a prime such that $p \equiv 1, 7 \pmod{8}$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{\sqrt{2}}{2}\right) \equiv \begin{cases} 2x(\frac{x}{3}) \pmod{p} & \text{if } p = x^2 + 6y^2 \equiv 1,7 \pmod{24}, \\ 0 \pmod{p} & \text{if } p \equiv 17,23 \pmod{24} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k} \equiv \begin{cases} 4x^2 \pmod{p} & \text{if } p = x^2 + 6y^2 \equiv 1,7 \pmod{24}, \\ 0 \pmod{p^2} & \text{if } p \equiv 17,23 \pmod{24}. \end{cases}$$

Proof. From Theorem 3.1 and Lemma 4.6 we deduce that

$$P_{\left[\frac{p}{3}\right]}\left(\frac{\sqrt{2}}{2}\right) \equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (-15 + 6\sqrt{2})n + 24 - 14\sqrt{2}}{p}\right)$$

$$\equiv \begin{cases} -2x(\frac{2x}{3}) = 2x(\frac{x}{3}) \pmod{p} & \text{if } p = x^2 + 6y^2 \equiv 1,7 \pmod{24}, \\ 0 \pmod{p} & \text{if } p \equiv 17,23 \pmod{24}. \end{cases}$$

Taking m=216 and $t=\frac{\sqrt{2}}{2}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Remark 4.3 For any prime p > 3, Z.W. Sun conjectured that ([Su1, A14])

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{216^k} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p = x^2 + 6y^2 \equiv 1,7 \pmod{24}, \\ 8x^2 - 2p \pmod{p^2} & \text{if } p = 2x^2 + 3y^2 \equiv 5,11 \pmod{24}, \\ 0 \pmod{p^2} & \text{if } p \equiv 13,17,19,23 \pmod{24}. \end{cases}$$

Conjecture 4.1. Let p be a prime such that $p \equiv 5, 11, 13, 19 \pmod{24}$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{\sqrt{2}}{2}\right) \equiv \begin{cases} 0 \pmod{p} & \text{if } p \equiv 13, 19 \pmod{24}, \\ -2x(\frac{x}{3})\sqrt{2} \pmod{p} & \text{if } p = 2x^2 + 3y^2 \equiv 5, 11 \pmod{24}. \end{cases}$$

Lemma 4.7. Let p be a prime with $p \equiv \pm 1 \pmod{5}$. Then

$$\begin{split} &\sum_{n=0}^{p-1} \left(\frac{n^3 + (-15 + 12\sqrt{5})n + 42 - 28\sqrt{5}}{p} \right) \\ &= \begin{cases} 2x(\frac{2x}{3}) & \text{if } p \equiv 1,4 \text{ (mod 15) } \text{ and so } p = x^2 + 15y^2, \\ 0 & \text{if } p \equiv 11,14 \text{ (mod 15)}. \end{cases} \end{split}$$

Proof. From [I, Proposition 3.3] we know that the elliptic curve defined by the equation $y^2 = x^3 + (105 + 48\sqrt{5})x - 784 - 350\sqrt{5}$ has complex multiplication by the order of discriminant -15. Since $4p = u^2 + 60v^2$ implies $2 \mid u$ and $p = (\frac{u}{2})^2 + 15v^2$, by (4.4) and [I, Theorem 3.1] we have

$$\begin{split} &\sum_{n=0}^{p-1} \Big(\frac{n^3 + (105 + 48\sqrt{5})n - 784 - 350\sqrt{5}}{p}\Big) \\ &= \left\{ \begin{array}{l} 2x(\frac{2x}{3})(\frac{(1+\sqrt{5})/2}{p}) & \text{if } p \equiv 1,4 \text{ (mod 15) and so } p = x^2 + 15y^2, \\ 0 & \text{if } p \equiv 11,14 \text{ (mod 15)}. \end{array} \right. \end{split}$$

Observe that

$$\frac{-15 + 12\sqrt{5}}{105 + 48\sqrt{5}} = (\sqrt{5} - 2)^2 \quad \text{and} \quad \frac{42 - 28\sqrt{5}}{-784 - 350\sqrt{5}} = (\sqrt{5} - 2)^3.$$

Using Lemma 4.1 we see that

$$\sum_{n=0}^{p-1} \left(\frac{n^3 + (-15 + 12\sqrt{5})n + 42 - 28\sqrt{5}}{p} \right)$$

$$= \left(\frac{\sqrt{5} - 2}{p} \right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (105 + 48\sqrt{5})n - 784 - 350\sqrt{5}}{p} \right).$$

Note that $(\sqrt{5}-2)\frac{1+\sqrt{5}}{2}=(\frac{1-\sqrt{5}}{2})^2$. We then have $(\frac{\sqrt{5}-2}{p})=(\frac{(1+\sqrt{5})/2}{p})$. Now putting all the above together we obtain the result.

Theorem 4.6. Let p be a prime such that $p \equiv 1, 4 \pmod{5}$. Then

$$P_{[\frac{p}{3}]}(\sqrt{5}) \equiv \begin{cases} 2x(\frac{x}{3}) \pmod{p} & \text{if } p = x^2 + 15y^2 \equiv 1,4 \pmod{15}, \\ 0 \pmod{p} & \text{if } p \equiv 11,14 \pmod{15} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-27)^k} \equiv \begin{cases} 4x^2 \pmod{p} & \text{if } p = x^2 + 15y^2 \equiv 1, 4 \pmod{15}, \\ 0 \pmod{p^2} & \text{if } p \equiv 11, 14 \pmod{15}. \end{cases}$$

Proof. From Theorem 3.1 and Lemma 4.7 we deduce that

$$P_{\left[\frac{p}{3}\right]}(\sqrt{5}) \equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (-15 + 12\sqrt{5})n + 42 - 28\sqrt{5}}{p}\right)$$

$$\equiv \begin{cases} -2x(\frac{2x}{3}) = 2x(\frac{x}{3}) \pmod{p} & \text{if } p = x^2 + 15y^2 \equiv 1, 4 \pmod{15}, \\ 0 \pmod{p} & \text{if } p \equiv 11, 14 \pmod{15}. \end{cases}$$

Taking m=-27 and $t=\sqrt{5}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Conjecture 4.2. Let p be an odd prime such that $p \equiv 2, 7, 8, 13 \pmod{15}$. Then

$$P_{[\frac{p}{3}]}(\sqrt{5}) \equiv \left\{ \begin{array}{ll} 0 \ (\text{mod} \ p) & \mbox{ if } p \equiv 7,13 \ (\text{mod} \ 15), \\ 2x(\frac{x}{3})\sqrt{5} \ (\text{mod} \ p) & \mbox{ if } p = 5x^2 + 3y^2 \equiv 2,8 \ (\text{mod} \ 15). \end{array} \right.$$

Lemma 4.8. Let p be a prime such that $p \equiv \pm 1 \pmod{5}$. Then

$$\begin{split} &\sum_{n=0}^{p-1} \Big(\frac{n^3 + (-300 + 108\sqrt{5})n - 2520 + 1042\sqrt{5}}{p} \Big) \\ &= \left\{ \begin{array}{ll} (\frac{2\sqrt{5}}{p})(\frac{x}{3})x & \textit{if } p \equiv 1,4 \pmod{15} \ \textit{and so } 4p = x^2 + 75y^2, \\ 0 & \textit{if } p \equiv 11,14 \pmod{15}. \end{array} \right. \end{split}$$

Proof. From [I, p.134] we know that the elliptic curve defined by the equation $y^2 = x^3 + (-2160 + 408\sqrt{5})x + 42130 - 10472\sqrt{5}$ has complex multiplication by the order of discriminant -75. By (4.4) and [I, Theorem 3.1] we have

$$\sum_{n=0}^{p-1} \left(\frac{n^3 + (-2160 + 408\sqrt{5})n + 42130 - 10472\sqrt{5}}{p} \right)$$

$$= \begin{cases} \left(\frac{-25 - 13\sqrt{5}}{p} \right) \left(\frac{x}{3} \right) x & \text{if } p \equiv 1, 4 \pmod{15} \text{ and so } 4p = x^2 + 75y^2, \\ 0 & \text{if } p \equiv 11, 14 \pmod{15}. \end{cases}$$

Observe that

$$\frac{-2160 + 408\sqrt{5}}{-300 + 108\sqrt{5}} = \left(-\frac{7 + \sqrt{5}}{2}\right)^2 \quad \text{and} \quad \frac{42130 - 10472\sqrt{5}}{-2520 + 1042\sqrt{5}} = \left(-\frac{7 + \sqrt{5}}{2}\right)^3.$$

Using Lemma 4.1 we see that

$$\sum_{n=0}^{p-1} \left(\frac{n^3 + (-2160 + 408\sqrt{5})n + 42130 - 10472\sqrt{5}}{p} \right)$$

$$= \left(\frac{-(7 + \sqrt{5})/2}{p} \right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (-300 + 108\sqrt{5})n - 2520 + 1042\sqrt{5}}{p} \right).$$

Since

$$2(-7-\sqrt{5})(-25-13\sqrt{5}) = 120+58\sqrt{5} = 2\sqrt{5}(29+12\sqrt{5}) = 2\sqrt{5}(3+2\sqrt{5})^2$$

from the above we deduce the result.

Theorem 4.7. Let p be a prime such that $p \equiv 1, 4 \pmod{5}$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{9}{20}\sqrt{5}\right) \equiv \begin{cases} -(\frac{x}{3})x \pmod{p} & \text{if } p \equiv 1,4 \pmod{15} \text{ and so } 4p = x^2 + 75y^2, \\ 0 \pmod{p} & \text{if } p \equiv 11,14 \pmod{15} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-8640)^k} \equiv \begin{cases} x^2 \pmod{p} & \text{if } p \equiv 1, 4 \pmod{15} \text{ and so } 4p = x^2 + 75y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 11, 14 \pmod{15}. \end{cases}$$

Proof. From Theorem 3.1 and Lemma 4.8 we deduce that

$$\begin{split} P_{\left[\frac{p}{3}\right]}\left(\frac{9}{20}\sqrt{5}\right) &\equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + \frac{27 - 15\sqrt{5}}{\sqrt{5}}n + \frac{521 - 252\sqrt{5}}{20}}{p}\right) \\ &= -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{\left(\frac{n}{2\sqrt{5}}\right)^3 + \frac{27 - 15\sqrt{5}}{\sqrt{5}} \cdot \frac{n}{2\sqrt{5}} + \frac{521 - 252\sqrt{5}}{20}}{p}\right) \\ &= -\left(\frac{p}{3}\right) \left(\frac{2\sqrt{5}}{p}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + \left(-300 + 108\sqrt{5}\right)n - 2520 + 1042\sqrt{5}}{p}\right) \\ &\equiv \left\{ \begin{array}{l} -\left(\frac{x}{3}\right)x \pmod{p} & \text{if } p \equiv 1, 4 \pmod{15} \text{ and so } 4p = x^2 + 75y^2, \\ 0 \pmod{p} & \text{if } p \equiv 11, 14 \pmod{15}. \end{array} \right. \end{split}$$

Taking m = -8640 and $t = \frac{9}{4\sqrt{5}}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Remark 4.4 In [S2] the author conjectured that for any prime p > 5,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-8640)^k} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } 3 \mid p-1, \ p = x^2 + 3y^2 \text{ and } 5 \mid xy, \\ p - 2x^2 \pm 6xy \pmod{p^2} & \text{if } 3 \mid p-1, \ p = x^2 + 3y^2, \ 5 \nmid xy \\ & \text{and } x \equiv \pm y, \pm 2y \pmod{5}, \\ 0 \pmod{p^2} & \text{if } 3 \mid p-2. \end{cases}$$

This is equivalent to

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-8640)^k} \equiv \begin{cases} x^2 - 2p \pmod{p^2} & \text{if } p \equiv 1, 4 \pmod{15} \text{ and so } 4p = x^2 + 75y^2, \\ 2p - 3x^2 \pmod{p^2} & \text{if } p \equiv 7, 13 \pmod{15} \text{ and so } 4p = 3x^2 + 25y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Conjecture 4.3. For any prime p > 5 we have

$$\sum_{k=0}^{p-1} \frac{9k+1}{(-8640)^k} {2k \choose k}^2 {3k \choose k} \equiv p\left(\frac{p}{15}\right) \pmod{p^3}.$$

Conjecture 4.4. Let p be a prime such that $p \equiv 7, 13, 17, 23 \pmod{30}$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{9}{20}\sqrt{5}\right) \equiv \begin{cases} 5y(\frac{y}{3}) \pmod{p} & \textit{if } p \equiv 7,13 \pmod{30} \ \textit{and so } 4p = 3x^2 + 25y^2, \\ 0 \pmod{p} & \textit{if } p \equiv 17,23 \pmod{30}. \end{cases}$$

Let $b \in \{17, 41, 89\}$ and $f(b) = -12^3, -48^3, -300^3$ according as b = 17, 41, 89. In [Su1, Conjectures A20, A22 and A23], Z.W. Sun conjectured that for any odd prime $p \neq 3, b$, (4.5)

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{f(b)^k} \equiv \begin{cases} x^2 - 2p \pmod{p^2} & \text{if } (\frac{p}{3}) = (\frac{p}{b}) = 1 \text{ and so } 4p = x^2 + 3by^2, \\ 2p - 3x^2 \pmod{p^2} & \text{if } (\frac{p}{3}) = (\frac{p}{b}) = -1 \text{ and so } 4p = 3x^2 + by^2, \\ 0 \pmod{p^2} & \text{if } (\frac{p}{3}) = -(\frac{p}{b}). \end{cases}$$

Now we partially solve (4.5).

Theorem 4.8. Let p be an odd prime such that $(\frac{17}{p}) = 1$. Then

$$P_{[\frac{p}{3}]}\Big(\frac{\sqrt{17}}{4}\Big) \equiv \left\{ \begin{array}{ll} -(\frac{x}{3})x \pmod{p} & \textit{if } p \equiv 1 \pmod{3} \ \textit{and so } 4p = x^2 + 51y^2, \\ 0 \pmod{p} & \textit{if } p \equiv 2 \pmod{3}. \end{array} \right.$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-12)^{3k}} \equiv \begin{cases} x^2 \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 51y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Proof. From [I, p.134] we know that the elliptic curve defined by the equation $y^2 = x^3 - (60 + 12\sqrt{17})x - 210 - 56\sqrt{17}$ has complex multiplication by the order of discriminant -51. Thus, by (4.4) and [I, Theorem 3.1] we have

$$\begin{split} &\sum_{n=0}^{p-1} \Big(\frac{n^3 - (60 + 12\sqrt{17})n - 210 - 56\sqrt{17}}{p} \Big) \\ &= \left\{ \begin{array}{ll} (\frac{-2}{p})(\frac{x}{3})x & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 51y^2, \\ 0 & \text{if } p \equiv 2 \pmod{3}. \end{array} \right. \end{split}$$

It then follows from (1.3) and Theorem 3.1 that

$$\begin{split} P_{\left[\frac{p}{3}\right]}\left(\frac{\sqrt{17}}{4}\right) &= \left(\frac{p}{3}\right)P_{\left[\frac{p}{3}\right]}\left(-\frac{\sqrt{17}}{4}\right) \equiv -\sum_{n=0}^{p-1}\left(\frac{n^3 + 3(-\sqrt{17} - 5)n + \frac{17}{4} + 22 + 7\sqrt{17}}{p}\right) \\ &= -\sum_{n=0}^{p-1}\left(\frac{(-\frac{n}{2})^3 - 3(5 + \sqrt{17})(-\frac{n}{2}) + \frac{105 + 28\sqrt{17}}{4}}{p}\right) \\ &= -\left(\frac{-2}{p}\right)\sum_{n=0}^{p-1}\left(\frac{n^3 - (60 + 12\sqrt{17})n - 210 - 56\sqrt{17}}{p}\right) \\ &\equiv \begin{cases} -(\frac{x}{3})x \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 51y^2, \\ 0 \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{split}$$

Taking $m = -12^3$ and $t = \frac{\sqrt{17}}{4}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Conjecture 4.5. Let p be an odd prime such that $(\frac{p}{17}) = -1$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{\sqrt{17}}{4}\right) \equiv \begin{cases} 0 \pmod{p} & \text{if } p \equiv 1 \pmod{3}, \\ -(\frac{y}{3})y\sqrt{17} \pmod{p} & \text{if } p \equiv 2 \pmod{3} \text{ and so } 4p = 3x^2 + 17y^2. \end{cases}$$

Theorem 4.9. Let p be an odd prime such that $(\frac{41}{p}) = 1$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5\sqrt{41}}{32}\right) \equiv \begin{cases} -(\frac{x}{3})x \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 123y^2, \\ 0 \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-48)^{3k}} \equiv \left\{ \begin{array}{ll} x^2 \pmod{p} & \textit{if } p \equiv 1 \pmod{3} \ \textit{and so } 4p = x^2 + 123y^2, \\ 0 \pmod{p^2} & \textit{if } p \equiv 2 \pmod{3}. \end{array} \right.$$

Proof. From [I,p.134] we know that the elliptic curve defined by the equation $y^2 = x^3 + (-960 + 120\sqrt{41})x - 13314 + 2240\sqrt{41}$ has complex multiplication by the order of discriminant -123. Thus, by [I, Theorem 3.1] we have

$$\begin{split} &\sum_{n=0}^{p-1} \left(\frac{n^3 - (960 - 120\sqrt{41})n - 13314 + 2240\sqrt{41}}{p} \right) \\ &= \left\{ \begin{array}{ll} \left(\frac{-2}{p} \right) (\frac{x}{3})x & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 123y^2, \\ 0 & \text{if } p \equiv 2 \pmod{3}. \end{array} \right. \end{split}$$

It then follows from (1.3) and Theorem 3.1 that

$$\begin{split} P_{\left[\frac{p}{3}\right]}\Big(\frac{5\sqrt{41}}{32}\Big) &\equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + 3(\frac{5}{8}\sqrt{41} - 5)n + (\frac{5\sqrt{41}}{16})^2 - 7 \cdot \frac{5\sqrt{41}}{8} + 22}{p}\right) \\ &= -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{(-\frac{n}{8})^3 + \frac{15}{8}(\sqrt{41} - 8)(-\frac{n}{8}) + \frac{6657 - 1120\sqrt{41}}{256}}{p}\right) \\ &= -\left(\frac{p}{3}\right) \left(\frac{-2}{p}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (960 - 120\sqrt{41})n - 13314 + 2240\sqrt{41}}{p}\right) \\ &\equiv \left\{\begin{array}{ll} -(\frac{x}{3})x \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 51y^2, \\ 0 \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{array}\right. \end{split}$$

Taking $m = -48^3$ and $t = \frac{5\sqrt{41}}{32}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Conjecture 4.6. Let p be an odd prime such that $(\frac{p}{41}) = -1$. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5\sqrt{41}}{32}\right) \equiv \left\{ \begin{array}{ll} 0 \; (\text{mod } p) & \text{if } p \equiv 1 \; (\text{mod } 3), \\ -(\frac{y}{3})y\sqrt{41} \; (\text{mod } p) & \text{if } p \equiv 2 \; (\text{mod } 3) \; \text{and so } 4p = 3x^2 + 41y^2. \end{array} \right.$$

Theorem 4.10. Let p > 5 be a prime such that $(\frac{89}{p}) = 1$. Then

$$P_{[\frac{p}{3}]}\Big(\frac{53\sqrt{89}}{500}\Big) \equiv \left\{ \begin{array}{ll} -(\frac{x}{3})x \; (\bmod \; p) & \textit{if } p \equiv 1 \; (\bmod \; 3) \; \textit{and so } 4p = x^2 + 267y^2, \\ 0 \; (\bmod \; p) & \textit{if } p \equiv 2 \; (\bmod \; 3). \end{array} \right.$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{(-300)^{3k}} \equiv \begin{cases} x^2 \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 267y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Proof. From [I, p.135] we know that the elliptic curve defined by the equation $y^2 =$ $x^{3} + (-37500 + 3180\sqrt{89})x + 3250002 - 371000\sqrt{89}$ has complex multiplication by the order of discriminant -267. Thus, by [I, Theorem 3.1] we have

$$\begin{split} &\sum_{n=0}^{p-1} \left(\frac{n^3 + (-37500 + 3180\sqrt{89})n + 3250002 - 371000\sqrt{89}}{p} \right) \\ &= \left\{ \begin{array}{ll} (\frac{2}{p})(\frac{x}{3})x & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 267y^2, \\ 0 & \text{if } p \equiv 2 \pmod{3}. \end{array} \right. \end{split}$$

It then follows from (1.3) and Theorem 3.1 that

$$\begin{split} P_{\left[\frac{p}{3}\right]}\Big(\frac{53\sqrt{89}}{500}\Big) &\equiv -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + 3(4 \cdot \frac{53}{500}\sqrt{89} - 5)n + 4(\frac{53}{500}\sqrt{89})^2 - 28 \cdot \frac{53\sqrt{89}}{500} + 22}{p}\right) \\ &= -\left(\frac{p}{3}\right) \sum_{n=0}^{p-1} \left(\frac{(\frac{n}{50})^3 + \frac{-1875 + 159\sqrt{89}}{125} \cdot \frac{n}{50} + \frac{1625001 - 185500\sqrt{89}}{250^2}\right) \\ &= -\left(\frac{p}{3}\right) \left(\frac{50}{p}\right) \sum_{n=0}^{p-1} \left(\frac{n^3 + (-37500 + 3180\sqrt{89})n + 3250002 - 371000\sqrt{89}}{p}\right) \\ &\equiv \left\{ \begin{array}{ll} -(\frac{x}{3})x \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } 4p = x^2 + 51y^2, \\ 0 \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{array} \right. \end{split}$$

Taking $m = -300^3$ and $t = \frac{53\sqrt{89}}{500}$ in Theorem 4.1 and then applying the above we deduce the remaining result.

Conjecture 4.7. Let p be an odd prime such that $(\frac{p}{89}) = -1$. Then

$$P_{\left[\frac{p}{3}\right]}\Big(\frac{53\sqrt{89}}{500}\Big) \equiv \left\{ \begin{array}{ll} 0 \ (\text{mod } p) & \text{if } p \equiv 1 \ (\text{mod } 3), \\ -(\frac{y}{3})y\sqrt{89} \ (\text{mod } p) & \text{if } p \equiv 2 \ (\text{mod } 3) \ and so \ 4p = 3x^2 + 89y^2. \end{array} \right.$$

In the end we pose the following conjectures.

Conjecture 4.8. Let p be a prime with $p \equiv \pm 1 \pmod{5}$. Then

$$\sum_{n=0}^{p-1} \left(\frac{n^3 + 3(-125 + 44\sqrt{5})n + 154(21 - 10\sqrt{5}}{p} \right)$$

$$= \begin{cases} 2x(\frac{2x}{3}) & \text{if } p \equiv 1, 4 \pmod{15} \text{ and so } p = x^2 + 15y^2, \\ 0 & \text{if } p \equiv 11, 14 \pmod{15}. \end{cases}$$

If Conjecture 4.8 is true, we may discuss the following conjecture in [S2]:

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{15^{3k}} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p \equiv 1, 4 \pmod{15} \text{ and so } p = x^2 + 15y^2, \\ 2p - 12x^2 \pmod{p^2} & \text{if } p \equiv 2, 8 \pmod{15} \text{ and so } p = 3x^2 + 5y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 7, 11, 13, 14 \pmod{15}. \end{cases}$$

Conjecture 4.9. For any prime p > 5 we have

$$\sum_{k=0}^{p-1} \frac{33k+4}{15^{3k}} {2k \choose k}^2 {3k \choose k} \equiv 4p \left(\frac{p}{3}\right) \pmod{p^3}.$$

Conjecture 4.10. Let p > 5 be a prime. Then

$$P_{\left[\frac{p}{3}\right]}\!\left(\frac{11\sqrt{5}}{25}\right) \equiv \left\{ \begin{array}{ll} 2x(\frac{x}{3}) \pmod{p} & \text{if } p \equiv 1,19 \pmod{30} \text{ and so } p = x^2 + 15y^2, \\ -2y(\frac{y}{3})\sqrt{5} \pmod{p} & \text{if } p \equiv 17,23 \pmod{30} \text{ and so } p = 3x^2 + 5y^2, \\ 0 \pmod{p} & \text{if } p \equiv 7,11,13,29 \pmod{30}. \end{array} \right.$$

In [S2], the author also conjectured that for any prime p > 3,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{3k}{k}}{1458^k} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p \equiv 1 \pmod{3} \text{ and so } p = x^2 + 3y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Conjecture 4.11. For any prime p > 3 we have

$$\sum_{k=0}^{p-1} \frac{15k+2}{1458^k} {2k \choose k}^2 {3k \choose k} \equiv (-1)^{\frac{p-1}{2}} 2p \pmod{p^3}.$$

Conjecture 4.12. Let p > 3 be a prime. Then

$$P_{\left[\frac{p}{3}\right]}\left(\frac{5\sqrt{3}}{9}\right) \equiv \begin{cases} (-1)^{\frac{p-1}{2}} 2x(\frac{x}{3}) \pmod{p} & \text{if } p \equiv 1 \pmod{3} \text{ and so } p = x^2 + 3y^2, \\ 0 \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

References

- [BE] B. C. Berndt and R. J. Evans, Sums of Gauss Eisenstein, Jacobi, Jacobsthal and Brewer, Illinois J. Math. 23 (1979), 374-437.
- [BEW] B.C. Berndt, R.J. Evans and K.S. Williams, *Gauss and Jacobi Sums*, John Wiley & Sons, New York, 1998.
- [BM] J. Brillhart and P. Morton, Class numbers of quadratic fields, Hasse invariants of elliptic curves, and the supersingular polynomial, J. Number Theory 106 (2004), 79-111.
- [C] D.A. Cox, Primes of the Form $x^2 + ny^2$: Fermat, Class Field Theory, and Complex Multiplication, Wiley, New York, 1989.
- [G] H.W. Gould, Combinatorial Identities, A Standardized Set of Tables Listing 500 Binomial Coefficient Summations, Morgantown, W. Va., 1972.
- [Gr] B.H. Gross, Minimal models for elliptic curves with complex multiplication, Compositio Math. 45 (1982), 155-164.
- [IR] K. Ireland and M. Rosen, A Classical Introduction to Modern Number Theory (2nd edition), Springer, New York, 1990, pp. 235-236.
- [I] N. Ishii, Trace of Frobenius endomorphism of an elliptic curve with complex multiplication, Bull. Austral. Math. Soc. **70** (2004), 125-142.
- [JM] A. Joux et F. Morain, Sur les sommes de caractères liées aux courbes elliptiques à multiplication complexe, J. Number Theory **55** (1995), 108-128.
- [LM] F. Leprévost and F. Morain, Revêtements de courbes elliptiques à multiplication complexe par des courbes hyperelliptiques et sommes de caractères, J. Number Theory 64 (1997), 165-182.
- [MOS] W. Magnus, F. Oberhettinger and R.P. Soni, Formulas and Theorems for the Special Functions of Mathematical Physics, 3rd. ed., Springer, New York, 1966, pp. 228-232.
- [M] F. Morain, Computing the cardinality of CM elliptic curves using torsion points, J. Théor. Nombres Bordeaux 19 (2007), 663-681.

- [Mo] E. Mortenson, Supercongruences for truncated $_{n+1}F_n$ hypergeometric series with applications to certain weight three newforms, Proc. Amer. Math. Soc. 133(2005), 321-330.
- [Mor] P. Morton, Explicit identities for invariants of elliptic curves, J. Number Theory 120 (2006), 234-271.
- [PV] R. Padma and S. Venkataraman, Elliptic curves with complex multiplication and a character sum, J. Number Theory **61** (1996), 274-282.
- [PR] J.C. Parnami and A.R. Rajwade, A new cubic character sum, Acta Arith. 40 (1982), 347-356.
- [RV] F. Rodriguez-Villegas, Hypergeometric families of Calabi-Yau manifolds. Calabi-Yau Varieties and Mirror Symmetry (Yui, Noriko (ed.) et al., Toronto, ON, 2001), 223-231, Fields Inst. Commun., 38, Amer. Math. Soc., Providence, RI, 2003.
- [RS] K. Rubin and A. Silverberg, *Point counting on reductions of CM elliptic curves*, J. Number Theory 129(2009), 2903-2923.
- [Si] A. Silverberg, Group order formulas for reductions of CM elliptic curves, in Proceedings of the Conference on Arithmetic, Geometry, Cryptography and Coding Theory, Contemporary Mathematics, 521, American Mathematical Society, Providence, RI, 2010, 107-120.
- [St] H.M. Stark, Counting points on CM elliptic curves, Rocky Mountain J. Math. 26 (1996), 1115-1138.
- [S1] Z.H. Sun, On the number of incongruent residues of $x^4 + ax^2 + bx$ modulo p, J. Number Theory 119 (2006), 210-241.
- [S2] Z.H. Sun, Congruences concerning Legendre polynomials, Proc. Amer. Math. Soc. 139 (2011), 1915-1929.
- [Su1] Z.W. Sun, Open conjectures on congruences, arXiv:0911.5665. http://arxiv.org/abs/0911.5665.
- [Su2] Z.W. Sun, On sums involving products of three binomial coefficients, preprint, arXiv:1012.3141. http://arxiv.org/abs/1012.3141.