

QUASI-UNIFORMITY OF MINIMAL WEIGHTED ENERGY POINTS ON COMPACT METRIC SPACES

D. P. HARDIN, E. B. SAFF, AND J. T. WHITEHOUSE

ABSTRACT. For a closed subset K of a compact metric space A possessing an α -regular measure μ with $\mu(K) > 0$, we prove that whenever $s > \alpha$, any sequence of weighted minimal Riesz s -energy configurations $\omega_N = \{x_{i,N}^{(s)}\}_{i=1}^N$ on K (for ‘nice’ weights) is quasi-uniform in the sense that the ratios of its mesh norm to separation distance remain bounded as N grows large. Furthermore, if K is an α -rectifiable compact subset of Euclidean space (α an integer) with positive and finite α -dimensional Hausdorff measure, it is possible to generate such a quasi-uniform sequence of configurations that also has (as $N \rightarrow \infty$) a prescribed positive continuous limit distribution with respect to α -dimensional Hausdorff measure.

1. INTRODUCTION

Let A be a compact infinite metric space with metric $d: A \times A \rightarrow [0, \infty)$ and let $\omega_N = \{x_i\}_{i=1}^N \subset A$ denote a configuration of $N \geq 2$ points in A . We are chiefly concerned with two ‘quality’ measures of ω_N ; namely, the *separation distance* of ω_N defined by

$$(1.1) \quad \delta(\omega_N) := \min_{1 \leq i \neq j \leq N} d(x_i, x_j),$$

and the *mesh norm* of ω_N with respect to A defined by

$$(1.2) \quad \rho(\omega_N, A) := \max_{y \in A} \min_{1 \leq i \leq N} d(y, x_i).$$

This quantity is also known as the *fill radius* or *covering radius* of ω_N relative to A . The optimal values of these quantities are also of interest and we consider, for $N \geq 2$, the *N -point best-packing distance on A* given by

$$\delta_N(A) := \max\{\delta(\omega_N) : \omega_N \subset A, |\omega_N| = N\},$$

and the *N -point mesh norm of A* given by

$$\rho_N(A) := \min\{\rho(\omega_N, A) : \omega_N \subset A, |\omega_N| = N\},$$

where $|S|$ denotes the cardinality of set S .

In the theory of approximation and interpolation (for example, by splines or radial basis functions (RBFs)), the separation distance is often associated with some

Date: May 26, 2018.

2000 Mathematics Subject Classification. Primary: 31C20, 65N50, 57N16; Secondary: 52A40, 28A78 .

Key words and phrases. Fill radius, Mesh-separation ratio, Best-packing, Optimal configurations, Covering radius, Minimal Riesz energy, Quasi-uniformity, Separation distance.

The research of all authors was supported, in part, by the U. S. National Science Foundation under grants DMS-0808093 and DMS-1109266.

measure of ‘stability’ of the approximation, while the mesh norm arises in the error of the approximation. In this context, the *mesh-separation ratio* (or *mesh ratio*)

$$\gamma(\omega_N, A) := \rho(\omega_N, A)/\delta(\omega_N),$$

can be regarded as a ‘condition number’ for ω_N relative to A . If $\{\omega_N\}_{N=2}^\infty$ is a sequence of N -point configurations such that $\gamma(\omega_N, A)$ is uniformly bounded in N , then the sequence is said to be *quasi-uniform on A* . Quasi-uniform sequences of configurations are important for a number of methods involving RBF approximation and interpolation (see [9, 15, 17, 19]).

We remark that in some cases it is easy to obtain positive lower bounds for the mesh-separation ratio. For example, if A is connected, then $\gamma(\omega_N, A) \geq 1/2$. Furthermore, letting

$$B(x, r) = \{y \in A : m(y, x) \leq r\}$$

be the closed ball in A with center x and radius r , then $\gamma(\omega_N, A) \geq \beta/2$ for any N -point configuration $\omega_N \subset A$ whenever A and $\beta \in (0, 1)$ have the property that for any $r \in (0, \text{diam}(A)]$ and any $x \in A$, the annulus $B(x, r) \setminus B(x, \beta r)$ is nonempty. The diameter of A is defined by

$$\text{diam}(A) := \max\{m(x, y) : x \in A, y \in A\}.$$

In this paper we consider the separation distance and mesh norm of finite point configurations in A that minimize certain weighted energy functionals. We call $w : A \times A \rightarrow [0, \infty)$ an *SLP weight on A* if it is symmetric and lower semi-continuous on $A \times A$ and is positive on the diagonal, $D(A)$, of $A \times A$. For $s > 0$ and a collection of $N \geq 2$ distinct points $\omega_N = \{x_1, \dots, x_N\} \subset A$, the (s, w) -energy of ω_N (also known as the *weighted Riesz s -energy*) is

$$(1.3) \quad E_s^w(\omega_N) := \sum_{\substack{i \neq j \\ j \neq i}} \frac{w(x_i, x_j)}{(x_i, x_j)^s} = \sum_{i=1}^N \sum_{\substack{j=1 \\ j \neq i}}^N \frac{w(x_i, x_j)}{(x_i, x_j)^s},$$

and we denote the *minimal N -point (s, w) -energy of A* by

$$(1.4) \quad \mathcal{E}_s^w(N, A) := \inf\{E_s^w(\omega_N) : \omega_N \subset A, |\omega_N| = N\}.$$

Since A is compact and the energy $E_s^w(\omega_N)$ is lower semi-continuous, there exists at least one N -point configuration $\omega_N^* \subset A$ such that $E_s^w(\omega_N^*) = \mathcal{E}_s^w(N, A)$. We refer to such an ω_N^* as an *N -point (s, w) -energy minimizing configuration on A* . The asymptotics as $N \rightarrow \infty$ of N -point (s, w) -energy minimizing configurations and their energies are investigated in [2, 10] for d -rectifiable sets $A \subset \mathbb{R}^p$ and $s > d$ (see further discussion in the next section).

In our results we shall require that A is either α -regular or upper α -regular as we next describe. For a positive Borel measure μ supported on A and $\alpha > 0$, we say that μ is *upper α -regular* if there is some finite constant C_0 such that

$$(1.5) \quad \mu(B(x, r)) \leq C_0 r^\alpha \quad (x \in A, 0 < r \leq \text{diam}(A)),$$

and we say that μ is *lower α -regular* if there is some positive constant c_0 such that

$$(1.6) \quad c_0^{-1} r^\alpha \leq \mu(B(x, r)) \quad (x \in A, 0 < r \leq \text{diam}(A)).$$

We shall refer to A as an *upper α -regular metric space* if there exists an upper α -regular measure $\bar{\mu}$ on A such that $\bar{\mu}(A) > 0$ and shall refer to A as a *lower α -regular*

metric space if there exists a lower α -regular measure $\underline{\mu}$ on A such that $\underline{\mu}(A) < \infty$. (Obviously, if A is upper α -regular then A has infinitely many points.) If A supports a measure that is both upper and lower α -regular, then we say that A is an α -regular *metric space*. If A is α -regular, then it is not difficult to show that the Hausdorff dimension of A , $\dim_{\mathcal{H}} A$, equals α (cf. [12, 16]). Furthermore, the α -dimensional Hausdorff measure of A , $\mathcal{H}_\alpha(A)$, is positive and finite.

Many of the constants appearing in this paper, either explicitly or implicitly involve the upper and lower regularity constants C_0 and c_0 appearing in (1.5) and (1.6). However, in certain cases we are interested in ‘local’ regularity estimates (i.e., for r small) which can substantially improve our explicit estimates for particular metric spaces of interest (e.g., A is the sphere S^d with the Euclidean metric). Specifically, if $\bar{\mu}$ is an upper α -regular measure, $\underline{\mu}$ is a lower α -regular measure and $r^* > 0$, we define

$$(1.7) \quad \begin{aligned} C_0(r^*) &:= \sup\{\bar{\mu}(B(x, r))/r^\alpha : x \in A, 0 < r \leq r^*\}, \\ c_0(r^*)^{-1} &:= \inf\{\underline{\mu}(B(x, r))/r^\alpha : x \in A, 0 < r \leq r^*\}. \end{aligned}$$

We note that both $C_0(r^*)$ and $c_0(r^*)$ are increasing in r^* , and we make the definitions

$$(1.8) \quad \begin{aligned} C_0(0) &:= \lim_{r^* \rightarrow 0^+} C_0(r^*), \\ c_0(0) &:= \lim_{r^* \rightarrow 0^+} c_0(r^*). \end{aligned}$$

Furthermore, if A is a compact (i.e., without boundary), C^1 , d -dimensional manifold and $\mu = \mathcal{H}_d$, then $C_0(0) \cdot c_0(0) = 1$. For the largest length scale of interest, with a slight abuse of notation, the global constants for $\bar{\mu}$ and $\underline{\mu}$, respectively, are related by $C_0 = C_0(\text{diam}(A))$ and $c_0 = c_0(\text{diam}(A))$.

One may obtain simple upper bounds for $\delta_N(A)$ (respectively, lower bounds for $\rho_N(A)$) in the case that A is lower (respectively, upper) α -regular. Specifically, if A is lower α -regular then there is a constant $c_A < \infty$ such that

$$(1.9) \quad \delta_N(A) \leq c_A N^{-1/\alpha}, \quad (N \geq 2),$$

while if A is upper α -regular then there is a constant $\tilde{c}_A > 0$ such that

$$(1.10) \quad \rho_N(A) \geq \tilde{c}_A N^{-1/\alpha}, \quad (N \geq 2).$$

The bound (1.9) is a consequence of the facts that the balls $\{B(x, \delta(\omega_N)/2) : x \in \omega_N\}$ are pairwise disjoint and that there exists a lower α -regular measure $\underline{\mu}$ with $\underline{\mu}(A) < \infty$. Similarly, if A is upper α -regular, then the bound (1.10) follows from the covering property of the balls $\{B(x, \rho(\omega_N, A)) : x \in \omega_N\}$ and the existence of an upper α -regular measure $\bar{\mu}$ with $\bar{\mu}(A) > 0$.

The main result of this paper, given in Theorem 5, is that a sequence of N -point (s, w) -energy minimizing configurations on an α -regular compact metric space A is quasi-uniform on A whenever $s > \alpha$. As an application, we deduce that, if $A \subset \mathbb{R}^p$ is d -rectifiable for some integer $0 < d \leq p$ with $\mathcal{H}_d(A) > 0$, then a quasi-uniform sequence of N -point configurations on A can be found that has a prescribed bounded positive density on A (see Corollary 6 and the discussion preceding it).

2. MAIN RESULTS

We first consider the separation distance of (s, w) -energy minimizing configurations on an upper α -regular compact metric space A . For these separation results, we consider symmetric weight functions w such that $\|w(\cdot, x)\|_{L_p(\mu)}$ is uniformly bounded on A for some $1 < p \leq \infty$. Here we use the standard notation,

$$\|f\|_{L_p(\mu)} := \begin{cases} (\int_A |f|^p d\mu)^{1/p}, & 1 \leq p < \infty, \\ \mu\text{-ess sup } |f|, & p = \infty, \end{cases}$$

where μ is a positive Borel measure and f is a Borel measurable function on A .

The following theorem extends a result [2, Theorem 4] to a more general class of weight functions and to more general compact metric spaces.

Theorem 1. *Let A be a compact, upper α -regular metric space with respect to $\bar{\mu}$ and let w be an SLP weight on A such that $\|w(\cdot, x)\|_{L_{p_0}(\bar{\mu})}$ is uniformly bounded on A for some $1 < p_0 \leq \infty$. Suppose $1 < p \leq p_0$, $s > \alpha(1 - 1/p)$, and $N \geq 2$. If ω_N^* is an N -point (s, w) -energy minimizing configuration on A , then*

$$(2.1) \quad \delta(\omega_N^*) \geq C_1 N^{-\left(\frac{1}{\alpha} + \frac{1}{sp}\right)} \quad (N \geq 2),$$

where C_1 is a constant independent of N indicated below in (3.13).

Taking w bounded and setting $p = \infty$ in Theorem 1 produces the following result.

Corollary 2. *Suppose A is a compact, upper α -regular metric space and w is a bounded SLP weight on A , and let $s > \alpha$. If ω_N^* is an N -point (s, w) -energy minimizing configuration on A , then*

$$(2.2) \quad \delta(\omega_N^*) \geq C_2 N^{-1/\alpha} \quad (N \geq 2),$$

where C_2 is a constant independent of N . Consequently,

$$(2.3) \quad \delta_N(A) \geq C_2 N^{-1/\alpha} \quad (N \geq 2).$$

For the unweighted case $w \equiv 1$, the constant C_2 satisfies

$$(2.4) \quad C_2 \geq \left[\frac{\bar{\mu}(A)}{C_0} \left(1 - \frac{\alpha}{s}\right) \right]^{1/\alpha} \left(\frac{\alpha}{s}\right)^{1/s},$$

where $C_0 = C_0(\text{diam}(A))$.

We note that if A in Corollary 2 is α -regular, then by inequality (1.9) we see that N -point (s, w) -energy minimizing configurations on A have the best possible order of separation as $N \rightarrow \infty$.

With respect to the separation constant of (2.4), if $d \geq 2$ and $A = \mathbb{S}^d$ with σ_d denoting the uniform probability distribution on \mathbb{S}^d , then we can get an explicit lower bound for C_2 by calculating the regularity constant C_0 . As stated in [13], for $x \in \mathbb{S}^d$, $0 \leq r \leq 2$, and

$$(2.5) \quad \gamma_d := \frac{\Gamma\left(\frac{d+1}{2}\right)}{\Gamma(d/2)\Gamma(1/2)},$$

there holds

$$\sigma_d(r) := \sigma_d(B(x, r)) = \gamma_d \int_{1-r^2/2}^1 (1-t^2)^{d/2-1} dt$$

from which it follows that

$$\sigma_d(r) \leq \frac{\gamma_d}{d} r^d,$$

and, as $r \rightarrow 0^+$,

$$\sigma_d(r) = \frac{\gamma_d}{d} r^d + \mathcal{O}(r^{d+2}).$$

Therefore, for the uniform probability distribution on \mathbb{S}^d , the global upper regularity constant is

$$(2.6) \quad C_0 = \sup_{0 < r \leq 2} \frac{\sigma_d(r)}{r^d} = \frac{\gamma_d}{d},$$

and when applied to (2.4) we obtain

$$(2.7) \quad C_2 \geq \left(\frac{d}{\gamma_d}\right)^{1/d} \left(1 - \frac{d}{s}\right)^{1/d} \left(\frac{d}{s}\right)^{1/s}.$$

With this lower bound for C_2 , (2.2) becomes

$$(2.8) \quad \delta(\omega_N^*) \geq \left(\frac{d}{\gamma_d}\right)^{1/d} \left(1 - \frac{d}{s}\right)^{1/d} \left(\frac{d}{s}\right)^{1/s} N^{-1/d} \quad (N \geq 2, s > d),$$

and, on letting $s \rightarrow \infty$, we deduce for the N -point best-packing distance

$$\delta_N(\mathbb{S}^d) \geq \left(\frac{d}{\gamma_d}\right)^{1/d} N^{-1/d} \quad (N \geq 2, s > d).$$

A less explicit lower bound for the separation constant of minimal energy points for $s > d$ on \mathbb{S}^d was obtained in [13, Corollary 4].

We next consider the mesh norm of (s, w) -energy minimizing configurations on an α -regular compact metric space A . In this case we require that the weight function w be bounded.

Theorem 3. *Let A be a compact, α -regular metric space with respect to the measure μ and $K \subset A$ be a compact set of positive μ -measure. Let w be a bounded SLP weight on K . If $s > \alpha$ and ω_N^* is an N -point (s, w) -energy minimizing configuration on K , then*

$$(2.9) \quad \rho(\omega_N^*, K) \leq C_3 N^{-1/\alpha} \quad (N \geq 2),$$

where C_3 is a constant independent of N given below in (3.41).

Theorem 3 substantially extends a result of [6] that holds for unweighted energy minimizing point configurations when $K \subset \mathbb{R}^p$ is restricted to be the finite union of bi-Lipschitz images of compact sets in \mathbb{R}^d .

We remark that for K and A as in Theorem 3, the set K need not inherit the lower α -regularity of A . However, since $\mu(K) > 0$, we do have that K is an upper α -regular metric space and, consequently, there is a constant $\tilde{c}_K > 0$ such that (1.10) holds with A replaced by K . Hence, the inequality (2.9) has the best possible order with respect to N .

Taking $w \equiv 1$ in Theorem 3 immediately yields the following.

Corollary 4. *Let A be a compact, α -regular metric space with respect to the measure μ and let $K \subset A$ be a compact set of positive μ -measure. Then there exists a constant C_4 such that*

$$\rho_N(K) \leq C_4 N^{-1/\alpha} \quad (N \geq 2).$$

Combining Corollary 2 and Theorem 3 we obtain our main result.

Theorem 5. *Let A be a compact, α -regular metric space with respect to the measure μ and let $K \subset A$ be a compact set of positive μ -measure. Furthermore, let w be a bounded SLP weight on K , and for $s > \alpha$ and $N \geq 2$, let ω_N^* be an N -point (s, w) -energy minimizing configuration on K . Then $\{\omega_N^*\}_{N=2}^\infty$ is quasi-uniform on K .*

We remark that there are α -regular sets A and values of $s < \alpha$ for which (unweighted) $(s, 1)$ -energy minimizing configurations on A have a mesh-separation ratio that goes to ∞ with N . One such example given in [4] is a ‘washer’ A obtained by revolving a certain rectangle about an axis parallel to one of its sides, where it turns out that for $s < 1/3$, the support of the limit distribution of the $(s, 1)$ -energy minimizing configurations on A omits an open subset of A . Also, for the logarithmic energy which corresponds to $s = 0$, it is shown in [11] that, for $w \equiv 1$, the support of the limit distribution of the log-energy minimizing configurations on a torus in \mathbb{R}^3 is only supported on the positive curvature portion of the torus, so that the mesh-separation ratio for such configurations is again unbounded as $N \rightarrow \infty$. Examples also abound in one dimension. For the logarithmic energy, it is well-known [21, Sections 6.7 and 6.21] that for $A = [-1, 1]$ and $w \equiv 1$ the minimum energy points are zeros of Jacobi orthogonal polynomials (together with ± 1) that have separation distance of precise order $1/N^2$ and mesh norm of precise order $1/N$, so that the mesh-separation ratio grows like N .

One of our main motivations for considering weighted minimum energy configurations is that for a large class of sets A one can design a weight function w so that a sequence of N -point (s, w) -energy minimizing configurations have a specified limiting density on A as $N \rightarrow \infty$. The following result is a consequence of Theorem 5 and [2, Corollary 2]. Recall that a set in \mathbb{R}^p is d -rectifiable if it is the Lipschitz image of a bounded set in \mathbb{R}^d .

Corollary 6. *Let $d \leq p$ and $A \subset \mathbb{R}^p$ be a compact, infinite set that is d -rectifiable and lower d -regular with respect to \mathcal{H}_d for some integer d . Suppose σ is a probability density on A that is continuous almost everywhere with respect to \mathcal{H}_d and is bounded above and below by positive constants. Let $s > d$ and $w : A \times A \rightarrow [0, \infty)$ be given by*

$$(2.10) \quad w(x, y) := (\sigma(x)\sigma(y))^{-s/2d}.$$

For $N \geq 2$, let ω_N^ be an N -point (s, w) -energy minimizing configuration on A . Then $\{\omega_N^*\}_{N=2}^\infty$ is quasi-uniform on A and the sequence of normalized counting measures associated with the ω_N^* 's converges weak-star (as $N \rightarrow \infty$) to $\sigma d\mathcal{H}_d$.*

For A an infinite, compact, metric space and $s > 0$, let ω_N^s be an N -point $(s, 1)$ -energy minimizing configuration on A . Furthermore, let ν_N be a cluster point (in the product topology on A^N) of ω_N^s as $s \rightarrow \infty$. As we now show, ν_N must be an N -point best-packing configuration on A , that is, $\delta(\nu_N) = \delta_N(A)$. For this purpose, let $\tilde{\omega}_N$ be an N -point best-packing configuration on A . Then we have

$$\delta(\omega_N^s)^{-s} \leq \mathcal{E}_s^1(N, A) \leq E_s^1(\tilde{\omega}_N) \leq N(N-1)\delta_N(A)^{-s},$$

and so

$$(N(N-1))^{-1/s}\delta_N(A) \leq \delta(\omega_N^s) \leq \delta_N(A),$$

which gives

$$(2.11) \quad \lim_{s \rightarrow \infty} \delta(\omega_N^s) = \delta_N(A).$$

Since $\omega_N^{s_j} \rightarrow \nu_N$ for some subsequence $s_j \rightarrow \infty$, it follows from (2.11) and continuity that $\delta(\nu_N) = \delta_N(A)$ and so ν_N is an N -point best-packing configuration on A .

In general, it is not true that a sequence of N -point best-packing configurations in A is quasi-uniform on A (e.g., if A is the classical $(1/3)$ -Cantor set in $[0,1]$ together with any point outside this interval). However, for A as in Theorem 5, it turns out that by using $(s, 1)$ -energy minimizing configurations on A and taking $s \rightarrow \infty$ we can construct a sequence of N -point best-packing configurations in A that is also quasi-uniform on A .

Theorem 7. *Let A be a compact, α -regular metric space with respect to the measure μ and let $K \subset A$ be a compact set of positive μ -measure. For $N \geq 2$, let ν_N be a cluster point of a family of N -point $(s, 1)$ -energy minimizing configurations on K as $s \rightarrow \infty$. Then $\{\nu_N\}_{N=2}^\infty$ is a sequence of N -point best-packing configurations on K that is also quasi-uniform on K .*

Furthermore, the mesh-separation ratios satisfy

$$(2.12) \quad \limsup_{N \rightarrow \infty} \gamma(\nu_N, K) \leq 2 \left(\frac{\mu(A)}{\mu(K)} \right)^{1/\alpha} [c_0(0) C_0(0)]^{1/\alpha},$$

where $c_0(0)$ and $C_0(0)$ are given in (1.8) for the set A .*

We note that the constant on the right-hand side of (2.12) is at least 2 per (1.7) and (1.8). One can also establish an analogous result concerning the existence of quasi-uniform sequences of *weighted* best-packing configurations (cf. [3]). We leave this extension to the reader.

In comparison with (2.12), we remark that one can construct examples of metric spaces A having n -point best-packing configurations with arbitrarily large mesh-separation ratio.

We conclude this section with further references to related results. Separation theorems for the case $s \leq d = \dim_{\mathcal{H}}(A)$ have been established only for rather special sets and values of s . Dahlberg [5] proved that (unweighted) optimal $((p - 2), 1)$ -energy configurations ω_N^* on A are *well-separated* (i.e., they satisfy $\delta(\omega_N^*) \geq CN^{-1/d}$ for some positive constant C) if $A \subset \mathbb{R}^p$ ($p \geq 3$) is a smooth $d = p - 1$ dimensional closed surface in \mathbb{R}^p that separates \mathbb{R}^p into two components. For the critical value $s = d$ and A a d -rectifiable subset of a smooth d -dimensional manifold in \mathbb{R}^p , it is shown in [2] that the following weaker separation result holds

$$(2.13) \quad \delta(\omega_N^*) \geq C(N \log N)^{-1/d},$$

for some positive constant C .

For the case that $A = \mathbb{S}^d$, the d -dimensional unit sphere in \mathbb{R}^{d+1} , well-separation was proved in [14] for the range of values $d - 1 < s < d$ and further extended by Dragnev and Saff [8] to the range $d - 2 < s < d$ with explicit estimates for the

**Added in proof:* In the manuscript [1], the first two authors together with A. Bondarenko have recently proved under more general conditions that the right-hand side of (2.12) can be replaced by 1.

separation constant C . Well-separation for $s = d - 2$ and $d \geq 3$ was established in [6].

Thus, for the important case of $A = \mathbb{S}^2$ it is known that optimal s -energy configurations on \mathbb{S}^2 are well-separated for all nonnegative values of $s \neq 2$ (well-separatedness for $s = 0$ was established in [18]; see also [7]); for the critical value $s = 2$, the only known separation results are of the weak form given in (2.13).

Much less is known with regard to covering (mesh norm) theorems in the case that $s \leq d$ (see [20, Sec. 1.3]).

3. PROOFS

In the proofs we shall need that an SLP weight w is bounded below in a neighborhood of the diagonal $D(A)$. Indeed, the positivity and lower semi-continuity of w on $D(A)$ and the compactness of A imply that there are positive numbers η and κ such that

$$(3.1) \quad w(x, y) \geq \eta \quad (x, y \in A, m(x, y) \leq \kappa).$$

Proof of Theorem 1. The initial part of this argument proceeds as in [13]. Let $N \geq 2$ be fixed and let $\omega_N^* = \{x_1, \dots, x_N\} \subset A$ be a fixed (s, w) -energy minimizing configuration in A . For $x \in A$ and $1 \leq i \leq N$, let

$$U_i(x) := \sum_{\substack{j=1 \\ j \neq i}}^N \frac{w(x, x_j)}{(x, x_j)^s}.$$

Since ω_N^* is a minimizing configuration we have the lower bound

$$(3.2) \quad U_i(x_i) \leq U_i(x) \text{ for all } x \in A.$$

Fix $r_1 \leq \text{diam}(A)$ such that

$$(3.3) \quad \bar{\mu} \left(\bigcup_{j=1}^N B(x_j, r_1) \right) \geq \bar{\mu}(A).$$

The radius r_1 can clearly be chosen independent of N , for example $r_1 = \text{diam}(A)$, and we note for future reference that it suffices to take $r_1 > \rho(\omega_N^*, A)$. For the rest of this proof we fix $r_1 = \text{diam}(A)$.

Now let $0 < \theta < 1$ and define

$$(3.4) \quad r_0 := \left(\frac{\theta \bar{\mu}(A)}{N C_0(r_1)} \right)^{1/\alpha},$$

where $C_0(r_1) = C_0$ is the upper regularity constant of $\bar{\mu}$ as in (1.7). We note that $r_0 < r_1$ as can be seen from the fact that $\bar{\mu}(A) \leq C_0(r_1)r_1^\alpha$.

For $B(x, r_0, r_1) := B(x, r_1) \setminus B(x, r_0)$, let

$$D := \bigcup_{j=1}^N B(x_j, r_0, r_1).$$

Using the upper regularity of $\bar{\mu}$ and (3.3) we see that

$$\bar{\mu}(D) \geq \bar{\mu}(A) - \sum_{j=1}^N \bar{\mu}(B(x_j, r_0)) \geq (1 - \theta)\bar{\mu}(A) > 0,$$

and thus by inequality (3.2) we have

$$(3.5) \quad U_i(x_i) \leq \frac{1}{\bar{\mu}(D)} \int_D U_i(x) d\bar{\mu}(x) \leq \frac{1}{(1 - \theta)\bar{\mu}(A)} \sum_{\substack{j=1 \\ j \neq i}}^N \int_{B(x_j, r_0, r_1)} \frac{w(x, x_j)}{(x, x_j)^s} d\bar{\mu}(x).$$

Applying Hölder's inequality with $1/q = 1 - 1/p$ we obtain

$$(3.6) \quad U_i(x_i) \leq \frac{1}{(1 - \theta)\bar{\mu}(A)} \sum_{\substack{j=1 \\ j \neq i}}^N \|w(\cdot, x_j)\|_{L_p(\bar{\mu})} \left(\int_{B(x_j, r_0, r_1)} \frac{1}{(x, x_j)^{sq}} d\bar{\mu}(x) \right)^{1/q}.$$

Converting the integral on the right-hand side of (3.6) to the appropriate integral of the distribution function, and noting that $sq > \alpha$ by assumption, we have

$$(3.7) \quad \begin{aligned} \int_{B(x_j, r_0, r_1)} \frac{1}{(x, x_j)^{sq}} d\bar{\mu}(x) &= \int_0^\infty \bar{\mu}(\{x \in B(x_j, r_0, r_1) : m(x_j, x)^{-sq} > t\}) dt \\ &\leq \int_{r_1^{-sq}}^{r_0^{-sq}} \bar{\mu}(B(x_j, t^{-1/sq})) dt \\ &\leq \frac{C_0(r_1) sq}{sq - \alpha} r_0^{\alpha - sq} \\ &= \frac{C_0(r_1) sq}{sq - \alpha} \left(\frac{\theta \bar{\mu}(A)}{N C_0(r_1)} \right)^{1 - (sq)/\alpha}, \end{aligned}$$

which, combined with (3.6), gives

$$(3.8) \quad \begin{aligned} U_i(x_i) &\leq \frac{\|w\|_{p, \infty}}{(1 - \theta)\bar{\mu}(A)} \left(\frac{C_0(r_1) sq}{sq - \alpha} \right)^{1/q} (N - 1) \left(\frac{\theta \bar{\mu}(A)}{N C_0(r_1)} \right)^{1/q - s/\alpha} \\ &< \frac{1}{\bar{\mu}(A)} \left(\frac{C_0(r_1)}{\bar{\mu}(A)} \right)^{s/\alpha} \left(\frac{\|w\|_{p, \infty}}{(1 - \theta)\theta^{s/\alpha - 1/q}} \right) \left(\frac{sq \bar{\mu}(A)}{sq - \alpha} \right)^{1/q} N^{1/p + s/\alpha}, \end{aligned}$$

where $\|w\|_{p, \infty} := \sup_{x \in A} \|w(\cdot, x)\|_{L_p(\bar{\mu})} < \infty$.

Choosing

$$(3.9) \quad \theta_0 := \frac{sq - \alpha}{sq - \alpha + \alpha q} = \left(\frac{s}{\alpha} - \frac{1}{q} \right) \left(\frac{s}{\alpha} + \frac{1}{p} \right)^{-1} < 1,$$

which minimizes the right-hand side of (3.8) with respect to θ , we obtain

$$(3.10) \quad U_i(x_i) \leq c_1 N^{s/\alpha + 1/p},$$

where after a bit of arithmetic we have

$$(3.11) \quad c_1 := \|w\|_{p, \infty} \left(\frac{C_0(r_1)}{\bar{\mu}(A)} \frac{s/\alpha + 1/p}{s/\alpha - 1/q} \right)^{s/\alpha} \left(\frac{s/\alpha + 1/p}{\bar{\mu}(A)} \right)^{1/p} (s/\alpha)^{1/q}.$$

Next, select the indices $1 \leq i_s \neq j_s \leq N$ so that $\delta(\omega_N^*) = (x_{i_s}, x_{j_s})$ and let κ and η be as in (3.1). If $\delta(\omega_N^*) \leq \kappa$, then

$$(3.12) \quad \frac{\eta}{\delta(\omega_N^*)^s} \leq \frac{w(x_{i_s}, x_{j_s})}{(x_{i_s}, x_{j_s})^s} \leq U_{i_s}(x_{i_s}) \leq c_1 N^{s/\alpha+1/p},$$

and therefore

$$\delta(\omega_N^*) \geq \left(\frac{\eta}{c_1} \right)^{1/s} N^{-\frac{1}{\alpha} - \frac{1}{sp}}.$$

Hence, (2.1) holds with

$$(3.13) \quad C_1 := \min\{\kappa, (\eta/c_1)^{1/s}\}.$$

□

We remark that for the case when $w \equiv 1$ and $p = \infty$, we can take $\kappa = \infty$, $\eta = 1$, and so from (3.13) we deduce the separation estimate

$$\delta(\omega_N^*) \geq C_2 N^{-1/\alpha} \quad (N \geq 2),$$

where

$$(3.14) \quad C_2 := \left[\frac{\bar{\mu}(A)}{C_0(r_1)} (1 - \alpha/s) \right]^{1/\alpha} (\alpha/s)^{1/s}, \quad r_1 = \text{diam}(A).$$

For the proof of Theorem 3, we utilize the following.

Lemma 8. *Let A be a compact, infinite, lower α -regular metric space with lower α -regular measure $\underline{\mu}$, $w : A \times A \rightarrow [0, \infty)$ be an SLP weight on A , and $s > \alpha$. Then there exists a positive integer N_0 independent of s , such that*

$$(3.15) \quad \mathcal{E}_s^w(N, A) \geq C_5 N^{1+s/\alpha} \quad (N \geq N_0),$$

where C_5 is a constant independent of N given below in (3.19).

Proof. Let κ and η be as in (3.1) and let $0 < r_2 \leq \kappa$. Since A is compact, there is some M such that the M -point best-packing distance satisfies

$$(3.16) \quad \delta_M(A) \leq r_2.$$

Let $N > M$ and let $\omega_N = \{x_1, \dots, x_N\} \subset A$ be an arbitrary N -point configuration of distinct points. For $1 \leq i \leq N$, let $y_i \in \omega_N$ be a fixed nearest neighbor to x_i in the configuration ω_N , and set

$$\delta_i := (x_i, y_i) = \min_{\substack{1 \leq j \leq N \\ j \neq i}} (x_i, x_j) > 0.$$

We assume an ordering on ω_N so that $\delta_i \leq \delta_{i+1}$ for $i = 1, \dots, N-1$. We note that $\omega_N \setminus \{x_1, \dots, x_{N-M}\}$ is of cardinality M and thus for all $i \leq N' := N - M$ we have that $\delta_i \leq r_2 \leq \kappa$.

The energy of ω_N then has the lower bound

$$(3.17) \quad \begin{aligned} E_s^w(\omega_N) &\geq \sum_{i=1}^{N'} \frac{w(x_i, y_i)}{\delta_i^s} \geq \sum_{i=1}^{N'} \eta \left(\frac{1}{\delta_i^\alpha} \right)^{s/\alpha} \geq \eta \left(\sum_{i=1}^{N'} \frac{1}{\delta_i^\alpha} \right)^{s/\alpha} (N')^{1-s/\alpha} \\ &\geq \eta \left(\sum_{i=1}^{N'} \delta_i^\alpha \right)^{-s/\alpha} (N')^{1+s/\alpha} = \eta 2^{-s} \left(\sum_{i=1}^{N'} \left(\frac{\delta_i}{2} \right)^\alpha \right)^{-s/\alpha} (N')^{1+s/\alpha}. \end{aligned}$$

where the last inequality in the first line follows from Jensen's inequality and the subsequent inequality follows from the harmonic-arithmetic mean inequality.

Let $\Lambda > 1$ and $N_0 := M\Lambda/(\Lambda - 1)$. Then $N' = N - M \geq \Lambda^{-1}N$ for $N \geq N_0$. Noting that the balls $B(x_i, \delta_i/2)$ are pairwise disjoint, we may apply the lower regularity of $\underline{\mu}$ (with regularity constant $c_0(r_2)$) to obtain

$$\begin{aligned}
 (3.18) \quad E_s^w(\omega_N) &\geq \eta 2^{-s} \left(c_0(r_2) \sum_{i=1}^{N'} \underline{\mu} \left(B(x_i, \frac{\delta_i}{2}) \right) \right)^{-s/\alpha} (N')^{1+s/\alpha} \\
 &\geq \frac{\eta}{(2^\alpha c_0(r_2) \underline{\mu}(A))^{s/\alpha}} (N')^{1+s/\alpha} \\
 &\geq \Lambda^{-1-s/\alpha} \frac{\eta}{(2^\alpha c_0(r_2) \underline{\mu}(A))^{s/\alpha}} N^{1+s/\alpha}
 \end{aligned}$$

Since (3.18) holds for arbitrary N -point configurations $\omega_N \subset A$ with $N \geq N_0$, we obtain that (3.15) holds with

$$(3.19) \quad C_5 := \Lambda^{-1-s/\alpha} \eta 2^{-s} (c_0(r_2) \underline{\mu}(A))^{-s/\alpha}.$$

We remark that N_0 depends on Λ and r_2 , but is independent of s . \square

Proof of Theorem 3. Appealing to the generality provided by Theorem 1 and Lemma 8, we can substantially extend and improve upon the arguments used in the proof of Theorem 3.6 in [6].

Let $\omega_N^* = \{x_1, \dots, x_N\}$ be an N -point (s, w) -energy minimizing configuration for the compact set K , and, for $y \in K$, consider the function

$$(3.20) \quad U(y) := \frac{1}{N} \sum_{i=1}^N \frac{w(y, x_i)}{(y, x_i)^s}.$$

For fixed $1 \leq j \leq N$, the function $U(y)$ can be decomposed as

$$(3.21) \quad U(y) = \frac{1}{N} \frac{w(y, x_j)}{(y, x_j)^s} + \frac{1}{N} \sum_{\substack{i=1 \\ i \neq j}}^N \frac{w(y, x_i)}{(y, x_i)^s},$$

and, since ω_N^* is a minimizing configuration on K , the point x_j minimizes the sum over $i \neq j$ on the right-hand side of equation (3.21). Thus for each fixed j and $y \in K$

$$(3.22) \quad U(y) \geq \frac{1}{N} \frac{w(y, x_j)}{(y, x_j)^s} + \frac{1}{N} \sum_{\substack{i=1 \\ i \neq j}}^N \frac{w(x_j, x_i)}{m(x_j, x_i)^s}.$$

Summing over j gives

$$(3.23) \quad NU(y) \geq \frac{1}{N} \sum_{j=1}^N \frac{w(y, x_j)}{(y, x_j)^s} + \frac{1}{N} \sum_{j=1}^N \sum_{\substack{i=1 \\ i \neq j}}^N \frac{w(x_j, x_i)}{m(x_j, x_i)^s}$$

$$(3.24) \quad = U(y) + \frac{1}{N} \mathcal{E}_s^w(N, K),$$

and thus

$$(3.25) \quad U(y) \geq \frac{1}{N(N-1)} \mathcal{E}_s^w(N, K) \geq \frac{\mathcal{E}_s^w(N, K)}{N^2} \quad (y \in K).$$

Since K is compact, there exists a point $y^* \in K$ such that

$$(3.26) \quad \min_{1 \leq i \leq N} (y^*, x_i) = \rho(\omega_N^*, K) =: \rho(\omega_N^*).$$

Using the fact that a function is lower semi-continuous if and only if it is the limit of an increasing sequence of continuous functions, it is not difficult to show that since w is a bounded SLP weight on K , it may be extended to a bounded SLP weight on A . Then, by Lemma 8, there are constants N_0 and $C_5 > 0$ such that

$$(3.27) \quad \mathcal{E}_s^w(N, K) \geq \mathcal{E}_s^w(N, A) \geq C_5 N^{1+s/\alpha} \quad (N \geq N_0).$$

We note that the constant C_5 of (3.27) does not depend on K , but rather on A (specifically on the lower regularity constant of A and on $\mu(A)$) as well as on the extended weight w .

Since (3.25) holds for the point y^* of (3.26), we combine (3.25) with (3.27) to obtain

$$(3.28) \quad U(y^*) \geq \frac{\mathcal{E}_s^w(N, K)}{N^2} \geq C_5 N^{s/\alpha-1} \quad (N \geq N_0).$$

Next we determine an upper bound for $U(y^*)$ using the α -regularity of the superset A . Since A is upper α -regular, we see that K is also because $\mu(K) > 0$. Hence, Corollary 2 applied to K implies that there is some $C_2 > 0$ such that $\delta(\omega_N^*) \geq C_2 N^{-1/\alpha}$ for $N \geq 2$. We note that the constant C_2 here depends on K , specifically $\mu(K)$.

Let \mathcal{N} consist of those $N \geq N_0$ such that

$$(3.29) \quad \rho(\omega_N^*) \geq \frac{C_2}{2} N^{-1/\alpha}.$$

If \mathcal{N} is empty (or finite) then we are done. Assuming that \mathcal{N} is nonempty, let $N \in \mathcal{N}$ be fixed.

For $0 < \epsilon < 1/2$, let

$$(3.30) \quad r_0 = r_0(N, \epsilon) := \epsilon C_2 N^{-1/\alpha}.$$

Note that any two of the balls $B(x_i, r_0) \subset A$, for $1 \leq i \leq N$, do not intersect since $r_0 < \delta(\omega_N^*)/2$.

For any $x \in B(x_i, r_0)$, inequalities (3.26) and (3.29) imply

$$(3.31) \quad \begin{aligned} (x, y^*) &\leq (x, x_i) + (x_i, y^*) \leq r_0 + (x_i, y^*) \\ &\leq 2\epsilon \rho(\omega_N^*) + (x_i, y^*) \leq (1 + 2\epsilon)(x_i, y^*). \end{aligned}$$

For fixed $1 \leq i \leq N$, using (3.31) and taking an average value on $B(x_i, r_0)$ we obtain

$$(3.32) \quad \begin{aligned} \frac{w(x_i, y^*)}{(x_i, y^*)^s} &\leq \frac{\|w\|_\infty (1 + 2\epsilon)^s}{\mu(B(x_i, r_0))} \int_{B(x_i, r_0)} \frac{d\mu(x)}{(x, y^*)^s} \\ &\leq \frac{\|w\|_\infty (1 + 2\epsilon)^s c_0(r_0)}{r_0^\alpha} \int_{B(x_i, r_0)} \frac{d\mu(x)}{(x, y^*)^s}, \end{aligned}$$

where $\|w\|_\infty$ denotes the sup-norm of w on $A \times A$ and $c_0(r_0)$ is the localized constant of (1.7) for the set A .

Inequality (3.29) and definition (3.30) imply $2\epsilon\rho(\omega_N^*) \geq r_0$ and thus, for $x \in B(x_i, r_0)$, we obtain

$$(3.33) \quad \begin{aligned} (x, y^*) &\geq (x_i, y^*) - (x, x_i) \geq (x_i, y^*) - r_0 \\ &\geq (x_i, y^*) - 2\epsilon\rho(\omega_N^*) \geq (1 - 2\epsilon)\rho(\omega_N^*). \end{aligned}$$

Inequality (3.33) implies

$$\bigcup_{i=1}^N B(x_i, r_0) \subset A \setminus B(y^*, (1 - 2\epsilon)\rho(\omega_N^*)),$$

and since the left-hand side is a disjoint union, averaging the inequalities of (3.32) we have

$$(3.34) \quad \begin{aligned} U(y^*) &\leq \frac{\|w\|_\infty (1 + 2\epsilon)^s c_0(r_0)}{N r_0^\alpha} \sum_{i=1}^N \int_{B(x_i, r_0)} \frac{d\mu(x)}{(x, y^*)^s} \\ &\leq \frac{\|w\|_\infty (1 + 2\epsilon)^s c_0(r_0)}{N r_0^\alpha} \int_{A \setminus B(y^*, (1-2\epsilon)\rho(\omega_N^*))} \frac{d\mu(x)}{(x, y^*)^s}. \end{aligned}$$

For fixed $\tau \geq 1$ we define the radius $R(N) := \tau(1 - 2\epsilon)\rho(\omega_N^*)$, and the constant

$$(3.35) \quad \tilde{C}_0(\tau) := C_0(R(N))(1 - \tau^{\alpha-s}) + C_0\tau^{\alpha-s}.$$

Note that if $\tau = 1$, then $\tilde{C}_0(1) = C_0$. (We retain τ as a parameter in our estimates as an option for the reader to optimize C_3 for a fixed s .) Now we break the integral on the right-hand side of (3.34) into two terms and proceed as in (3.7) to obtain

$$(3.36) \quad \begin{aligned} &\int_{A \setminus B(y^*, (1-2\epsilon)\rho(\omega_N^*))} \frac{d\mu(x)}{(x, y^*)^s} \\ &= \int_{B(y^*, (1-2\epsilon)\rho(\omega_N^*), R(N))} \frac{d\mu(x)}{(x, y^*)^s} + \int_{A \setminus B(y^*, R(N))} \frac{d\mu(x)}{(x, y^*)^s} \\ &\leq C_0(R(N)) \int_{R(N)^{-s}}^{[(1-2\epsilon)\rho(\omega_N^*)]^{-s}} t^{-\alpha/s} dt + C_0 \int_0^{R(N)^{-s}} t^{-\alpha/s} dt \\ &= \frac{\tilde{C}_0(\tau)}{(1 - \alpha/s)(1 - 2\epsilon)^{s-\alpha}} \rho(\omega_N^*)^{\alpha-s}. \end{aligned}$$

It is convenient to define the quantity

$$(3.37) \quad \beta(\epsilon) := \frac{\|w\|_\infty (1 + 2\epsilon)^s}{(1 - \alpha/s)(1 - 2\epsilon)^{s-\alpha} (\epsilon C_2)^\alpha},$$

and we note that for fixed $s > \alpha$ it is minimized as a function of ϵ for

$$(3.38) \quad \epsilon_0 := \frac{1}{2(2(s/\alpha) - 1)} < \frac{1}{2},$$

with minimal value

$$(3.39) \quad \beta_0 := \beta(\epsilon_0) = \frac{\|w\|_\infty}{(1 - \alpha/s)^{s-\alpha+1}} \left(\frac{4s}{\alpha C_2} \right)^\alpha.$$

Using ϵ_0 and combining inequality (3.34) with inequality (3.36) we obtain

$$(3.40) \quad U(y^*) \leq c_0(r_0)\beta_0\tilde{C}_0(\tau)\rho(\omega_N^*)^{\alpha-s}.$$

If $N \in \mathcal{N}$, then (3.40) and (3.28) imply

$$\rho(\omega_N^*) \leq \left[\frac{c_0(r_0)\beta_0\tilde{C}_0(\tau)}{C_5} \right]^{1/(s-\alpha)} N^{-1/\alpha}.$$

If $N \notin \mathcal{N}$, then either $N \leq N_0$ or $\rho(\omega_N^*) < \frac{C_2}{2}N^{-1/\alpha}$. Hence (2.9) holds with

$$(3.41) \quad C_3 := \max \left\{ \text{diam}(A)N_0^{1/\alpha}, \left[\frac{c_0(r_0)\beta_0\tilde{C}_0(\tau)}{C_5} \right]^{1/(s-\alpha)}, \frac{C_2}{2} \right\}.$$

We note that if $N > N_0$, then it suffices to take

$$(3.42) \quad C_3 = \max \left\{ \left[\frac{c_0(r_0)\beta_0\tilde{C}_0(\tau)}{C_5} \right]^{1/(s-\alpha)}, \frac{C_2}{2} \right\}$$

□

Proof of Theorem 7. Starting with Theorem 3 we shall employ a bootstrapping argument whereby the constants C_2 , C_5 , and subsequently C_3 are redefined so as to depend on N .

We begin by noting that if $s \geq 2\alpha$, then the constant C_3 of (3.41) has a uniform upper bound in s ; indeed, with $\kappa = \infty$, C_2 as defined in (3.14) and C_5 as defined in (3.19) (with $\eta = 1$), each of the three terms appearing in braces in (3.41) is uniformly bounded above. Thus there exists a constant C^* independent of $N \geq 2$ and of $s \geq 2\alpha$ such that $\rho(\omega_N^{(s)}, K) < C^*N^{-1/\alpha}$, where $\omega_N^{(s)}$ is any N -point $(s, 1)$ -energy minimizing configuration on K .

We next note that $C_0(0)$ of (1.8) is finite and positive, and utilizing the constant c_A of (1.9) we fix

$$(3.43) \quad C^{**} := \max \left\{ C^*, c_A, \left(\frac{\mu(K)}{C_0(0)} \right)^{1/\alpha} \right\},$$

and we now redefine the radius r_1 to be a function of N ,

$$(3.44) \quad r_1(N) := C^{**} N^{-1/\alpha} \quad (N \geq 2).$$

Returning to the proof of Theorem 1, we note that $r_1(N) > \rho(\omega_N^{(s)}, K)$, and so inequality (3.3) holds. Furthermore, by the choice of C^{**} we have that for $0 < \theta_0 < 1$ as in (3.9)

$$r_0(N) := \left(\frac{\theta_0 \mu(K)}{N C_0(0)} \right)^{1/\alpha} < r_1(N).$$

Taking $r_0 = r_0(N)$ in the proof and remembering that $q = 1$ in the current context, we see that with A replaced by K the penultimate term on right-hand side of (3.7) becomes

$$\frac{sC_0(r_1(N))}{s-\alpha} \left(\frac{\theta_0 \mu(K)}{N C_0(0)} \right)^{1-s/\alpha},$$

and thus

$$(3.45) \quad \int_{B(x_j, r_0(N), r_1(N))} \frac{d\mu(x)}{(x, x_j)^s} \leq \frac{sC_0(r_1(N))}{s-\alpha} \left(\frac{\theta_0 \mu(K)}{N C_0(0)} \right)^{1-s/\alpha} \\ \leq \frac{s}{s-\alpha} \left(\frac{\theta_0 \mu(K)}{N} \right)^{1-s/\alpha} C_0(r_1(N))^{s/\alpha},$$

where the last inequality follows from the fact that $C_0(0) \leq C_0(r_1(N))$ and $s > \alpha$.

For $w \equiv 1$, the constant C_2 of (3.14) with $r_1 = r_1(N)$ becomes

$$(3.46) \quad C_2(N) := \left(\frac{\alpha}{s} \right)^{1/s} \left(\frac{1-\alpha/s}{C_0(r_1(N))} \right)^{1/\alpha} \mu(K)^{1/\alpha},$$

where $C_0(r_1(N))$ is the local upper regularity constant of (1.7), and we have

$$\delta(\omega_N^{(s)}) \geq C_2(N) N^{-1/\alpha} \quad (N \geq 2, s \geq 2\alpha).$$

Furthermore, allowing the radius r_2 appearing in (3.16) to depend on $N \geq 2$ by taking $r_2 := r_1(N)$, we see via (1.9) and (3.43) that

$$r_1(N) \geq \delta_N(A) \quad (N \geq 2),$$

and there is no need to designate the integer M in the proof of Lemma 8. Thus we can take $\Lambda = 1$ in (3.19), and it follows (with $\eta = 1$) that

$$E_s^1(\omega_N^{(s)}) \geq C_5(N) N^{1+s/\alpha} \quad (N \geq 2, s \geq 2\alpha),$$

where

$$(3.47) \quad C_5(N) := \frac{1}{2^s [c_0(r_1(N)) \mu(A)]^{s/\alpha}}.$$

We remark that $C_2(N)$ clearly depends on the subset K , whereas $C_5(N)$ depends on the superset A .

We now return to the proof of Theorem 3 utilizing the constants $C_2(N)$ and $C_5(N)$. For β_0 as in (3.39), we see that

$$\rho(\omega_N^{(s)}, K) \leq C_3(N) N^{-1/\alpha} \quad (N \geq N_0, s \geq 2\alpha),$$

where N_0 is as in Lemma 8, and by (3.42) (choosing $\tau = 1$, so that $\tilde{C}_0(\tau) = C_0$)

$$(3.48) \quad C_3(N) := \max \left\{ \left[\frac{c_0(r_0) \beta_0 C_0}{C_5(N)} \right]^{1/(s-\alpha)}, \frac{C_2(N)}{2} \right\}.$$

With equations (3.46)-(3.48) in mind, we are ready to complete the proof of Theorem 7. The argument leading to equation (2.11) shows that ν_N is an N -point best-packing configuration on K for each $N \geq 2$. We now need to determine the limits of the constants $C_2(N)$ of (3.46) and $C_3(N)$ of (3.48) as $s \rightarrow \infty$. Fixing N in (3.46) yields

$$(3.49) \quad \lim_{s \rightarrow \infty} C_2(N) = \left(\frac{\mu(K)}{C_0(r_1(N))} \right)^{1/\alpha} =: \hat{C}_2(N).$$

Since $c_0(r_0)$ and C_0 are independent of s and $\lim_{s \rightarrow \infty} \beta_0^{1/(s-\alpha)} = 1$, it follows, that for fixed N

$$\begin{aligned}
 \lim_{s \rightarrow \infty} C_3(N) &= \max \left\{ \frac{\hat{C}_2(N)}{2}, \lim_{s \rightarrow \infty} C_5(N)^{1/(\alpha-s)} \right\} \\
 (3.50) \quad &= \max \left\{ \frac{1}{2} \left(\frac{\mu(K)}{C_0(r_1(N))} \right)^{1/\alpha}, 2[c_0(r_1(N))\mu(A)]^{1/\alpha} \right\} \\
 &:= \hat{C}_3(N)
 \end{aligned}$$

From the continuity of $\delta(\cdot)$ and $\rho(\cdot, K)$ on K^N we deduce that

$$\delta(\nu_N) \geq \hat{C}_2(N)N^{-1/\alpha} \quad \text{and} \quad \rho(\nu_N, K) \leq \hat{C}_3(N)N^{-1/\alpha} \quad (N \geq N_0).$$

Taking the ratio of these two quantities we have that

$$(3.51) \quad \frac{\rho(\nu_N, K)}{\delta(\nu_N)} \leq \frac{\hat{C}_3(N)}{\hat{C}_2(N)} = \max \left\{ \frac{1}{2}, 2 \left(\frac{\mu(A)}{\mu(K)} \right)^{1/\alpha} [c_0(r_1(N)) C_0(r_1(N))]^{1/\alpha} \right\},$$

and hence for $N \geq N_0$

$$(3.52) \quad \limsup_{N \rightarrow \infty} \frac{\rho(\nu_N, K)}{\delta(\nu_N)} \leq \max \left\{ \frac{1}{2}, 2 \left(\frac{\mu(A)}{\mu(K)} \right)^{1/\alpha} [c_0(0) C_0(0)]^{1/\alpha} \right\}$$

$$(3.53) \quad = 2 \left(\frac{\mu(A)}{\mu(K)} \right)^{1/\alpha} [c_0(0) C_0(0)]^{1/\alpha} < \infty.$$

Therefore, the sequence of configurations $\{\nu_N\}_{N=2}^\infty$ is quasi-uniform on K . \square

Acknowledgments. We thank the referees for their careful reading and detailed comments.

REFERENCES

- [1] A.V. Bondarenko, D.P. Hardin, E.B. Saff, *Mesh norms and separation distances for best-packing configurations*, (manuscript).
- [2] S.V. Borodachov, D.P. Hardin, E.B. Saff, *Asymptotics for discrete weighted minimal Riesz energy problems on rectifiable sets*, Trans. Amer. Math. Soc. **360** (2008), 1559–1580.
- [3] S.V. Borodachov, E.B. Saff, D.P. Hardin, *Asymptotics of densest weighted packing on rectifiable sets* (Russian) Mat. Sb. **199** (2008), 3–20; translation in Sb. Math. **199** (2008), 1579–1595.
- [4] J.S. Brauchart, D.P. Hardin, E.B. Saff, *The support of the limit distribution of optimal Riesz energy points on sets of revolution in \mathbb{R}^3* , J. Math. Phys. **48** (2007), 122901, 24 pp.
- [5] B.E.J. Dahlberg, *On the distribution of Fekete points*, Duke Math. J. **45** (1978), 537–542.
- [6] S.B. Damelin, V. Maymeskul; *On point energies, separation radius, and mesh norm for s -extremal configurations on compact sets in \mathbb{R}^n* , J. Complexity **21** (2005), 845–863.
- [7] P.D. Dragnev, *On the separation of logarithmic points on the sphere*, Approximation Theory X: Abstract and Classical Analysis (C. K. Chui, L. L. Schumaker, and J. Stöckler, eds.) (2002) 137–144.
- [8] P.D. Dragnev, E.B. Saff, *Riesz spherical potentials with external fields and minimal energy points separation*, Potential Anal. **26** (2007), 139–162.
- [9] E.J. Fuselier, G.B. Wright, *Stability and error estimates for vector field interpolation and decomposition on the sphere with RBFs*, SIAM J. Numer. Anal. **47** (2009), 3213–3239.
- [10] D.P. Hardin, E.B. Saff, *Minimal Riesz energy point configurations for rectifiable d -dimensional manifolds*, Adv. Math. **193** (2005), 174–204.
- [11] D.P. Hardin, E.B. Saff, H. Stahl, *Support of the logarithmic equilibrium measure on sets of revolution in \mathbb{R}^3* , J. Math. Phys., **48** (2007), 022901, 14 pp.

- [12] J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer-Verlag New York, Universitext (2001)
- [13] A.B.J. Kuijlaars, E.B. Saff, *Asymptotics for minimal discrete energy on the sphere*, Trans. Amer. Math. Soc. **350** (1998), 523-538.
- [14] A.B.J. Kuijlaars, E.B. Saff, X. Sun, *On a separation of minimal Riesz energy points on spheres in Euclidean spaces*, J. Comp. Appl. Math. **199**: (2007) 172-180.
- [15] Q.T. Le Gia, I.H. Sloan, H. Wendland, *Multiscale analysis in Sobolev spaces on the sphere*, *SIAM J. Numer. Anal.* 48 (2010), 20652090. 65J10 (46E35 46N40)
- [16] P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces; fractals and rectifiability*, Cambridge Studies in Advanced Math. vol. **44**, Cambridge University Press (1995).
- [17] I. Pesenson, *A sampling theorem on homogeneous manifolds*, Trans. Amer. Math. Soc. **352** (2000), 4257-4269.
- [18] E.A. Rakhmanov, E.B. Saff, Y.M. Zhou, *Electrons on the sphere*. Comput. Methods Funct. Theory, 293-309, Ser. Approx. Decompos., 5, World Sci. Publ., River Edge, NJ, 1995. 78A30 .
- [19] R. Schaback, *Error estimates and condition numbers for radial basis function interpolation*, Adv. Comput. Math. **3** (1995), 251-264.
- [20] A. Schürmann, *Computational Geometry of Positive Definite Quadratic Forms*, University Lecture Series vol. 48, Amer. Math. Soc., Providence, RI (2009).
- [21] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Colloq. Publ. **23**, Amer. Math. Soc., Providence, RI, 4th edition (1975).

D. P. HARDIN, E. B. SAFF, AND J. T. WHITEHOUSE: CENTER FOR CONSTRUCTIVE APPROXIMATION, DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240, USA

E-mail address: Doug.Hardin@Vanderbilt.Edu

E-mail address: Edward.B.Saff@Vanderbilt.Edu

E-mail address: Tyler.Whitehouse@gmail.com