Extension of vertex operator algebra $V_{\widehat{H}_{4}}(\ell,0)$

Cuipo Jiang¹ and Song Wang²

Department of Mathematics, Shanghai Jiaotong University, Shanghai 200240 China

Abstract

We classify the irreducible restricted modules for the affine Nappi-Witten Lie algebra H_4 with some natural conditions. It turns out the representation theory of H_4 is quite different from the theory of representations of Heisenberg algebras. We also study the extension of the vertex operator algebra $V_{\hat{H}_4}(\ell,0)$ by the even lattice L. We give the structure of the extension $V_{\widehat{H}_4}(\ell,0)\otimes\mathbb{C}[\widehat{L}]$ and its irreducible modules via irreducible representations of $V_{\hat{H}_4}(\ell, 0)$ viewed as a vertex algebra.

2000MSC:17B69

1 Introduction

The Nappi-Witten algebra H_4 is a four-dimensional Lie algebra with the basis $\{a, b, c, d\}$ and the following Lie bracket:

$$
[a, b] = c
$$
, $[d, a] = a$, $[d, b] = -b$, $[c, d] = [c, a] = [c, b] = 0$.

There exists a non-degenerate symmetric bilinear form $($, $)$ on H_4 defined by

 $(a, b) = 1, (c, d) = 1, \text{ otherwise, } (,) = 0.$

The associated affine Nappi-Witten Lie algebra is

$$
\widehat{H}_4 = H_4 \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C} \mathbf{k}.
$$

The study of the affine Nappi-Witten algebra \widehat{H}_4 was initiated by the theory of Wess-Zumino-Novikov-Witten (WZNW) models based on non-abelian non-semisimple groups [31, 27]. WZNW models corresponding to abelian or compact semisimple groups have been studied extensively. The importance of non-abelian non-semisimple types was first noticed in [8, 27]. It was shown in [27] that the NW model corresponding to a central extension of the two-dimensional Euclidean groups describes the spectrum and scattering of strings moving in a gravitational wave background. In [19], irreducible representations with unitary base for the affine NW algebra were constructed through four free fields.

¹Supported by China NSF grants 10931006,10871125, the Innovation Program of Shanghai Municipal Education Commission (11ZZ18) and a grant of Science and Technology Commission of Shanghai Municipality (No. 09XD1402500)(E-mail: cpjiang@sjtu.edu.cn).

 ${}^{2}E$ -mail: wangsong025@sina.com.

In [1], Verma type modules for the affine Nappi-Witten algebra \widehat{H}_4 were investigated. The associated vertex operator algebra $V_{\hat{H}_4}(\ell,0)$ for a given complex number ℓ and its representations were also studied in [1]. More results on the NW model can be found in [4, 5, 6, 7].

It is known that every restricted \hat{H}_4 -module of level ℓ carries a unique module structure for the vertex algebra $V_{\hat{H}_4}(\ell,0)$, extending the action of H_4 in a canonical way and that every module of the vertex algebra $V_{\widehat{H}_{4}}(\ell,0)$ is naturally a restricted H_{4} -module of level ℓ . This leads us to study restricted modules for the affine Lie algebra H_4 . On the other hand, let $L = \mathbb{Z}c \otimes \mathbb{Z}d$, then L is a non-degenerate even lattice. Motivated by [22], [23], [24], [25] and [11], we consider the extension $V_{\hat{H}_4}(\ell,0)[L]$ of $V_{\hat{H}_4}(\ell,0)$ by L. Some ideas we use come from these papers. For the simple vertex operator algebra $V_{\hat{\mathfrak{g}}}(\ell,0)$ and an irreducible highest weight $V_{\hat{\mathfrak{g}}}(\ell,0)$ -module W with \mathfrak{g} a finite-dimensional simple Lie algebra and ℓ a positive integer, or with $\mathfrak g$ abelian and $\ell = 1$, it is known that $W^{(\alpha)}$ is still an irreducible highest weight module for the affine Lie algebra $\hat{\mathfrak{g}}$ (cf. [11], [22], [23], [24], [25]), where $\alpha \in \mathfrak{h}$ such that $\alpha(0)$ acts semisimply on V with integral eigenvalues and \mathfrak{h} is a Cartan subalgebra of **g**. But for $V_{\hat{H}_4}(\ell, 0)$, it turns out it is quite different. For general $\alpha \in L$ and an irreducible highest weight H_4 -module W, $W^{(\alpha)}$ is not a highest weight module of H_4 anymore. We characterize $W^{(\alpha)}$ for every irreducible highest weight module of \hat{H}_4 .

The paper is organized as follows: In Section 2, we recall the various modules, admissible modules and ordinary modules for a vertex operator algebra. We also give some known related results on the affine Nappi-Witten algebra and the associated vertex operator algebra. In section 3, we characterize three types of restricted H_4 -modules and their irreducible quotients. Section 4 is devoted to the extension $V_{\widehat{H}_{4}}(\ell,0)[L]$ and its irreducible modules.

2 Preliminaries

In this section, we briefly review the definitions of vertex operator algebra and various modules (cf. [3], [9], [12], [16], [36], [20], [35]). We also give some known results related to the Nappi-Witten algebra H_4 .

A vertex operator algebra V is a Z-graded vector space $V = \bigoplus_{n \in \mathbb{Z}} V_n$ equipped with a linear map $Y: V \to (\text{End}V)[[z, z^{-1}]], a \mapsto Y(a, z) = \sum_{n \in \mathbb{Z}}$ $a_n z^{-n-1}$ for $a \in V$ such that dim V_n is finite for all integer n and that $V_n = 0$ for sufficiently small n (see [16]). There are two distinguished vectors, the vacuum vector $1 \in V_0$ and the Virasoro element $\omega \in V_2$. By definition $Y(1, z) = id_V$, and the component operators $\{L(n)|n \in \mathbb{Z}\}\$ of $Y(\omega, z) = \sum$ $\overline{n\varepsilon}\mathbb{Z}$ $L(n)z^{-n-2}$ give a representation of the Virasoro algebra on V with central charge c. Each homogeneous subspace V_n ($n \geq 0$) is an eigenspace for $L(0)$ with eigenvalue n .

Definition 2.1. A weak V-module is a vector space M equipped with a linear map

$$
Y_M: V \to (\text{End}M)[[z, z^{-1}]]
$$

$$
v \mapsto Y_M(v, z) = \sum_{n \in \mathbb{Z}} v_n z^{-n-1}, v_n \in \text{End}M
$$

satisfying the following conditions:

- 1) $v_n w = 0$ for $n >> 0$ where $v \in V$ and $w \in M$
- 2) $Y_M(1, z) = \text{Id}_M$
- 3) The Jacobi identity holds:

$$
z_0^{-1}\delta\left(\frac{z_1 - z_2}{z_0}\right)Y_M(u, z_1)Y_M(v, z_2) - z_0^{-1}\delta\left(\frac{z_2 - z_1}{-z_0}\right)Y_M(v, z_2)Y_M(u, z_1)
$$

=
$$
z_2^{-1}\delta\left(\frac{z_1 - z_0}{z_2}\right)Y_M(Y(u, z_0)v, z_2).
$$
 (2.1)

Definition 2.2. An admissible V-module is a weak V-module which carries a \mathbb{Z}_+ -grading $M = \bigoplus_{r=0}^{\infty} M(n)$, such that if $v \in V_r$, then $v_m M(n) \subseteq M(n + r - m - 1)$. $n{\in}\mathbb{Z}_+$

Definition 2.3. An ordinary V-module is a weak V-module which carries a \mathbb{C} -grading $M = \bigoplus M_{\lambda}$, such that:

- $\lambda\overline{\in}\mathbb{C}$ 1) $dim(M_\lambda) < \infty$,
- 2) $M_{\lambda+n} = 0$ for fixed λ and $n \ll 0$,

3) $L(0)w = \lambda w = \text{wt}(w)w$ for $w \in M_\lambda$, where $L(0)$ is the component operator of $Y_M(\omega, z) = \sum$ n∈Z $L(n)z^{-n-2}$.

Remark 2.4. It is easy to see that an ordinary V-module is an admissible one. If W is an ordinary V -module, we simply call W a V -module.

We now introduce the Nappi-Witten algebra and some known related results. The Nappi-Witten Lie algebra H_4 is a 4-dimensional vector space

$$
H_4 = \mathbb{C}a \oplus \mathbb{C}b \oplus \mathbb{C}c \oplus \mathbb{C}d
$$

equipped with the bracket relations

$$
[a, b] = c
$$
, $[d, a] = a$, $[d, b] = -b$, $[c, d] = [c, a] = [c, b] = 0$.

and a non-degenerate symmetric bilinear form $($, $)$ on H_4 defined by

$$
(a, b) = 1
$$
, $(c, d) = 1$, otherwise, (,) = 0.

The Nappi-Witten algebra H_4 is has the following triangular decomposition:

$$
H_4 = H_4^+ \oplus H_4^0 \oplus H_4^- = H_4^{\geq 0} \oplus H_4^-,
$$

where

$$
H_4^+ = \mathbb{C}a, \quad H_4^0 = \mathbb{C}c \oplus \mathbb{C}d, \quad H_4^- = \mathbb{C}b.
$$

For $\lambda \in (H_4^0)^*$, the highest weight module (Verma module) of H_4 is defined by

$$
M(\lambda) = U(H_4) \otimes_{U(H_4^{\geq 0})} \mathbb{C}_{\lambda},
$$
\n(2.2)

where \mathbb{C}_{λ} is the 1-dimensional $H_4^{\geq 0}$ $\frac{1}{4}^{\infty}$ -module, on which $h \in H_4^0$ acts as multiplication by $\lambda(h)$ and H_4^+ acts as 0. For convenience, we denote $M(\lambda)$ by $M(\mathfrak{c},\mathfrak{d})$ when $\lambda(c) = \mathfrak{c}$ and $\lambda(d) = d.$

We have the following lemma from [28].

Lemma 2.5. For $c, d \in \mathbb{C}$, $M(c, d)$ is irreducible if and only if $c \neq 0$. If $c = 0$, then the irreducible quotient $L(\mathbf{d}) = L(0, \mathbf{d}) = \mathbb{C}v_{\mathbf{d}}$ such that

$$
av_{\mathbf{d}} = bv_{\mathbf{d}} = cv_{\mathbf{d}} = 0, \quad dv_{\mathbf{d}} = dv_{\mathbf{d}}.
$$
\n(2.3)

Let $M = V(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu})$ for $\tilde{\lambda}, \tilde{\mu}, \tilde{\nu} \in \mathbb{C}$, where $V(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu})$ is the H_4 -module defined as follows: $V(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}) = \bigoplus$ n∈Z $\mathbb{C}v_n$ such that

$$
dv_n = (\tilde{\lambda} + n)v_n, \ cv_n = \tilde{\mu}v_n, \ av_n = -\tilde{\mu}v_{n+1}, \ bv_n = (\tilde{\lambda} + \tilde{\nu} + n)v_{n-1}.
$$
 (2.4)

The following result was obtained by M. Willard [30].

Theorem 2.6. All irreducible weight modules of H_4 with finite-dimensional weight spaces can be classified into the following classes:

- (1) Irreducible modules $L(\mathbf{d})$ for $\mathbf{d} \in \mathbb{C}$.
- (2) Irreducible highest weight modules $M(\lambda)$ or irreducible lowest weight modules $M^-(\mu)$.
- (3) Intermediate series modules $V(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu})$ defined as (2.4) such that $\tilde{\mu} \neq 0$ and $\tilde{\lambda} + \tilde{\nu} \notin$ Z.

 \Box

 \Box

To the pair $(H_4, (,))$, let \widehat{H}_4 be the associated *affine Nappi-Witten Lie algebra* with the underlying vector space

$$
\widehat{H}_4 = H_4 \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C} \mathbf{k} \tag{2.5}
$$

equipped with the bracket relations

$$
[h_1 \otimes t^m, h_2 \otimes t^n] = [h_1, h_2] \otimes t^{m+n} + m(h_1, h_2) \delta_{m+n,0} \mathbf{k}, \; [\hat{H}_4, \mathbf{k}] = 0,\tag{2.6}
$$

for $h_1, h_2 \in H_4$ and $m, n \in \mathbb{Z}$. It is clear that \widehat{H}_4 has the Z-grading:

$$
\widehat{H}_4 = \coprod_{n \in \mathbb{Z}} \widehat{H}_4^{(n)},
$$

where

$$
\widehat{H}_4^{(0)} = H_4 \oplus \mathbb{C} \mathbf{k}, \quad \text{and} \quad \widehat{H}_4^{(n)} = H_4 \otimes t^n \quad \text{for } n \neq 0.
$$

Then one has the following graded subalgebras of \widehat{H}_4 :

$$
\widehat{H}_4^{(\pm)} = \coprod_{n>0} \widehat{H}_4^{(\pm n)}, \quad \widehat{H}_4^{(\geq 0)} = \coprod_{n\geq 0} \widehat{H}_4^{(n)} = \widehat{H}_4^{(+)} \oplus H_4 \oplus \mathbb{C} \mathbf{k}.
$$

Let M be an H_4 -module and $\ell \in \mathbb{C}$. Regard M as an $\widehat{H}_4^{(\geq 0)}$ -module with $\widehat{H}_4^{(+)}$ acting trivially and **k** as the scalar ℓ . The induced H_4 -module $V_{\hat{H}_4}(\ell, M)$ is defined by

$$
V_{\widehat{H}_4}(\ell, M) = \text{Ind}_{\widehat{H}_4^{(2\ 0)}}^{\widehat{H}_4}(M) = U(\widehat{H}_4) \otimes_{U(\widehat{H}_4^{(2\ 0)})} M. \tag{2.7}
$$

Let $M = L(\mathbf{d}) = \mathbb{C}v_{\mathbf{d}}$ for $\mathbf{d} \in \mathbb{C}$ be the one-dimensional irreducible H_4 -module defined as in (2.3). Denote

$$
V_{\widehat{H}_4}(\ell, d) = U(\widehat{H}_4) \otimes_{U(\widehat{H}_4^{(\geq 0)})} \mathbb{C}v_d.
$$
\n(2.8)

We have the following result:

Theorem 2.7 ([1]). For $\ell, d \in \mathbb{C}$, the \widehat{H}_4 -module $V_{\widehat{H}_4}(\ell, d)$ is irreducible if and only if $\ell \neq 0$. Furthermore, if $\ell = 0$, then the irreducible quotient of $V_{\widehat{H}_{4}}(\ell, d)$ is isomorphic to the one-dimensional \hat{H}_4 -module $\mathbb{C}v_{\rm d}$.

For $h \in H_4$, we define the generating function

$$
h(x) = \sum_{n \in \mathbb{Z}} (h \otimes t^n) x^{-n-1} \in \widehat{H}_4[[x, x^{-1}]].
$$

Then the defining relations (2.6) can be equivalently written as

$$
[h_1(x_1), h_2(x_2)] = [h_1, h_2](x_2) x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) - (h_1, h_2) \frac{\partial}{\partial x_1} x_2^{-1} \delta\left(\frac{x_1}{x_2}\right) \mathbf{k}.
$$
 (2.9)

Given an \hat{H}_4 -module W, let $h(n)$ denote the operator on W corresponding to $h \otimes t^n$ for $h \in H_4$ and $n \in \mathbb{Z}$. We shall use the notation for the action of $h(x)$ on W:

$$
h_W(x) = \sum_{n \in \mathbb{Z}} h(n)x^{-n-1} \in (\text{End}W)[[x, x^{-1}]].
$$
\n(2.10)

Definition 2.8. Let W be a restricted \widehat{H}_4 -module in the sense that for every $h \in H_4$ and $w \in W$, $h(n)w = 0$ for n sufficiently large. We say that the H_4 -module W is of level ℓ if the central element **k** acts as a scalar ℓ in \mathbb{C} .

Let ℓ be a complex number. Consider the induced module defined as $(2.8)(let d = 0)$:

$$
V_{\widehat{H}_{4}}(\ell,0) = U(\widehat{H}_{4}) \otimes_{U(\widehat{H}_{4}^{(\leq 0)})} v_{0}.
$$

Set

$$
\mathbf{1} = v_0 \in V_{\widehat{H}_4}(\ell, 0).
$$

Then

$$
V_{\hat{H}_4}(\ell,0) = \coprod_{n\geq 0} V_{\hat{H}_4}(\ell,0)_{(n)},
$$

where $V_{\hat{H}_4}(\ell,0)_{(n)}$ is spanned by the vectors

$$
h_{-m_1}^{(1)}\cdots h_{-m_r}^{(r)}\mathbf{1}
$$

for $r \geq 0$, $h^{(i)} \in H_4$, $m_i \geq 1$, with $n = m_1 + \cdots + m_r$. It is clear that $V_{\hat{H}_4}(\ell, 0)$ is a restricted H_4 -module of level ℓ . We can regard H_4 as a subspace of $V_{\hat{H}_4}(\ell,0)$ through the map

$$
H_4 \to V_{\widehat{H}_4}(\ell, 0), \quad h \mapsto h(-1)\mathbf{1}.
$$

In fact, $H_4 = V_{\hat{H}_4}(\ell, 0)_{(1)}$.

Theorem 2.9 (cf. [20, 26]). Let ℓ be any complex number. Then there exists a unique vertex algebra structure $(V_{\hat{H}_4}(\ell,0),Y,1)$ on $V_{\hat{H}_4}(\ell,0)$ such that 1 is the vacuum vector and

$$
Y(h, x) = h(x) \in (\text{End}V_{\hat{H}_4}(\ell, 0))[[x, x^{-1}]]
$$

for $h \in H_4$. For $r \geq 0$, $h^{(i)} \in H_4$, $n_i \in \mathbb{Z}_+$, the vertex operator map for this vertex algebra structure is given by

$$
Y(h^{(1)}(-n_1)\cdots h^{(r)}(-n_r)\mathbf{1},x) = \int_{0}^{\infty} \partial^{(n_1-1)}h^{(1)}(x)\cdots \partial^{(n_r-1)}h^{(r)}(x) \big|_{0}^{\infty} 1,
$$

where

$$
\partial^{(n)} = \frac{1}{n!} \left(\frac{d}{dx} \right)^n
$$

◦ ◦ ◦ \int_{α}^{∞} is the normal-ordering operation, and 1 is the identity operator on $V_{\widehat{H}_{4}}(\ell,0)$.

Proposition 2.10 (cf. [20]). Any module W for the vertex algebra $V_{\widehat{H}_4}(\ell,0)$ is naturally a restricted \hat{H}_4 -module of level ℓ , with $h_W(x) = Y_W(h,x)$ for $h \in H_4$. Conversely, any restricted H₄-module W of level ℓ is naturally a $V_{\widehat{H}_{4}}(\ell,0)$ -module as vertex algebra with

$$
Y_W(h^{(1)}(-n_1)\cdots h^{(r)}(-n_r)\mathbf{1},x) = \frac{6}{9}\partial^{(n_1-1)}h_W^{(1)}(x)\cdots \partial^{(n_r-1)}h_W^{(r)}(x)\frac{6}{9}\mathbf{1}_W,
$$

for $r \geq 0, h^{(i)} \in \widehat{H}_4, n_i \in \mathbb{Z}_+$. Furthermore, for any $V_{\widehat{H}_4}(\ell, 0)$ -module W, the $V_{\widehat{H}_4}(\ell, 0)$ submodules of W coincide with the \widehat{H}_4 -submodules of W.

Let ℓ be a non-zero complex number. Set

$$
\omega = \frac{1}{\ell} (a(-1)b(-1)\mathbf{1} + c(-1)d(-1)\mathbf{1}) - \frac{1}{2\ell}c(-2)\mathbf{1} - \frac{1}{2\ell^2}c(-1)c(-1)\mathbf{1} \quad (2.11)
$$

and define operators $L(n)$ for $n \in \mathbb{Z}$ by

$$
Y(\omega, x) = \sum_{n \in \mathbb{Z}} \omega_n x^{-n-1} = \sum_{n \in \mathbb{Z}} L(n) x^{-n-2}.
$$

Then we have the following results from [1].

Theorem 2.11. Let ℓ be a complex number such that $\ell \neq 0$. Then the vertex algebra $V_{\widehat{H}_{4}}(\ell,0)$ constructed in Theorem 2.9 is a vertex operator algebra of central charge 4 with ω defined in (2.11) as the conformal vector. The Z-grading on $V_{\hat{H}_4}(\ell,0)$ is given by $L(0)$ eigenvalues. Moreover, $H_4 = V_{\widehat{H}_4}(\ell,0)_{(1)}$, which generates $V_{\widehat{H}_4}(\ell,0)$ as a vertex algebra, and

$$
[h(n), L(m)] = nh(m+n), \quad \text{for } h \in H_4, \ m, n \in \mathbb{Z}.
$$

Theorem 2.12. For $\ell \neq 0$ and $d \in \mathbb{C}$, the \widehat{H}_4 -module $V_{\widehat{H}_4}(\ell, d)$ is naturally an irreducible ordinary module for the vertex operator algebra $V_{\widehat{H}_{4}}(\ell,0)$. Furthermore, the modules $V_{\hat{H}_4}(\ell, d)$ exhaust all the irreducible ordinary $V_{\hat{H}_4}(\ell, 0)$ -modules up to equivalence.

Note that the central charge of the vertex operator algebra $V_{\hat{H}_4}(\ell,0)$ for $\ell \neq 0$ is independent of ℓ . We have the following result.

Theorem 2.13. Let $\ell \neq 0$ be any nonzero complex number and let $\sqrt{\ell}$ be one of the two square roots of ℓ . Then the vertex operator algebra $V_{\widehat{H}_{4}}(\ell,0)$ is isomorphic to $V_{\widehat{H}_{4}}(1,0)$ under the following linear map

$$
\rho: V_{\widehat{H}_4}(\ell,0) \to V_{\widehat{H}_4}(1,0)
$$

$$
d(-m_{41}) \cdots d(-m_{4i_4})c(-m_{31}) \cdots c(-m_{3i_3})
$$

\n
$$
a(-m_{21}) \cdots a(-m_{2i_2})b(-m_{11}) \cdots b(-m_{1i_1})\mathbf{1}
$$

\n
$$
\rightarrow (\sqrt{\ell})^{i_1+i_2} \ell^{i_3} d(-m_{41}) \cdots d(-m_{4i_4})c(-m_{31}) \cdots c(-m_{3i_3})
$$

\n
$$
a(-m_{21}) \cdots a(-m_{2i_2})b(-m_{11}) \cdots b(-m_{1i_1})\mathbf{1},
$$

where $i_k \geq 0$, $m_{kj} \geq 1$, $m_{k1} \leq m_{k2} \leq \cdots \leq m_{ki_k}$, $k = 1, \cdots, 4, j = 1, 2, \cdots, i_k$.

Following Theorem 2.13, for the rest of the paper, we always assume that $\ell = 1$.

3 Restricted modules for the affine Lie algebra \hat{H}_4

In this section, we will discuss the structures of restricted \hat{H}_4 -modules under some conditions which are natural from the point of view of representations of vertex algebras. We first have the following lemma.

Lemma 3.1. Let W be an irreducible \widehat{H}_4 -module, then c(0) and **k** act as scalars on W.

Proof: Note that \widehat{H}_4 has countable basis and W is irreducible. So W is of countable dimension over \mathbb{C} . Then the lemma follows from the fact that $c(0)$ and **k** are in the center of \widehat{H}_4 and $\text{Hom}_{\widehat{H}_4}(W, W) = \mathbb{C}$.

Let $\mathfrak{h} = \mathbb{C}c \oplus \mathbb{C}d$ and $\hat{\mathfrak{h}} = \mathfrak{h} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}k$ be the associated subalgebra of the affine Lie algebra \hat{H}_4 .

We now introduce a subalgebra $\widehat{H}_4(m, \epsilon)$ of \widehat{H}_4 for given $m \in \mathbb{Z}$ and $\epsilon \in \{0, 1\}$ as follows:

$$
\widehat{H}_4(m,\epsilon) = (\bigoplus_{n\geq -m} \mathbb{C}a \otimes t^n) \bigoplus (\bigoplus_{n\geq m+\epsilon} \mathbb{C}b \otimes t^n) \bigoplus ((\mathbb{C}c \bigoplus \mathbb{C}d) \otimes \mathbb{C}[t]) \bigoplus \mathbb{C}\mathbf{k}.
$$
 (3.1)

Let $0 \neq \alpha = \text{d}c + \text{d}c \in \mathfrak{h}^*$. We define a one-dimensional $\widehat{H}_4(m, \epsilon)$ -module $\mathbb{C}_{\alpha} = \mathbb{C}1_{\alpha}$ by

$$
(a\otimes t^s)\mathbf{1}_{\alpha}=(b\otimes t^r)\mathbf{1}_{\alpha}=(c\otimes t^l)\mathbf{1}_{\alpha}=(d\otimes t^l)\mathbf{1}_{\alpha}=0, \ s\geq -m, \ r\geq m+\epsilon, \ l\geq 1,
$$

 $c\mathbf{1}_{\alpha} = c\mathbf{1}_{\alpha}, \ d\mathbf{1}_{\alpha} = d\mathbf{1}_{\alpha}, \ k\mathbf{1}_{\alpha} = \mathbf{1}_{\alpha}.$

Then we get a generalized Verma type module $V_{\hat{H}_4}(1,(m,\epsilon), \mathbf{1}_{\alpha})$ as follows:

$$
V_{\widehat{H}_4}(1,(m,\epsilon),\mathbf{1}_{\alpha})=U(\widehat{H}_4)\otimes_{U(\widehat{H}_4(m,\epsilon))}\mathbb{C}_{\alpha}.
$$
\n(3.2)

If $m = \mathfrak{c} \in \mathbb{Z}$ and $\epsilon = 0$, we briefly denote $V_{\widehat{H}_4}(1,(m,\epsilon), \mathbf{1}_{\alpha})$ by $V_{\widehat{H}_4}(1,\mathfrak{c}, \mathbf{1}_{\alpha})$. Note that

$$
V_{\hat{H}_4}(1,0,\mathbf{1}_{\alpha})=V_{\hat{H}_4}(1,\mathbf{d}).
$$

We first consider the module $V_{\hat{H}_4}(1,\mathbb{C},\mathbf{1}_{\alpha})$ for $\mathbb{C} \in \mathbb{Z}$. We give the first main result of this section as follows.

Theorem 3.2. Let $0 \neq \alpha = \text{d}c + \text{d}c \in \mathfrak{h}^*$ such that $\mathfrak{c} \in \mathbb{Z}$. Then the generalized Verma type H_4 -module $V_{\hat{H}_4}(1, \mathbb{c}, \mathbf{1}_{\alpha})$ is irreducible.

Proof: It is clear that $\sigma : \tilde{H}_4 \to \tilde{H}_4$ defined by $\sigma(a \otimes t^n) = b \otimes t^n$, $\sigma(b \otimes t^n) = a \otimes t^n$, $\sigma(c \otimes t^n) = -c \otimes t^n$, $\sigma(d \otimes t^n) = -d \otimes t^n$, $n \in \mathbb{Z}$, $\sigma(\mathbf{k}) = \mathbf{k}$ is an automorphism of \widehat{H}_4 . So we may assume that $c \ge 0$. By PBW theorem and the definition of $V_{\hat{H}_4}(1, c, \mathbf{1}_{\alpha})$,

$$
\mathbf{1}_{\alpha}, d(m_{41})\cdots d(m_{4i_4})c(m_{31})\cdots c(m_{3i_3})a(m_{21})\cdots a(m_{2i_2})b(m_{11})\cdots b(m_{1i_1})\mathbf{1}_{\alpha},
$$

such that $i_k \geq 0$, $m_{k1} \leq m_{k2} \leq \cdots \leq m_{ki_k}$, m_{4j} , $m_{3j} \leq -1$, $m_{2j} \leq -\varepsilon - 1$, $m_{1j} \leq \varepsilon - 1$, $k = 1, \dots, 4, j = 1, 2, \dots, i_k$, is a basis of $V_{\widehat{H}_4}(1, \infty, \mathbf{1}_{\alpha})$.

For $n \in \mathbb{Z}_+$, let

$$
p_n = c(n), \quad q_n = d(-n).
$$

Then

$$
[p_m, q_n] = m\delta_{m,n}\mathbf{k}.
$$

Therefore

$$
\mathfrak{s} = \mathop{\oplus}_{n \in \mathbb{Z}_+} (\mathbb{C}p_n \oplus \mathbb{C}q_n) \oplus \mathbb{C}{\bf k}
$$

is a Heisenberg algebra and $V_{\widehat{H}_{4}}(1, \mathbb{c}, 1_\alpha)$ is an $\mathfrak{s}\text{-module}$ such that $\bf k$ acts as 1. Since every highest weight $\mathfrak s$ -module generated by one element with k acting as a non-zero scalar is irreducible, it follows that $V_{\widehat H_4}(1,\mathbb{c},\mathbf{1}_\alpha)$ can be decomposed into a direct sum of irreducible highest weight modules of $\boldsymbol{\mathfrak{s}}$ with the highest weight vectors

$$
\mathbf{1}_{\alpha},c(m_{31})\cdots c(m_{3i_3})a(m_{21})\cdots a(m_{2i_2})b(m_{11})\cdots b(m_{1i_1})\mathbf{1}_{\alpha},
$$

such that $i_k \geq 0$, $m_{k1} \leq m_{k2} \leq \cdots \leq m_{ki_k}$, $m_{3j} \leq -1$, $m_{2j} \leq -\mathfrak{c} - 1$, $m_{1j} \leq \mathfrak{c} - 1$, $k = 1, 2, 3, j = 1, 2, \cdots, i_k$. For convenience, we denote the set of such highest weight vectors by N . Set

$$
\widehat{H}_4(\mathbf{c},+)=(\bigoplus_{n\geq -\mathbf{c}}\mathbb{C}a\otimes t^n)\bigoplus(\bigoplus_{n\geq \mathbf{c}}\mathbb{C}b\otimes t^n)\bigoplus ((\mathbb{C}c\bigoplus\mathbb{C}d)\otimes \mathbb{C}t[t])
$$

Let U be a non-zero submodule of $V_{\hat{H}_4}(1, \mathbb{c}, 1)$. It suffices to prove that $1_{\alpha} \in U$. Let u be a non-zero element in U. Since as an \mathfrak{s} -module, U is a direct sum of irreducible highest weight $\mathfrak{s}\text{-modules}$, we may assume that

$$
u=\sum_{i=1}^s a_i c(m_{31}^{(i)})\cdots c(m_{3i_3}^{(i)})a(m_{21}^{(i)})\cdots a(m_{2i_2}^{(i)})b(m_{11}^{(i)})\cdots b(m_{1i_1}^{(i)})b(n_{i1})\cdots b(n_{ir_i})\mathbf{1}_{\alpha},
$$

where $0 \neq a_i \in \mathbb{C}, i_k \geq 0, m_{k1}^{(i)} \leq m_{k2}^{(i)} \leq \cdots \leq m_{ki}^{(i)}$ $\binom{(i)}{ki_k}, m_{3j}^{(i)} \leq -1, m_{2j}^{(i)} \leq -\mathbb{c} - 1,$ $m_{1j}^{(i)} \leq -1, 0 \leq n_{i1}, \dots, n_{ir_i} \leq \epsilon - 1, i = 1, 2, \dots, s, k = 1, 2, 3, j = 1, 2, \dots, i_k$, and $\sum_{j=1}^{1} m_{1j}^{(1)} \leq \sum_{j=1}^{2} m_{1j}^{(2)} \leq \cdots \leq \sum_{j=1}^{s_1} m_{1j}^{(s)}$ $_{1j}^{\left(s\right) }\text{. }$ Let

$$
u^{(i)} = c(m_{31}^{(i)}) \cdots c(m_{3i_3}^{(i)}) a(m_{21}^{(i)}) \cdots a(m_{2i_2}^{(i)}) b(m_{11}^{(i)}) \cdots b(m_{1i_1}^{(i)}) b(n_{i1}) \cdots b(n_{ir_i}) \mathbf{1}_{\alpha}, i = 1, 2, \cdots, s.
$$

Note that

$$
a(-m_{11_1}^{(1)}) \cdots a(-m_{12}^{(1)})a(-m_{11}^{(1)})u^{(1)}
$$

= $q \prod_{j=1}^{1_1} (c-m_{1j}^{(1)})c(m_{31}^{(1)}) \cdots c(m_{31_3}^{(i)})a(m_{21}^{(1)}) \cdots a(m_{21_2}^{(i)})b(n_{i1}) \cdots b(n_{ir_i})\mathbf{1}_{\alpha}$
 $\neq 0,$

for some $0 \neq q \in \mathbb{Z}$ and for the case $b(m_{11}^{(i)}) \cdots b(m_{1i_1}^{(i)})$ $\binom{i}{1}$ \neq $b(m_{11}^{(1)}) \cdots b(m_{11}^{(1)})$ $\binom{11}{111}$, we have

$$
a(-m_{11_1}^{(1)})\cdots a(-m_{12}^{(1)})a(-m_{11}^{(1)})u^{(i)} = 0.
$$

We may further assume that

$$
u=\sum_{i=1}^s a_i c(m_{31}^{(i)})\cdots c(m_{3i_3}^{(i)})a(m_{21}^{(i)})\cdots a(m_{2i_2}^{(i)})b(n_{i1})\cdots b(n_{ir_i})\mathbf{1}_{\alpha},
$$

and

$$
u^{(i)} = c(m_{31}^{(i)}) \cdots c(m_{3i_3}^{(i)}) a(m_{21}^{(i)}) \cdots a(m_{2i_2}^{(i)}) b(n_{i1}) \cdots b(n_{ir_i}) \mathbf{1}_{\alpha}, \ i = 1, 2, \cdots, s,
$$

where $0 \neq a_i \in \mathbb{C}, i_k \geq 0, m_{k1}^{(i)} \leq m_{k2}^{(i)} \leq \cdots \leq m_{ki}^{(i)}$ $\left(\begin{smallmatrix} (i)\ iki \ k i_k, \end{smallmatrix} \right) \left. m^{(i)}_{2j} \right| \leq -1, \ m^{(i)}_{2j} \leq -\mathbb{c} \left. -1, \right.$ $0 \le n_{i1} \le \cdots \le n_{ir_i} \le c-1, i = 1, 2, \cdots, s, k = 1, 2, j = 1, 2, \cdots, i_k \text{ and } \sum_{j=1}^{1_2} m_{2j}^{(1)} \le$

$$
\sum_{j=1}^{2_2} m_{2j}^{(2)} \leq \dots \leq \sum_{j=1}^{s_2} m_{2j}^{(s)}.
$$
 Note also that
\n
$$
b(-m_{21_2}^{(1)}) \dots + b(-m_{22}^{(1)})b(-m_{21}^{(1)})u^{(1)}
$$
\n
$$
= q_1 \prod_{j=1}^{1_1} (-\mathbf{c} - m_{2j}^{(1)}) c(m_{31}^{(1)}) \dots + c(m_{31_3}^{(i)}) b(n_{i1}) \dots + b(n_{ir_i}) \mathbf{1}_{\alpha}
$$
\n
$$
\neq 0,
$$

for some $0 \neq q_1 \in \mathbb{Z}$ and for the case $a(m_{21}^{(i)}) \cdots a(m_{2i_2}^{(i)})$ $a_{2i_2}^{(i)}$ $\neq a(m_{21}^{(1)}) \cdots b(m_{21}^{(1)})$ $\binom{1}{21_2}$, we have

$$
b(-m_{21_2}^{(1)})\cdots b(-m_{22}^{(1)})b(-m_{21}^{(1)})u^{(i)}=0.
$$

We may also assume that

$$
u=\sum_{i=1}^s a_i c(m_{i1})\cdots c(m_{ip_i})b(n_{i1})\cdots b(n_{ir_i})\mathbf{1}_{\alpha},
$$

where $0 \neq a_i \in \mathbb{C}, p_i, r_i \geq 0, m_{i1} \leq m_{i2} \leq \cdots \leq m_{ip_i} \leq -1, 0 \leq n_{i1} \leq \cdots \leq n_{ir_i} \leq \epsilon - 1,$ $i = 1, 2, \dots, s$. Note that $[d, b] = -b$, so we may assume that

$$
r_1=r_2=\cdots=r_s=r.
$$

If $r = 0$, then by the fact that $[c(p), d(q)] = p\delta_{p+q,0}k$, it is easy to see that $\mathbf{1}_{\alpha} \in U$. If $r \neq 0$. Assume that

$$
(n_{i1}, n_{i2}, \cdots, n_{ir}) \le (n_{i+1,1}, n_{i+1,2} \cdots, n_{i+1,r}), i = 1, 2, \cdots, r
$$

in the sense that

$$
n_{i1} \leq n_{i2} \leq \cdots \leq n_{ir},
$$

and

$$
n_{i1} = n_{i+1,1}, \dots, n_{ik} = n_{i+1,k}, n_{i,k+1} < n_{i+1,k+1}, \text{ for some } k.
$$

Then

$$
a(-n_{1r})\cdots a(-n_{11})c(m_{11})\cdots c(m_{1p_1})b(n_{11})\cdots b(n_{1r}) = kc(m_{11})\cdots c(m_{1p_1})\mathbf{1}_{\alpha} \neq 0
$$

and for the case that $(n_{11}, \dots, n_{1r}) < (n_{i1}, \dots, n_{ir}),$ we have

$$
a(-n_{1r})\cdots a(-n_{11})c(m_{i1})\cdots c(m_{ip_i})b(n_{i1})\cdots b(n_{ir})1_{\alpha}=0.
$$

So

$$
a(-n_{1r})\cdots a(-n_{11})u=\sum_{i=1}^{r_1}c_ic(m_{11})\cdots c(m_{1p_1})\mathbf{1}_{\alpha}\in U,
$$

for some $r_1 \le r$ and $c_i \neq 0, i = 1, 2, \dots, r_1$. From this we deduce that $\mathbf{1}_{\alpha} \in U$.

Next we consider the second class of generalized Verma type module $V_{\hat{H}_4}(1,(m,\epsilon),\mathbf{1}_{\alpha})$ for $\epsilon = 1$. Let $0 \neq \alpha = \text{d}c + \text{d} \in \mathfrak{h}^*$, $(m, 1) \in \mathbb{Z}^2$, and let $V_{\hat{H}_4}(1, (m, 1), \mathbf{1}_{\alpha})$ be the generalized Verma type \widehat{H}_4 -module defined by (3.2). For $n \in \mathbb{Z}$, define

$$
\deg a(-n - m) = n - 1, \ \deg b(-n + m) = n + 1,
$$

$$
\deg c(-n) = n, \ \deg d(-n) = n, \ \ \deg k = 0.
$$

Then $U(\widehat{H}_4)$ is Z-graded. So we get an N-gradation of $V_{\widehat{H}_4}(1,(m,1), 1)$ by defining $\deg \mathbf{1}_{\alpha} = 0$. Let

$$
\widehat{H}_4((m,1),+) = (\bigoplus_{n \ge -m} \mathbb{C}a \otimes t^n) \bigoplus (\bigoplus_{n \ge m+1} \mathbb{C}b \otimes t^n) \bigoplus ((\mathbb{C}c \bigoplus \mathbb{C}d) \otimes \mathbb{C}t[t])
$$
 (3.3)

A homogeneous element u in $V_{\widehat{H}_4}(1,(m,1), \mathbf{1}_{\alpha})$ is called a homogeneous singular vector if

 $x \cdot u = 0.$

for all $x \in \mathcal{U}(\widehat{H}_{4}(m, 1), +)$. It is obvious that the module generated by a homogeneous singular vector is a proper submodule. The following is the second main result of this section.

Theorem 3.3. For $0 \neq \alpha = \text{d}c + \text{d}c \in \mathfrak{h}^*$ and $(m, 1) \in \mathbb{Z}^2$, the \widehat{H}_4 -module $V_{\widehat{H}_4}(1, (m, 1), \mathbf{1}_{\alpha})$ is irreducible if and only if $c \notin \mathbb{Z}$. Furthermore, we have

- (i) If $\mathfrak{c} = m$, then $u = b(m) \mathbf{1}_{\alpha}$ is a singular vector and $V_{\hat{H}_4}(1,(m,1), \mathbf{1}_{\alpha})/U(H_4)u$ is irreducible and isomorphic to $V_{\widehat{H}_{4}}(1, \infty, \mathbf{1}_{\alpha})$.
- (ii) If $c + n = m$ for some $n \in \mathbb{Z}_+ \setminus \{0\}$, then all the linearly independent singular vectors are

$$
u^{k} = \left[\sum_{\lambda \in P(n)} (a_{\lambda}c(-\lambda)b(m) + \sum_{i=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_{i}, \lambda_{i}}c(-\lambda \setminus \lambda_{i})b(-\lambda_{i}+m)\right]^{k} \mathbf{1}_{\alpha},
$$

for all $k \in \mathbb{Z}_+$, satisfying

$$
a_{\lambda}q_i\lambda_i - b_{\lambda\setminus\lambda_i,\lambda_i} = 0, \quad i = 1, 2, \cdots, k_{\lambda},
$$

$$
b_{\lambda\setminus\lambda_i,\lambda_i}(-n+\lambda_i) + \sum_{j=1}^{k_{\lambda}} b_{\lambda\setminus\lambda_i,\lambda_j,\lambda_i+\lambda_j} = 0;
$$

(iii) If $c = m + n$ for some $n \in \mathbb{Z}_+ \setminus \{0\}$, then all the linearly independent singular vectors are

$$
u^{k} = \left[\sum_{\lambda \in P(n)} \sum_{i=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_{i}, \lambda_{i}} c(-\lambda \setminus \lambda_{i}) a(-\lambda_{i} - m)\right]^{k} \mathbf{1}_{\alpha},
$$

for all $k \in \mathbb{Z}_+$, satisfying

$$
q_i\lambda_i c_{\lambda \setminus \lambda_j, \lambda_j} + c_{\lambda \setminus \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0, \ i, j = 1, 2, \cdots, k_\lambda, i \neq j,
$$

$$
c_{\lambda \setminus \lambda_i, \lambda_i}(-n + \lambda_i) - \sum_{j=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0, \ i = 1, 2, \cdots, k_{\lambda},
$$

where in (ii) and (iii), $P(n)$ is the set of all partitions of weight n,

$$
\lambda^{(k)} = (\lambda, \lambda, \cdots, \lambda) \in \mathbb{Z}_+^k, \quad \lambda = (\lambda_1^{(q_1)}, \lambda_2^{(q_2)}, \cdots, \lambda_{k_\lambda}^{(q_{k_\lambda})})
$$

such that
$$
\sum_{i=1}^{k_{\lambda}} q_i \lambda_i = n
$$
, and $\lambda \setminus \lambda_i = (\lambda_1^{(q_1)}, \cdots, \lambda_i^{(q_i-1)}, \cdots, \lambda_{k_{\lambda}}^{(q_{k_{\lambda}})})$ and if $\lambda = \lambda_i$,
then $c(-\lambda \setminus \lambda_i)b(-\lambda_i + m) = b(-\lambda_i + m)$, $c(-\lambda \setminus \lambda_i)a(-\lambda_i - m) = a(-\lambda_i - m)$.

Proof: If $c = m$, then it is easy to see that $b(m) \mathbf{1}_{\alpha}$ is a singular vector. Then (i) follows from Theorem 3.2.

We now assume that $\mathbb{c} \neq m$.

Let $\mathfrak s$ be the infinite-dimensional Heisenberg algebra defined in the proof of Theorem 3.2. Then as an s-module, $V_{\hat{H}_4}(1,(m,1), \mathbf{1}_{\alpha})$ is a direct sum of irreducible highest weight s-modules with k acting as 1 and highest weight vectors in

$$
\mathcal{N}_{\alpha} = \{b(m)^{i}\mathbf{1}_{\alpha}, c(-\lambda)a(-\mu)b(-\nu)b(m)^{i}\mathbf{1}_{\alpha}, y_{1}(-\lambda)y_{2}(-\mu)b(m)^{i}\mathbf{1}_{\alpha},
$$

 $x(-\lambda)b(m)^{i} \mathbf{1}_{\alpha} \mid i \geq 0, (y_1, y_2) \in \{(a, b), (c, a), (c, b)\}, x \in \{a, b, c\}, \lambda, \mu, \nu \in \mathcal{P}\},\$

where for a partition $\lambda = (\lambda_1, \cdots, \lambda_r)$,

$$
d(-\lambda) = d(-\lambda_1)d(-\lambda_2)\cdots d(-\lambda_r),
$$

\n
$$
c(-\lambda) = c(-\lambda_1)c(-\lambda_2)\cdots c(-\lambda_r),
$$

\n
$$
a(-\lambda) = a(-\lambda_1 - m)a(-\lambda_2 - m)\cdots a(-\lambda_r - m),
$$

\n
$$
b(-\lambda) = b(-\lambda_1 + m)b(-\lambda_2 + m)\cdots b(-\lambda_r + m).
$$

Suppose that $u \in V_{\hat{H}_4}(1,(m,1), \mathbf{1}_{\alpha})$ is a singular vector such that

$$
U(\widehat{H}_4^+) = 0.
$$

Since $V_{\hat{H}_4}(1,(m,1), \mathbf{1}_{\alpha})$ is a direct sum of irreducible highest weight s-modules and u is homogeneous, we may assume that

$$
u = \sum_{i=1}^{3} u_i,
$$

where

$$
u_1 = \sum_{j=1}^{l_1} a_{1j} c(-\lambda^{(1j)}) a(-\mu^{(1j)}) b(-\nu^{(1j)}) b(m)^{k_{1j}} \mathbf{1}_{\alpha},
$$

for $(\lambda^{(1j)}, \mu^{(1j)}, \nu^{(1j)}) \succ (\lambda^{(1,j+1)}, \mu^{(1,j+1)}, \nu^{(1,j+1)}), j = 1, \cdots, l_1 - 1,$

$$
u_2 = u_{21} + u_{22} + u_{23},
$$

where

$$
u_{21} = \sum_{j=1}^{l_2} a_{2j}c(-\lambda^{(2j)})a(-\mu^{(2j)})b(m)^{k_{2j}}\mathbf{1}_{\alpha}, \ u_{22} = \sum_{j=1}^{l_3} a_{3j}c(-\lambda^{(3j)})b(-\nu^{(3j)})b(m)^{k_{3j}}\mathbf{1}_{\alpha},
$$

$$
u_{23} = \sum_{j=1}^{l_4} a_{4j}a(-\mu^{(4j)})b(-\nu^{(4j)})b(m)^{k_{4j}}\mathbf{1}_{\alpha},
$$

for $(\lambda^{(2j)}, \mu^{(2j)}) \succ (\lambda^{(2,j+1)}, \mu^{(2,j+1)}), j = 1, \cdots, l_2-1; (\lambda^{(3j)}, \nu^{(3j)}) \succ (\lambda^{(3,j+1)}, \mu^{(3,j+1)}), j =$ $1, \cdots, l_3-1; (\mu^{(4j)}, \nu^{(4j)}) \succ (\mu^{(4,j+1)}, \nu^{(4,j+1)}), j=1, \cdots, l_4-1, \text{ and}$

$$
u_3 = u_{31} + u_{32} + u_{33},
$$

where

$$
u_{31} = \sum_{j=1}^{l_5} a_{5j}c(-\lambda^{(5j)})b(m)^{k_{5j}}\mathbf{1}_{\alpha},
$$

\n
$$
u_{32} = \sum_{j=1}^{l_6} a_{6j}a(-\mu^{(6j)})b(m)^{k_{6j}}\mathbf{1}_{\alpha}, u_{33} = \sum_{j=1}^{l_7} a_{7j}b(-\nu^{(7j)})b(m)^{k_{7j}}\mathbf{1}_{\alpha},
$$

\n
$$
u_{3j+1} = 1 \dots l_7 = 1 \dots l_8 = 1 \dots l_9 = 1 \dots l_8 = 1 \dots l_9 =
$$

for $\lambda^{(5j)} \succ \lambda$ $^{(5,j+1)}, j = 1, \cdots, l_5 - 1; \mu$ 1; $\mu^{(0)} > \mu$ $^{(6,j+1)}, j=1,\cdots, l_6-1; \nu$ $^{(7j)}\succ\nu$ \cdot , J $1, \cdots, l_7-1.$

Case 1 $\in \mathcal{L}$ Z. One can prove that $u = 0$ by the same method used in the proof of Theorem 3.2. Therefore $V_{\widehat{H}_4}(1,(m,1), \alpha)$ is irreducible.

Case 2 $\mathbb{c} \in \mathbb{Z}$ and $\mathbb{c} \neq m$.

Since $au = 0$, it is easy to see that $k_{ij} = 0$, for $i = 4, 6, 7$. If $u_{33} \neq 0$, we consider $a(\nu_1^{(71)}$ $\binom{(71)}{1}u$. Since no monomial of $a(\nu_1^{(71)})$ $a_1^{(71)})u_1, a(\nu_1^{(71)})$ $u_1^{(71)}$) u_2 , $a(\nu_1^{(71)})$ $a_1^{(71)})u_{31}$ and $a(\nu_1^{(71)})$ $x_1^{(1)}$) u_{32} is of the form $b(-\eta)$, where $\eta \in \mathcal{P}$, we deduce that

$$
x + \nu_1^{(71)} - m = 0.
$$

Similarly, considering $a(\nu_{r_{7p}}^{(7p)})u$, where $\nu_{r_{7p}}^{(7p)} = min{\nu_{r_{7j}}^{(7j)}}, j = 1, 2, \cdots, l_7$, we have

$$
\mathbb{c} + \nu_{r_{7p}}^{(7p)} - m = 0.
$$

Then we deduce that $l_7 = 1$ and $u_{33} = a_{71}(b(-\nu_1^{(71)})$ $\binom{(71)}{1}$ ^{k₃}**1**_α such that

$$
c + \nu_1^{(71)} - m = 0, \quad k_3(\nu_1^{(71)} + 1) = \deg u.
$$

If $u_{32} \neq 0$, then by the fact that $b(\mu_1^{(61)})$ $\binom{0}{1} u = 0$, we have

$$
c - (\mu_1^{(61)} + m) = 0.
$$

Therefore

$$
\mu_1^{(61)} + \nu_1^{(71)} = 0,
$$

which is impossible since $\mu_1^{(61)} + \nu_1^{(71)} > 0$. We deduce that $u_{32} = 0$ or $u_{33} = 0$.

Subcase 1 $u_{33} \neq 0$, $u_{32} = 0$.

By the fact that $b(\mu_1^{(41)})$ $j_1^{(41)})u = b(\mu_1^{(41)})$ $b(\mu_1^{(41)})u_{31} = b(\mu_1^{(41)})$ $b(\mu_1^{(41)})u_{33} = b(\mu_1^{(41)})$ $\binom{41}{1}u_{22} = 0$, and both $b(\mu_1^{(41)}$ $_1^{(41)})u_1$ and $b(\mu_1^{(41)})$ $a_{1}^{(41)})u_{21}$ contain no monomials of the forms $a(-\lambda)b(-\mu)$ and $b(-\mu)$, where $\lambda, \mu \in \mathcal{P}$, we have

$$
b(\mu_1^{(41)})u_{23} = \sum_{j=1}^{l_4} a_{4j} \sum_{p=1}^{r_{4j}} \delta_{\mu_1^{(41)}, \mu_p^{(4j)}}(-\mathbf{c} + \mu_1^{(41)} + m)
$$

$$
\cdot a(-\mu^{(4j)})a(-\mu_p^{(4j)})b(-\nu^{(4j)})\mathbf{1}_{\alpha} = 0,
$$

where $a(-\mu_p^{(4j)})$ means this factor is deleted. So if $u_{23} \neq 0$, then $-c + \mu_1^{(41)} + m = 0$, which is impossible since $\mathfrak{c} + \nu_1^{(71)} - m = 0$. This proves that $u_{23} = 0$. Let $Y = b(\mu_1^{(21)}$ $\binom{(21)}{1},$ then $Yu = Yu_1 + Yu_{21} = 0$. If $u_{21} \neq 0$, comparing Yu_1 and Yu_{21} , we have

$$
-\mathbf{c} + \mu_1^{(21)} + m = 0,
$$

which is not true. So $u_{21} = 0$. Similarly, $u_1 = 0$. Therefore

$$
u = u_{22} + u_{31} + u_{33}.
$$

By the fact that $a(\nu_1^{(31)})$ $(v^{(31)})u = 0$, we can easily deduce that $v^{(3j)} \preceq v^{(71)}$, $j = 1, \dots, l_3$. Similarly, we have $\lambda^{(3j)} \leq \nu^{(71)}$, $\lambda^{(5k)} \leq \nu^{(71)}$, $j = 1, \dots, l_3$, $k = 1, \dots, l_5$. If $\nu_1^{(71)} = 1$, then $x + 1 - m = 0$, $u_{33} = a_{71}b(-1)^{k_3}1_\alpha$ and

$$
u_{22} = \sum_{j=1}^{l_3} a_{3j}c(-1)^{p_{3j}}b(-1)^{q_{3j}}b(m)^{k_{3j}}\mathbf{1}_{\alpha}, \ u_{31} = \sum_{j=1}^{l_5} a_{5j}c(-1)^{p_{5j}}b(m)^{k_{5j}}\mathbf{1}_{\alpha}.
$$

Since $d(1)u = 0$, we have

$$
u_{22} + u_{31} + u_{33} = k(\sum_{j=0}^{k_3} C_{k_3}^j c(-1)^{k_3-j} b(-1+m)^j b(m)^{k_3-j} \mathbf{1}_{\alpha}),
$$

for some $k \in \mathbb{C}$. Let

$$
u = \sum_{j=0}^{k_3} C_{k_3}^j c(-1)^{k_3-j} b(-1+m)^j b(m)^{k_3-j} \mathbf{1}_{\alpha} = (c(-1)b(m) + b(-1+m))^{k_3} \mathbf{1}_{\alpha}.
$$

Then we have

$$
a(-m)(c(-1)b(m) + b(-1+m)) = (c(-1)b(m) + b(-1+m))a(-m).
$$

Thus

$$
a(-m)u=0.
$$

It is clear that $a(r - m)u = b(r + m + 1)u = c(r + 1)u = d(r + 1)u = 0$, for $r \ge 0$. We prove that u is a non-zero singular vector for each $k_3 \in \mathbb{N}$.

Assume that $\nu_1^{(71)} > 1$. If $\nu_1^{(3j)} = 1$ for all $j = 1, 2, \dots, l_3$. Then $\nu_1^{(71)} = 2$. So $x+2-m=0$. In fact, if $\nu_1^{(71)} > 2$, then $a(\nu_1^{(71)}-1-m)u_{22} = a(\nu_1^{(71)}-1-m)u_{31} = 0$, $a(\nu_1^{(71)} - 1 - m)u_{33} \neq 0$, a contradiction. If $k_3 > 1$, then $a(1-m)u_{31} = 0$, and $a(1-m)u_{33}$ contains factor $b(-2 + m)$, but no monomial of $a(1 - m)u_{22}$ contains factor $b(-2 + m)$. So $u_{33} = u_{22} = 0$. Then $u_{31} = 0$ and we deduce that $u = 0$. If $k_3 = 1$, then $u_{33} =$ $a_{71}b(-2+m)\mathbf{1}_{\alpha}$ and one can easily deduce that

$$
u = (c(-2)b(m) + c(-1)^{2}b(m) + 2c(-1)b(-1+m) + 2b(-2+m))\mathbf{1}_{\alpha}
$$

is a singular vector. Generally, for $\nu_1^{(71)} = 2$, we have $\varepsilon + 2 - m = 0$ and

$$
u = (c(-2)b(m) + c(-1)^{2}b(m) + 2c(-1)b(-1+m) + 2b(-2+m))^{k}\mathbf{1}_{\alpha}, \quad k \ge 1.
$$

Now assume that $\nu_1^{(71)} = n > 2$ and $l(\nu^{(71)}) = 1$. Note that $\varepsilon + n - m = 0$. Let $1 \leq p \leq l_3$ be such that $l(\nu^{(3p)}) = max\{l(\nu^{(3j)}), j = 1, 2, \cdots, l_3\}$ and if $l(\nu^{(3q)}) = l(\nu^{(3p)}),$ then $\nu^{(3p)} \succeq \nu^{(3q)}$. If $l(\nu^{(3p)}) > 1$, then $d(\lambda^{(3p)})u \neq 0$, a contradiction. So $l(\nu^{(3j)}) = 1$ for all $j = 1, 2, \dots, l_3$. It follows that $k_{3j} = 0, j = 1, \dots, l_3$ since $a(-m)u = 0$. For $x \in \mathbb{Z}_+, k \in \mathbb{Z}_+$, denote $(x, \dots, x) \in \mathbb{N}^k$ by $x^{(k)}$. Then

$$
u = \sum_{\lambda \in P(n)} [a_{\lambda}c(-\lambda)b(m) + \sum_{i=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_i, \lambda_i}c(-\lambda \setminus \lambda_i)b(-\lambda_i)]\mathbf{1}_{\alpha},
$$

where $P(n)$ is the set of all partitions of weight $n, \lambda = (\lambda_1^{(q_1)})$ $\lambda_1^{(q_1)},\lambda_2^{(q_2)},\cdots,\lambda_{k_\lambda}^{(q_{k_\lambda})}$ $\binom{(4k\lambda)}{k\lambda}$ such that $\sum_{i=1}^{k_{\lambda}} q_i \lambda_i = n$, and $\lambda \setminus \lambda_i = (\lambda_1^{(q_1)})$ $\lambda_1^{(q_1)},\cdots,\lambda_i^{(q_i-1)},\cdots,\lambda_{k_\lambda}^{(q_{k_\lambda})}$ $\left(\begin{array}{c}\n\langle \Psi_{k}\rangle \\
k_{\lambda}\end{array}\right)$ and if $\lambda = \lambda_i$, then $c(-\lambda\setminus\lambda_i)b(-\lambda_i+m) = b(-\lambda_i+m)$. By the fact that $d(r+1)u = a(r-m)u = b(r+m+1)u =$ 0 for $r \in \mathbb{Z}_+$, and $c + n - m = 0$, we deduce that $a_{\lambda}, b_{\lambda \setminus \lambda_i, \lambda_i}, i = 1, 2, \cdots, k_{\lambda}$ are uniquely determined by the following equations up to a non-zero scalar.

$$
a_{\lambda}q_i\lambda_i - b_{\lambda\setminus\lambda_i,\lambda_i} = 0, \ i = 1,2,\cdots,k_{\lambda},\tag{3.4}
$$

$$
b_{\lambda \setminus \lambda_i, \lambda_i}(-n + \lambda_i) + \sum_{j=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0.
$$
 (3.5)

It is easy to check that the u determined by (3.4) and (3.5) is indeed a non-zero singular vector. Generally

$$
u = \left[\sum_{\lambda \in P(n)} (a_{\lambda}c(-\lambda)b(m) + \sum_{i=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_i, \lambda_i}c(-\lambda \setminus \lambda_i)b(-\lambda_i))\right]^k \mathbf{1}_{\alpha}, \ k = 1, 2, \cdots
$$

satisfying (3.4) and (3.5) are all the linearly independent singular vectors.

Subcase 2 $u_{33} = 0$, $u_{32} \neq 0$. Then $-c + \mu_1^{(61)} + m = 0$. Similar to the proof for Subcase 1, we can deduce that $u_1 = u_{22} = u_{23} = 0$, $l_6 = 1$, $\mu^{(61)} = (\mu_1^{(61)}$ $\mathcal{L}_1^{(61)}, \cdots, \mathcal{L}_1^{(61)}$ $_1^{\text{(01)}}$ and $k_{2i} = k_{5j} = 0, i = 1, 2, \cdots, l_2, j = 1, 2, \cdots, l_5.$

We first assume that $l(\mu^{(61)}) = 1$ and $\mu_1^{(61)} = n$. Then it is easy to see that $l(\mu^{(2j)}) = 1$, $j = 1, 2, \dots, l_2, u_{21} = 0$. Therefore

$$
u = \sum_{\lambda \in P(n)} \sum_{i=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_i, \lambda_i} c(-\lambda \setminus \lambda_i) a(-\lambda_i)) \mathbf{1}_{\alpha},
$$

where $P(n)$ is the set of all partitions of weight $n, \lambda^{(k)} = (\lambda, \lambda, \dots, \lambda) \in \mathbb{Z}_+^k$, $\lambda = (\lambda_1^{(q_1)}, \lambda_2^{(q_2)})$ $\binom{(q_1)}{1},$ $\lambda_2^{(q_2)}$ $\lambda_2^{(q_2)}, \cdots, \lambda_{k_\lambda}^{(q_{k_\lambda})}$ $\binom{(q_{k_{\lambda}})}{k_{\lambda}}$ such that $\sum_{i=1}^{k_{\lambda}} q_i \lambda_i = n$, and $\lambda \setminus \lambda_i = (\lambda_1^{(q_1)})$ $\lambda_1^{(q_1)},\cdots,\lambda_i^{(q_i-1)},\cdots,\lambda_{k_\lambda}^{(q_{k_\lambda})}$ $\binom{q_{k_{\lambda}}}{k_{\lambda}}$. By $d(n)u = b(n)u = 0$ for $n \ge 1$, we deduce that

$$
q_i \lambda_i c_{\lambda \lambda_j, \lambda_j} + c_{\lambda \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0, \ i, j = 1, 2, \cdots, k_\lambda, i \neq j,
$$
\n(3.6)

$$
c_{\lambda \setminus \lambda_i, \lambda_i}(-n + \lambda_i) - \sum_{j=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0, \ i = 1, 2, \cdots, k_{\lambda}.
$$
 (3.7)

Actually, u is uniquely determined by (3.6) and (3.7).

Next for the case that $l(\mu^{(61)}) > 1$ and $\mu_1^{(61)} = n$, one can easily deduce that

$$
u = \left[\sum_{\lambda \in P(n)} \sum_{i=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_i, \lambda_i} c(-\lambda \setminus \lambda_i) a(-\lambda_i)\right]^k \mathbf{1}_{\alpha}, \ k \in \mathbb{Z}_+
$$

satisfying (3.6) and (3.7) for some $k \in \mathbb{N}$.

Conversely, for $k \in \mathbb{Z}$, $k \geq 1$, it is easy to check that

$$
u = \left[\sum_{\lambda \in P(n)} \sum_{i=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_i, \lambda_i} c(-\lambda \setminus \lambda_i) a(-\lambda_i)\right]^k \mathbf{1}_{\alpha}, \ k \in \mathbb{Z}_+
$$

satisfying (3.6) and (3.7) is a singular vector. The proof of the theorem is complete. \square **Corollary 3.4.** (1) If $c + n = m$ for some $n \in \mathbb{Z}_+$, then

$$
L_{\widehat{H}_4}(1,(m,1),\mathbf{1}_{\alpha})=V_{\widehat{H}_4}(1,(m,1),\mathbf{1}_{\alpha})/U(\widehat{H}_4)u^1
$$

is irreducible, where

$$
u^{1} = \sum_{\lambda \in P(n)} (a_{\lambda}c(-\lambda)b(m) + \sum_{i=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_{i}, \lambda_{i}}c(-\lambda \setminus \lambda_{i})b(-\lambda_{i}))\mathbf{1}_{\alpha},
$$

satisfying (3.4) and (3.5) ;

(2) If $c - n = m$ for some $n \in \mathbb{Z}_+$, then

$$
L_{\hat{H}_4}(1,(m,1),\mathbf{1}_{\alpha})=V_{\hat{H}_4}(1,(m,1),\mathbf{1}_{\alpha})/U(\hat{H}_4)u^1
$$

is irreducible, where

$$
u^{1} = \sum_{\lambda \in P(n)} \sum_{i=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_{i}, \lambda_{i}} c(-\lambda \setminus \lambda_{i}) a(-\lambda_{i})) \mathbf{1}_{\alpha},
$$

satisfying (3.6) and (3.7) .

Proof: We only prove (1), since the proof of (2) is quite similar. Let

$$
V' = U(\widehat{H}_4)u^1.
$$

By Theorem 3.3 (ii), all the homogeneous singular vectors

$$
u^{k} = \left[\sum_{\lambda \in P(n)} (a_{\lambda}c(-\lambda)b(m) + \sum_{i=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_{i}, \lambda_{i}}c(-\lambda \setminus \lambda_{i})b(-\lambda_{i}))\right]^{k} \mathbf{1}_{\alpha}, \ k = 1, 2, \cdots
$$

lie in V'. Now let U/V' be a proper submodule of $V_{\hat{H}_4}(1,(m,1), 1_\alpha)/V'$, where U is a proper submodule of $V_{\hat{H}_4}(1,(m,1), \mathbf{1}_{\alpha})$. Recall that

$$
V_{\widehat{H}_4}(1,(m,1),\mathbf{1}_{\alpha})=\bigoplus_{j=1}^{\infty}V_{\widehat{H}_4}(1,(m,1),\mathbf{1}_{\alpha})_j.
$$

Let

$$
v = \sum_{i} v_i \in U,
$$

where $0 \neq v_i \in V_{\widehat{H}_4}(1, (m, 1), \mathbf{1}_{\alpha})_i$, be such that $|v| = \sum_i$ i i is the minimal. Then

$$
\widehat{H}_4((m,1), +)v_i = 0,
$$

for each i, where $\hat{H}_4((m, 1), +)$ is the same as (3.3). By Theorem 3.3 (ii), $v_i \in V'$. So $U = V'$. This proves that $V_{\hat{H}_4}(1, (m, 1), \mathbf{1}_{\alpha})/V'$ is irreducible.

Finally we introduce the so-called third class of generalized Verma type modules of \widehat{H}_4 . For $m \in \mathbb{Z}$, let $\widehat{H}_4(m, 0)$ be the subalgebra of \widehat{H}_4 defined by (3.1) for the case $\epsilon = 0$, that is,

$$
\widehat{H}_4(m,0)=(\bigoplus_{n\geq -m}\mathbb{C}a\otimes t^n)\bigoplus(\bigoplus_{n\geq m}\mathbb{C}b\otimes t^n)\bigoplus((\mathbb{C}c\bigoplus\mathbb{C}d)\otimes\mathbb{C}[t])\bigoplus\mathbb{C}\mathbf{k}.
$$

Set

$$
\widehat{H}_4(m,0)^{+} = (\bigoplus_{n \ge -m+1} \mathbb{C}a \otimes t^n) \bigoplus (\bigoplus_{n \ge m+1} \mathbb{C}b \otimes t^n) \bigoplus ((\mathbb{C}c \bigoplus \mathbb{C}d) \otimes \mathbb{C}t[t]).
$$
 (3.8)

For $\tilde{\lambda}, \tilde{\mu}, \tilde{\nu} \in \mathbb{C}$, let

$$
V(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, m) = \bigoplus_{n \in \mathbb{Z}} \mathbb{C}v_n
$$

be an infinite dimensional module of $\widehat{H}_4(m, 0)$ defined by

$$
\widehat{H}_4(m,0)^+v_n = 0, \ \mathbf{k} \cdot v_n = v_n \quad n \in \mathbb{Z},
$$

$$
c(0)v_n = (\tilde{\mu} + m)v_n, \ a(-m)v_n = -\tilde{\mu}v_{n+1},
$$

$$
d(0)v_n = (\tilde{\lambda} + n)v_n, \ b(m)v_n = (\tilde{\lambda} + \tilde{\nu} + n)v_{n-1}.
$$

The following lemma is obvious.

Lemma 3.5. $V(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, m)$ is an irreducible $\hat{H}_4(m, 0)$ -module if and only if $\tilde{\mu} \neq 0$ and $\lambda + \tilde{\nu} \notin \mathbb{Z}$.

Let

$$
V_{\widehat{H}_4}(\widetilde{\lambda}, \widetilde{\mu}, \widetilde{\nu}, m) = U(\widehat{H}_4) \otimes_{U(\widehat{H}_4(m, 0))} V(\widetilde{\lambda}, \widetilde{\mu}, \widetilde{\nu}, m)
$$

be the induced \widehat{H}_4 -module.

For $n \in \mathbb{Z}$, define

$$
\deg a(-n - m) = n, \ \deg b(-n + m) = n, \deg c(-n) = n, \ \deg d(-n) = n, \ \ \deg k = 0.
$$

Then $U(\widehat{H}_4)$ is Z-graded. So we get an N-gradation of $V_{\widehat{H}_4}(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, m)$ by defining deg $v_n =$ 0 for $n \in \mathbb{Z}$. A homogeneous vector u in $V_{\hat{H}_4}(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, m)$ is called a homogeneous singular vector if

$$
\hat{H}_4(m,0)^+(u) = 0.
$$

Similar to Theorem 3.3, we have the following last main result of this section.

Theorem 3.6. Let $\tilde{\lambda}, \tilde{\mu}, \tilde{\nu} \in \mathbb{C}$ be such that $\tilde{\lambda} + \tilde{\nu} \notin \mathbb{Z}$, $\tilde{\mu} \neq 0$. Then the \hat{H}_4 -module $V_{\widehat{H}_{4}}(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, m)$ is irreducible if and only if $\tilde{\mu} + n \neq 0$ for all $n \in \mathbb{Z}$. Furthermore, we have

(i) If $\tilde{\mu} + n = 0$ for some $n \in \mathbb{Z}_+$, then all the linearly independent homogeneous singular vectors are

$$
u^{(k,j)} = \left[\sum_{\lambda \in P(n)} (a_{\lambda}c(-\lambda)b(m) + \sum_{i=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_i, \lambda_i}c(-\lambda \setminus \lambda_i)b(-\lambda_i+m))\right]^{k}v_j, \ j \in \mathbb{Z}, \ k = 1, 2, \cdots
$$

satisfying

$$
a_{\lambda}q_i\lambda_i - b_{\lambda\setminus\lambda_i,\lambda_i} = 0, \ i = 1,2,\cdots,k_{\lambda},\tag{3.9}
$$

$$
b_{\lambda \setminus \lambda_i, \lambda_i}(\tilde{\mu} + \lambda_i) + \sum_{j=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0; \qquad (3.10)
$$

The submodules generated by $u^{(k,i)}$ and $u^{(k,j)}$ for $i, j \in \mathbb{Z}$, $i \neq j$ are isomorphic to each other.

(ii) If $\tilde{\mu} - n = 0$ for some $n \in \mathbb{Z}_+$, then all the homogeneous singular vectors are

$$
u^{(k,j)} = \left[\sum_{\lambda \in P(n)} \sum_{i=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_i, \lambda_i} c(-\lambda \setminus \lambda_i) a(-\lambda_i)\right]^k v_j, \ j \in \mathbb{Z}, \ k = 1, 2, \dots
$$

satisfying

$$
q_i \lambda_i c_{\lambda \lambda_j, \lambda_j} + c_{\lambda \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0, \ i, j = 1, 2, \cdots, k_\lambda, i \neq j,
$$
 (3.11)

$$
c_{\lambda \setminus \lambda_i, \lambda_i}(-\tilde{\mu} + \lambda_i) - \sum_{j=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_i, \lambda_j, \lambda_i + \lambda_j} = 0, \ i = 1, 2, \cdots, k_{\lambda}, \tag{3.12}
$$

and the submodules generated by $u^{(k,i)}$ and $u^{(k,j)}$ for $i, j \in \mathbb{Z}$, $i \neq j$ are isomorphic to each other, where in (i) and (ii), $P(n)$ is the set of all partitions of weight n,

$$
\lambda^{(k)} = (\lambda, \lambda, \cdots, \lambda) \in \mathbb{Z}_+^k, \ \lambda = (\lambda_1^{(q_1)}, \ \lambda_2^{(q_2)}, \cdots, \lambda_{k_\lambda}^{(q_{k_\lambda})})
$$

such that $\sum_{i=1}^{k_{\lambda}} q_i \lambda_i = n$, and $\lambda \setminus \lambda_i = (\lambda_1^{(q_1)})$ $\lambda_1^{(q_1)},\cdots,\lambda_i^{(q_i-1)},\cdots,\lambda_{k_\lambda}^{(q_{k_\lambda})}$ $\lambda_{k_{\lambda}}^{(\Psi_{k_{\lambda}})}$ and if $\lambda = \lambda_i$, then $c(-\lambda \setminus \lambda_i)b(-\lambda_i) = b(-\lambda_i)$, $c(-\lambda \setminus \lambda_i)a(-\lambda_i) = a(-\lambda_i)$.

Corollary 3.7. (1) If $\tilde{\mu} + n = m$ for some $n \in \mathbb{Z}_+$, then

$$
L_{\widehat{H}_{4}}(\widetilde{\lambda}, \widetilde{\mu}, \widetilde{\nu}, m) = V_{\widehat{H}_{4}}(\widetilde{\lambda}, \widetilde{\mu}, \widetilde{\nu}, m) / U(\widehat{H}_{4}) u^{1}
$$

is irreducible, where

$$
u^{1} = \sum_{\lambda \in P(n)} (a_{\lambda}c(-\lambda)b(m) + \sum_{i=1}^{k_{\lambda}} b_{\lambda \setminus \lambda_{i}, \lambda_{i}}c(-\lambda \setminus \lambda_{i})b(-\lambda_{i}))v_{0},
$$

satisfying (3.9) and (3.10) ;

(2) If $\tilde{\mu} - n = m$ for some $n \in \mathbb{Z}_+$, then

$$
L_{\widehat{H}_4}(\widetilde{\lambda}, \widetilde{\mu}, \widetilde{\nu}, m) = V_{\widehat{H}_4}(\widetilde{\lambda}, \widetilde{\mu}, \widetilde{\nu}, m) / U(\widehat{H}_4) u^1
$$

is irreducible, where

$$
u^{1} = \sum_{\lambda \in P(n)} \sum_{i=1}^{k_{\lambda}} c_{\lambda \setminus \lambda_{i}, \lambda_{i}} c(-\lambda \setminus \lambda_{i}) a(-\lambda_{i})) v_{0},
$$

satisfying (3.11) and (3.12) .

4 Extension of the vertex operator algebra $V_{\widehat{H}_{4}}(1,0)$

Set

$$
\mathfrak{h}=\mathbb{C}c(-1)\mathbf{1}\oplus \mathbb{C}d(-1)\mathbf{1}.
$$

Then

$$
L(n)h = \delta_{n,0}h, \quad h(n)h' = (h, h')\delta_{n,1}\mathbf{1} \quad \text{for } n \in \mathbb{Z}_+, \quad h, h' \in \mathfrak{h}
$$

and $($, $)$ restricted to $\mathfrak h$ is a non-degenerate symmetric bilinear form. Thus we can identify h with its dual h^{*}. One can easily see that for any $h \in \mathfrak{h}$, $h(0)$ acts semisimply on $V_{\hat{H}_4}(1,0)$. So

$$
V_{\widehat{H}_4}(1,0) = \bigoplus_{\alpha \in \mathfrak{h}} V_{\widehat{H}_4}(1,0)^{(0,\alpha)},
$$

where

$$
V_{\widehat{H}_4}(1,0)^{(0,\alpha)} = \{ v \in V_{\widehat{H}_4}(1,0) | h(0)v = (\alpha, h)v \text{ for } h \in \mathfrak{h} \}.
$$

Set

$$
P = \{ \alpha \in \mathfrak{h} | V_{\widehat{H}_4}(1,0)^{(0,\alpha)} \neq 0 \}
$$

According to $[22]$, P equipped with the bilinear form $($, $)$ is a lattice. Then

 $P = \mathbb{Z}c$

Set

$$
L=\mathbb{Z} c\oplus \mathbb{Z} d
$$

Then for any $\alpha \in L$, $\alpha(0)$ acting on $V_{\hat{H}_4}(1,0)$ has only integral eigenvalues. This amounts to $L \subset P^0$, where

 $P^0 = \{ h \in \mathfrak{h} | (h, \alpha) \in \mathbb{Z} \text{ for } \alpha \in P \}$

Let $\epsilon: L \times L \rightarrow {\pm 1}$ be a 2-cocycle on L such that

$$
\epsilon(c,c) = 1, \quad \epsilon(c,d) = -1, \quad \epsilon(d,d) = 1,
$$

and

$$
\epsilon(\alpha,\beta)\epsilon(\beta,\alpha)^{-1} = (-1)^{(\alpha,\beta)+(\alpha,\alpha)(\beta,\beta)} = (-1)^{(\alpha,\beta)} \text{ for } \alpha,\beta \in L.
$$

For $\alpha \in L$, let

$$
E^{\pm}(\alpha, z) = exp\left(\sum_{n=1}^{\infty} \frac{\alpha(\pm n)}{\pm n} z^{\mp n}\right), \quad \Delta(\alpha, z) = z^{\alpha(0)} E^{+}(-\alpha, -z).
$$

Let $(W, Y(\cdot, z))$ be any (irreducible) weak $V_{\hat{H}_4}(1, 0)$ -module, then by Proposition 2.9 of [22], $(W^{(\alpha)}, Y_{\alpha}(\cdot, z)) = (W, Y(\Delta(\alpha, z), z))$ carries the structure of an (irreducible) weak $V_{\hat{H}_4}(1,0)$ -module. We expand $Y_{\alpha}(u,z)$ for $u \in V_{\hat{H}_4}(1,0)$ as follows:

$$
Y_{\alpha}(u,z)) = \sum_{n \in \mathbb{Z}} u_{(\alpha,n)} z^{-n-1}.
$$

We have the following result.

Proposition 4.1. (1) Let $k \in \mathbb{Z}$ and $\alpha = kc \in L$, then $V_{\hat{H}_4}(1,0)^{(\alpha)} \cong V_{\hat{H}_4}(1,\text{d})$ with $d = k$ and $V_{\widehat{H}_4}(1,0)^{(\alpha)}$ is a simple current of $V_{\widehat{H}_4}(1,0)$.

(2) Let $m_1, m_2 \in \mathbb{Z}, m_2 \neq 0$ and $\alpha = m_1c + m_2d$. Then $V_{\hat{H}_4}(1,0)^{(\alpha)}$ is isomorphic to the generalized Verma type module $V_{\hat{H}_4}(1, m_2, \mathbf{1}_{\alpha})$.

 $(3) V_{\hat{H}_4}(1,0)^{(\alpha)}$ is irreducible for each $\alpha \in L$ and for $\alpha, \beta \in L$, $V_{\hat{H}_4}(1,0)^{(\alpha)} \cong V_{\hat{H}_4}(1,0)^{(\beta)}$ if and only if $\alpha = \beta$.

Proof: For $\alpha = kc$, let W be any irreducible weak $V_{\hat{H}_4}(1,0)$ -module. By Lemma 3.1, $c(0)$ acts as a scalar on W. Then by Theorem 2.15 of [23] (see also Theorem 2.13 of [22]), $V_{\hat{H}_4}(1,0)^{(\alpha)}$ is a simple current of $V_{\hat{H}_4}(1,0)$. By Theorem 2.12, $W \cong V_{\hat{H}_4}(1,d)$ for some $d \in \mathbb{C}$. From the definition of $Y_{\alpha}(\cdot, z) = Y(\Delta(\alpha, z) \cdot, z)$, it is easy to check that $W^{(\alpha)} \cong V_{\widehat{H}_4}(1, d+k)$. Therefore (1) holds.

For (2), by the definition of Y_α , we have

$$
Y_{\alpha}(a(-1)\mathbf{1},z)\mathbf{1} = z^{m_2}Y(a(-1)\mathbf{1},z)\mathbf{1}, Y_{\alpha}(b(-1)\mathbf{1},z)\mathbf{1} = z^{-m_2}Y(b(-1)\mathbf{1},z)\mathbf{1},
$$

\n
$$
Y_{\alpha}(c(-1)\mathbf{1},z)\mathbf{1} = Y(c(-1)\mathbf{1} + z^{-1}m_2\mathbf{1},z)\mathbf{1},
$$

\n
$$
Y_{\alpha}(d(-1)\mathbf{1},z)\mathbf{1} = Y(d(-1)\mathbf{1} + z^{-1}m_1\mathbf{1},z)\mathbf{1},
$$

So we have

$$
(a(-1)\mathbf{1})_{(\alpha,n)}\mathbf{1} = 0, \quad (b(-1)\mathbf{1})_{(\alpha,m)}\mathbf{1} = 0, \text{ for } n \ge -m_2, m \ge m_2,
$$

$$
(c(-1)\mathbf{1})_{(\alpha,n)}\mathbf{1} = (d(-1)\mathbf{1})_{(\alpha,n)}\mathbf{1} = 0, \text{ for } n \ge 1,
$$

$$
(c(-1)\mathbf{1})_{(\alpha,0)}\mathbf{1} = m_2\mathbf{1}, \quad (d(-1)\mathbf{1})_{(\alpha,0)}\mathbf{1} = m_1\mathbf{1}.
$$

Therefore as an \hat{H}_4 -module, $V_{\hat{H}_4}(1,0)^{(\alpha)}$ is a quotient of the generalized Verma type module $V_{\hat{H}_4}(1,m_2,\mathbf{1}_{\alpha})$. By Theorem 3.2, $V_{\hat{H}_4}(1,0)^{(\alpha)}$ is isomorphic to $V_{\hat{H}_4}(1,m_2,\mathbf{1}_{\alpha})$.

(3) follows from Theorem 3.2 and (1)-(2). \square

Furthermore, we have

Proposition 4.2. Let $\alpha = m_1c + m_2d \in L$, $\beta = n_1c + n_2d \in \mathfrak{h}^*$ and $m \in \mathbb{Z}$. Then (1)

$$
V_{\hat{H}_4}(1, (m, 1), \mathbf{1}_{\beta})^{(\alpha)} \cong V_{\hat{H}_4}(1, (m + m_2, 1), \mathbf{1}_{\alpha + \beta}),
$$

$$
L_{\hat{H}_4}(1, (m, 1), \mathbf{1}_{\beta})^{(\alpha)} \cong L_{\hat{H}_4}(1, (m + m_2, 1), \mathbf{1}_{\alpha + \beta}).
$$

(2) Let $\tilde{\lambda}, \tilde{\mu}, \tilde{\nu} \in \mathbb{C}$ be such that $\tilde{\lambda} + \tilde{\nu} \notin \mathbb{Z}, \tilde{\mu} \neq 0$. Then

$$
V_{\hat{H}_4}(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, m)^{(\alpha)} \cong V_{\hat{H}_4}(\tilde{\lambda} + m_1, \tilde{\mu}, \tilde{\nu}, m + m_2),
$$

$$
L_{\hat{H}_4}(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, m)^{(\alpha)} \cong L_{\hat{H}_4}(\tilde{\lambda} + m_1, \tilde{\mu}, \tilde{\nu}, m + m_2).
$$

For convenience, we give the following definition.

Definition 4.3. Let F be the category of irreducible restricted \hat{H}_4 -modules W of level 1 such that $d(0)$ is semisimple, and there exists an integer m such that $h(n)$, $a(-m+n)$ and $b(m + n)$ for $n \geq 1$, $h \in \mathfrak{h}$ are locally nilpotent on W.

The following lemma is easy to prove.

Lemma 4.4. Let $W \in \mathcal{F}$. Then there exist $m \in \mathbb{Z}$ and a non-zero element w in W such that

$$
h(n)w = a(-m+n)w = b(m+n)w = 0, \forall n \ge 1, \forall h \in \mathfrak{h}.
$$

We are now in a position to state the first main result of this section.

Theorem 4.5. Let $W \in \mathcal{F}$ be such that $c(0)$ acts on W as \mathbf{c} . Denote $\alpha = \mathrm{d}c + \mathrm{d}c$, then W is isomorphic to one of the following irreducible modules:

$$
V_{\widehat{H}_{4}}(1,m,\boldsymbol{1}_{\alpha}),V_{\widehat{H}_{4}}(1,(m,1),\boldsymbol{1}_{\alpha}),\ V_{\widehat{H}_{4}}(\tilde{\lambda},\tilde{\mu},\tilde{\nu},m)\ L_{\widehat{H}_{4}}(1,(m,1),\boldsymbol{1}_{\alpha}),\ L_{\widehat{H}_{4}}(\tilde{\lambda},\tilde{\mu},\tilde{\nu},m)
$$

given in Theorems 3.2, 3.3 and 3.6, Corollary 3.4 and Corollary 3.7 respectively.

Proof: Let $W \in \mathcal{F}$, by the definition of \mathcal{F} and Lemma 4.4, there exist an integer m and a non-zero element u in W such that

$$
c(0)u = \mathbf{c}u, \ d(0)u = \mathbf{d}u,
$$

for some $d \in \mathbb{C}$, and

$$
h(n)u = a(-m+n)u = b(m+n)u = 0, \forall n \ge 1, \forall h \in \mathfrak{h}.
$$

We first assume that $m = 0$. Then

$$
h(n)u = a(n)u = b(n)u = 0, \forall n \ge 1, \forall h \in \mathfrak{h}.
$$
\n
$$
(4.1)
$$

Let W^0 be the subspace of W consisting of elements satisfying (4.1). Then W^0 is a module of H_4 . Since W is irreducible, it follows that W^0 is an irreducible H_4 -module. By Theorem 2.6, Theorems 3.2, 3.3 and 3.6, Corollaries 3.4 and 3.7, we have

$$
W \cong V_{\hat{H}_4}(1, d), \text{ if } c = 0,
$$

$$
W \cong V_{\hat{H}_4}(1, (0, 1), \mathbf{1}_{\alpha}) \text{ or } V_{\hat{H}_4}(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, 0), \text{ if } 0 \neq c \notin \mathbb{Z},
$$

and

Next we assume that $m \neq 0$. Note that W can be viewed as a module for the vertex algebra $V_{\hat{H}_4}(1,0)$. Let (\cdot, z) be the corresponding operator. Let $\beta = -md$. By the proof of Proposition 4.1, $(W^{(\beta)} = W, Y_{\beta}(\cdot, z))$ is a module of $V_{\widehat{H}_{4}}(1,0)$ satisfying

 $W \cong L_{\widehat{H}_4}(1,(0,1), \mathbf{1}_{\alpha})$ or $L_{\widehat{H}_4}(\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, 0)$, if $0 \neq \alpha \in \mathbb{Z}$.

$$
(x(-1)\mathbf{1})_{(\beta,n)}u=0, \ \forall \ x\in H_4, \ \forall \ n\in \mathbb{Z}_+,
$$

$$
(c(-1){\bf 1})_{(\beta,0)}u=(c-m)u,\ (d(-1){\bf 1})_{(\beta,0)}u={\rm d} u.
$$

Therefore $W^{(\beta)}$ is isomorphic to one of the following module:

 $V_{\widehat{H}_{4}}(1,0,1_{\alpha+\beta}),\ V_{\widehat{H}_{4}}(1,(0,1),1_{\alpha+\beta}),\ V_{\widehat{H}_{4}}(\tilde{\lambda},\tilde{\mu},\tilde{\nu},0),\ L_{\widehat{H}_{4}}(1,(0,1),1_{\alpha+\beta}),\ L_{\widehat{H}_{4}}(\tilde{\lambda},\tilde{\mu},\tilde{\nu},0).$ This means that W is isomorphic to one of the irreducible modules:

$$
V_{\hat{H}_4}(1,m,\mathbf{1}_{\alpha}),\ V_{\hat{H}_4}(1,(m,1),\mathbf{1}_{\alpha}),\ V_{\hat{H}_4}(\tilde{\lambda},\tilde{\mu},\tilde{\nu},m),\ L_{\hat{H}_4}(1,(m,1),\mathbf{1}_{\alpha}),\ L_{\hat{H}_4}(\tilde{\lambda},\tilde{\mu},\tilde{\nu},m).
$$

Set

$$
V_{\widehat{H}_4}(1,0)[L] = \mathbb{C}[L] \otimes V_{\widehat{H}_4}(1,0).
$$

For $\alpha, \beta \in L$, $u, v \in V_{\hat{H}_4}(1,0)$, we define a vertex operator map Y on $V_{\hat{H}_4}(1,0)[L]$ by $Y(e^{\alpha} \otimes u, z)(e^{\beta} \otimes v) = \epsilon(\alpha, \beta)e^{\alpha+\beta} \otimes z^{(\alpha, \beta)}E^{-}(-\alpha, z)Y(\Delta(\beta, z)u, z)E^{+}(-\alpha, z)(-z)^{\alpha(0)}v.$ From [22], we have

Theorem 4.6. $V_{\widehat{H}_4}(1,0)[L]$ equipped with the vertex operator map Y is a vertex algebra.

It is easy to see that for each $\alpha \in L$, $\mathbb{C}e^{\alpha} \otimes V_{\widehat{H}_4}(1,0)$ is isomorphic to $V_{\widehat{H}_4}(1,0)^{(\alpha)}$ as $V_{\widehat{H}_{4}}(1,0)$ -modules. So we also denote

$$
V_{\hat{H}_4}(1,0)^{(\alpha)} = \mathbb{C}e^{\alpha} \otimes V_{\hat{H}_4}(1,0).
$$

It is obvious that $L(0)$ has integral eigenvalues on $V_{\hat{H}_4}(1,0)[L]$. In fact, for

$$
v = e^{k_1 c + k_2 d} \otimes d(-m_{11}) \cdots d(-m_{1i_1}) c(-m_{21}) \cdots c(-m_{2i_2}) a(-m_{31})
$$

$$
\cdots a(-m_{3i_3}) b(-m_{41}) \cdots b(-m_{4i_4}) \mathbf{1},
$$

$$
L(0)v = (k_1 k_2 + \sum_{k=1}^4 \sum_{j=1}^{i_k} m_{kj} + k_2 (i_3 - i_4)) v.
$$

Moreover, the same proof of Theorem 3.5 in [11] gives that for $v \in V_{\hat{H}_4}(1,0)[L]$,

$$
[L(-1), Y(v, z)] = \frac{d}{dz}Y(v, z).
$$

Thus $V_{\hat{H}_4}(1,0)[L]$ is a vertex operator algebra except the two grading restrictions.

Theorem 4.7. Let W be a module for the vertex algebra $V_{\hat{H}_4}(1,0)$ such that for each $\alpha \in L$, $\alpha(0)$ acts on W semisimply with only integral eigenvalues. Set

$$
W[L] = \mathbb{C}[L] \otimes W.
$$

For $\alpha, \beta \in L$, $v \in V_{\hat{H}_4}(1,0)$, $w \in W$, we define

$$
Y_{W[L]}(e^{\alpha}\otimes v,z)(e^{\beta}\otimes w)=\epsilon(\alpha,\beta)e^{\alpha+\beta}\otimes z^{(\alpha,\beta)}E^{-}(-\alpha,z)Y_{W}(\triangle(\beta,z)v,z)E^{+}(-\alpha,z)(-z)^{\alpha(0)}w.
$$

Then $W[L]$ carries the structure of a $V_{\hat{H}_4}(1,0)[L]$ -module. Furthermore, if W is irreducible, then $W[L]$ is irreducible. In particular, $V_{\hat{H}_4}(1,0)[L]$ is a simple vertex operator algebra.

Lemma 4.8. Let U be an irreducible $V_{\hat{H}_4}(1,0)[L]$ -module. Then

(1) There exists an element w in U such that for all $\alpha \in L$ and $n \geq 1$,

$$
\alpha(n)w=0.
$$

(2) For $\alpha \in L$, $\alpha(0)$ acts on U semisimply with only integral eigenvalues and all $\alpha(n)$ for $n \geq 1$ are locally nilpotent.

Proof: Note that $V_{\hat{H}_4}(1,0)[L]$ contains the lattice vertex algebra V_L and $V_{\hat{H}_4}(1,0)[L]$ is simple. By Lemma 3.15 in [12], there exists an element w in U such that for all $\alpha \in L$, and $n \geq 1$,

$$
\alpha(n)w=0.
$$

Let α be any element in L. Since $V_{\hat{H}_4}(1,0)[L]$ is simple, similar to the argument in [9] (see also [25]), from [10], we have $Y(e^{\alpha}, z)w \neq 0$. Then there exists $k \in \mathbb{Z}$ such that $e^{\alpha}(k)w \neq 0$ and $e^{\alpha}(n)w = 0$ for $n > k$. By the fact that $\frac{d}{dz}Y(e^{\alpha}, z) = \alpha(z)Y(e^{\alpha}, z)$, we have

$$
e^{\alpha}(k)\alpha(0)w = (-k-1)e^{\alpha}(k)w.
$$

Since

$$
[\alpha(n), e^{\beta}(r)] = (\alpha, \beta)e^{\beta}(n+r),
$$

we have

$$
\alpha(0)e^{\alpha}(k)w = (-k - 1 + (\alpha, \alpha))e^{\alpha}(k)w.
$$

Note that $(\alpha, \alpha) \in \mathbb{Z}$, so $-k-1 + (\alpha, \alpha) \in \mathbb{Z}$. Then (2) follows from Lemma 4.4 and the fact that $e^{\alpha}(k)w$ generates W by operators $x(n), e^{\beta}(n)$, for $x \in H_4$, $\beta \in L$, and $[d, a] = a$, $[d, b] = -b.$

Theorem 4.9. Let W be an irreducible $V_{\hat{H}_4}(1,0)[L]$ -module, then $W \cong \mathbb{C}[L] \otimes U$, where U is an irreducible module of $V_{\hat{H}_4}(1,0)$ such that for $\alpha \in L$, $\alpha(0)$ acts on U semisimply with only integral eigenvalues and all $\alpha(n)$ with $n \geq 1$ are locally nilpotent.

Proof: By Theorem 3.16 of $[12]$, W viewed as a V_L -module is completely reducible and each simple weak module of V_L is $V_{L+\lambda}$, for some $\lambda \in L^{\circ}$, where L° is the dual of L and V_L is the vertex algebra associated with the even lattice L . Therefore there exist an element $u \in W$ and $\lambda \in L^{\circ}$ such that for $\alpha \in L$,

$$
\alpha(n)u = 0, \ \alpha(0)u = (\alpha, \lambda)u,
$$

and

$$
e^{\alpha}(-(\alpha,\lambda)-1)u \neq 0, e^{\alpha}(-(\alpha,\lambda)-1+n)u = 0,
$$

for $n \in \mathbb{Z}_+$. Since

$$
[\alpha(n), e^{\beta}(r)] = (\alpha, \beta)e^{\beta}(n+r),
$$

it follows that

$$
\alpha(n)e^{\alpha}(-(\alpha,\lambda)-1)u=0
$$

for $n \in \mathbb{Z}_+$ and

$$
\alpha(0)e^{\alpha}(-(\alpha,\lambda)-1)u = (\alpha,\alpha+\lambda)e^{\alpha}(-(\alpha,\lambda)-1)u.
$$

Let U be the $V_{\widehat{H}_4}(1,0)$ -module generated by u. Then $\mathbb{C}[L]\otimes U$ is a module of $V_{\widehat{H}_4}(1,0)[L]$. For $\alpha \in L$, let

$$
E_{\alpha} = E^-(\alpha, z)Y_W(e^{\alpha}, z)E^+(\alpha, z)(-z)^{-\alpha(0)}.
$$

Then as the proof of Theorem 3.10 in [25], we have

$$
\frac{d}{dz}E_{\alpha}=0,
$$

and

$$
E_{\alpha}E_{\beta} = \epsilon(\alpha, \beta)E_{\alpha+\beta}.
$$

Then it is easy to see that W is linearly spanned by $\{E_\alpha w \mid \alpha \in L, w \in U\}.$

Define $\psi: \mathbb{C}[L] \otimes U \to W$ by

$$
\psi(e^{\alpha} \otimes w) = E_{\alpha} \otimes w,
$$

for $\alpha \in L, w \in U$. Quite similar to the proof of Theorem 3.10 in [25], we have

$$
\psi(Y_{U[L]}(h(-1)\mathbf{1},z)(e^{\alpha} \otimes w)) = Y_W(h(-1)\mathbf{1},z)\psi(e^{\alpha} \otimes w),
$$

$$
\psi(Y_{U[L]}(e^{\alpha},z)(e^{\beta} \otimes w)) = Y_W(e^{\alpha},z)\psi(e^{\alpha} \otimes w),
$$

for $h \in \mathfrak{h}$, $\alpha, \beta \in L$, $w \in U$. Furthermore, for $\alpha = m_1c + m_2d \in L$, $w \in U$, we have

$$
Y_{U[L]}(a(-1)\mathbf{1}, z)(e^{\alpha} \otimes w) = z^{m_2} e^{\alpha} \otimes Y_{U[L]}(a(-1)\mathbf{1}, z)w,
$$

\n
$$
Y_W(a(-1)\mathbf{1}, z)E_{\alpha}w = z^{m_2} E_{\alpha} Y_W(a(-1)\mathbf{1}, z)w,
$$

\n
$$
Y_{U[L]}(b(-1)\mathbf{1}, z)(e^{\alpha} \otimes w) = z^{-m_2} e^{\alpha} \otimes Y_{U[L]}(b(-1)\mathbf{1}, z)w,
$$

\n
$$
Y_W(b(-1)\mathbf{1}, z)E_{\alpha}w = z^{-m_2} E_{\alpha} Y_W(b(-1)\mathbf{1}, z)w.
$$

So

$$
\psi(Y_{U[L]}(a(-1)\mathbf{1},z)(e^{\alpha} \otimes w)) = Y_W(a(-1)\mathbf{1},z)\psi(e^{\alpha} \otimes w),
$$

$$
\psi(Y_{U[L]}(b(-1)\mathbf{1},z)(e^{\alpha} \otimes w)) = Y_W(b(-1)\mathbf{1},z)\psi(e^{\alpha} \otimes w).
$$

This means that ψ is a module isomorphism. Hence $W \cong \mathbb{C}[L] \otimes U$. Since W is irreducible, it follows that U is irreducible. it follows that U is irreducible.

Acknowledgements We would like to thank the referee for valuable comments and suggestions.

References

- [1] Yixin Bao, Cuipo Jiang and Yufeng Pei, Representations of affine Nappi-Witten algebras (preprint)
- [2] K. Bardakci and M. B. Halpern, Phys. Rev. D3 (1971) 2493.
- [3] R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, Proc. Natl. Acad. Sci. USA **83** (1986) 3068-3071.
- [4] Y. Cheung, L. Freidel and K. Savvidy, Strings in gravimagnetic fields, Journal of High Energy Physics 054 (2004) 1-48.
- [5] G. DAppollonio and T. Quella, The abelian cosets of the Heisenberg group, Journal of High Energy Physics 045 (2007).
- [6] G. DAppollonio and T. Quella, The diagonal cosets of the Heisenberg group, Journal of High Energy Physics 060 (2008).
- [7] G. DAppollonio and E. Kiritsis, String interactions in gravitational wave backgrounds, Nucl. Phys. B 674 (2003), 80-170.
- [8] J. Distler, talk at the Strings 93 Conference, Berkeley, May 1993.
- [9] C. Dong, Vertex algebras associated with even lattices, J. Algebra 160 (1993), 245-265.
- [10] C. Dong and J. Lepowsky, Generalized Vertex Algebras and Relative Vertex Operators, Progress in Math. Vol. 112, Birkhäuser, Boston 1993.
- [11] C. Dong, H. Li and G. Mason, Simple currents and extensions of vertex operator algebras, Commun. Math. Phys. 180 (1996), 671-707.
- [12] C. Dong, H. Li and G. Mason, Regularity of rational vertex operator algebras, Advances. in Math. 132 (1997), 148-166.
- [13] B. Feigin and E. Frenkel, A new class of representations of affine Kac-Moody algebras, Russ. Math. Surv.43 (1988), 221-222.
- [14] J. M. Figueroa-O'Farrill and S. Stanciu, Nonsemisimple Sugawara constructions, Phys. Lett. B (1994) 327-340.
- [15] E. Frenkel and D. Ben-Zvi, Vertex algebras and algebraic curves, Mathematical Surveys and Monographs 88, AMS, 2001.
- [16] I. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the monster, Pure and Appl. Math. 134, Academic Press, Boston, 1988.
- [17] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, Duke Math. J. 66 (1992), 123-168.
- [18] V. Kac, Infinite-Dimensional Lie Algebras, (1990) 3rd ed. Cambridge, UK: Cambridge University Press.
- [19] E. Kiritsis and C. Kounnas, String propagation in gravitational wave backgrounds, Phys. Lett. B 594 (1994), 368-374.
- [20] J. Lepowsky, H. Li, Introduction to Vertex Operator Algebras and Their Representations, Progress in Math. 227, Birkhäuser, Boston, 2003.
- [21] H. Li, Local systems of vertex operators, vertex superalgebras and modules, J. Pure Appl. Alg. 109 (1996), 143-195.
- [22] H. Li, Certain extensions of vertex operator algebras of affine type, Commun. Math. Phys. **217** (2001), 653-696.
- [23] H. Li, The physics superselection principle in vertex operator algebra theory, J. $Alg.$ 196 (1997), 73-114.
- [24] H. Li, Extension of vertex operator algebras by a self-dual simple module, J. Alg. 187 (1997), 237-267.
- [25] H. Li and Q. Wang, On vertex operator algebras and their modules associated with even alttices, $arXiv:0802.0173v1$.
- [26] B. Lian, On the classification of simple vertex operator algebras, Commun. Math. Phys. 163 (1994), 307-357.
- [27] C. Nappi, E. Witten, Wess-Zumino-Witten model based on a nonsemisimple group, Phys. Rev. Lett. 23 (1993), 3751-3753.
- [28] R. V. Moody and A. Pianzola, Lie algebras with triangular decompositions, Canadian Mathematical Society Series of Monographs and Advanced Texts. John Wiley & Sons Inc., New York, 1995.
- [29] M. Szczesny, Wakimoto modules for twisted affine Lie algebras, Math. Res. Lett. 9 (2002), 433C448.
- [30] M. Willard, Lie Theory and Special Functions, Academic Press, Inc., New York, 1968.
- [31] E. Witten, Non-abelian bosonization in two-dimensions, Commun. Math. Phys. 92 (1984), 455-472.
- [32] M. Wakimoto, Fock representations of the affine Lie algebra $A_1^{(1)}$ $_1^{(1)}$, Commun. Math. Phys. 104 (1986), 605-609.
- [33] S. Wang, Construction of the vertex operator algebra associated to finite nondegenerate solvable Lie algebra, Acta Math. Sinica 48 (2005), 867-878.
- [34] W. Wang, Rationality of Virasoro vertex operator algebras, Internat. Math. Res. Notices 7 (1993), 197-211.
- [35] X. Xu, Introduction to Vertex Operator Superalgebras and Their Modules, Kluwer Academic Publishers, Dordrecht/Boston/London, 1998.
- [36] Y. Zhu, Modular invariance of characters of vertex operator algebras, J. Amer. Math. Soc. **9** (1996), 237-301.