MEASURABLE TIME-RESTRICTED SENSITIVITY

DOMENICO AIELLO, HANSHENG DIAO, ZHOU FAN, DANIEL O. KING, JESSICA LIN, AND CESAR E. SILVA

ABSTRACT. We develop two notions of time-restricted sensitivity to initial conditions for measurable dynamical systems, where the time before divergence of a pair of paths is at most an asymptotically logarithmic function of a measure of their initial distance. In the context of finite measure-preserving transformations on a compact space, we relate these notions to the metric entropy of the system. We examine one of these notions for classes of non-measurepreserving, nonsingular transformations.

1. INTRODUCTION

Sensitivity has been widely studied as a characterization of chaos for topological dynamical systems, see e.g. [BBC⁺92], [GW93], [AAB96]. Recently, sensitivity has been explored in the context of other measurable dynamical properties, such as weak mixing and entropy, for a finite measure-preserving transformation equipped with a metric of full support [ABC02], [HYW04], [CJ05]. More recently, in [JKL⁺08], the authors introduced a measure-theoretic version of sensitivity, invariant under measurable isomorphism, for nonsingular transformations. This has been further studied in [GII⁺08].

In broad terms, sensitivity asserts that for any point x in the space, there exists another arbitrarily close point y such that at some future positive time n = n(x, y)the points $T^n(x)$ and $T^n(y)$ are separated by some predetermined distance. (Let us call this n a sensitive time.) This simplest notion of sensitivity does not assert anything about the sensitive time other than its existence. A refinement of this definition, called strong sensitivity, was introduced in [ABC02], where for each point x the set of sensitive times n(x, y), for some y, is co-finite. Strong sensitivity was also studied in the measurable context in [JKL⁺08]. An alternative refinement of (topological) sensitivity was studied in [Moo07], where the set of sensitive times is required to be syndetic. After completing this paper we learned of [HLYar], where the authors study extensions of pairwise sensitivity of for finite invariant measures in topological dynamical systems.

In this paper, we are interested in placing a quantitative, asymptotic bound on the sensitive time, restricting the first sensitive time of a point x and a point in an ε -ball around x to be at most asymptotically logarithmic in the measure of that ε -ball. We develop two notions of measurable sensitivity that restrict the sensitive time in this way, show that these notions are related to positive metric entropy for finite measure-preserving systems, and explore one of the notions in the context

Date: June 15, 2018.

¹⁹⁹¹ Mathematics Subject Classification. Primary 37A05; Secondary 37F10.

Key words and phrases. Measure-preserving, ergodic, sensitive dependence.

of (non-measure-preserving) nonsingular transformations. An outstanding problem in nonsingular ergodic theory is the lack of a theory of entropy, see e.g. [DS09]; our definitions of restricted sensitivity are related to entropy in the finite measurepreserving, and can be formulated in the context of nonsingular transformations and be thought of as an approach to positive nonsingular entropy.

In Section 2, we define the notions of restricted sensitivity, extending $[JKL^+08]$, and restricted pairwise sensitivity, extending [CJ05]. We prove that under mild conditions restricted pairwise sensitivity implies restricted sensitivity, and we explore these notions for Bernoulli shifts. In Section 3, we consider the setting of finite measure-preserving transformations on a compact space and prove that restricted pairwise sensitivity implies positive metric entropy and positive metric entropy for a continuous, ergodic transformation implies restricted sensitivity. We explore a quantitative relationship between the entropy of a Bernoulli shift and the asymptotic bound on the sensitive time. In Section 4, we explore restricted sensitivity in the context of (non-measure-preserving) nonsingular transformations. We construct a class of nonsingular transformations, that includes type III (i.e., not admitting an equivalent σ -finite invariant measure) transformations, that are restricted sensitive as well as a class of nonsingular rank-one transformations that are not. It is wellknown that finite measure-preserving rank-one transformations have zero entropy; it would be interesting to know if all nonsingular rank-one transformations are not restrictive sensitive.

1.1. Acknowledgements. This paper is based on research by the Ergodic Theory group of the 2008 SMALL summer research project at Williams College. Support for the project was provided by National Science Foundation REU Grant DMS - 0353634 and the Bronfman Science Center of Williams College.

2. TIME-RESTRICTED NOTIONS OF SENSITIVITY

Throughout, let $(X, \mathcal{S}(X), \mu)$ denote a standard probability space. We also consider a metric d on X, and assume that d is measurable so that the d-balls are measurable. It will be convenient for us to consider two compatibility conditions between d and μ . Let us say that d is μ -compatible if all nonempty open d-balls have positive μ -measure. The topology of a μ -compatible metric is separable [JKL⁺08]. Let us say that d is μ -regular if for all $x \in X$, there exists c > 0 such that

$$c\mu(\overline{B_r}(x)) \le \mu(B_r(x))$$

for all r > 0. In particular, if all *d*-balls are μ -continuity sets, then *d* is μ -regular. Recall the following definitions of measurable sensitivity:

Definition 2.1. [JKL⁺08] A nonsingular dynamical system $(X, S(X), \mu, T)$ is said to be **measurably sensitive** if whenever a dynamical system $(X_1, S(X_1), \mu_1, T_1)$ is measurably isomorphic to $(X, S(X), \mu, T)$ and d is a μ_1 -supported metric on X_1 , then there exists $\delta > 0$ such that for all $x \in X_1$ and all $\varepsilon > 0$, there exists $n \in \mathbb{N}_0$ such that

$$\mu_1\{y \in B_{\varepsilon}(x) : d(T_1^n x, T_1^n y) > \delta\} > 0.$$

Definition 2.2. [CJ05] A measure-preserving transformation T on a probability space (X, μ) equipped with a metric d is **pairwise sensitive** if there exists $\delta > 0$

such that for μ^{\otimes^2} -a.e. $(x,y) \in X \times X$, there exists $n \in \mathbb{N}_0$ such that

$$d(T^n x, T^n y) > \delta.$$

In this paper, we study the following time-restricted modifications of these sensitivity notions:

Definition 2.3. A measurable, nonsingular transformation T on the probability space $(X, \mathcal{S}(X), \mu)$ equipped with the metric d is **restricted sensitive** if for a.e. $x \in X$, there exist $\delta > 0$ and a > 0 such that for every $\varepsilon > 0$, there exists $n \in \mathbb{N}_0$, $n \leq -a \log \mu B_{\varepsilon}(x)$, with

$$\mu \{ y \in B_{\varepsilon}(x) : d(T^n x, T^n y) > \delta \} > 0.$$

Definition 2.4. A measurable, nonsingular transformation T on the probability space $(X, \mathcal{S}(X), \mu)$ equipped with the metric d is **restricted pairwise sensitive** if there exists $\delta > 0$ and a > 0 such that for μ^{\otimes^2} -a.e. $(x, y) \in X \times X$, there exists $n \in \mathbb{N}_0, n \leq -a \log \mu B_{d(x,y)}(x)$ such that $d(T^n x, T^n y) > \delta$.

Let us call δ and a from these definitions a **sensitivity constant** and an **asymptotic rate**, respectively. For both Definitions 2.3 and 2.4, if δ is a sensitivity constant, then any $\delta' < \delta$ is also a sensitivity constant, and if a is an asymptotic rate, than any a' > a is also an asymptotic rate. Note that we allow the sensitivity constant and asymptotic rate to vary over $x \in X$ in Definition 2.3 but assume that they are constant across $x \in X$ in Definition 2.4. Finally, note that the condition $d(T^nx, T^ny) > \delta$ is true for n = 0 when $d(x, y) > \delta$, so that it suffices to check $\varepsilon \leq \delta$ in Definition 2.3 and pairs of points (x, y) with $d(x, y) \leq \delta$ in Definition 2.4.

Restricted pairwise sensitivity is a stronger notion than restricted sensitivity, in the following sense:

Proposition 2.5. Suppose d is μ -regular. If a transformation T is restricted pairwise sensitive, then T is restricted sensitive.

Proof. Let $\delta > 0$ and a > 0 be such that for a.e. $x \in X$, for a.e. $y \in X$, there exists $n \leq -a \log \mu B_{d(x,y)}(x)$ with $d(T^n x, T^n y) > \delta$. Consider any such $x \in X$, let c > 0 be such that $c\mu(\overline{B_r}(x)) \leq \mu(B_r(x))$ for all r > 0, and consider any $\varepsilon \leq \delta$. Let $\varepsilon_M = \sup\{\varepsilon' < \varepsilon : \mu(B_{\varepsilon'}(x)) < \mu(B_{\varepsilon}(x))\}$. If $\mu(B_{\varepsilon_M}(x)) = \mu(B_{\varepsilon}(x))$, then choose $\varepsilon' < \varepsilon_M$ sufficiently close to ε_M such that $\mu(B_{\varepsilon}(x) \setminus B_{\varepsilon'}(x)) > 0$ and $\mu(B_{\varepsilon'}(x)) = \mu(B_{\varepsilon}(x))$, otherwise if $\mu(B_{\varepsilon_M}(x)) < \mu(B_{\varepsilon}(x))$, then we must have $\mu(\overline{B_{\varepsilon_M}}(x)) = \mu(B_{\varepsilon}(x))$, and we may choose $\varepsilon' = \varepsilon_M$ so that $\mu(B_{\varepsilon}(x) \setminus B_{\varepsilon'}(x)) > 0$ and $\mu(B_{\varepsilon'}(x)) \geq c\mu(\overline{B_{\varepsilon'}}(x)) = \mu(B_{\varepsilon}(x))$ by the definition of c.

For a.e. $y \in B_{\varepsilon}(x) \setminus B_{\varepsilon'}(x)$, there exists $n \leq -a \log \mu B_{d(x,y)}(x)$ such that $d(T^nx, T^ny) > \delta$. We note that $\mu B_{d(x,y)}(x) \geq \mu B_{\varepsilon'}(x) \geq c\mu B_{\varepsilon}(x)$, so $n \leq -a \log c\mu B_{\varepsilon}(x) = -a \log \mu B_{\varepsilon}(x) - a \log c$. As we may choose the pairwise sensitivity constant δ so that $\mu B_{\varepsilon}(x) \leq \mu B_{\delta}(x) < 1$, we may take \tilde{a} so that $-a \log \mu B_{\varepsilon}(x) - a \log c \leq -\tilde{a} \log \mu B_{\varepsilon}(x)$ for all $\varepsilon \leq \delta$. Hence T is restricted sensitive with sensitivity constant δ and sensitivity function \tilde{a} .

As an application of these notions of restricted sensitivity and restricted pairwise sensitivity, let us consider the standard one-sided and two-sided Bernoulli shift transformations. **Example 2.6.** Consider the space $\Sigma_N^+ = \prod_{i=0}^{\infty} \{1, \ldots, N\}$ with its product σ -algebra and the probability measure $\mu = \bigotimes_{i=0}^{\infty} \mu'$, where μ' is a probability measure on $\{1, \ldots, N\}$ of full support. Consider the metric $d(\sigma, \tau) = 2^{-I(\sigma,\tau)}$ where $I(\sigma, \tau) = \min\{i \ge 0 : \sigma_i \ne \tau_i\}$. Let T be the one-sided Bernoulli shift transformation $T((\sigma_0, \sigma_1, \sigma_2, \ldots)) = (\sigma_1, \sigma_2, \sigma_3, \ldots)$. Let $p = \max_k \mu'(k), \ \delta = \frac{1}{2}$, and $a = -\frac{1}{\log p}$. Consider any two points $\sigma, \tau \in \Sigma_N^+$ and suppose that $I(\sigma, \tau) = n$. Then $\mu B_{d(\sigma,\tau)}(\sigma) \le p^n$, so $-a \log \mu B_{d(\sigma,\tau)}(\sigma) \ge n$. Since $d(T^n\sigma, T^n\tau) = 1 > \delta$, T is restricted pairwise sensitive.

Example 2.7. Consider the space $\Sigma_N = \prod_{i=-\infty}^{\infty} \{1, \ldots, N\}$ with its product σ -algebra and the probability measure $\mu = \bigotimes_{i=-\infty}^{\infty} \mu'$, where μ' is a probability measure on $\{1, \ldots, N\}$ of full support. Consider the metric $d(\sigma, \tau) = 2^{-I(\sigma, \tau)}$, where $I(\sigma, \tau) = \min\{|i| : \sigma_i \neq \tau_i\}$. Let T be the two-sided Bernoulli shift transformation such that $T(\sigma)_i = \sigma_{i+1}$. Let $p_k = \mu'(k)$ for each $k = 1, \ldots, N$, and consider any $\delta < 1$ and any a > 0. For any $\sigma \in \Sigma_N$, choose an integer $k_1 > 0$ such that $2^{-k_1} < \delta$ and let $P = \prod_{i=-k_1+1}^{k_1-1} p_{\sigma_i}$. Choose an integer $k_2 > k_1$ such that $2^{-(k_2+a\log P)} < \delta$. Consider any τ in the cylinder set $[\bar{\sigma}_{-k_1}\sigma_{-k_1+1}\ldots\sigma_{0}\ldots\sigma_{k_2}]$, where $\bar{\sigma}_{-k_1}$ is some symbol not equal to σ_{-k_1} . Then by the construction of k_1 and k_2 , for all $n \leq -a\log P = -a\log \mu B_{d(\sigma,\tau)}(\sigma)$, $d(T^n\sigma,T^n\tau) \leq \max(2^{-k_1},2^{-(k_2-n)}) \leq \max(2^{-k_1},2^{-(k_2+a\log P)}) < \delta$. Hence T is not restricted pairwise sensitive.

On the other hand, let $p = \max_k \sigma'(k)$, $\delta = \frac{1}{4}$, and $a = -\frac{1}{2\log p}$. Consider any $\sigma \in \Sigma_N$ and any ball $B(\sigma) = [\sigma_{-n} \dots \sigma_0 \dots \sigma_n]$. We have that $\mu B(\sigma) \leq p^{2n+1}$, so $n < -a \log \mu B(\sigma)$. Since $d(T^n \sigma, T^n \tau) = \frac{1}{2} > \delta$ for all $\tau \in [\sigma_{-n} \dots \sigma_n \overline{\sigma}_{n+1}] \subset [\sigma_{-n} \dots \sigma_n]$ where $\overline{\sigma}_{n+1}$ is some symbol not equal to σ_{n+1} , T is restricted sensitive.

Thus we have examples of transformations that are restricted pairwise sensitive and transformations that are restricted sensitive but not restricted pairwise sensitive. One can also easily construct transformations that are not restricted sensitive but that are measurably sensitive according to Definition 2.1; examples of such transformations include the rank-one cutting and stacking transformations that we will study in Section 4.

3. Sensitivity and Entropy for Measure-Preserving Transformations

In the setting of finite measure-preserving transformations on a compact metric space, the notion of metric entropy measures the rate at which a transformation disorganizes the space. As measurable time-restricted notions of sensitivity convey that points separate from one another rapidly on a local level, it is natural to explore the connection between notions of measurable time-restricted sensitivity and the metric entropy of a dynamical system.

The following theorem and its immediate corollary show that, under mild conditions on μ and d, restricted pairwise sensitivity implies positive metric entropy in the context of finite measure-preserving transformations on a compact metric space:

Theorem 3.1. Suppose μ is a nonatomic probability measure on X and d is μ compatible and μ -regular. Let T be a measure-preserving transformation on X. Let $\mathcal{A} = \{A_1, \ldots, A_k\}$ be a partition of X such that diam $A_i < \delta$ for all i, and let

 $h_{\mu}(T, \mathcal{A})$ be the metric entropy of T with respect to \mathcal{A} . If T is restricted pairwise sensitive with asymptotic rate a, then $h_{\mu}(T, \mathcal{A}) \geq \frac{1}{a}$.

Proof. Suppose T is restricted pairwise sensitive with sensitivity constant δ and asymptotic rate a. Take $x \in X$ such that for a.e. $y \in X$, there exists $n \leq -a \log \mu B_{d(x,y)}(x)$ with $d(T^n x, T^n y) > \delta$. Let $C_n(x)$ denote the element of the partition $\bigvee_{i=0}^{n} T^{-i} \mathcal{A}$ containing x. If $y \in C_n(x)$, then $T^i x$ and $T^i y$ are in the same element of \mathcal{A} for all $i \leq n$, so $d(T^i x, T^i y) < \delta$ for all $i \leq n$.

Suppose c > 0 is such that $e^{-c}\mu(\overline{B_r}(x)) \leq \mu(B_r(x))$ for all r > 0. (The existence of this c is given by μ -regularity.) Since μ is nonatomic, $\mu(B_r(x)) \to 0$ as $r \to 0$. For each integer $n \geq 1$, we may take ε_n such that $e^{-(n-1)c} > \mu(B_{\varepsilon_n}(x)) \geq e^{-nc}$, for otherwise there must exist ε' with $\mu(\overline{B_{\varepsilon'}}(x)) \geq e^{-(n-1)c}$ and $\mu(B_{\varepsilon'}(x)) < e^{-nc}$, contradicting the definition of c. If $y \notin B_{\varepsilon_{\lfloor \frac{n}{ac} \rfloor}}(x)$, then $\mu B_{d(x,y)}(x) \geq B_{\varepsilon_{\lfloor \frac{n}{ac} \rfloor}}(x) \geq e^{-\lfloor \frac{n}{ac} \rfloor c}$, so for a.e. $y \notin B_{\varepsilon_{\lfloor \frac{n}{ac} \rfloor}}(x)$, there exists $i \leq -a \log \mu B_{d(x,y)}(x) \leq n$ such that $d(T^ix, T^iy) > \delta$, and so $y \notin C_n(x)$. Hence $C_n(x) \subset B_{\varepsilon_{\lfloor \frac{n}{ac} \rfloor}}(x) \mod \mu$, so $\mu C_n(x) \leq \mu B_{\varepsilon_{\lfloor \frac{n}{ac} \rfloor}}(x) < e^{(\lfloor \frac{n}{ac} \rfloor - 1)c}$. Letting $h(x) = \liminf_{n \to \infty} -\frac{1}{n+1} \log \mu C_n(x)$, we have

$$h(x) \ge \liminf_{n \to \infty} -\frac{c}{n+1} \left(\left\lfloor \frac{n}{ac} \right\rfloor - 1 \right) = \frac{1}{a}.$$

This holds for a.e. $x \in X$, so by Fatou's lemma,

$$h_{\mu}(T,\mathcal{A}) = \lim_{n \to \infty} \int -\frac{1}{n+1} \log \mu C_n(x) \, d\mu \ge \int h(x) \, d\mu \ge \frac{1}{a}.$$

Corollary 3.2. Suppose (X, d) is compact, μ is nonatomic, and d is μ -compatible and μ -regular. Let T be a measure-preserving transformation on X. If T is restricted pairwise sensitive with asymptotic rate a, then $h_{\mu}(T) \geq \frac{1}{a}$.

Example 2.7 shows that a converse to this theorem is not true, for the two-sided Bernoulli shift has positive entropy but is not restricted pairwise sensitive. The following result shows, however, that the implication in this direction is true if we replace restricted pairwise sensitivity with restricted sensitivity, under assumptions of ergodicity and continuity of T:

Theorem 3.3. Suppose (X, d) is compact and μ is nonatomic. Let T be a continuous, ergodic, measure-preserving transformation on X. If $h_{\mu}(T) > \frac{1}{a}$, then T is restricted sensitive with asymptotic rate a for all $x \in X$.

Proof. Suppose T is not restricted sensitive with asymptotic rate a over all $x \in X$. Then there exists a positive measure set A such that for any $x \in A$ and $\delta > 0$, there exists $\varepsilon(\delta) \leq \delta$ such that $\mu \{y \in B_{\varepsilon(\delta)}(x) : d(T^n x, T^n y) > \delta\} = 0$ for all $n \leq -a \log \mu B_{\varepsilon(\delta)}(x)$. For each n, let

$$C(x, n, \delta) = \{ y \in X : d(T^i x, T^i y) \le \delta \text{ for all } 0 \le i < n \}.$$

Then $B_{\varepsilon(\delta)}(x) \subset C(x, \left[-a \log \mu B_{\varepsilon(\delta)}(x)\right], \delta) \mod \mu$, so

$$\mu B_{\varepsilon(\delta)}(x) \le \mu C(x, \left\lceil -a \log \mu B_{\varepsilon(\delta)}(x) \right\rceil, \delta).$$

As μ is nonatomic, $\lim_{\delta\to\infty} -a\log\mu B_{\varepsilon(\delta)}(x) = \infty$. Then using the Brin-Katok Theorem, there exists an $x \in A$ such that the metric entropy is given by

$$\begin{split} h_{\mu}(T) &= \liminf_{\delta \to 0} \liminf_{n \to \infty} -\frac{1}{n} \log \mu C(x, n, \delta) \\ &\leq \liminf_{m \to \infty} \liminf_{n \to \infty} \frac{-\log \mu C(x, \lceil -a \log \mu B_{\varepsilon(\frac{1}{n})}(x) \rceil, \frac{1}{m})}{\lceil -a \log \mu B_{\varepsilon(\frac{1}{n})}(x) \rceil} \\ &\leq \liminf_{n \to \infty} \frac{-\log \mu C(x, \lceil -a \log \mu B_{\varepsilon(\frac{1}{n})}(x) \rceil}{\lceil -a \log \mu B_{\varepsilon(\frac{1}{n})}(x) \rceil} \\ &\leq \liminf_{n \to \infty} \frac{-\log \mu B_{\varepsilon(\frac{1}{n})}(x)}{\lceil -a \log \mu B_{\varepsilon(\frac{1}{n})}(x) \rceil} \\ &\leq \lim_{n \to \infty} \frac{-\log \mu B_{\varepsilon(\frac{1}{n})}(x)}{\lceil -a \log \mu B_{\varepsilon(\frac{1}{n})}(x) \rceil} \\ &\leq \frac{1}{a}, \end{split}$$

where in the third line we have used the inequality $\mu C(x, n, \delta) \leq \mu C(x, n, \delta')$ if $\delta \leq \delta'$. This gives the desired contradiction, and thus T is restricted sensitive with asymptotic rate a.

The conditions of ergodicity and continuity on T in the above proof were required for the use of the Brin-Katok Theorem. Note that ergodicity of T is necessary for any result of this type, because positive entropy is a global property of the system whereas sensitivity is a condition that must hold locally at almost every point. (Indeed, if we consider the disjoint union of a restricted sensitive system with positive entropy and a non-restricted sensitive system with zero entropy, the resulting system would have positive entropy but not be restricted sensitive.) That the continuity of T is necessary is not clear, and we suspect that a result of this type holds true without the continuity condition.

Regarding the converse of this theorem, the following example constructs an ergodic, measure-preserving transformation which is restricted sensitive but has zero measure-theoretic entropy:

Example 3.4. Let X be a disjoint union of two copies of [0, 1/2], labeled I_1 and I_2 , equipped with their Borel sigma algebras \mathcal{B}_1 and \mathcal{B}_2 and Lebesgue measures. Consider the σ -algebra on X given by $\mathcal{S} = \{S_1 \cup S_2 : S_1 \in \mathcal{B}_1 \text{ and } S_2 \in \mathcal{B}_2\}$ and probability measure μ on X given by $\mu(A) = \lambda(A \cap I_1) + \lambda(A \cap I_2)$. Define a metric d on X by d(x, y) = |x - y| if x and y are in the same copy of [0, 1/2], and d(x, y) = 2 if not.

Let $A \subset [0, 1/2]$ be a Borel set of Lebesgue measure $\lambda(A) = \frac{1}{4}$ with the property that for any subinterval K of [0, 1/2], $\lambda(K \cap A) > 0$ and $\lambda(K \cap A^c) > 0$. (For a construction of such a set, see Appendix B of [Sil08].) By Theorem 3.4.23 in [Sri98], there exist measure-preserving Borel isomorphisms $\phi : A \to [0, 1/4)$ and $\psi :$ $A^c \to [0, 1/4)$. Let A_1 and A_2 be the copies of A inside I_1 and I_2 , respectively, and let ϕ_1, ψ_1 and ϕ_2, ψ_2 be copies of the maps ϕ, ψ on I_1, I_2 . Define a transformation $T : X \to X$ by

$$T(x) = \begin{cases} \psi_1^{-1} \circ \phi_1(x) & \text{for } x \in A_1 \\ \phi_2^{-1} \circ \psi_1(x) & \text{for } x \in A_1^c \cap I_1 \\ \psi_2^{-1} \circ \phi_2(x) & \text{for } x \in A_2 \\ \phi_1^{-1} \circ R \circ \psi_2(x) & \text{for } x \in A_2^c \cap I_2, \end{cases}$$

where $R : [0, 1/4) \to [0, 1/4)$ is an irrational rotation. Note that T is finite measurepreserving. Also, note that when T^4 is restricted to any one of the four "segments" $A_1, A_1^c \cap I_1, A_2$, or $A_2^c \cap I_2$, it is isomorphic to an irrational rotation. We claim that T is ergodic and restricted sensitive and that $h_{\mu}(T) = 0$.

To see that T is ergodic, let $C \subset X$ and $D \subset X$ have positive measure. Then we can find positive measure subsets $C^* \subset C$ and $D^* \subset D$ each of which is completely contained inside one of the four segments A_1 , $A_1^c \cap I_1$, A_2 , and $A_2^c \cap I_2$. Let k < 4 be the integer such that $T^k(C^*)$ and D^* are in the same segment. Then, since T^4 is isomorphic to an irrational rotation when restricted to that segment, there exists n a multiple of 4 such that $\mu(T^{k+n}(C^*) \cap D^*) > 0$. Therefore, $\mu(T^{k+n}(C) \cap D) > 0$.

To see that T is restricted sensitive, choose $\delta = 1$ and a = 2 for all $x \in X$. For any $x \in I_1$ and $\varepsilon \leq 1$, by construction, $\mu(B_{\varepsilon}(x) \cap A_1) > 0$ and $\mu(B_{\varepsilon}(x) \cap (A_1^c \cap I_1)) > 0$, so $\mu\{y \in B_{\varepsilon}(x) : d(Tx, Ty) > \delta\} > 0$ and we note that $n = 1 < -2\log \frac{1}{2} \leq -a\log \mu B_{\varepsilon}(x)$. A similar argument holds for any $x \in I_2$, so T is restricted sensitive.

To see that T has zero entropy, simply note that $h_{\mu}(T^4) = 0$ because T^4 is isomorphic to a disjoint union of 4 irrational rotations. This implies that $h_{\mu}(T) = 0$.

Theorem 3.1 places a quantitative lower bound on the entropy of a system using the asymptotic rate a in Definition 2.4. In the context of restricted sensitivity, we may consider the minimal asymptotic rate $a_T^*(x) = \inf_{\delta(x)} \inf_{a(x)} a(x)$ of a restricted sensitive transformation T as a function of $x \in X$, where the infimums are taken over all sensitivity constants and asymptotic rates at the point x. Theorem 3.3 then implies that if a > 0 is such that $a_T^*(x) > a$ for a.e. $x \in X$, then $\frac{1}{a} \ge h_{\mu}(T)$. This function $a_T^*(x)$ is well-defined over a.e. $x \in X$, and it is in fact measurable under mild conditions on μ and d:

Proposition 3.5. Suppose (X, d) is separable and μ is a Borel probability measure. Then for any restricted sensitive transformation T on X, its minimal asymptotic rate function $a_T^*: X \to \mathbb{R}$ is measurable.

We defer the proof of this technical proposition to Appendix A. The following proposition computes a_T^* for the one-sided Bernoulli shift transformation of Example 2.6 and shows that the upper bound on the entropy of the Bernoulli shift from Theorem 3.3 is tight:

Proposition 3.6. Let $(\Sigma_N^+, \mathcal{B}, \mu)$, d, and T be as in Example 2.6. Then

$$\frac{1}{a_T^*(\sigma)} = h_\mu(T)$$

for a.e. $\sigma \in \Sigma_N^+$.

Proof. Take $\sigma \in \Sigma_N^+$ for which the restricted sensitivity condition holds with sensitivity constant $\delta(\sigma) > 0$ and asymptotic rate $a(\sigma) > 0$. Let $c(\sigma) \ge 0$ be the integer such that $2^{-c(\sigma)} > \delta(\sigma) \ge 2^{-c(\sigma)-1}$. For each $i = 1, \ldots, N$, let $k_i^{(n)}(\sigma)$ be the number of occurrences of the symbol i in σ_0 through σ_{n-1} . For any ball $B(\sigma) = [\sigma_0, \sigma_1, \sigma_2, \ldots, \sigma_{n-1}]$, $\min\{k : \mu(\tau \in B(\sigma) : d(T^k\sigma, T^k\tau) > \delta(\sigma)) > 0\} = n - c(\sigma)$. Hence we must have that $n - c(\sigma) \le -a(\sigma)\log(p_1^{k_1^{(n)}(\sigma)} \ldots p_N^{k_N^{(n)}(\sigma)})$ for all $n > c(\sigma)$, so for fixed $\delta(\sigma)$,

$$\inf a(\sigma) = \sup_{n > c(\sigma)} -(n - c(\sigma)) \left(\log(p_1^{k_1^{(n)}(\sigma)} \dots p_N^{k_N^{(n)}(\sigma)}) \right)^{-1}$$

Thus

$$\begin{aligned} a_{T}^{*}(\sigma) &= \inf_{\delta(\sigma)} \sup_{n > c(\sigma)} -(n - c(\sigma)) \left(\log(p_{1}^{k_{1}^{(n)}(\sigma)} \dots p_{N}^{k_{N}^{(n)}(\sigma)}) \right)^{-1} \\ &= \left(\sup_{c \ge 0} \inf_{n > c} \frac{n}{n - c} \cdot \frac{1}{n} \sum_{i=1}^{N} -k_{i}^{(n)}(\sigma) \log p_{i} \right)^{-1} \\ &= \left(\lim_{c \to \infty} \inf_{n > c} \frac{n}{n - c} \cdot \frac{1}{n} \sum_{i=1}^{N} -k_{i}^{(n)}(\sigma) \log p_{i} \right)^{-1}. \end{aligned}$$

By the Birkhoff Ergodic Theorem, for a.e. $\sigma \in \Sigma_N^+$,

$$\lim_{n \to \infty} \frac{k_i^{(n)}(\sigma)}{n} = \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^N \chi_{[i]} \left(T^i(\sigma) \right) p_i,$$

 \mathbf{SO}

$$\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{N} -k_i^{(n)}(\sigma) \log p_i = \sum_{i=1}^{N} -p_i \log p_i = h_{\mu}(T).$$

Then

As ε we

$$\frac{1}{a_T^*(\sigma)} \le \lim_{c \to \infty} \lim_{n \to \infty} \frac{n}{n-c} \cdot \frac{1}{n} \sum_{i=1}^N -k_i^{(n)}(\sigma) \log p_i = h_\mu(T).$$

For any $\varepsilon > 0$, there exists c sufficiently large so that $\frac{1}{n} \sum_{i=1}^{N} -k_i^{(n)}(\sigma) \log p_i \ge h_{\mu}(T) - \varepsilon$ for all n > c; hence

$$\frac{1}{a_T^*(\sigma)} \ge \lim_{c \to \infty} \lim_{n \to \infty} \frac{n}{n-c} \cdot (h_\mu(T) - \varepsilon) = h_\mu(T) - \varepsilon.$$

as arbitrary, $\frac{1}{a_T^*(\sigma)} = h_\mu(T).$

4. Restricted Sensitivity for Nonsingular Transformations

In the preceding section, the notions of restricted sensitivity and restricted pairwise sensitivity were used to examine measure-preserving transformations, for which there is a well-developed theory of metric entropy. These notions can be applied as well to nonsingular transformations; let us consider restricted sensitivity in this section. We show that a general class of nonsingular rank-one transformations (including measure-preserving rank-one transformations) are not restricted sensitive, and we construct a class of nonsingular type III transformations that are restricted sensitive.

Let us recall the definition of rank-one transformations. This class is known to contain finite measure-preserving mixing transformations [Orn72] and type III power weakly mixing nonsingular transformations [AFS01]. A nonsingular transformation $T: (X, \mu) \to (X, \mu)$ is **type III** if there are no σ -finite measures invariant under T that are equivalent to μ . The first example of a type III transformation was rank-one [Orn60]. By [JKL⁺08], it follows that the class of rank-one transformations further includes strong measurably sensitive finite measure-preserving transformations and measurably sensitive type III transformations.

We give the cutting and stacking definition of these transformations and follow the notation of [CS04] and [DS09]. Our presentation includes nonsingular transformations. A **column** consists of a finite ordered collection of disjoint intervals in

8

 \mathbb{R} . Each interval is called a **level**, and the levels may be of different lengths. The **height** of the column is the number of levels in the column. Each column defines an associated **column map**, defined on all levels except the top, by mapping each interval of the column to the next interval in the column by the unique orientation-preserving affine map that takes one interval to the other. Hence the column map is defined on all but the last level.

A rank-one nonsingular transformation is specified by a sequence of integers $\{r_n \geq 2\}$, a sequence of functions $s_n : \{0, \ldots, r_n - 1\} \rightarrow \mathbb{N}_0$, and a sequence $\{p_n\}$ of probability vectors on $\{0, \ldots, r_n - 1\}$. In the case of a measure-preserving transformations the probability vectors are all uniform, i.e., $p_n(i) = 1/r_n$ for all $i \in \{0, \ldots, r_n - 1\}$. We now describe the inductive procedure that constructs a sequence of columns C_n . Start by letting C_0 consist of a single interval. Assume that column $C_n = \{I_{n,i}\}_{i=0}^{h_n-1}$ of height h_n has been constructed. This we have a column map T_n , where $T_n(I_{n,i}) = I_{n,i+1}$ for $i \neq h_n - 1$. To construct column C_{n+1} subdivide C_n into r_n subcolumns by cutting each level $I_{n,i}$ into r_n subintervals or **sublevels**, $\{I_{n,i}^{[j]}\}_{j=0}^{r_n-1}$, (where $I_{n,i}^{[0]}$ is the leftmost sublevel and $I_{n,i}^{[r_n-1]}$ is the rightmost) whose lengths are in the proportions

$$p_n(0): p_n(1): \dots : p_n(r_n-1).$$

(For example, if $r_n = 1$, and $p_n(0) = 1/3$, $p_n(1) = 2/3$, then every level is cut in the proportions 1/3 : 2/3.) Then the **subcolumns** of C_n are $C_n^{[j]} = \{I_{n,i}^{[j]}\}_{i=0}^{h_n-1}$. By preserving the order on the levels, each subcolumn, $C_n^{[j]}$, is a column in its own right with the associated map $T_n^{[j]}$ which is the restriction of T_n to $C_n^{[j]}$. The next step is to place new intervals the size of the top sublevel above each subcolumn by adding $s_{n,j}$ levels above $C_n^{[j]}$; these new intervals are called **spacer levels**. To obtain the next column then we **stack** the resulting subcolumns with spacers right on top of left yielding the new column C_{n+1} with height

$$h_{n+1} = r_n h_n + \sum_{j=0}^{r_n - 1} s_n(j).$$

Let S_n denote the **union of spacer levels** added to C_n , the collection of levels in C_{n+1} that are not sublevels of levels in C_n ; hence, $C_{n+1} = C_n \sqcup S_n$, and denote by $S_n^{[j]}$ the collection of spacers added over $C_n^{[j]}$, so that $S_n = \bigcup_{j=0}^{r_n-1} S_n^{[j]}$. The associated column map T_{n+1} restricts to T_n on the levels in C_n . Let X be the union of all the levels in all columns. We assume that as $n \to \infty$ the maximal length of the intervals in C_n converges to 0, so we may define a transformation T of (X, μ) by

$$T(x) := \lim_{n \to \infty} T_n(x).$$

One can verify that T is well-defined and invertible a.e. and that it is nonsingular and ergodic. T is measure-preserving if all the probability vectors p_n are uniform, and $\mu(X) < \infty$ if and only if the total measure of the added spacers is finite.

The following proposition shows that nonsingular rank-one transformations constructed in this way are not restricted sensitive if there is a uniform lower bound on the elements of the probability vectors $\{p_n\}$.

Proposition 4.1. Let T be a nonsingular rank-one transformation on [0, 1) with the Euclidean metric d and Lebesgue measure λ . Suppose that column C_n is divided

into r_n subcolumns with proportions $p_n(0), \ldots, p_n(r_n-1)$. If there exists c > 0 such that $p_n(j) \ge c$ for all n and j, then T is not restricted sensitive.

Proof. For each n, let us further divide the leftmost subcolumn $C_n^{[0]}$ of column C_n into three equal subcolumns, labeled from left to right as $C_n^{[0],1}$, $C_n^{[0],2}$, and $C_n^{[0],3}$. Let $S_n = C_n \setminus C_n^{[0],2}$ and let $S = \bigcup_{k=0}^{\infty} \bigcap_{n=k}^{\infty} S_n$. As $\lambda S_n \leq (1 - \frac{c}{3})\lambda C_n \leq 1 - \frac{c}{3}$, $\lambda(\bigcap_{n=k}^{\infty} S_n) \leq 1 - \frac{c}{3}$ for each k and thus $\lambda S \leq 1 - \frac{c}{3}$. So the complement of S in [0,1) has positive measure.

For any $x \notin S$, there is an increasing sequence $\{n_k\}$ such that $x \in C_{n_k} \setminus S_{n_k}$. Let h_n and w_n be the height and the width of the smallest level, respectively, of column C_n . For any $\delta > 0$ and a > 0, there exists n_k sufficiently large such that $w_{n_k} < \delta$ and $a(n_k \log \frac{1}{c} + \log \frac{3}{2w_0}) < 2^{n_k-1}$. By the construction of S_{n_k-1} , if h is the smallest number such that $T^h(x)$ is in the highest level of C_{n_k} and w is the distance from x to the closer of the two endpoints of the level containing x in C_{n_k} , then $h \ge \frac{h_{n_k}}{2}$ and $w \ge \frac{w_{n_k}}{3}$. We also note that $w_{n_k} \ge c^{n_k} w_0$ and $h_{n_k} \ge 2^{n_k}$. Consider the ball $B_w(x)$. We note that

$$-a \log \lambda B_w(x) = -a \log 2w \le -a \log \frac{2w_{n_k}}{3} \le -a \log \frac{2c^{n_k}w_0}{3}$$
$$= a \left(n_k \log \frac{1}{c} + \log \frac{3}{2w_0} \right) < 2^{n_k - 1} \le \frac{h_{n_k}}{2} \le h.$$

Hence, for all $y \in B_w(x)$, $T^n y \in B_w(T^n x)$ for all $n \leq -a \log \lambda B_w(x)$, so

$$d(T^n x, T^n y) < w < \delta.$$

Hence T is not restricted sensitive.

This proposition addresses a large class of nonsingular rank-one transformations. As measure-preserving transformations are those for which the probability vectors p_n are uniform, measure-preserving transformations for which r_n is bounded above over all n satisfy the conditions of this proposition. In fact, the argument in the above proof can be modified to hold for all measure-preserving rank-one transformations:

Proposition 4.2. If T is a measure-preserving rank-one transformation on [0, 1)with the Euclidean metric d and Lebesque measure λ , then T is not restricted sensitive.

Proof. As the proof is very similar to that for Proposition 4.1, we will highlight the modifications required. Let us divide each subcolumn $C_n^{[j]}$ of column C_n into three equal subcolumns, labeled from left to right as $C_n^{[j],1}$, $C_n^{[j],2}$, and $C_n^{[j],3}$. Let

$$S_n = \bigcup_{j=\lceil \frac{r_n-1}{2}\rceil}^{r_n-1} C_n^{[j]} \cup \bigcup_{j=0}^{\lceil \frac{r_n-1}{2}\rceil-1} C_n^{[j],1} \cup \bigcup_{j=0}^{\lceil \frac{r_n-1}{2}\rceil-1} C_n^{[j],3},$$

and let $S = \bigcup_{k=0}^{\infty} \bigcap_{n=k}^{\infty} S_n$ as before. We have $\lambda S_n \leq \frac{8}{9} \lambda C_n \leq \frac{8}{9}$ for all n, so the complement of S in [0, 1) has positive measure.

The rest of the proof is the same as for Proposition 4.1, except that for any $\delta > 0$ and a > 0, we choose n_k sufficiently large such that $w_{n_k} < \delta$ and $a \log \frac{3h_{n_k}}{2w_0h_0} < \frac{h_{n_k}}{2}$.

As $w_{n_k}h_{n_k} \ge w_0h_0$ for all k, we use the bound $w_{n_k} \ge \frac{w_0h_0}{h_{n_k}}$ in place of the bounds $w_{n_k} \ge c^{n_k}w_0$ and $h_{n_k} \ge 2^{n_k}$ from the proof of Proposition 4.1.

We end by constructing type III nonsingular transformations that are restricted sensitive. The construction is a consequence of the following proposition:

Proposition 4.3. Let T be a transformation on a probability space (X, μ) with metric d_X and S be a transformation on a probability space (Y, ν) with metric d_Y , and let d_Y be ν -supported. If T is restricted sensitive, then the transformation $T \times S$ on $X \times Y$ (with the product σ -algebra and product measure) is restricted sensitive under the metric given by

$$d((x_1, y_1), (x_2, y_2)) = \max \left\{ d_X(x_1, x_2), d_Y(y_1, y_2) \right\}.$$

Proof. Suppose T is restricted sensitive at $x \in X$ with sensitivity constant $\delta > 0$ and asymptotic rate a > 0. For any $\varepsilon > 0$, suppose $n \ge$ is such that

$$\mu \left\{ x' \in B_{\varepsilon}^{d_X}(x) : d_X(T^n(x), T^n(x')) > \delta \right\} > 0.$$

Then for any $y \in Y$,

$$(\mu \times \nu) \left\{ (x', y') \in B^d_{\varepsilon}(x, y) : d((T \times S)^n(x, y), (T \times S)^n(x', y')) > \delta \right\}$$

$$\geq \nu(B^{d_Y}_{\varepsilon}(y)) \mu \left\{ x' \in B^{d_X}_{\varepsilon}(x) : d_X(T^n(x), T^n(x')) > \delta \right\} > 0.$$

As ν is a probability measure, $(\mu \times \nu)B^d_{\varepsilon}(x,y) \leq \mu B^{d_X}_{\varepsilon}(x)$ for any $\varepsilon > 0$, so $-a\log(\mu \times \nu)B^d_{\varepsilon}(x,y) \geq -a\log\mu B^{d_X}_{\varepsilon}(x)$. Hence $T \times S$ is restricted sensitive at (x,y) with the same sensitivity constant δ and asymptotic rate a. This holds for a.e. $x \in X$ and all $y \in Y$, so $T \times S$ is restricted sensitive.

Corollary 4.4. Let T be a measure-preserving mixing (or mildly mixing) transformation on a probability space (X, μ) with metric d_X , and suppose that T is restricted sensitive. Let S be a type III nonsingular conservative ergodic invertible transformation on a probability space (Y, ν) with metric d_Y . Then $T \times S$ on $X \times Y$ with the metric d given in Proposition 4.3 is a type III conservative ergodic transformation that is restricted sensitive.

Proof. By Proposition 5.4 and Theorem 5.2 of [HS98] the transformation $T \times S$ is conservative ergodic and type III. Proposition 4.3 implies that $T \times S$ is restricted sensitive.

References

- [AAB96] Ethan Akin, Joseph Auslander, and Kenneth Berg, When is a transitive map chaotic?, Convergence in ergodic theory and probability (Columbus, OH, 1993), Ohio State Univ. Math. Res. Inst. Publ., vol. 5, de Gruyter, Berlin, 1996, pp. 25–40. MR MR1412595 (97i:58106)
- [ABC02] Christophe Abraham, Gérard Biau, and Benoît Cadre, Chaotic properties of mappings on a probability space, J. Math. Anal. Appl. 266 (2002), no. 2, 420–431. MR MR1880515 (2002j:37009)
- [AFS01] Terrence Adams, Nathaniel Friedman, and Cesar E. Silva, Rank-one power weakly mixing non-singular transformations, Ergodic Theory Dynam. Systems 21 (2001), no. 5, 1321–1332. MR MR1855834 (2002h:37013)
- [BBC⁺92] J. Banks, J. Brooks, G. Cairns, G. Davis, and P. Stacey, On Devaney's definition of chaos, Amer. Math. Monthly 99 (1992), no. 4, 332–334. MR MR1157223 (93d:54059)
- [CJ05] Benoît Cadre and Pierre Jacob, On pairwise sensitivity, J. Math. Anal. Appl. 309 (2005), no. 1, 375–382. MR MR2154050 (2006b:28026)

- [CS04] Darren Creutz and Cesar E. Silva, Mixing on a class of rank-one transformations, Ergodic Theory Dynam. Systems 24 (2004), no. 2, 407–440. MR MR2054050 (2005a:37011)
- [DS09] Alexandre I. Danilenko and Cesar E. Silva, Ergodic theory: Nonsingular transformations, Encyclopedia of Complexity and System Science, vol. Part 5, Springer, 2009, pp. 3055–3083.
- [GII⁺08] Ilya Grigoriev, Nathaniel Ince, Cătălin M. Iordan, Amos Lubin, and Cesar E. Silva, On μ-compatible metrics and measurable sensitivity, preprint. (2008).
- [GW93] Eli Glasner and Benjamin Weiss, Sensitive dependence on initial conditions, Nonlinearity 6 (1993), no. 6, 1067–1075. MR MR1251259 (94j:58109)
- [HLYar] Wen Huang, Ping Lu, and Xiangdong Ye, Measure-theoretical sensitivity and equicontinuity, Israel J. Math. (to appear), 1–51.
- [HS98] Jane Hawkins and Cesar E. Silva, Characterizing mildly mixing actions by orbit equivalence of products, New York J. Math. 3A (1997/98), no. Proceedings of the New York Journal of Mathematics Conference, June 9–13, 1997, 99–115 (electronic). MR MR1611117 (99b:28019)
- [HYW04] Lianfa He, Xinhua Yan, and Lingshu Wang, Weak-mixing implies sensitive dependence, J. Math. Anal. Appl. 299 (2004), no. 1, 300–304. MR MR2091290 (2005i:37005)
- [JKL⁺08] Jennifer James, Thomas Koberda, Kathryn Lindsey, Cesar E. Silva, and Peter Speh, Measurable sensitivity, Proc. Amer. Math. Soc. 136 (2008), no. 10, 3549–3559.
- [Moo07] T. K. Subrahmonian Moothathu, Stronger forms of sensitivity for dynamical systems, Nonlinearity 20 (2007), no. 9, 2115–2126. MR MR2351026
- [Orn60] Donald S. Ornstein, On invariant measures, Bull. Amer. Math. Soc. 66 (1960), 297– 300. MR MR0146350 (26 #3872)
- [Orn72] _____, On the root problem in ergodic theory, Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971), Vol. II: Probability theory (Berkeley, Calif.), Univ. California Press, 1972, pp. 347–356. MR MR0399415 (53 #3259)
- [Sil08] C. E. Silva, Invitation to ergodic theory, Student Mathematical Library, vol. 42, American Mathematical Society, Providence, RI, 2008. MR MR2371216
- [Sri98] S.M. Srivastava, A Course on Borel Sets, Springer, 1998.

APPENDIX A. MEASURABILITY OF THE MINIMUM ASYMPTOTIC RATE

The proof of Proposition 3.5 requires the following lemmas:

Lemma A.1. Let (X,d) be a metric space with a probability measure μ . For all r > 0, the function $f(x) = \mu B_r(x)$ is lower semi-continuous.

Proof. Note that for any $x \in X$,

$$\lim_{n \to \infty} \mu B_{r-\frac{1}{n}}(x) = \mu \left(\bigcup_{n=1}^{\infty} B_{r-\frac{1}{n}}(x) \right) = \mu B_r(x)$$

Hence for any $\varepsilon > 0$, there exists n such that $\mu B_r(x) - \mu B_{r-\frac{1}{n}}(x) < \varepsilon$. For any $y \in B_{\frac{1}{n}}(x)$, $B_{r-\frac{1}{n}}(x) \subset B_r(y)$, so $\mu B_{r-\frac{1}{n}}(x) \le \mu B_r(y)$ and thus $\mu B_r(x) - \mu B_r(y) < \varepsilon$. Hence f is lower semi-continuous.

Lemma A.2. Suppose (X, d) is a separable metric space, μ is a Borel probability measure, and T is a transformation on X. Let $\delta > 0$, $\varepsilon > 0$, and $n \ge 0$, and let $A_{n,\varepsilon,\delta} = \{x \in X : \mu\{y \in B_{\varepsilon}(x) : d(T^nx, T^ny) > \delta\} > 0\}$. Then $A_{n,\varepsilon,\delta}$ is measurable.

Proof. By Lusin's Theorem, for each integer k > 0, there exists a closed set $F_k \subset X$ such that $\mu(X \setminus F_k) < \frac{1}{k}$ and $T|_{F_k} : F_k \to X$ is continuous. Let $E_k = X \setminus \bigcup_{i=0}^{n-1} T^{-i}(X \setminus F_k)$, so T^n is continuous on E_k , and define $f_k : E_k \to [0,1]$ as $f_k(x) = \mu\{y \in B_{\varepsilon}(x) \cap E_k : d(T^n x, T^n y) \geq \delta\}.$ We claim that f_k is lower-semicontinuous. Suppose $f_k(x) > 0$ for some $x \in E_k$, and let c > 0. There exists $\eta > 0$ such that $\mu\{y \in B_{\varepsilon}(x) \cap E_k : d(T^n x, T^n y) \ge \delta + \eta\} > f_k(x) - \frac{c}{2}$. Since $T|_{E_k}$ is continuous, there exists r > 0 such that $d(T^n x, T^n x') < \eta$ for all $x' \in B_r(x) \cap E_k$. We may choose r sufficiently small so that $\mu B_{\varepsilon - r}(x) > \mu B_{\varepsilon}(x) - \frac{c}{2}$. Then for all $x' \in B_r(x) \cap E_k$, $\mu(B_{\varepsilon}(x) \cap B_{\varepsilon}(x')) \ge \mu B_{\varepsilon - r}(x) > \mu B_{\varepsilon}(x) - \frac{c}{2}$. Thus

$$f_{k}(x') = \mu \{ y \in B_{\varepsilon}(x') \cap E_{k} : d(T^{n}x', T^{n}y) \geq \delta \}$$

$$\geq \mu \{ y \in B_{\varepsilon}(x') \cap E_{k} : d(T^{n}x, T^{n}y) \geq \delta + \eta \}$$

$$\geq \mu \{ y \in B_{\varepsilon}(x') \cap B_{\varepsilon}(x) \cap E_{k} : d(T^{n}x, T^{n}y) \geq \delta + \eta \}$$

$$\geq \mu \{ y \in B_{\varepsilon}(x) \cap E_{k} : d(T^{n}x, T^{n}y) \geq \delta + \eta \} - \mu (B_{\varepsilon}(x) \cap E_{k} \setminus B_{\varepsilon}(x'))$$

$$\geq f_{k}(x) - \frac{c}{2} - \frac{c}{2}$$

$$= f_{k}(x) - c.$$

Hence f_k is lower-semicontinuous on E_k , so f_k is measurable since μ is a Borel measure. Then $f_k^{-1}(0,1] = \{x \in X : \mu\{y \in B_{\varepsilon}(x) \cap E_k : d(T^nx,T^ny) \ge \delta\} \ge 0\}$ is measurable.

Let $S_{n,\varepsilon,\delta} = \{x \in X : \mu\{y \in B_{\varepsilon}(x) : d(T^nx,T^ny) \geq \delta\} > 0\}$. We note that $f_k^{-1}(0,1] \subset S_{n,\varepsilon,\delta}$ for all k. Since $\mu E_k^c < \frac{n}{k}$, for each $x \in S_{n,\varepsilon,\delta}$, if there exists k such that $\mu\{y \in B_{\varepsilon}(x) : d(T^nx,T^ny) \geq \delta\} > \frac{n}{k}$, then $f_k(x) > 0$. Hence $x \in f_k^{-1}(0,1]$ for some k, so $S_{n,\varepsilon,\delta} = \bigcup_{k=1}^{\infty} f_k^{-1}(0,1]$, which is measurable. Finally, $A_{n,\varepsilon,\delta} = \bigcup_{\delta' < \delta} S_{n,\varepsilon,\delta'}$, which is measurable.

Proof of Proposition 3.5. For any $\varepsilon > 0$ and $\delta > 0$, define $n_{\varepsilon,\delta} : X \to \mathbb{Z}$ as $n_{\varepsilon,\delta}(x) = \min\{n \ge 0 : \mu\{y \in B_{\varepsilon}(x) : d(T^nx, T^ny) > \delta\} > 0\}$. For any $n \ge 0$, let $A_{n,\varepsilon,\delta} = \{x \in X : \mu\{y \in B_{\varepsilon}(x) : d(T^nx, T^ny) > \delta\} > 0\}$. Define $S_{n,\varepsilon,\delta} = A_{n,\varepsilon,\delta} \setminus \bigcup_{i=0}^{n-1} A_{i,\varepsilon,\delta}$. Then $n_{\varepsilon,\delta}(x) = \sum_{n=0}^{\infty} n\chi_{S_{n,\varepsilon,\delta}}(x)$. By Lemma A.2, $A_{n,\varepsilon,\delta}$ is measurable for all n, so $n_{\varepsilon,\delta}$ is measurable. For any $\delta > 0$, consider

$$\bar{a}_{\delta}(x) = \inf\{a > 0 : \forall \varepsilon > 0, \exists n \le -a \log \mu B_{\varepsilon}(x) \text{ s.t.} \\ \mu\{y \in B_{\varepsilon}(x) : d(T^n x, T^n y) > \delta\} > 0\},\$$

as an extended real-valued function with $\bar{a}_{\delta}(x) = \infty$ if the set of such a > 0 is empty (i.e. if δ is not a sensitivity constant for $x \in X$). Then $\bar{a}_{\delta}(x) = \sup_{\varepsilon > 0} \frac{n_{\varepsilon,\delta}(x)}{-\log \mu B_{\varepsilon}(x)}$. By Lemma A.1, $-\log \mu B_{\varepsilon}(x)$ is upper semi-continuous and thus measurable since μ is a Borel measure, so $\frac{n_{\varepsilon,\delta}(x)}{-\log \mu B_{\varepsilon}(x)}$ is measurable.

Let us fix x and δ and consider $n_{\varepsilon,\delta}(x)$ and $-\log \mu B_{\varepsilon}(x)$ as monotonically decreasing functions of ε . We note that both functions are left-continuous because $\mu B_{\varepsilon}(x)$ is left-continuous in ε . For each n, let $\varepsilon_{\delta,n} = \max\{\varepsilon > 0 : n_{\varepsilon,\delta}(x) = n\}$. Then $n_{\varepsilon,\delta}(x)$ as a function of ε is constant on each of the intervals $(\varepsilon_{n,\delta}, \varepsilon_{n+1,\delta}]$ for all n, so $\frac{n_{\varepsilon,\delta}(x)}{-\log \mu B_{\varepsilon}(x)}$ as a function of ε is monotonically increasing on these intervals. Hence $\bar{a}_{\delta}(x) = \sup_{n\geq 0} \lim_{\varepsilon \to \varepsilon_{n,\delta}^{-1}} \left(\frac{n_{\varepsilon,\delta}(x)}{-\log \mu B_{\varepsilon}(x)}\right)$, which is measurable. Finally, we note that for any fixed x, $\bar{a}_{\delta}(x)$ is monotonically increasing in δ , so $a_T^* = \inf_{\delta>0} \bar{a}_{\delta} = \lim_{\delta \to 0} \bar{a}_{\delta}$ as extended real-valued functions, which is measurable.

(Domenico Aiello) WILLIAMS COLLEGE, MA 01267, USA *E-mail address*: Domenico Aiello <09da@williams.edu>

(Hansheng Diao) MASSACHUSETTS INSTITUTE OF TECHNOLOGY, MA 02139, USA $E\text{-}mail\ address:\ \texttt{hansheng@mit.edu}$

(Zhou Fan) HARVARD COLLEGE, MA 02138, USA E-mail address: Zhou Fan <zhoufan@fas.harvard.edu>

(Daniel King) WILLIAMS COLLEGE, MA 01267, USA *E-mail address*: Dan King <09dok@williams.edu>

(Jessica Lin) Courant Institute of Mathematical Sciences, New York University, NY 10012, USA

 $E\text{-}mail\ address: \texttt{jessicalin}\texttt{Q}\texttt{nyu}\texttt{.}\texttt{edu}$

(Cesar Silva) Department of Mathematics, Williams College, Williamstown, MA 01267, USA

 $E\text{-}mail \ address: \texttt{csilva@williams.edu}$

14