Bases for Riemann-Roch spaces of one point divisors on an optimal tower of function fields

Francesco Noseda[∗] Gilvan Oliveira† Luciane Quoos[∗]

Abstract

For applications in algebraic geometric codes, an explicit description of bases of Riemann-Roch spaces of divisors on function fields over finite fields is needed. We give an algorithm to compute such bases for one point divisors, and Weierstrass semigroups over an optimal tower of function fields. We also explicitly compute Weierstrass semigroups till level eight.

MSC2010 Subject Classification Numbers: 14H05, 14H55, 14G50, 94B05. Keywords: Riemann-Roch spaces, tower of function fields, Weierstrass semigroup, algebraic geometric codes.

1 Introduction

Algebraic geometric codes are defined by means of Riemann-Roch spaces of divisors on function fields over finite fields. In practice, for applications in coding theory, one needs an explicit description of bases of such spaces. The problem of computing these bases is a hard one, and it is addressed, for instance, in [\[7\]](#page-13-0), [\[8\]](#page-13-1), [\[9\]](#page-14-0), [\[10\]](#page-14-1) and [\[13\]](#page-14-2). In this correspondence we give an algorithm to compute bases and Weierstrass semigroups over an optimal tower of function fields.

We use the language of function fields as in [\[11\]](#page-14-3). Let \mathbb{F}_q be the finite field with q elements, and consider a tower of function fields $\mathcal{T} = (T_j)_{j\geq 0}$ over \mathbb{F}_q . Let $g(T_j)$ denote the genus of T_j/\mathbb{F}_q and $N(T_j)$ its number of \mathbb{F}_q -rational points. The well-known Drinfeld-Vlădut bound (see [\[12\]](#page-14-4)) guarantees that the limit $\lambda(\mathcal{T}) := \lim_{j \to \infty} N(T_j)/g(T_j)$ satisfies the inequality:

$$
0 \leq \lambda(\mathcal{T}) \leq \sqrt{q} - 1.
$$

The tower T is said to be good if $\lambda(T) > 0$, and optimal if $\lambda(T) = \sqrt{q} - 1$. In the literature there are many explicit descriptions of several good and optimal towers (see $[1]$, $[2]$, $[4]$ and $[5]$), from which algebraic geometric codes can be constructed having parameters that attain the Tsfasman-Vlădut-Zink bound (see $[11]$).

In this paper we consider the tower $\mathcal{T} = (T_j)_{j\geq 0}$ over the finite field \mathbb{F}_{p^2} in odd characteristic. This tower is defined recursively by $T_0 = \mathbb{F}_{p^2}(x_0)$ and, for $j \geq 0$, $T_{j+1} = T_j(x_{j+1})$, where the function x_{j+1} satisfies the relation:

$$
x_{j+1}^2 = \frac{x_j^2 + 1}{2x_j}.
$$

This tower was introduced and proven to be optimal in [\[3\]](#page-13-6). Let P^j_∞ be the unique pole of the function x_0 in T_j . For each $s \in \mathbb{N}$ fix the divisors sP^j_{∞} , and let $D^j = P_1 + \cdots + P_N$ be the sum of all rational places in T_j that lie over a set Ω of points in T_0 that completely split in the tower, see [\[3\]](#page-13-6). We get a sequence of algebraic geometric codes defined, for $j \geq 0$, as:

$$
\mathcal{C}_L(sP_\infty^j, D^j) = \{ (z(P_1), \cdots, z(P_N)) \mid z \in L(sP_\infty^j) \} \subseteq \mathbb{F}_{p^2}^N
$$

^{*}Instituto de Matemática, UFRJ, CP 68530, Cidade Universitária, CEP: 21941-909, Rio de Janeiro/RJ - Brasil (noseda@im.ufrj.br, luciane@im.ufrj.br).

[†]Departamento de Matemática, CCE/UFES, Goiabeiras, CEP: 29075-910, Vitória/ES - Brasil (jgilvanol@gmail.com).

where the Riemann-Roch space $L(sP_{\infty}^j)$ is defined by:

 $L(sP_{\infty}^j) = \{z \in T_j | \text{ the divisor of } z \text{ satisfies } (z) \ge -sP_{\infty}^j\}.$

The knowledge of explicit bases of the Riemann-Roch spaces allows to construct the matrices of such codes. The main result of this paper, Theorem [3.1,](#page-6-0) is an algorithm to compute such bases. The central idea is to apply results of [\[6\]](#page-13-7) (see Theorem [2.9](#page-5-0) in Section [2](#page-1-0) below) to decompose the vector space $L(sP^j_{\infty})$ in T_j as a direct sum of Riemann-Roch spaces of divisors at the lower level T_{j-1} , and continue this way till the rational function field T_0 , where the bases can be easily computed. In order this process to be performed, the divisors we get at each level $k < j$ should be invariant for the action of the Galois group of T_k/T_{k-1} . Unfortunately this condition is not always satisfied and the procedure has to be suitably modified, as done in Sections [3.1](#page-7-0) and [3.2.](#page-9-0)

As a consequence of the main result we get an algorithm to compute the Weierstrass semigroups:

 $H(P_{\infty}^j) = \{ s \in \mathbb{N} \mid \exists z \in T_j \text{ s.t. the pole divisor of } z \text{ satisfies } (z)_{\infty} = s P_{\infty}^j \}$

at the totally ramified points P_{∞}^{j} , Theorem [3.2.](#page-6-1) As an application, we also explicitly present the semigroups till level eight, Section [3.3.](#page-12-0)

The ramification structure of the tower and the computation of the genus is presented in Section [2.](#page-1-0)

Acknowledgments. The authors would like to express their gratitude to A. Garcia for suggesting the subject and for valuable discussions, and to P. Zingano for useful help in the computational part.

2 Preliminaries

The object of study is an asymptotically optimal tower of functions fields defined over the finite field $K = \mathbb{F}_{p^2}$ with p^2 elements, where p is an odd prime. This tower is recursively defined by: $T_0 = K(x_0)$ and, for $j \geq 0$, $T_{j+1} = T_j(x_{j+1})$, where the function x_{j+1} satisfies the relation:

$$
x_{j+1}^2 = \frac{x_j^2 + 1}{2x_j}.
$$

The field K is the full constant field of T_j/K for any $j \geq 0$. Moreover, the extension T_{j+1}/T_j is a Kummer extension of degree 2. For the general theory of function fields we refer to [\[11\]](#page-14-3).

For $\alpha \in K \cup \{\infty\}$ we denote by P^0_α the unique place of T_0/K such that $x_0(P^0_\alpha) = \alpha$. We start by studying the ramification structure of the tower above the places P^0_α with $\alpha \in \{\infty, 0, \pm i, \pm 1\}$, where $i^2 = -1$; this is the content of Lemma [2.2.](#page-2-0) We shall consider the results of the lemma, and its consequences, to be known. Nonetheless, we are not aware of a complete written account of the ramification structure, and the results we need are beyond what is explicitly written about it in [\[3\]](#page-13-6), [\[4\]](#page-13-4), and [\[11\]](#page-14-3).

Lemma 2.1 For all $j \geq 0$, let P be a place of T_j/K and Q be a place of T_{j+1}/K that lies over P. Then:

(a) We have:

(b) We have:

$$
x_{j+1}(Q) = \infty \qquad \Longleftrightarrow \qquad x_j(P) \in \{\infty, 0\}
$$

\n
$$
x_{j+1}(Q) \in \{1, -1\} \qquad \Longleftrightarrow \qquad x_j(P) = 1.
$$

\n
$$
x_{j+1}(Q) = 0 \qquad \Longleftrightarrow \qquad x_j(P) = 1.
$$

\n
$$
x_{j+1}(Q) \in \{i, -i\} \qquad \Longleftrightarrow \qquad x_j(P) = -1.
$$

(c) If $x_j(P) \in \{1, -1\}$ then $e(Q|P) = f(Q|P) = 1$. In both cases there are exactly two places in T_{j+1}/K over P, and x_{j+1} takes one of the values admitted by item (b) in one place, and the other value in the other place.

PROOF: It follows from the theory of Kummer extensions of function fields, see [\[11\]](#page-14-3).

Lemma 2.2 Denote $\mathcal{R} := \{P_{\infty}^0, P_0^0, P_i^0, P_{-i}^0, P_1^0, P_{-1}^0\}$. Then, for any $j \ge 0$ the following holds.

(a) For $\alpha \in \{\infty, 0, \pm i\}$ there is a unique place P^j_α in T_j/K over P^0_α . If $j \geq 1$ then $e(P^j_\alpha | P^{j-1}_\alpha) = 2$. Moreover $v_{P_{\infty}^j}(x_j) = -1$ and:

$$
v_{P_0^j}(x_j) = \begin{cases} 1 & \text{if } j = 0 \\ -1 & \text{if } j \ge 1 \end{cases} \qquad v_{P_{\pm i}^j}(x_j) = \begin{cases} 0 & \text{if } j = 0 \\ 1 & \text{if } j = 1 \\ -1 & \text{if } j \ge 2. \end{cases}
$$

(b) For $\beta \in \{\pm 1\}$ there is a unique place P^j_β in T_j/K with $x_j(P^j_\beta) = \beta$. If $j \ge 1$, we have: $P^j_{\pm 1} | P^{j-1}_1$ and $e(P_{\pm 1}^j | P_1^{j-1}) = 1.$

(c) Let r with $0 \le r \le \lfloor (j-3)/2 \rfloor$; notice that $j \ge 3$ and $r < j-1$. Denote by $Q = Q_i^j$ a place in T_j/K that lie over P_{-1}^r . Then the sum of the degrees of such places is 2^{r+2} . Moreover, $e(Q_r^j|Q_r^{j-1})=2$ and $v_Q(x_j) = -1.$

(d) Let r with $\lfloor (j-1)/2 \rfloor \le r \le j-2$; notice that $j \ge 2$ and $r \ge 0$. Denote by $Q = Q_r^j$ a place in T_j/K that lie over P_{-1}^r . Then the sum of the degrees of such places is 2^{j-r} . Moreover, we have $e(Q_r^j|Q_r^{j-1})=1$ and:

$$
v_Q(x_j) = \begin{cases} -2^{2r-j+2} & \text{if } r \le j-3\\ 2^{j-2} & \text{if } r = j-2. \end{cases}
$$

(e) For $\beta \in {\pm i}$ there is a unique place Q_{β}^{j} in T_j/K with $x_j(Q_{\beta}^j) = \beta$. If $j \ge 1$, we have: $Q_{\pm i}^j | P_{-1}^{j-1}$ and $e(Q_{\pm i}^j | P_{-1}^{j-1}) = 1$, and $Q_{\pm i}^j$ are the unique places of T_j/K that lie over P_{-1}^{j-1} .

(f) The places of T_j/K defined in items (a) to (e) are the unique places over R. With the exception of $Q_{\pm i}^0 = P_{\pm i}^0$, they are all distinct.

- (g) The function x_j has no zeros nor poles in T_j outside the set of places over \mathcal{R} .
- (h) The places $Q_{\pm i}^j$ defined in item (e) satisfy $v_{Q_{\pm i}^j}(x_j^2 + 1) = 2^j$.

PROOF: The proof is by induction on j. For $j = 0$ the thesis follows by the properties of the rational function field. Let $j \geq 0$, and assume the thesis is true for j.

(a) Use Lemma [2.1](#page-1-1) to deduce $v_{P_{\infty,0}^{j+1}}(x_{j+1}) = -1$ and $e(P_{\infty,0}^{j+1}|P_{\infty,0}^j) = 2$ for $j \ge 0$, and to deduce $v_{P_{\pm i}^{j+1}}(x_{j+1}) = -1$ and $e(P_{\pm i}^{j+1}|P_{\pm i}^j) = 2$ for $j \ge 1$. For $j = 0$ the assertions $v_{P_{\pm i}^1}(x_1) = 1$ and $e(P_{\pm i}^1 | P_{\pm i}^0) = 2$ follows by $v_{P_{\pm i}^0}(x_0^2 + 1) = 1$ and by the same Lemma.

(b) It follows by Lemma [2.1.](#page-1-1)

 \Box

(c) Let r be such that $0 \le r \le \lfloor (j-2)/2 \rfloor$, which implies $j \ge 2$ and $r < j$. Let Q be a place in T_{j+1}/K that lies over P_{-1}^r . The image P of Q in T_j/K lies over P_{-1}^r as well. Applying Lemma [2.1,](#page-1-1) the thesis follows if we prove the claim that: $v_P(x_j) = -1$, and the sum of the degrees of such places P is 2^{r+2} . First, if $r \leq (i-3)/2$, then the claim follows by induction hypothesis (c).

Second, if $r > |(j-3)/2|$ then j is even and $r = (j-2)/2$, i.e., $j = 2r+2$. Notice that $r = |(j-1)/2| \leq$ $j-2$. Then the result follows by induction hypothesis (d) .

(d) Let r with $|j/2| \leq r \leq j-1$, which implies $j \geq 1$ and $r \geq 0$. Let Q be a place in T_{j+1}/K that lies over P_{-1}^r . The image P of Q in T_j/K lies over P_{-1}^r as well.

First case: $r \leq j - 3$. Then $\lfloor (j - 1)/2 \rfloor \leq r \leq j - 2$ and we can apply induction (d) to deduce $v_P(x_j) = -2^{2r-j+2}$, and the sum of the degrees of such places P is 2^{j-r} . Since in fact $(j-1)/2 \leq r$, then $2r-j+2 \geq 1$ and $v_P(x_j)$ is even. Apply Lemma [2.1](#page-1-1) to deduce $e(Q|P) = 1$ and $v_Q(x_{j+1}) = -2^{2r-(j+1)+2}$. The claim on the sum of the degrees follows.

Second case: $r = j - 2$. Then $\lfloor (j - 1)/2 \rfloor \le r \le j - 2$ and we can apply induction (d) to deduce $v_P(x_j) = 2^{j-2}$, and the sum of the degrees of such places P is 2^{j-r} . Since $\lfloor j/2 \rfloor \leq j-2$, then $j \geq 3$. Hence $v_P(x_j)$ is even. Apply Lemma [2.1](#page-1-1) to deduce $e(Q|P) = 1$ and $v_Q(x_{j+1}) = -2^{j-3} = -2^{2r-(j+1)+2}$. The claim on the sum of the degrees follows.

Third case: $r = j - 1$. In this case, by induction (e) and (h), the place P is one of the places $Q_{\pm i}^j$, and we have: $v_Q(x_j^2 + 1) = 2^j$ and $v_Q(x_j) = 0$. By Lemma [2.1](#page-1-1) it follows that $e(Q|P) = 1$ and $v_Q(x_{j+1}) = 2^{j-1} = 2^{(j+1)-2}$. The result on the sum of the degrees follows.

(e) By Lemma [2.1](#page-1-1) and induction (b).

(f) Any place Q of T_{j+1}/K that lies over R lies over a place P of T_j/K that lies over R. By induction (f) , any such a place P is one of the places of T_i/K defined by items (a) to (e). The thesis follows by observing that if a place of T_{j+1}/K lies above a place of T_j/K defined by (a) to (e), then it is itself defined by (a) to (e) .

(g) We have to show that if Q is a place of T_{j+1} that does not lie over R then $v_Q(x_{j+1}) = 0$. Let P the image of Q in T_j . Since P does not lie over R, by induction (g) we have $v_P(x_j) = 0$. By the relation between x_{j+1} and x_j it follows that we have to show that $v_P(x_j^2 + 1) = 0$. Necessarily $v_P(x_j^2 + 1) \ge 0$. If it happened that $v_P(x_j^2 + 1) > 0$ then $x_j(P) = \pm i$ and by induction (e) the place P would lie over R.

(h) Let's analyse zeros and poles of $x_{j+1}^2 + 1$ in T_{j+1}/K . The unique zeros are $Q_{\pm i}^{j+1}$, by what was already proven for (e). Since the two places $Q_{\pm i}^{j+1}$ are permuted by the action of the unique non-trivial automorphism of T_{j+1}/T_j , then the valuations of $x_{j+1}^2 + 1$ at these two places are equal, say equal to v. We have to show that $v = 2^{j+1}$. By (g) and (a) to (e) the poles of x_{j+1} are the following: P_{∞}^{j+1} , P_0^{j+1} 0 (order 1); $P_{\pm i}^{j+1}$ if $j \ge 1$ (order 1); the places Q_r^{j+1} for r with $0 \le r \le j-2$ (order 1 if $r \le \lfloor (j-2)/2 \rfloor$, the sum of the degrees of such places is 2^{r+2} ; order 2^{2r-j+1} if $r > \lfloor (j-2)/2 \rfloor$, the sum of the degrees of such places is 2^{j+1-r}). We can compute:

$$
\deg(x_{j+1})_{\infty} = 1 + \sum_{n=0}^{j} 2^n = 2^{j+1}.
$$

. В последните постание и производите на селото на приема в селото на селото на селото на селото на селото на
Селото на селото на

Then $0 = \deg(x_{j+1}^2 + 1) = 2v - 2\deg(x_{j+1})_\infty$, and we conclude $v = 2^{j+1}$

Definition 2.3 For any $j \geq 0$ and for any r with $0 \leq r \leq j$, denote by:

$$
D^j_r:=\sum_{Q^j\mid P^r_{-1}}Q^j
$$

the divisor given by the sum of the places in T_j that lie over P_{-1}^r . Notice that $D_j^j = P_{-1}^j$. Extend the definition of D_r^j for $r = -2, -1$ by $D_{-2}^j := P_0^j$ and $D_{-1}^j := P_i^j + P_{-i}^j$.

The following corollary is a consequence of Lemma [2.2.](#page-2-0)

Corollary 2.4 For any $j \geq 0$ and any r with $-2 \leq r \leq j$ we have:

$$
\deg(D_r^j) = \begin{cases} 2^{j-r} & \text{if } j \le 2r+2 \\ 2^{r+2} & \text{if } j \ge 2r+2. \end{cases}
$$

Moreover, if $j \ge 1$ then, for any r with $-2 \le r \le j$, the places in D_r^j ramifies in T_j/T_{j-1} if and only if $j \geq 2r + 3$.

Proposition 2.5 The divisors of x_j and $1 + x_j$ in T_j are given by the following formulae: $(x_0) = -P^0_{\infty} + P^0_{0}$, $(x_1) = -P^1_{\infty} - P^1_{0} + D^1_{-1}$, and:

$$
(x_j) = -P_{\infty}^j - \sum_{r=-2}^{\lfloor \frac{j-3}{2} \rfloor} D_r^j - \sum_{r=\lfloor \frac{j-1}{2} \rfloor}^{j-3} 2^{2r-j+2} D_r^j + 2^{j-2} D_{j-2}^j \quad \text{if} \quad j \ge 2,
$$

 $(1+x_0) = -P^0_{\infty} + P^0_{-1}, (1+x_1) = -P^1_{\infty} - P^1_0 + 2P^1_{-1}, and$

$$
(1+x_j) = -P_{\infty}^j - \sum_{r=-2}^{\lfloor \frac{j-3}{2} \rfloor} D_r^j - \sum_{r=\lfloor \frac{j-1}{2} \rfloor}^{j-3} 2^{2r-j+2} D_r^j + 2^j P_{-1}^j \quad \text{if} \quad j \ge 2.
$$

PROOF: We apply Lemma [2.2.](#page-2-0) The claim on the divisor of x_j follows. The function $1 + x_j$ has the same poles as x_j ; moreover, it has P_{-1}^j as unique zero. By the proof of item (h) of the lemma we have: $\deg(1+x_j)_{\infty} = \deg(x_j)_{\infty} = 2^j$. Hence, the coefficient of P_{-1}^j of $(1+x_j)$ is 2^j .

 \Box

Proposition 2.6 For $j \geq 0$ the genus g_j of T_j/K is given by:

$$
g_j = \begin{cases} (2^{\frac{j+2}{2}} - 1)(2^{\frac{j}{2}} - 1) & \text{if } j \text{ is even} \\ (2^{\frac{j+1}{2}} - 1)^2 & \text{if } j \text{ is odd.} \end{cases}
$$

PROOF: By the Riemann-Hurwitz Theorem we have that for any $j \geq 0$:

$$
g_{j+1} = 2g_j - 1 + \frac{1}{2}R_j
$$

where $R_j = \sum_P \deg P$, and the sum is over all places of T_j/K such that $v_P((x_j^2 + 1)/2x_j)$ is odd. The last condition is satisfied in two cases: $v_P(x_j) = 0$ and $v_P(x_j^2 + 1)$ is odd; or $v_P(x_j)$ is odd. By Lemma [2.2,](#page-2-0) the first case is realized exactly by $P_{\pm i}^j$ if $j = 0$. The second one is realized by the places: P_{∞}^j , P_0^j ; $P_{\pm i}^j$ if $j \geq 1$; Q_r^j with $0 \leq r \leq \lfloor (j-3)/2 \rfloor$ (the sum of the degrees of such places is 2^{r+2}); Q_r^j with $\lfloor (j-1)/2 \rfloor \le r \le j-2$ and: (a) $r = j-2$ and $j-2 = 0$; or (b) $r \le j-3$ and $2r - j + 2 = 0$ (the sum of the degrees of such places is 2^{j-r}).

If j is odd then cases (a) and (b) above do not contribute, and we have $R_j = 1 + \sum_{n=0}^{(j+1)/2} 2^n = 2^{\frac{j+3}{2}}$. If j is even then case (a) contributes with $r = 0$ if $j = 2$ (the sum of the degrees is 4), and case (b) contributes with $r = (j-2)/2$ if $j \ge 4$ (the sum of the degrees is 2^{r+2}). Hence, $R_j = 1 + \sum_{n=0}^{(j+2)/2} 2^n = 2^{\frac{j+4}{2}}$.

From this we deduce the following recursive relations:

$$
g_{j+2} = \begin{cases} 4g_j + 3 \cdot 2^{\frac{j+2}{2}} - 3 & \text{if } j \text{ is even} \\ 4g_j + 2^{\frac{j+5}{2}} - 3 & \text{if } j \text{ is odd.} \end{cases}
$$

Since $g_0 = 0$, from which $g_1 = 1$, then the thesis follows by induction.

Next, we state a particular case of Theorem 2.2 [\[6\]](#page-13-7), by H. Maharaj, that will play a central role in the construction of a basis of the Riemann-Roch space $L(sP_{\infty}^j)$. In order to state the theorem we need the notions of invariant divisor and of restriction of a divisor; this is the content of the next two remarks.

Remark 2.7 We will be dealing with divisors in T_i/K of the form:

$$
D = \alpha_{\infty} P_{\infty}^j + \sum_{r=-2}^{j-1} \alpha_r D_r^j + \alpha_j P_{-1}^j
$$

where the coefficients α_{∞} , α_r and α_j are integers. If $j \geq 1$ then such a divisor is invariant under the action of the Galois group of T_j/T_{j-1} (briefly, invariant in T_j/T_{j-1}) if and only if $\alpha_j = 0$. Indeed, because the unique non-trivial automorphism of T_j/T_{j-1} sends x_j to $-x_j$, the place P_{-1}^j is sent to P_1^j , while the divisors P^j_{∞} and D^j_r with $-2 \le r \le j-1$ are invariant.

Remark 2.8 Let $j \geq 1$, and $D = \alpha_{\infty} P_{\infty}^{j} + \sum_{r=-2}^{j-1} \alpha_{r} D_{r}^{j}$ be a divisor of T_{j}/K . Then, applying the definition given in [\[6\]](#page-13-7), and applying Corollary [2.4,](#page-4-0) the restriction of D to T_{j-1} is given by:

$$
D_{|T_{j-1}} = \left\lfloor \frac{\alpha_{\infty}}{2} \right\rfloor P_{\infty}^{j-1} + \sum_{r=-2}^{\left\lfloor \frac{j-3}{2} \right\rfloor} \left\lfloor \frac{\alpha_r}{2} \right\rfloor D_r^{j-1} + \sum_{r=\left\lfloor \frac{j-1}{2} \right\rfloor}^{j-1} \alpha_r D_r^{j-1}
$$

Theorem 2.9 Let $j \ge 1$ and D be a divisor in T_j/K that is invariant in T_j/T_{j-1} . Then:

$$
L(D) = L(D_{|T_{j-1}}) \oplus L([D + (x_j)]_{|T_{j-1}})x_j.
$$

PROOF: This is Theorem 2.2 of [\[6\]](#page-13-7) applied to the Kummer extension T_j/T_{j-1} .

In the next remark we explain how to compute the power series expansion of the generators x_k around P_{-1}^j , for $k \leq j$, with respect to a suitably chosen local parameter. This will be used in the sequel to make bases of $L(sP^j_{\infty})$ and the Weierstrass semigroups $H(P^j_{\infty})$ computable.

Remark 2.10 The function $t := 1 - x_0$ is a local parameter around P_{-1}^j , for any $j \ge 1$. Indeed, $v_{P_1^0}(1-x_0)=1$ and $e(P_{-1}^j|P_1^0)=1$. Notice that, for the same reason, t is a local parameter around P_1^j as well, for any $j \geq 0$.

We show how to compute the power series expansion up to any order $\varepsilon \geq 1$ of x_j around P_1^j , for any $j \geq 0$. Assume to have the expansion: $x_j = 1 + \sum_{k=1}^{\varepsilon} a_k t^k + O(t^{\varepsilon+1})$ around P_1^j . Then we can compute the expansion $(x_j^2 + 1)/2x_j = 1 + \sum_{k=1}^{\varepsilon} b_k t^k + \widetilde{O(t^{\varepsilon+1})}$. Since $e(P^{j+1}|P^j) = 1$ then the same expansion holds around P_1^{j+1} . Let the expansion of x_{j+1} around P_1^{j+1} be given by: $x_{j+1} = 1 + \sum_{k=1}^{\varepsilon} c_k t^k + O(t^{\varepsilon+1}),$ where the $c_k \in K$ have to be computed. By the relation: $x_{j+1}^2 = (x_j^2 + 1)/2x_j$ it follows that:

$$
1 + \sum_{k=1}^{\varepsilon} \left(2c_k + \sum_{l=1}^{k-1} c_l c_{k-l} \right) t^k + O(t^{\varepsilon+1}) = 1 + \sum_{k=1}^{\varepsilon} b_k t^k + O(t^{\varepsilon+1}).
$$

Hence, the unknown coefficients c_k can be computed by induction on k from the formulae:

$$
c_k = \frac{1}{2} \left(b_k - \sum_{l=1}^{k-1} c_l c_{k-l} \right) \qquad 1 \le k \le \varepsilon.
$$

We also remark that for $j \ge 1$ and k with $0 \le k \le j$ the expansion of x_k around P_{-1}^j is: the one given above if $k < j$ (since $e(P_{-1}^j | P_1^k) = 1$); minus the one given above if $j = k$ (since we have to take the other determination of the square root of $(x_j^2 + 1)/2x_j$.

.

3 Hermitian basis of the Riemann-Roch space $L(sP^j_\infty)$

We state the main result of the paper: Theorem [3.1.](#page-6-0) This gives bases of the spaces $L(sP^j_{\infty})$, as j and s vary, in a constructive way. As a corollary of the main theorem we get a constructive way to compute the Weierstrass semigroups $H(P_{\infty}^j)$: this is the content of Theorem [3.2.](#page-6-1) We recall that a basis of a Riemann-Roch space is Hermitian (with respect to P_{∞}^j) if its elements have distinct pole order at P_{∞}^j .

Theorem 3.1 There exists an algorithm that, for all $j \geq 0$ and all m with $0 \leq m < 2^{j}$, constructs integers $c_m^{(j)}$ and functions $w_m^{(j)} \in T_j$ s.t. for all $s \in \mathbb{Z}$ the family parametrized by m and l:

> $x_0^l w_m^{(j)}$ (i) $0 \le m < 2^j$ $0 \le l \le \left\lfloor \frac{s-m}{2^j} \right\rfloor - c_m^{(j)} := l_m(s)$

is a Hermitian basis of $L(sP^j_\infty)$. (When the value of s is such that $l_m(s)$ is negative for a given m, it is understood that the m-th family does not contribute to the basis.)

PROOF: The proof is by induction on j and uses Theorems [2.9](#page-5-0) and [3.4.](#page-7-1) For $j = 0$, put $c_0^{(0)} := 0$ and $w_0^{(0)} := 1.$

Let $j \geq 1$. We apply Theorem [2.9](#page-5-0) and get, for any $s \in \mathbb{Z}$:

$$
L(sP_\infty^j)=L([sP_\infty^j]_{|T_{j-1}})\oplus L([sP_\infty^j+(x_j)]_{|T_{j-1}})x_j.
$$

Since $[sP^j_{\infty}]|_{T_{j-1}} = \left[\frac{s}{2}\right]P^{j-1}_{\infty}$ (see Remark [2.8\)](#page-5-1), then the induction hypothesis gives a basis of the first direct summand:

$$
x_0^l w_m^{(j-1)} \qquad \qquad 0 \le m < 2^{j-1} \qquad 0 \le l \le \left\lfloor \frac{s-2m}{2^j} \right\rfloor - c_m^{(j-1)}
$$

where we used the fact that: $\lfloor (\lfloor s/2 \rfloor - m)/2^{j-1} \rfloor = \lfloor (s - 2m)/2^{j} \rfloor$.

A basis of $L([sP^j_{\infty}+(x_j)]_{|T_{j-1}})$ is constructed by Theorem [3.4.](#page-7-1) The desired basis of $L(sP^j_{\infty})$ is obtained by defining, for $0 \leq m < 2^j$:

$$
c_m^{(j)} := \begin{cases} c_{m/2}^{(j-1)} & \text{if m is even} \\ \tilde{c}_m^{(j-1)} & \text{if m is odd} \end{cases} \qquad w_m^{(j)} := \begin{cases} w_{m/2}^{(j-1)} & \text{if m is even} \\ \tilde{w}_m^{(j-1)} x_j & \text{if m is odd.} \end{cases}
$$

The fact that the basis is Hermitian follows by Corollary [3.3.](#page-7-2)

The knowledge of the dimension of $L(sP^j_{\infty})$ for fixed j and s variable, allows to compute the Weierstrass semigroup $H(P_{\infty}^{j})$. Because of the structure of the basis given in the above theorem, we get an elegant and efficient way to recover the semigroup.

Theorem 3.2 For any $j \geq 0$ the Weierstrass semigroup $H(P_{\infty}^{j})$ can be recovered from the coefficients $c_m^{(j)}$ given in Theorem [3.1](#page-6-0) as follows. For any $s \in \mathbb{Z}$, let $q(s)$ and $m(s)$ be quotient and rest of the division of s by 2^j , i.e., $s = 2^j q(s) + m(s)$ with $0 \le m(s) < 2^j$. Then:

$$
s\in H(P_\infty^j) \qquad \iff \qquad q(s)\ge c_{m(s)}^{(j)}.
$$

PROOF: Fix $j \geq 0$. We define a function $\sharp : \mathbb{Z} \to \mathbb{Z}$ by: $\sharp(L) := L + 1$ if $L \geq -1$, and $\sharp(L) :=$ 0 if $L \le -1$. By counting the elements of the basis of $L(sP^j_{\infty})$ given in Theorem [3.1](#page-6-0) we get that for any $s \in \mathbb{Z}$: dim $L(sP_{\infty}^j) = \sum_{m=0}^{2^j-1} \sharp(l_m(s))$. Define $\Delta(s) := \dim L(sP_{\infty}^j) - \dim L((s-1)P_{\infty}^j) =$ $\sum_{m=0}^{2^{j}-1} [\sharp(l_{m}(s)) - \sharp(l_{m}(s-1))]$. By writing $s-m=2^{j}q+r$ with $0 \leq r < 2^{j}$, we can deduce that for all m and s:

$$
l_m(s) - l_m(s-1) = \left\lfloor \frac{s-m}{2^j} \right\rfloor - \left\lfloor \frac{s-1-m}{2^j} \right\rfloor = \begin{cases} 1 & \text{if } m = s \mod 2^j \\ 0 & \text{if } m \neq s \mod 2^j \end{cases}
$$

.

It follows that $\Delta(s) = \sharp(l_{m(s)}(s)) - \sharp(l_{m(s)}(s-1))$. Then, for any $s \in \mathbb{Z}$:

$$
s \in H(P_{\infty}^{j})
$$
 \iff $\Delta(s) = 1$ \iff $l_{m(s)}(s) \ge 0$ \iff $q(s) - c_{m(s)}^{(j)} \ge 0.$

Corollary 3.3 With the same notation of Theorem [3.1](#page-6-0) we have that the pole divisor of the function $w_m^{(j)}$ is:

$$
(w_m^{(j)})_{\infty} = (2^j c_m^{(j)} + m) P_{\infty}^j.
$$

Moreover, the set:

$$
\{2^j\}\cup\{2^j c_m^{(j)}+m\,|\,0
$$

generates $H(P_{\infty}^j)$.

PROOF: Fix m and define $s := 2^{j}c_m^{(j)} + m$. Since $l_m(s-1) = -1$ and $l_m(s) = 0$ then $w_m^{(j)} \in \mathbb{R}$ $L(sP^j_\infty)\backslash L((s-1)P^j_\infty)$. The function x_0 has a unique pole of order 2^j at P^j_∞ , then we are done by Theorem [3.2.](#page-6-1) (We remark that this is not a minimal set of generators in general.)

Thanks to Theorem [2.9,](#page-5-0) the construction of a basis of $L(sP_{\infty}^j)$ was reduced, in the proof of Theorem [3.1,](#page-6-0) to the construction of a basis of $L([sP^j_{\infty}+(x_j)]_{|T_{j-1}})$. We state the main technical result of the paper.

Theorem 3.4 There exists an algorithm that, for all $j \ge 1$ and all odd m with $0 < m < 2^j$, constructs integers $\tilde{c}_m^{(j-1)}$ and functions $\tilde{w}_m^{(j-1)} \in T_{j-1}$ s.t. for all $s \in \mathbb{Z}$ the family parametrized by m and l:

> $x_0^l \tilde{w}_m^{(j-1)}$ $\begin{array}{ll} (j-1) \\ m \end{array}$ $m \text{ odd}$ $0 < m < 2^j$ $0 \leq l \leq \left\lfloor \frac{s-m}{2^j} \right\rfloor - \tilde{c}_m^{(j-1)}$

is a basis of $L([sP^j_{\infty}+(x_j)]_{|T_{j-1}})$.

PROOF: The result is a direct consequence of Proposition [3.7.](#page-9-1) Item (ii) applied with $k = j - 1$ gives a basis of $L(A_1^{j-1}(s)) := L([sP^j_{\infty} + (x_j)]_{|T_{j-1}})$. Thanks to item (i) we can reorder the families in such a way to get the basis in the stated form.

 \Box

3.1 Basis of $L([sP^j_{\infty}+(x_j)]_{|T_{j-1}})$: first part.

This subsection and the next one are dedicated to the proof of Theorem [3.4.](#page-7-1) We fix once and for all an integer $j \geq 1$. All of what will be defined in the sequel will depend on j but we will not indicate this fact in the notation.

In this subsection we will construct divisors $A_n^k(s)$ and $B_n^k(s)$ at level $k \leq j-1$. In the next subsection suitable bases of the Riemann-Roch spaces of these divisors will be constructed.

Proposition 3.5 There is an algorithm to construct divisors $A_n^k(s)$ and $B_n^k(s)$, $s \in \mathbb{Z}$, and integers a_n^k , b_n^k , α_n^k , γ_n^k , and δ_n^k , labeled by integers k and n s.t. $0 \le k \le j-1$ and $1 \le n \le 2^{j-k-1}$, in such a way that the following conditions are satisfied.

- (i) $A_1^{j-1}(s) = [sP^j_{\infty} + (x_j)]_{|T_{j-1}}.$
- (ii) The divisors $A_n^k(s)$ and $B_n^k(s)$ have the form:

(a)
$$
A_n^k(s) = \left[\frac{s + a_n^k}{2^{j-k}} \right] P_{\infty}^k + \sum_{m=-2}^{k-1} \alpha_{n,m}^k D_m^k + \alpha_n^k P_{-1}^k
$$

(b)
$$
B_n^k(s) = \left[\frac{s + b_n^k}{2^{j-k}} \right] P_{\infty}^k + \sum_{m=-2}^{k-1} \beta_{n,m}^k D_m^k.
$$

- (iii) $B_n^k(s) = A_n^k(s) + \gamma_n^k(1 + x_k) + \delta_n^k P_{-1}^k$.
- (iv) If $k < j 1$, and $\lceil \cdot \rceil$ is the roof function, then:

$$
A_n^k(s) = \begin{cases} [B_{\lceil n/2 \rceil}^{k+1}(s)]_{\lceil T_k} & \text{if } n \text{ is odd} \\ [B_{\lceil n/2 \rceil}^{k+1}(s) + (x_{k+1})]_{\lceil T_k} & \text{if } n \text{ is even.} \end{cases}
$$

- (v) $-\alpha_n^k = \gamma_n^k 2^k + \delta_n^k$ and $0 \leq \delta_n^k < 2^k$.
- (vi) For k fixed, all the odd integers modulo 2^{j-k} appear exactly once: (a) in the sequence of the coefficients a_n^k , as n varies, and (b) in the sequence of the coefficients b_n^k , as n varies.

PROOF: The proof is by descending induction on k , and it is given in three steps. In step 1 we define the divisors A^{j-1} . In step 2, for any k, we define the divisors B^k given the divisors A^k . In step 3, for any $k < j - 1$, we define the divisors A^k given the divisors B^{k+1} .

Step 1. For $k = j - 1$ define, for all $s \in \mathbb{Z}$, $A_1^{j-1}(s) := [sP^j_{\infty} + (x_j)]_{|T_{j-1}}$, so that (i) is satisfied by definition. Define the coefficients $a_1^{j-1} := -1$ and $\alpha_1^{j-1} := 0$. Since the divisor (x_j) at level j has coefficient -1 at P_{∞}^{j} , see Proposition [2.5,](#page-4-1) then (ii, a) is satisfied. Moreover, (vi, a) is trivially satisfied.

Step 2. Fix k with $0 \le k \le j - 1$, and assume to have defined divisors $A_n^k(s)$ and integers a_n^k and α_n^k
s.t. (ii, a) and (vi, a) are satisfied for the given k. Define γ_n^k and δ_n^k by dividing $-\alpha_n^k$ by 2^k , is satisfied for the given k by definition. Define $B_n^k(s)$ by (iii) , so that (iii) is satisfied for the given k by definition. We will define coefficients b_n^k such that (ii, b) and (vi, b) are satisfied for the given k.

By the form of the divisor $(1 + x_k)$, see Proposition [2.5,](#page-4-1) item (ii, b) follows by taking care of the coefficients of $B_n^k(s)$ at P_{∞}^k and P_{-1}^k . Since the coefficient of $(1+x_k)$ at P_{-1}^k is 2^k then by (v) it is easily seen that the coefficient of $B_n^k(s)$ at P_{-1}^k is zero as desired. Since the coefficient of $(1+x_k)$ at P_{∞}^k is -1 then the coefficient of $B_n^k(s)$ at P_{∞}^k is given by $\lfloor (s + a_n^k - 2^{j-k} \gamma_n^k)/2^{j-k} \rfloor$. We define $b_n^k := a_n^k - 2^{j-k} \gamma_n^k$
and this concludes with items (ii, b) and (vi, b) for the given k, the latter because $b_n^k = a_n$

Step 3. Fix k with $0 \le k < j - 1$, and assume to have defined divisors $B_n^{k+1}(s)$ and integers b_n^{k+1} s.t. (ii, b) and (vi, b) are satisfied for $k + 1$ in place of k. We will define divisors $A_n^k(s)$ and integers a_n^k and α_n^k s.t. (ii, a) and (vi, a) are satisfied for k.

Define $A_n^k(s)$ by (iv) , so that (iv) is satisfied for k. Let:

$$
a_n^k := \begin{cases} b_{\lceil n/2 \rceil}^{k+1} & \text{if } n \text{ is odd} \\ b_{\lceil n/2 \rceil}^{k+1} - 2^{j-k-1} & \text{if } n \text{ is even.} \end{cases}
$$

By Proposition [2.5,](#page-4-1) the coefficient of the divisor (x_{k+1}) at P_{∞}^{k+1} is equal to -1 , then (ii, a) is satisfied for k. In order to prove (vi, a) we argue as follows. For $n = 1, ..., 2^{j-k-1}$ with n odd, the number $\lceil n/2 \rceil$ takes all the values $1, ..., 2^{j-(k+1)-1}$ exactly once. The same is true taking the even values of n. It follows that the set $\{a_n^k | n = 1, ..., 2^{j-k}\}$ coincides with the set: $\{b_n^{k+1}, b_n^{k+1} - 2^{j-k-1} | n = 1, ..., 2^{j-(k+1)-1}\}$. Since (v_i, b) holds with $k+1$ in place of k by hypothesis, then the b_n^{k+1} 's take exactly once all the odd residues modulo $2^{j-(k+1)}$. The conclusion (vi, a) for the given k follows.

The next corollary will be used in the next subsection to construct bases of the divisors given in the above proposition.

Corollary 3.6 For all integers k, n, s with $0 \le k < j - 1$ and $1 \le n \le 2^{j-k-2}$ we have:

$$
L(B_n^{k+1}(s)) = L(A_{2n-1}^k(s)) \oplus L(A_{2n}^k(s))x_{k+1}.
$$

PROOF: It follows by item (ii, b) of Proposition [3.5](#page-7-3) (applied with $k+1$ in place of k) that $B_n^{k+1}(s)$ is invariant in T_{k+1}/T_k . The thesis of the corollary follows by Theorem [2.9](#page-5-0) and item (iv) of Proposition [3.5.](#page-7-3) \Box

3.2 Basis of $L([sP^j_{\infty}+(x_j)]_{|T_{j-1}})$: second part.

In this subsection we will complete the proof of Theorem [3.4](#page-7-1) by computing bases of the Riemann-Roch spaces $L(A_n^k(s))$ and $L(B_n^k(s))$ of the divisors constructed in Subsection [3.1.](#page-7-0) The integer $j \ge 1$ will be fixed once and for all.

Proposition 3.7 There is an algorithm to construct functions w_m^k , $z_m^k \in T_k$ and integers b_m , c_m^k , d_m^k
labeled by integers k and m s.t. $0 \le k \le j - 1$ and $1 \le m \le 2^{j-1}$, in such a way that the following conditions are satisfied.

- (i) For all $m, 0 \leq b_m < 2^j$. Moreover, as m varies, the coefficients b_m take exactly once all the odd values modulo 2^j .
- (ii) For all k, for all n with $1 \le n \le 2^{j-k-1}$, and for all $s \in \mathbb{Z}$, the family parametrized by m and l:

$$
x_0^l w_m^k \qquad (n-1)2^k + 1 \le m \le n2^k \qquad 0 \le l \le \left\lfloor \frac{s-b_m}{2^j} \right\rfloor - c_m^k
$$

is a basis of $L(A_n^k(s))$.

(iii) For all k, for all n with $1 \le n \le 2^{j-k-1}$, and for all $s \in \mathbb{Z}$, the family parametrized by m and l:

$$
x_0^l z_m^k \qquad (n-1)2^k + 1 \le m \le n2^k \qquad 0 \le l \le \left\lfloor \frac{s-b_m}{2^j} \right\rfloor - d_m^k
$$

is a basis of $L(B_n^k(s))$.

PROOF: The proof is by induction on k , and it is given in three steps. In step 1 we construct bases of the spaces $L(B^0)$. In step 2, for any k, we construct bases of the spaces $L(A^k)$ given bases of the spaces $L(B^k)$. In step 3, for any $k < j - 1$, we construct bases of the spaces $L(B^{k+1})$ given bases of the spaces $L(A^k)$.

Step 1. For $k = 0$, and $n = 1, ..., 2^{j-1}$, and $s \in \mathbb{Z}$ we have, by item (ii, b) of Proposition [3.5:](#page-7-3)

$$
B_n^0(s) = \left[\frac{s + b_n^0}{2^j} \right] P_{\infty}^0 + \beta_{n, -2}^0 P_0^0 + \beta_{n, -1}^0 D_{-1}^0.
$$

Define, for $m = 1, ..., 2^{j-1}, z_m^0 := x_0^{-\beta_{m,-2}^0} (1 + x_0^2)^{-\beta_{m,-1}^0}$. Define b_m by a division by 2^j , that is: $-b_m^0 = q_m 2^j + b_m$ and $0 \leq b_m < 2^j$, and define $d_m^0 := q_m - \beta_{m,-2}^0 - 2\beta_{m,-1}^0$. Then item (i) is satisfied thanks to item (v_i, b) of Proposition [3.5](#page-7-3) for $k = 0$. Since T_0 is the rational function field, then the family parametrized by l:

$$
x_0^l z_m^0 = x_0^{l-\beta_{m,-2}^0} (1+x_0^2)^{-\beta_{m,-1}^0} \qquad 0 \le l \le \left\lfloor \frac{s+b_m^0}{2^j} \right\rfloor + \beta_{m,-2}^0 + 2\beta_{m,-1}^0 = \left\lfloor \frac{s-b_m}{2^j} \right\rfloor - d_m^0
$$

is a basis of $L(B_n^0(s))$. Then *(iii)* is satisfied for $k = 0$.

Step 2. Fix k with $0 \le k \le j-1$. Assume that, for $m = 1, ..., 2^{j-1}$, the elements d_m^k and z_m^k have already been defined in such a way that item (iii) is satisfied for the given k. We will define, for $m = 1, ..., 2^{j-1}$, elements c_m^k and w_m^k in such a way that item (ii) is satisfied for k.

Fix n with $1 \le n \le 2^{j-k-1}$. For any $s \in \mathbb{Z}$, let $C_n^k(s) := A_n^k(s) + \gamma_n^k(1+x_k)$. Define $c_m^k := d_{m,\delta}$ and $\tilde{w}_n^k := z_{m,\delta}$, for $m = (n-1)2^k + 1, ..., n2^k$, where $d_{m,\delta}$ and $z_{m,\delta}$ are constructed in Lemma [3.8.](#page-10-0) Then, for any $s \in \mathbb{Z}$ the family parametrized by m and l:

$$
x_0^l \tilde{w}_m^k \qquad (n-1)2^k + 1 \le m \le n2^k \qquad 0 \le l \le \left\lfloor \frac{s-b_m}{2^j} \right\rfloor - c_m^k
$$

is a basis of $L(C_n^k(s))$. Then, defining $w_m^k := \tilde{w}_m^k (1 + x_k)^{\gamma_m^k}$, the family parametrized by m and l:

$$
x_0^l w_m^k \qquad (n-1)2^k + 1 \le m \le n2^k \qquad 0 \le l \le \left\lfloor \frac{s - b_m}{2^j} \right\rfloor - c_m^k
$$

is a basis of $L(A_n^k(s))$ for any $s \in \mathbb{Z}$, as desired.

Step 3. Fix k with $0 \le k < j-1$. Assume that, for $m = 1, ..., 2^{j-1}$, the elements c_m^k and w_m^k have already been defined in such a way that item (ii) is satisfied for the given k. We will define, for $m = 1, ..., 2^{j-1}$, elements d_m^{k+1} and z_m^{k+1} in such a way that item *(iii)* is satisfied with $k+1$ in place of k.

Define $d_m^{k+1} := c_m^k$. For $n = 1, ..., 2^{j-k-2}$, and $m = (n-1)2^{k+1} + 1, ..., n2^{k+1}$, denote:

$$
z_m^{k+1} := \begin{cases} w_m^k & \text{if } m \le (2n-1)2^k \\ w_m^k x_{k+1} & \text{if } m > (2n-1)2^k. \end{cases}
$$

Then, (iii) with $k+1$ in place of k follows by Corollary [3.6.](#page-8-0)

The following lemma is needed in the proof of Proposition [3.7.](#page-9-1)

Lemma 3.8 The notation of step 2 in the proof of Proposition [3.7](#page-9-1) is in order. In particular, integers k and n are fixed with $0 \leq k \leq j-1$ and $1 \leq n \leq 2^{j-k-1}$ and will be dropped from the notation. The simplified notation is $C(s) = B(s) - \delta P$, where $\delta = \delta_n^k$ is the coefficient of $P := P_{-1}^k$ in item (iii) of Proposition [3.5.](#page-7-3) Then, there is an algorithm to construct functions $z_{m,\varepsilon} \in T_k$ and integers $d_{m,\varepsilon}$ parametrized by integers m and ε with $0 \leq \varepsilon \leq \delta$ and $(n-1)2^k + 1 \leq m \leq n2^k$ in such a way that for any $\varepsilon = 0, \ldots, \delta$, and for any $s \in \mathbb{Z}$ the family parametrized by m and l:

$$
x_0^l z_{m,\varepsilon} \qquad (n-1)2^k + 1 \le m \le n2^k \qquad 0 \le l \le \left\lfloor \frac{s - b_m}{2^j} \right\rfloor - d_{m,\varepsilon} \qquad (3.1)
$$

is a basis of $L(B(s) - \varepsilon P)$.

PROOF: The proof is by induction on ε . For $\varepsilon = 0$, define $d_{m,0} := d_m^k$ and $z_{m,0} := z_m^k$, where d_m^k and z_m^k are given in step 2 of the proof of Proposition [3.7.](#page-9-1)

Fix ε with $0 \leq \varepsilon < \delta$, and suppose to have already defined $d_{m,\varepsilon}$ and $z_{m,\varepsilon}$ giving a basis of $L(B(s)-\varepsilon P)$ for any s. We will define integers $d_{m,\varepsilon+1}$ and elements $z_{m,\varepsilon+1}$ that give a basis of $L(B(s) - (\varepsilon+1)P)$ for any s.

Define, for any m and s, $l_m(s) := \left\lfloor \frac{s-b_m}{2^j} \right\rfloor - d_{m,\varepsilon}$. By Lemma [3.9](#page-11-0) there is a unique bijection:

$$
\sigma: \{i \mid 1 \le i \le 2^k\} \longrightarrow \{m \mid (n-1)2^k + 1 \le m \le n2^k\}
$$

s.t. for any $s \in \mathbb{Z}$ we have $l_{\sigma(1)}(s) \leq l_{\sigma(2)}(s) \leq \ldots \leq l_{\sigma(2^k)}(s)$. Next, we prove the following two facts that will be needed in the sequel:

(i) For any $m = (n-1)2^k + 1, ..., n2^k$ we have $v_P(z_{m,\varepsilon}) \ge \varepsilon$.

(ii) There exists m s.t. $v_P(z_{m,\varepsilon}) = \varepsilon$.

Indeed, we will use the fact that formula [\(3.1\)](#page-10-1) is valid for any $s \in \mathbb{Z}$ (for the given ε). In particular, for s big enough we have $l_m(s) \ge 0$ for any m. It follows, taking $l = 0$ for any m, that $z_{m,\varepsilon} \in L(B(s) - \varepsilon P)$; hence, $v_P(z_{m,\varepsilon}) \geq \varepsilon$, and (i) is proved. Since, by Proposition [3.5](#page-7-3) (ii, b) , deg $B(s) \to +\infty$ as $s \to +\infty$, then, applying the Riemann-Roch Theorem, it follows that for s big enough we have $L((B(s)-\varepsilon P)-P) \neq$ $L(B(s) - \varepsilon P)$; thus, it cannot happen that $v_P(z_{m,\varepsilon}) > \varepsilon$ for any m (use that $v_P(x_0) \geq 0$, and Remark [3.10\)](#page-11-1), and item (ii) is proved.

The function $t := 1 - x_0$ is a local parameter around P: since δ is strictly positive then $k > 0$, and we can apply Remark [2.10.](#page-5-2)

For $z \in T_k$ with $v_P(z) \geq \varepsilon$, we denote by $z(P) \in K$ the order ε coefficient in the power series expansion of z around P with respect to t: $z = z(P)t^{\epsilon} + O(t^{\epsilon+1})$. Notice that $z(P) = 0$ if and only if $v_P(z) > \varepsilon$. Denote:

$$
I := \max\{i \mid 1 \le i \le 2^k \text{ and } v_P(z_{\sigma(i),\varepsilon}) = \varepsilon\}.
$$

Notice that $z_{\sigma(I),\varepsilon}(P) \neq 0$. The constructibility of $z_{m,\varepsilon}(P)$ follows by Remark [2.10](#page-5-2) and by the fact that the functions $z_{m,\varepsilon}$ are expressed in terms of the generators x_i : in fact, they are constructed starting from the explicit functions z_m^0 of step 1 of the proof of Proposition [3.7.](#page-9-1)

We are now in the position to define $z_{m,\epsilon+1}$ and $d_{m,\epsilon+1}$. For any m with $(n-1)2^k + 1 \leq m \leq n2^k$ define:

$$
z_{m,\varepsilon+1} := \begin{cases} z_{m,\varepsilon} - \frac{z_{m,\varepsilon}(P)}{z_{\sigma(I),\varepsilon}(P)} z_{\sigma(I),\varepsilon} & \text{if } m \neq \sigma(I) \\ z_{\sigma(I),\varepsilon}(1-x_0) & \text{if } m = \sigma(I) \end{cases}
$$

and:

$$
d_{m,\varepsilon+1} := \begin{cases} d_{m,\varepsilon} & \text{if } m \neq \sigma(I) \\ d_{\sigma(I),\varepsilon} + 1 & \text{if } m = \sigma(I). \end{cases}
$$

We are left to prove that for any $s \in \mathbb{Z}$ the $z_{m,\varepsilon+1}$'s and the $d_{m,\varepsilon+1}$'s give a basis of $L(B(s) - (\varepsilon + 1))$ as in the statement. We isolate the proof of this fact in Lemma [3.11.](#page-11-2)

 \Box

 \Box

The following lemmas and remark are needed in the proof of Lemma [3.8.](#page-10-0)

Lemma 3.9 With the notation of the proof of Lemma [3.8,](#page-10-0) there exists a unique bijection:

$$
\sigma: \{i \mid 1 \le i \le 2^k\} \longrightarrow \{m \mid (n-1)2^k + 1 \le m \le n2^k\}
$$

s.t., for any $s \in \mathbb{Z}$, we have $l_{\sigma(1)}(s) \leq l_{\sigma(2)}(s) \leq \ldots \leq l_{\sigma(2^k)}(s)$.

PROOF: We put a lexicographical order on $\mathbb{Z} \times \mathbb{Z}$ by declaring: $(d, b) \leq (d', b')$ if $d < d'$ or $(d =$ d' and $b \leq b'$). Since as m varies the b_m 's are all distinct, then there exists a unique bijection σ as in the statement s.t.: $(d_{\sigma(1),\varepsilon},b_{\sigma(1)})>(d_{\sigma(2),\varepsilon},b_{\sigma(2)})>...>(d_{\sigma(2^k),\varepsilon},b_{\sigma(2^k)})$. We have to show that such a σ has the desired property. For this it's enough to show that if (d, b) , $(d', b') \in \mathbb{Z} \times \mathbb{Z}$ with $0 \leq b, b' < 2^j$ and $(b, d) \ge (b', d')$ then, for all $s \in \mathbb{Z}$:

$$
\left\lfloor \frac{s-b}{2^j} \right\rfloor - d \le \left\lfloor \frac{s-b'}{2^j} \right\rfloor - d'.
$$

We leave the easy proof to the reader.

Remark 3.10 With the notation of the proof of Lemma [3.8,](#page-10-0) for any $s \in \mathbb{Z}$ and for any $z \in L(B(s) - \varepsilon P)$ we have that $v_P(z) \geq \varepsilon$ and:

$$
z \in L(B(s) - (\varepsilon + 1)P)
$$
 \iff $v_P(z) > \varepsilon$ \iff $z(P) = 0.$

Lemma 3.11 With the notation of the proof of Lemma [3.8,](#page-10-0) for any $s \in \mathbb{Z}$ the family parametrized by m and l:

$$
x_0^l z_{m,\varepsilon+1} \qquad (n-1)2^k + 1 \le m \le n2^k \qquad 0 \le l \le \left\lfloor \frac{s-b_m}{2^j} \right\rfloor - d_{m,\varepsilon+1}
$$

is a basis of $L(B(s) - (\varepsilon + 1)P)$.

PROOF: Fix $s \in \mathbb{Z}$. We simplify the notation by writing, for any i with $1 \leq i \leq 2^k$:

$$
z_i = z_{\sigma(i),\varepsilon}, \quad d_i = d_{\sigma(i),\varepsilon}, \quad \bar{z}_i = z_{\sigma(i),\varepsilon+1}, \quad \bar{d}_i = d_{\sigma(i),\varepsilon+1}
$$

and:

$$
l_i = l_{\sigma(i)}(s),
$$
 $\bar{l}_i = \bar{l}_{\sigma(i)}(s)$ $D = B(s) - \varepsilon P.$

With the new notation we have that $l_1 \leq l_2 \leq \ldots \leq l_{2^k}$, and:

$$
\bar{z}_i = \begin{cases}\nz_i - \frac{z_i(P)}{z_I(P)} z_I & \text{if } i \neq I \\
z_I(1 - x_0) & \text{if } i = I\n\end{cases}\n\qquad\n\bar{l}_i = \begin{cases}\nl_i & \text{if } i \neq I \\
l_I - 1 & \text{if } i = I.\n\end{cases}
$$

Recall that for any i we have $v_P(z_i) \geq \varepsilon$, and I is the biggest index with $v_P(z_I) = \varepsilon$.

What is left to prove is the following: if the family parametrized by i and $l: x_0^l z_i$ with $1 \leq i \leq 2^k$ and $0 \leq l \leq l_i$ is a basis of $L(D)$, then the family $x_0^l \overline{z}_i$ with $0 \leq i \leq 2^k$ and $0 \leq l \leq \overline{l}_i$ is a basis of $L(D-P)$.

If $L(D - P) = L(D)$, we start by observing that if i is such that $l_i \geq 0$ then $v_P(z_i) > \varepsilon$ and $z_i(P) = 0$: this is a consequence of the fact that $z_i \in L(D) = L(D-P)$ and Remark [3.10.](#page-11-1) Notice that such values of i are exactly the ones that contribute to the basis of $L(D)$, and that, in particular, $l_I < 0$ since $v_P(z_I) = \varepsilon$. It follows that, for i with $l_i \geq 0$, we have $\overline{z}_i = z_i$ and $\overline{l}_i = l_i$. We conclude that the family $x_0^l \overline{z}_i$ is exactly the same as the family $x_0^l z_i$, and we are done.

If $L(D - P) \subsetneq L(D)$, we start by observing that the codimension is 1 and there exists i with $l_i \geq 0$ and $v_P(z_i) = \varepsilon$. Indeed, if for any i with $l_i \geq 0$ we had $v_P(z_i) > \varepsilon$, then, by Remark [3.10,](#page-11-1) we would have that any element $x_0^l z_i$ that contributes to the basis of $L(D)$ would belong to $L(D - P)$, which would imply $L(D - P) = L(D)$, contrary to the assumption. By the maximality of I we have that $l_I \ge 0$. In particular, the family $x_0^l \bar{z}_i$ has one element less than the family $x_0^l z_i$. We are left to prove that all the elements of $x_0^l \bar{z}_i$ belong to $L(D - P)$ and that they are linearly independent over K.

Let's prove that for any i and for any l with $0 \leq l \leq \bar{l}_i$ then $x_0^l \bar{z}_i \in L(D - P)$.

First, consider the case $i = I$. Then $x_0^l \overline{z}_I = x_0^l z_I - x_0^{l+1} z_I$. Since $\overline{l}_I = l_I - 1$ and $0 \le l \le \overline{l}_I$ then both $x_0^l z_I$ and $x_0^{l+1} z_I$ belongs to $L(D)$, hence $x_0^l \overline{z}_I$ belongs as well. Since $v_P(x_0^l \overline{z}_I) = v_P(x_0^l z_I(1-x_0)) =$ $v_P(z_I) + v_P(1-x_0) = \varepsilon + 1 > \varepsilon$, then we can apply Remark [3.10](#page-11-1) to conclude $x_0^l \overline{z}_I \in L(D-P)$.

Second, consider the case in which $i \neq I$ and $v_P(z_i) > \varepsilon$. Then $z_i(P) = 0$. It follows that $x_0^l \overline{z}_i = x_0^l z_i$. Now, $x_0^l z_i \in L(D - P)$ by $x_0^l z_i \in L(D)$ since $0 \leq l \leq \overline{l}_i = l_i$, and by application of Remark [3.10.](#page-11-1)

Third, consider the case $i \neq I$ and $v_P(z_i) = \varepsilon$. Then $i < I$ and $\overline{l}_i = l_i \leq l_I$. It follows that for $0 \leq l \leq \overline{l}_i$ we have $x_0^l \bar{z}_i = x_0^l z_i - (z_i(P)/z_I(P)) x_0^l z_I \in L(D)$ since both $x_0^l z_i$ and $x_0^l z_I$ belongs to $L(D)$. Moreover, it's clear that $\bar{z}_i(P) = 0$ so that $v_P(x_0^l \bar{z}_i) = v_P(\bar{z}_i) > \varepsilon$, and we can apply Remark [3.10](#page-11-1) to conclude $x_0^l \overline{z}_i \in L(D-P).$

Now, let $\alpha_{i,l} \in K$ s.t. $\sum_{i=1}^{2^k}$ $\sum_{i=1}^{2^k} \sum_{l=0}^{\bar{l}_i} \alpha_{i,l} x_0^l \bar{z}_i = 0$. Write the set $\mathcal{I} := \{i \mid 1 \leq i \leq 2^k\}$ as a disjoint union:

$$
\mathcal{I} = \{I\} \cup \mathcal{I}_0 \cup \mathcal{I}_1
$$

where $\mathcal{I}_0 := \{i \in \mathcal{I} \mid i \neq I \text{ and } v_P(z_i) = \varepsilon\}$ and $\mathcal{I}_1 := \{i \in \mathcal{I} \mid v_P(z_i) > \varepsilon\}$, and split the summation:

$$
\sum_{i=1}^{2^k} \sum_{l=0}^{\bar{l}_i} \alpha_{i,l} x_0^l \bar{z}_i = \sum_{i \in \mathcal{I}_0} \sum_{l=0}^{l_i} \alpha_{i,l} x_0^l \left(z_i - \frac{z_i(P)}{z_I(P)} z_I \right) + \sum_{i \in \mathcal{I}_1} \sum_{l=0}^{l_i} \alpha_{i,l} x_0^l z_i + \sum_{l=0}^{l_I - 1} \alpha_{I,l} (x_0^l z_I - x_0^{l+1} z_I)
$$

where we used the definitions of \overline{z}_i and of \overline{l}_i . (We are using that for $i \in \mathcal{I}_1$ we have $\overline{z}_i = z_i$.) For any $i \in \mathcal{I}$, all the $x_0^l z_i$ appearing in the sum above are with $0 \leq l \leq l_i$. (For the $x_0^l z_l$ in the first summation in the right-hand side, we are using that for $i \in \mathcal{I}_0$ we have $l_i \leq l_i$, by the maximality of I.) Since such elements $x_0^l z_i$ are linearly independent, then it follows that for $i \neq I$ we have $\alpha_{i,l} = 0$. It follows that $\sum_{l=0}^{l_I} (\alpha_{I,l} - \alpha_{I,l-1}) x_0^l z_I = 0$, where in case $l_I \geq 0$ we put $\alpha_{I,-1} := \alpha_{I,l_I} := 0$. (In case $l_I < 0$ there are no coefficients $\alpha_{I,l}$ at all, and we are done.) By the linear independence of the $x_0^l z_I$'s, it follows that $\alpha_{I,l-1} = \alpha_{I,l}$ for any l. Then, by induction on l, all the coefficient $\alpha_{I,l}$ vanish, and the proof of the linear independence is complete.

 \Box

3.3 Weierstrass semigroups at P^j_{∞} up to level 8

As an application of Theorems [3.1](#page-6-0) and [3.2](#page-6-1) we implemented an algorithm in Scilab (www.scilab.org) to compute the Weierstrass semigroups $H^j := H(P^j_{\infty})$ at P^j_{∞} for any $0 \le j \le 8$.

We list the elements of the semigroup H^j by making a list of the non-gap intervals: a_1-b_1, a_2-b_2, \ldots a_n-b_n , a_{n+1} -∞. This means that:

$$
H^j = \bigcup_{k=1}^n [a_k, b_k] \cup [a_{n+1}, \infty).
$$

If $a_k = b_k$ we write a_k-b_k as a_k . In the following table g_j denotes the genus of T_j/K .

	g_i	H^{j}
0	θ	$0-\infty$
$\mathbf{1}$		$0: 2-\infty$
\mathfrak{D}	3	$0: 3-4: 6-\infty$
3	9	0; 6; 8; 11-12; 14- ∞
4	21	0; 12; 15-16; 22-24; 27-32; 34- ∞
5	49	0; 24; 30-32; 44; 46-48; 53-56; 58-64; 68-72; 74- ∞
6	105	$0; 48; 60; 62-64; 88; 92; 94-96; 103; 106-112;$
		115-128; 135-136; 138-144; 147- ∞
	225	0; 96; 120; 124; 126-128; 176; 184; 188; 190-192; 206;
		212-216; 218; 220-224; 230-232; 234-240; 242-256;
		263; 269-272; 276-280; 282-288; 291; 293- ∞

For $j = 8$ we have $g_8 = 465$ and the non-gap intervals are:

References

- [1] Garcia A., Bezerra J. A tower with non-Galois steps which attains the Drinfeld-Vlădut bound, J. Number Theory, 106, no. 1, 142-154, (2004).
- [2] Garcia A., Stichtenoth H., A tower of Artin-Schreier extensions of function Fields attaining the Drinfeld-Vlădut bound, Invent. Math., 121, 211-222, (1995).
- [3] Garcia A., Stichtenoth H., On tame towers over finite fields, J. Reine Angew. Math., 557, 53-80, (2003).
- [4] Garcia A., Stichtenoth H., Explicit towers of function fields over finite fields, in "Topics in Geometry, Coding Theory and Cryptography", Springer, (2007).
- [5] Li W., Maharaj H., Stichtenoth H., Elkies N. D. New optimal tame towers of function fields over small finite fields Algorithmic Number Theory, LNCS, 2369, 233-249, (2002).
- [6] Maharaj H., Code construction on fiber products of Kummer curves, IEEE Transactions on Information Theory, 50, no. 9, 2169-2173, (2004).
- [7] Maharaj H., Explicit constructions of algebraic-geometric codes, IEEE Transactions on Information Theory, 51, no. 2, 714-722, (2005).
- [8] Maharaj H., Hu X., On the qth power algorithm, Finite Fields and Their Applications, 14, 1068-1082, (2008).
- [9] Pellikaan R., Stichtenoth H., Torres F., Weierstrass semigroups in an asymptotically good tower of function fields Finite Fields and Their Applications, 4, 4, 381-392, (1998).
- [10] Shum W. K., Aleshnikov I., Kumar P. V., Stichtenoth H., Deolalikar V. A low complexity algorithm for the construction of algebraic-geometric codes better than the Gilbert-Varshamov bound, IEEE Transactions on Information Theory, 47, no. 6, 2225-2241, (2001).
- [11] Stichtenoth H., Algebraic Function Fields and Codes, Second Edition, Springer, 2009.
- [12] Vlădut S. G., Drinfel'd V.G., Number of points of an algebraic curve, Functional Analysis and Its Applications, 17, 53-54, (1983).
- [13] Voss C., Høholdt T., An explicit construction of a sequence of codes attaining the Tsfasman-Vlădut-Zink bound. The first steps, IEEE Transactions on Information Theory, 43, no. 1, (1997).