

The Number of Multistate Nested Canalizing Functions *

David Murrugarra and Reinhard Laubenbacher
Virginia Bioinformatics Institute and Mathematics Department
Virginia Polytechnic Institute and State University

November 9, 2018

Abstract

Identifying features of molecular regulatory networks is an important problem in systems biology. It has been shown that the combinatorial logic of such networks can be captured in many cases by special functions called nested canalizing in the context of discrete dynamic network models. It was also shown that the dynamics of networks constructed from such functions has very special properties that are consistent with what is known about molecular networks, and that simplify analysis. It is important to know how restrictive this class of functions is, for instance for the purpose of network reverse-engineering. This paper contains a formula for the number of such functions and a comparison to the class of all functions. In particular, it is shown that, as the number of variables becomes large, the ratio of the number of nested canalizing functions to the number of all functions converges to zero. This shows that the class of nested canalizing functions is indeed very restrictive, indicating that molecular net-

works have very special properties. The principal tool used for this investigation is a description of these functions as polynomials and a parameterization of the class of all such polynomials in terms of relations on their coefficients.

1 Introduction

A central problem of molecular systems biology is to understand the structure and dynamics of molecular networks, such as gene regulatory, signaling, or metabolic networks. Some progress has been made in elucidating general design principles of such networks. For instance, in [9] it was shown that certain graph theoretic motifs appear far more often in the topology of regulatory network graphs than would be expected at random. In [6, 7, 10] it was shown that a certain type of Boolean regulatory logic has the kind of dynamic properties one would expect from molecular networks. And in [2] it was shown that logical rules that appear in published Boolean models of regulatory networks are overwhelmingly of this type.

These rules, so-called *nested canalizing* rules, are a special case of *canalizing* rules, which are reminiscent of Waddington's concept of cana-

*This research was supported by the National Science Foundation under Grant Nr. CMMI-0908201.

lyzation in gene regulation [14]. Nested canalizing Boolean rules were shown in [4] to be identical with the class of unate cascade functions, which have been studied extensively in computer engineering. They represent exactly the class of Boolean functions that result in binary decision diagrams of shortest average path length [1]. This in itself has interesting implications for information processing in molecular networks. One consequence of this result is that a formula derived earlier for the number of unate cascade functions of a given number of variables [12] applies to give a formula for the number of nested canalizing Boolean functions, described in [4]. A formula for the number of canalizing Boolean functions had been given in [5].

Many molecular networks cannot be described using the Boolean framework, since more than one threshold for a molecular species might be required to represent different modes of action. There are several frameworks available for multistate discrete models, such as so-called logical models, Petri nets, and agent-based models. It has been shown in [13] and [3] that all these model types can be translated into the general and mathematically well-founded framework of polynomial dynamical systems over a finite number system. In [10] the concept of nested canalizing logical rule has been generalized to such polynomial systems. It has been shown there, furthermore, that a large proportion of rules in multistate discrete models are indeed nested canalizing, showing that this concept captures an important feature of the regulatory logic of molecular networks.

As was pointed out in [5] and [4], knowing the number of nested canalizing rules for a given number of input variables and for a given number of possible variable states is important because on the one hand it provides an estimate of

how plausible it is that such rules have evolved as regulatory principles and, on the other hand, provides an estimate of how restrictive the set of rules is. The latter is important, for instance, for the reverse-engineering of networks. If the set of rules is sufficiently restrictive, then the reverse-engineering problem, which is almost always underdetermined due to limited data, becomes more tractable when restricted to reverse-engineering networks consisting of nested canalizing functions. In this paper we present a formula for the number of nested canalizing functions in a given number of variables and show that the ratio of nested canalizing functions and all multistate functions converges to zero as the number of variables increases. We follow the approach in [4] and solve the problem within the framework of polynomial dynamical systems, which makes it possible to frame it as a problem of counting solutions to a system of polynomial equations.

2 Nested Canalizing Functions

As mentioned in the previous section, it is possible to view most discrete models within the framework of dynamical systems over a finite number system, or finite field. For our purposes we will use the finite fields $\mathbb{F}_p = \{0, 1, \dots, p-1\}$, p an arbitrary prime number, otherwise known as \mathbb{Z}/p , the integers modulo p . Furthermore, we will assume that \mathbb{F}_p is totally ordered under the canonical order, that is, its elements are arranged in linear increasing order, $\mathbb{F}_p = \{0 < 1 < \dots < p-1\}$. Let $\mathbb{F} = \mathbb{F}_p$ for some prime p . We first recall the general definition of a nested canalizing function in variables x_1, \dots, x_n from [10]. The underlying idea is as follows: A rule is nested canalizing, if there exists a variable x

such that, if x receives certain inputs, then it by itself determines the value of the function. If x does not receive these certain inputs, then there exists another variable y such that, if y receives certain other inputs, then it by itself determines the value of the function; and so on, until all variables are exhausted.

Definition 2.1. Let $S_i \subset \mathbb{F}, i = 1, \dots, n$, be subsets that satisfy the property that each S_i is a proper, nonempty subinterval of \mathbb{F} ; that is, every element of \mathbb{F} that lies between two elements of S_i in the chosen order is also in S_i . Furthermore, we assume that the complement of each S_i is also a subinterval, that is, each S_i can be described by a threshold s_i , with all elements of S_i either larger or smaller than s_i . Let σ be a permutation on $\{1, \dots, n\}$.

- The function $f : \mathbb{F}^n \rightarrow \mathbb{F}$ is a nested canalyzing function in the variable order $x_{\sigma(1)}, \dots, x_{\sigma(n)}$ with canalyzing input sets $S_1, \dots, S_n \subset \mathbb{F}$ and canalyzing output values $b_1, \dots, b_n, b_{n+1} \in \mathbb{F}$ with $b_n \neq b_{n+1}$ if it can be represented in the form

$$f(x_1, \dots, x_n) = \begin{cases} b_1 & \text{if } x_{\sigma(1)} \in S_1 \\ b_2 & \text{if } x_{\sigma(1)} \notin S_1, x_{\sigma(2)} \in S_2 \\ \vdots & \\ b_n & \text{if } x_{\sigma(1)} \notin S_1, \dots, x_{\sigma(n)} \in S_n \\ b_{n+1} & \text{if } x_{\sigma(1)} \notin S_1, \dots, x_{\sigma(n)} \notin S_n \end{cases}$$

- The function $f : \mathbb{F}^n \rightarrow \mathbb{F}$ is a nested canalyzing function if it is a nested canalyzing function in some variable order $x_{\sigma(1)}, \dots, x_{\sigma(n)}$ for some permutation σ on $\{1, \dots, n\}$.

It is straightforward to verify that, if $p = 2$, that is $\mathbb{F} = \{0, 1\}$, then we recover the definition in [6] of a Boolean nested canalyzing rule.

As mentioned above, several important classes of multistate discrete models can be represented in the form of a dynamical system $f : \mathbb{F}^n \rightarrow \mathbb{F}^n$, so that the concept of a nested canalyzing rule defined in this way has broad applicability.

3 Polynomial form of nested canalyzing functions

We now use the fact that any function $f : \mathbb{F}^n \rightarrow \mathbb{F}$ can be expressed as a polynomial in n variables [8, p. 369]. In this section we determine the polynomial form of nested canalyzing functions. That is, we will determine relationships among the coefficients of a polynomial that make it nested canalyzing. We follow the approach in [4]. Let B_n be the set of functions from \mathbb{F}^n to \mathbb{F} , i.e., $B_n = \{f : \mathbb{F}^n \rightarrow \mathbb{F}\}$. The set B_n is endowed with an addition and multiplication that is induced from that of \mathbb{F} , which makes it into a ring. Let I be the ideal of the ring of polynomials $\mathbb{F}[x_1, \dots, x_n]$ generated by the polynomials $\{x_i^p - x_i\}$ for all $i = 1, \dots, n$, where p is the number of elements in \mathbb{F} . There is an isomorphism between B_n and the quotient ring $\mathbb{F}[x_1, \dots, x_n]/I$ which is also isomorphic to

$$R = \left\{ \sum_{\substack{(i_1, \dots, i_n) \\ i_t \in \mathbb{F} \\ t=1, \dots, n}} C_{i_1 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \right\}$$

Now we use this identification to study nested canalyzing functions as elements of R .

Given a subset S of \mathbb{F} , we will denote by Q_S the indicator function of the complement of S , i.e., for $x_0 \in \mathbb{F}$, let

$$Q_S(x_0) = \begin{cases} 0 & \text{if } x_0 \in S \\ 1 & \text{if } x_0 \notin S \end{cases}$$

We will derive the polynomial form for $Q_S(x)$ in Lemma 8.2. The following theorem gives the polynomial form of a nested canalyzing function.

Theorem 3.1. *Let f be a function in R . Then the function f is nested canalyzing in the variable order x_1, \dots, x_n with canalyzing input sets S_1, \dots, S_n and canalyzing output values b_1, \dots, b_n, b_{n+1} with $b_n \neq b_{n+1}$, if and only if it has the polynomial form*

$$f(x_1, \dots, x_n) = \sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=1}^{n-j} Q_{S_i}(x_i) \right\} + b_1 \quad (3.1)$$

where Q_{S_i} is defined as in Lemma 8.2.

Proof. Let f be a nested canalyzing function as in Definition 2.1, and let

$$g(x_1, \dots, x_n) = \sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=1}^{n-j} Q_{S_i}(x_i) \right\} + b_1$$

Since g has the right form to be in R , we can use the isomorphism between B_n and R , to reduce the proof to showing that

$$g(a_1, \dots, a_n) = f(a_1, \dots, a_n)$$

for all $(a_1, \dots, a_n) \in \mathbb{F}^n$.

If $a_1 \in S_1$, then $Q_{S_1}(a_1) = 0$, therefore

$$g(a_1, \dots, a_n) = b_1 \text{ whenever } a_1 \in S_1.$$

If $a_1 \notin S_1$ and $a_2 \in S_2$, then $Q_{S_1}(a_1) = 1$ and $Q_{S_2}(a_2) = 0$, therefore

$$g(a_1, \dots, a_n) = (b_2 - b_1) + b_1 = b_2.$$

Iterating this process, if $a_1 \notin S_1, a_2 \notin S_2, \dots, a_n \in S_n$, then $Q_{S_1}(a_1) = 1, Q_{S_2}(a_2) = 1, \dots$, and $Q_{S_n}(a_n) = 0$, therefore

$$g(a_1, \dots, a_n) = (b_n - b_{n-1}) + \dots + (b_2 - b_1) + b_1 = b_n.$$

Finally, if $a_1 \notin S_1, \dots, a_n \notin S_n$, then $Q_{S_1}(a_1) = 1, Q_{S_2}(a_2) = 1, \dots$, and $Q_{S_n}(a_n) = 1$. Therefore,

$$g(a_1, \dots, a_n) = (b_{n+1} - b_n) + \dots + (b_2 - b_1) + b_1 = b_{n+1}$$

This completes the proof. \square

Theorem 3.2. *Let f be a function in R . Then the function f is nested canalyzing in the variable order $x_{\sigma(1)}, \dots, x_{\sigma(n)}$ with canalyzing input sets S_1, \dots, S_n and canalyzing output values b_1, \dots, b_n, b_{n+1} with $b_n \neq b_{n+1}$, if and only if it has the polynomial form*

$$f(x_1, \dots, x_n) = \sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=1}^{n-j} Q_{S_{\sigma(i)}}(x_i) \right\} + b_1$$

where $Q_{S_{\sigma(i)}}$ is defined as in Lemma 8.2.

Proof. The proof is very similar to the proof of Theorem 3.1. \square

4 The algebraic variety of nested canalyzing functions

Here we derive a parametrization for the coefficients of any nested canalyzing function. We will use this parametrization to derive a formula to compute the number nested canalyzing functions for a given number of variables within a finite field in the next section.

Recall that elements of $B_n = \{f : \mathbb{F}^n \rightarrow \mathbb{F}\}$ can be seen as elements of

$$R = \left\{ \sum_{\substack{(i_1, \dots, i_n) \\ i_t \in \mathbb{F} \\ t=1, \dots, n}} C_{i_1 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \right\}$$

Now, as a vector space over \mathbb{F} , R is isomorphic to \mathbb{F}^{p^n} via the correspondence

$$\sum_{\substack{(i_1, \dots, i_n) \\ i_t \in \mathbb{F} \\ t=1, \dots, n}} C_{i_1 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \leftrightarrow (\dots, C_{i_1 \dots i_n}, \dots)$$

We will identify the set of nested canalizing functions in R with a subset V^{ncf} of \mathbb{F}^{p^n} by imposing relations on the coordinates of its elements. We are going to use the following notation:

Notation 4.1. For $r \in \mathbb{F}$ and for $S \subset \mathbb{F}$,

$$C_{[r]} = C_{r \dots r}$$

$$C_{[r] \setminus \{i\}}^j = C_{r \dots j \dots r} \text{ where the value } j$$

goes in the i th position.

$$C_{[r] \setminus \{1, \dots, j\}}^{i_1, \dots, i_j} = C_{i_1, \dots, i_j, r \dots r}$$

where the values i_1, \dots, i_j go in the

$1, \dots, j$ positions, respectively.

$$C_{i_1 \dots i_{n-j}} = C_{i_1 \dots i_{n-j} 0 \dots 0}, \text{ i.e., } i_{n-j} \neq 0 \text{ and}$$

$$i_s = 0 \text{ for all } s > n - j.$$

$$S^c = \mathbb{F} \setminus S, \text{ i.e., } S^c \text{ denote the complement of } S.$$

Theorem 4.2. Let $f \in B_n$ be given by

$$f(x_1, \dots, x_n) = \sum_{\substack{(i_1, \dots, i_n) \\ i_t \in \mathbb{F} \\ t=1, \dots, n}} C_{i_1 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n} \quad (4.1)$$

The polynomial $f(x_1, \dots, x_n)$ is a nested canalizing function in the variable order x_1, \dots, x_n with canalizing input sets S_1, \dots, S_n and canalizing output values b_1, \dots, b_{n+1} if and only if its coefficients satisfy the following equations:

$$C_{i_1 \dots i_{n-\mu}} = C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} \prod_{j=1}^{n-\mu} C_{[p-1]}^{-1} C_{[p-1] \setminus \{j\}}^{i_j} \quad (4.2)$$

for $\mu = 0, \dots, n - 1$, where

$$C_{[p-1]} = (b_{n+1} - b_n)(p - 1)^n \prod_{i=1}^n |S_i^c|, \quad (4.3)$$

$$C_{[p-1] \setminus \{j\}}^{i_j} = (p - 1) |S_j^c|^{-1} \left(\sum_{r \in S_j^c} r^{p-1-i_j} \right) C_{[p-1]}, \quad (4.4)$$

for $i_j \neq 0, p - 1$, and

$$C_{[p-1] \setminus \{j\}}^0 = (p - 1) |S_j^c|^{-1} Q_{s_j}(0) C_{[p-1]}, \quad (4.5)$$

for $j = 1, \dots, n - 1$.

$$C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} = \left[\prod_{i=1}^{n-\mu} (p - 1) |S_i^c| \right] \sum_{j=0}^{\mu} \left\{ B_{n-j} \prod_{i=n-\mu+1}^{n-j} Q_{s_j}(0) \right\}, \quad (4.6)$$

$$C_{[0]\setminus\{1\}}^{p-1} C_{[p-1]\setminus\{1\}}^0 = C_{[p-1]} \left(C_{[0]} - b_1 \right) \quad (4.7)$$

where $Q_{S_j}(0)$ is defined as in Lemma 8.2 and $B_{n-j} = (b_{n-j+1} - b_{n-j})$ for $j = 1, \dots, n-1$.

The proof of Theorem 4.2 is given in Appendix 2. We now need to provide a similar parametrization for functions that are nested canalyzing with respect to an arbitrary variable ordering.

Theorem 4.3. *Let $f \in B_n$ given by*

$$f(x_1, \dots, x_n) = \sum_{\substack{(i_1, \dots, i_n) \\ i_t \in \mathbb{F} \\ t=1, \dots, n}} C_{i_1 \dots i_n} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n}. \quad (4.8)$$

The polynomial $f(x_1, \dots, x_n)$ is a nested canalyzing function in the variable order $x_{\sigma(1)}, \dots, x_{\sigma(n)}$ with canalyzing input sets S_1, \dots, S_n and canalyzing output values b_1, \dots, b_{n+1} if and only if,

$$C_{i_1 \dots i_{n-\mu}} = C_{[0]\setminus\{\sigma(i_1) \dots \sigma(i_{n-\mu})\}}^{p-1, \dots, p-1} \prod_{j=1}^{n-\mu} C_{[p-1]}^{-1} C_{[p-1]\setminus\{\sigma(j)\}}^{i_j} \quad (4.9)$$

for $\mu = 0, \dots, n-1$, where

$$C_{[p-1]} = (b_{n+1} - b_n) (p-1)^n \prod_{i=1}^n |S_{\sigma(i)}^c|, \quad (4.10)$$

$$C_{[p-1]\setminus\{\sigma(j)\}}^{i_j} = (p-1) |S_{\sigma(j)}^c|^{-1} \left(\sum_{r \in S_{\sigma(j)}^c} r^{p-1-i_j} \right) C_{[p-1]}, \quad (4.11)$$

for $i_j \neq 0, p-1$, and

$$C_{[p-1]\setminus\{\sigma(j)\}}^0 = (p-1) |S_{\sigma(j)}^c|^{-1} Q_{S_{\sigma(j)}}(0) C_{[p-1]}, \quad (4.12)$$

for $j = 1, \dots, n-1$.

$$C_{[0]\setminus\{\sigma(i_1) \dots \sigma(i_{n-\mu})\}}^{p-1, \dots, p-1} = \left[\prod_{i=1}^{n-\mu} (p-1) |S_{\sigma(i)}^c| \right] \sum_{j=0}^{\mu} \left\{ B_{n-j} \prod_{i=n-\mu+1}^{n-j} Q_{S_{\sigma(i)}}(0) \right\}, \quad (4.13)$$

$$C_{[0]\setminus\{\sigma(1)\}}^{p-1} C_{[p-1]\setminus\{\sigma(1)\}}^0 = C_{[p-1]} \left(C_{[0]} - b_1 \right) \quad (4.14)$$

where $Q_{S_{\sigma(j)}}(0)$ is defined as in Lemma 8.2 and

$$B_{n-j} = (b_{n-j+1} - b_{n-j}) \text{ for } j = 1, \dots, n-1.$$

Proof. The proof follows the same line of reasoning used for the proof of Theorem 4.2. \square

Remark 4.1. Notice from Equation 4.10 that a nested canalyzing function $f(x_1, \dots, x_n)$ with canalyzing input sets S_1, \dots, S_n and canalyzing output values b_1, \dots, b_n, b_{n+1} is also a nested canalyzing function in the same variable order with canalyzing input sets S_1, \dots, S_n^c and canalyzing output values b_1, \dots, b_{n+1}, b_n . In fact, Equation 4.10 implies that

$$\begin{aligned}
C_{[p-1]} &= (b_{n+1} - b_n)(p-1)^n \prod_{i=1}^n |S_{\sigma(i)}^c| \\
&= (b_n - b_{n+1}) |S_{\sigma(n)}| (p-1)^n \prod_{i=1}^{n-1} |S_{\sigma(i)}^c|
\end{aligned}$$

Note that $-|S_{\sigma(n)}^c| = |S_{\sigma(n)}| \pmod{p}$.

Remark 4.2. For every nonzero $b \in \mathbb{F}$ and for every nested canalizing function $f(x_1, \dots, x_n)$ in the variable order $x_{\sigma(1)}, \dots, x_{\sigma(n)}$ with canalizing input sets S_1, \dots, S_n and canalizing output values b_1, \dots, b_{n+1} , $f(x_1, \dots, x_n) + b$ is also a nested canalizing function in the same variable order and with the same canalizing input sets S_1, \dots, S_n and canalizing output values $b_1 + b, \dots, b_{n+1} + b$. In fact, for S_1, \dots, S_n and $b_1 + b, \dots, b_{n+1} + b$ Equations 4.9 - 4.13 stay the same and Equation 4.14 becomes

$$C_{[0] \setminus \{\sigma(1)\}}^{p-1} C_{[p-1] \setminus \{\sigma(1)\}}^0 = C_{[p-1]} \left(C_{[0]} - (b_1 + b) \right)$$

5 Number of nested canalizing functions

Here we derive a formula to compute the number of nested canalizing functions in a given number of variables n and a given finite field \mathbb{F} with p elements.

Let us denote the number of distinct nested canalizing functions in n variables by $NCF(n)$ and the number of distinct nested canalizing functions that can be written as a product of r nested canalizing functions by $RNCF(n, r)$ (R for reducible). It is clear from Formula 3.1 that any nested canalizing function can be written as a product of at most n nested canalizing functions. Hence $RNCF(n, r) = 0$ for all

$r > n$. We will denote the number of distinct nested canalizing functions in n variables that cannot be written as a product of two or more nested canalizing functions by $INCF(n)$ (I for irreducible) and the number of distinct nested canalizing functions in n variables that can be written as a product of two or more nested canalizing functions by $RNCF(n)$. Then

$$RNCF(n) = \sum_{r=2}^n RNCF(n, r)$$

The following lemma relates $RNCF(n)$ and $INCF(n)$.

Lemma 5.1. *For each natural number n , we have $INCF(n) = (p-1)RNCF(n)$.*

Proof. From Remark 4.2, for each $f \in RNCF(n)$, there are $p-1$ functions in $INCF(n)$. Conversely, Let $f \in INCF(n)$. From formula 3.1,

$$\begin{aligned}
f(x_1, \dots, x_n) &= \\
&\sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=1}^{n-j} Q_{S_i}(x_i) \right\} + b_1
\end{aligned}$$

Let $b = p - b_1$. Hence

$$\begin{aligned}
f(x_1, \dots, x_n) + b &= \\
&\sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=1}^{n-j} Q_{S_i}(x_i) \right\} \\
&= Q_{S_1}(x_1) \left[\sum_{j=0}^{n-2} \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=2}^{n-j} Q_{S_i}(x_i) \right\} \right. \\
&\quad \left. + (b_2 - b_1) \right]
\end{aligned}$$

Therefore, any element of $INCF(n)$ can be obtained from an element of $RNCF(n)$. \square

The following theorem gives us a formula to compute the number of nested canalyzing functions for a given number of variables n .

Theorem 5.2. *The number of nested canalyzing functions in n variables, denoted by $NCF(n)$, is given by*

$$NCF(n) = pRNCF(n),$$

where

$$RNCF(1) = (p-1)^2,$$

$$RNCF(2) = 4(p-1)^4,$$

and, for $n \geq 3$,

$$RNCF(n) =$$

$$\sum_{r=2}^{n-1} \binom{n}{r-1} 2^{r-1} (p-1)^r RNCF(n-r+1) + 2^{n-1} (p-1)^{n+1} (2+n(p-2)).$$

Proof. From Remark 4.2, in order to calculate $NCF(1)$ it is enough to calculate the number of nested canalyzing functions with canalyzing output values $(0, b_2)$, with $b_2 \neq 0$, and then multiply by p , because of the isomorphism between the sets of vectors $\{(0, b_2)\}$ and $\{(b_1, b_2)\}$ with $b_2 \neq b_1$. This isomorphism is given by

$$(0, b_2) \xrightarrow{+(b_1, b_1)} (b_1, b_1 + b_2)$$

for $b_1 = 1, \dots, p-1$. For output values of the form $(0, b_2)$, Formula 3.1 gives us

$$f(x_1) = b_2 Q_{S_1}(x_1)$$

There are $p-1$ choices for b_2 and $p-1$ choices for $Q_{S_1}(x_1)$. Note that we do not consider $2(p-1)$ choices for $Q_{S_1}(x_1)$ because a nested canalyzing

function with canalyzing input set S_1 and canalyzing output values $(0, b_2)$ is also a nested canalyzing function with canalyzing input set S_1^c and canalyzing output values $(b_2, 0)$ (see remark 4.1). Therefore

$$RNCF(1) = (p-1)^2,$$

Similarly, in order to calculate $NCF(2)$, it is enough to calculate the number of nested canalyzing functions with canalyzing output values $(0, b_2, b_3)$, with $b_3 \neq b_2$, and then multiply by p , because of the isomorphism between the sets of vectors $\{(0, b_2, b_3)\}$ and $\{(b_1, b_2, b_3)\}$ given by

$$(0, b_2, b_3) \xrightarrow{+(b_1, b_1, b_1)} (b_1, b_1 + b_2, b_1 + b_3)$$

for $b_1 = 1, \dots, p-1$. For output values of the form $(0, 0, b_3)$ formula 3.1 gives us

$$f(x_1) = b_3 Q_{S_1}(x_1) Q_{S_2}(x_2)$$

There are $p-1$ choices for b_3 and $2(p-1)$ choices for each of the $Q_{S_i}(x_i)$ for $i = 1, 2$. Therefore, for $(0, 0, b_3)$ there are $4(p-1)^3$ functions.

For $(0, b_2, b_3)$, where $b_2 \neq 0$, $b_3 \neq 0$, and $b_3 \neq b_2$, Formula 3.1 give us

$$f(x_1) = Q_{S_1}(x_1) \left((b_3 - b_2) Q_{S_2}(x_2) + b_2 \right)$$

There are $2(p-1)$ choices for $Q_{S_1}(x_1)$, $p-1$ choices for b_2 , $p-2$ choices for b_3 , and $p-1$ choices for $Q_{S_2}(x_2)$. Note that we do not consider $2(p-1)$ choices for $Q_{S_2}(x_2)$ because a nested canalyzing function with canalyzing input sets S_1, S_2 and canalyzing output values $(0, 0, b_3)$ is also a nested canalyzing function with canalyzing input sets S_1, S_2^c and canalyzing output values $(0, b_3, 0)$ (see Remark 4.1). Therefore, there are $2(p-1)^3(p-2)$ functions. If we count

the number of functions after permuting the variables, we get

$$RNCF(2) = 4(p-1)^3 + \binom{2}{1}2(p-1)^3(p-2),$$

Simplifying the formula above we get

$$RNCF(2) = 4(p-1)^4.$$

For $n \geq 3$ let us compute the number of distinct nested canalyzing functions that can be written as a product of n nested canalyzing functions, $RNCF(n, n)$. From Formula 3.1 it is clear that a nested canalyzing function can be written as a product of n nested canalyzing functions if and only if the output values must have the form $(0, \dots, 0, b_n, b_{n+1})$, where $b_{n+1} \neq b_n$. First consider the case where $b_n = 0$, i.e. the output values have the form $(0, \dots, 0, b_{n+1})$, with $b_{n+1} \neq 0$, for which we have:

$$f(x_1, \dots, x_n) = b_{n+1}Q_{S_1}(x_1) \dots Q_{S_n}(x_n)$$

There are $p-1$ choices for b_{n+1} and $2(p-1)$ choices for each of the $Q_{S_i}(x_i)$ for $i = 1, \dots, n$. Therefore, there are $2^n(p-1)^{n+1}$ functions. Note that if we permute the variables, we will still get the same functions.

For $(0, \dots, 0, b_n, b_{n+1})$, where $b_{n+1} \neq 0$, $b_n \neq 0$, and $b_{n+1} \neq b_n$, Formula 3.1 gives us

$$f(x_1, \dots, x_n) = Q_{S_1}(x_1) \dots Q_{S_{n-1}}(x_{n-1}) \left\{ (b_{n+1} - b_n)Q_{S_n}(x_n) + b_n \right\}$$

There are $2(p-1)$ choices for each $Q_{S_i}(x_i)$ for $i = 1, \dots, n-1$, $p-1$ choices for b_n , $p-2$ choices for b_{n+1} , and $p-1$ choices for $Q_{S_n}(x_n)$. Note that we do not consider $2(p-1)$ choices for

$Q_{S_n}(x_n)$ because a nested canalyzing function with canalyzing input sets S_1, \dots, S_n and canalyzing output values $(0, \dots, 0, b_n, b_{n+1})$ is also a nested canalyzing function with canalyzing input set $S_1, \dots, S_{n-1}, S_n^c$ and canalyzing output values $(0, \dots, 0, b_{n+1}, b_n)$ (see Remark 4.1). Therefore, there are $2^{n-1}(p-1)^{n+1}(p-2)$ such functions. If we count the number of functions after permuting the variables, we get

$$\begin{aligned} RNCF(n, n) &= 2^n(p-1)^{n+1} + \binom{n}{1}2^{n-1}(p-1)^{n+1}(p-2) \\ &= 2^{n-1}(p-1)^{n+1}(2 + n(p-2)). \end{aligned}$$

Let us now compute the number of distinct nested canalyzing functions that can be written as a product of $n-r$ nested canalyzing functions, $RNCF(n, n-r)$ for $r = 1, \dots, n-2$. From Formula 3.1, it is clear that a nested canalyzing function can be written as a product of $n-1$ nested canalyzing functions if and only if the output values have the form $(0, \dots, 0, b_{n-r}, \dots, b_n, b_{n+1})$, where $b_{n-r} \neq 0$ and $b_{n+1} \neq b_n$. In this case we have:

$$f(x_1, \dots, x_n) = \prod_{i=1}^{n-r-1} Q_{S_i}(x_i) \sum_{j=0}^r \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=n-r}^{n-j} Q_{S_i}(x_i) + b_{n-r} \right\}$$

There are $2(p-1)$ choices for each $Q_{S_i}(x_i)$ for $i = 1, \dots, n-r-1$ and $INCF(r+1)$ functions for the summation part. Note that here we do have repetition coming from Remark 4.1 because $b_{n-r} \neq 0$. Therefore, there are $2^{n-r-1}(p-1)^{n-r-1}INCF(r+1)$ such functions. If we count

the number of functions after permuting the variables, we get

$$RNCF(n, n-r) = \binom{n}{r+1} 2^{n-r-1} (p-1)^{n-r-1} INCF(r+1). \quad (5.1)$$

Now since

$$RNCF(n) = \sum_{r=2}^n RNCF(n, r) = \sum_{r=0}^{n-2} RNCF(n, n-r),$$

we have

$$RNCF(n) = \sum_{r=1}^{n-2} RNCF(n, n-r) + RNCF(n, n).$$

Now, replacing equation 5.1 in the previous formula, we have

$$RNCF(n) = \sum_{r=1}^{n-2} \binom{n}{r+1} 2^{n-r-1} (p-1)^{n-r-1} INCF(r+1) + RNCF(n, n).$$

If we make the change of variable $\mu = n-r$, then $\mu-1 = n-r-1$ and $r+1 = n-\mu+1$. Therefore,

$$RNCF(n) = \sum_{\mu=2}^{n-1} \binom{n}{n-\mu+1} 2^{\mu-1} (p-1)^{\mu-1} INCF(n-\mu+1) + RNCF(n, n).$$

But since,

$$\binom{n}{n-(\mu-1)} = \binom{n}{\mu-1},$$

we have

$$RNCF(n) = \sum_{\mu=2}^{n-1} \binom{n}{\mu-1} 2^{\mu-1} (p-1)^{\mu-1} INCF(n-\mu+1) + RNCF(n, n),$$

From Lemma 5.1,

$$RNCF(n) = \sum_{\mu=2}^{n-1} \binom{n}{\mu-1} 2^{\mu-1} (p-1)^{\mu} RNCF(n-\mu+1) + RNCF(n, n),$$

where

$$RNCF(n, n) = 2^{n-1} (p-1)^{n+1} (2 + n(p-2)). \quad (5.2)$$

Finally, again using Lemma 5.1

$$NCF(n) = RNCF(n) + INCF(n) = pRCF(n)$$

This completes the proof. \square

Example 5.3 (Boolean case). *Jarrah et. al. [4] show that the class of Boolean nested canalizing functions is identical to the class of unate cascade functions. Sasao and Kinoshita [12] found a recursive formula for the number of unate cascade functions. Therefore, the same formula can be used to compute the number of Boolean nested canalizing functions. Below is the formula originally given by Sasao and Kinoshita [12] which is a particular case of our formula in Theorem 5.2, namely when $p = 2$:*

$$NCF(n) = 2E(n),$$

Table 1: Number of nested analyzing functions for $p = 3$ and $n = 1, \dots, 8$

n	NCF(n)
1	12
2	192
3	5568
4	219648
5	10834944
6	641335296
7	44288360448
8	3495313145856

Table 2: Number of nested analyzing functions for $p = 5$ and $n = 1, \dots, 8$

n	NCF(n)
1	80
2	5120
3	547840
4	78561280
5	14082703360
6	3029304606720
7	760232846295040
8	218043057365319680

where

$$E(1) = 2, E(2) = 4,$$

and

$$E(n) = \sum_{r=2}^{n-1} \binom{n}{r-1} 2^{r-1} E(n-r+1) + 2^n.$$

Tables 1 - 2 show the number of nested analyzing functions for $p = 3, 5$ and $n = 1, \dots, 8$.

6 Asymptotic properties of $NCF(n)$

In this section we examine the asymptotic properties of the formula given in Theorem 5.2, i.e. we want to know the behavior of $NCF(n)$ as n becomes large.

First we derive the following inequalities:

Lemma 6.1. For fixed n and $r = 2, \dots, n-1$, we have

$$2^{r-1}(p-1)^r RNCF(n-r+1) \leq RNC(n)$$

Proof. It is a direct consequence of the formula given at Theorem 5.2. \square

Lemma 6.2. For all natural numbers n , we have

$$RNCF(n, n) \leq 2^{2n}(p-1)^{n+2}.$$

Proof. From equation 5.2,

$$\begin{aligned} RNCF(n, n) &= 2^{n-1}(p-1)^{n+1}(2+n(p-2)) \\ &\leq 2^{n-1}(p-1)^{n+1}(2+2^n(p-1)) \\ &\leq 2^{n-1}(p-1)^{n+1}(2^{n+1}(p-1)) \\ &\leq 2^{2n}(p-1)^{n+2} \end{aligned}$$

\square

Lemma 6.3. For all natural numbers $n \geq 3$, we have

$$RNCF(n) \leq 2^{n(n-1)}(p-1)^{2n}.$$

Proof. We prove this by induction over n . First

note that for $n = 3$,

$$\begin{aligned}
RNCF(3) &= \binom{3}{1} 2^1 (p-1)^2 RNCF(2) \\
&\quad + 2^2 (p-1)^4 (2 + 3p - 6) \\
&= 24(p-1)^6 + 4(p-1)^4 (3p-4) \\
&\leq 2^5 (p-1)^6 + 4(p-1)^4 (3(p-1) - 1) \\
&\leq 2^5 (p-1)^6 + 2^5 (p-1)^6 \\
&= 2^6 (p-1)^6 = 2^{3(3-1)} (p-1)^{2(3)}.
\end{aligned}$$

Now, assume

$$RNCF(n) \leq 2^{n(n-1)} (p-1)^{2n}.$$

Then

$$\begin{aligned}
RNCF(n+1) &= \\
&\sum_{r=2}^n \binom{n+1}{r-1} 2^{r-1} (p-1)^r RNCF(n-r+2) \\
&\quad + RNCF(n+1, n+1) \\
&= \binom{n+1}{1} 2(p-1)^2 RNCF(n) \\
&\quad + \sum_{r=3}^n \binom{n+1}{r-1} 2^{r-1} (p-1)^r RNCF(n-r+2) \\
&\quad + RNCF(n+1, n+1)
\end{aligned}$$

From Lemma 6.1 we have

$$\begin{aligned}
&\sum_{r=3}^n \binom{n+1}{r-1} 2^{r-1} (p-1)^r RNCF(n-r+2) \\
&= \sum_{r=2}^{n-1} \binom{n+1}{r} 2^r (p-1)^{r+1} RNCF(n-r+1) \\
&\leq \sum_{r=2}^{n-1} \binom{n+1}{r} 2(p-1) RNCF(n)
\end{aligned}$$

Therefore

$$\begin{aligned}
RNCF(n+1) &= \\
&\leq \binom{n+1}{1} 2(p-1)^2 RNCF(n) \\
&\quad + \sum_{r=2}^{n-1} \binom{n+1}{r} 2(p-1) RNCF(n) \\
&\quad + RNCF(n+1, n+1) \\
&\leq 2(p-1)^2 RNCF(n) \sum_{r=1}^{n-1} \binom{n+1}{r} \\
&\quad + RNCF(n+1, n+1)
\end{aligned}$$

Using the inductive hypothesis and Lemma 6.2

we have

$$\begin{aligned}
RNCF(n+1) &= \\
&\leq 2(p-1)^2 2^{n(n-1)} (p-1)^{2n} \sum_{r=1}^{n-1} \binom{n+1}{r} \\
&\quad + 2^{2(n+1)} (p-1)^{n+3} \\
&= 2^{n(n-1)+1} (p-1)^{2(n+1)} \sum_{r=1}^{n-1} \binom{n+1}{r} \\
&\quad + 2^{2(n+1)} (p-1)^{n+3}
\end{aligned}$$

Now since

$$\sum_{r=1}^{n-1} \binom{n+1}{r} = 2^{n+1} - n - 3$$

we have

$$\begin{aligned}
RNCF(n+1) &\leq \\
&2^{n(n-1)+1} (p-1)^{2(n+1)} (2^{n+1}) \\
&\quad + 2^{2(n+1)} (p-1)^{n+3} \\
&\leq 2^{n^2+2} (p-1)^{2(n+1)} \\
&\quad + 2^{n^2+n-1} (p-1)^{2(n+1)} \\
&\leq 2^{(n+1)n} (p-1)^{2(n+1)}.
\end{aligned}$$

Note that $n-1 \geq 2$. This completes our proof. \square

Let us denote the number of all possible functions on n variables by $\psi(n)$. The following theorem show that the set of all nested canalyzing functions is an increasingly smaller subset of the set of all functions.

Theorem 6.4. *The ratio $NCF(n)/\psi(n)$ converges to 0 as n becomes large.*

Proof.

$$\begin{aligned}
\frac{NCF(n)}{\psi(n)} &= \frac{NCF(n)}{p^{p^n}} = \frac{pRNCF(n)}{p^{p^n}} \\
&\leq \frac{p(2^{n(n-1)}(p-1)^{2n})}{p^{p^n}} \\
&\leq \frac{2^{n(n-1)} p^{2n+1}}{p^{p^n}} \\
&\leq \frac{p^{n(n-1)} p^{2n+1}}{p^{p^n}} \\
&= \frac{p^{n^2+n+1}}{p^{p^n}} \rightarrow 0 \text{ as } n \rightarrow \infty
\end{aligned}$$

because the exponential function p^n grows much faster than the quadratic function $n^2 + n + 1$ as n becomes large. \square

7 Discussion

The concept of a nested canalyzing rule has been shown to be a useful approach to elucidating design principles for molecular regulatory networks, and such rules appear very frequently in published network models. But how much of a restriction do such rules impose on the regulatory logic of the network, that is, how “special” are such rules? It was shown in [10] that networks with nested canalyzing rules have very special dynamic properties. In this paper we have shown that nested canalyzing rules do indeed make up a very small subset of all possible rules. In particular, we provide an explicit formula for the number of such rules for a given number of variables.

This was done by translating the problem into the mathematical context of polynomial functions over finite fields and the language of alge-

braic geometry. The formula we provide uses a parametric description of the class of all nested canalyzing polynomials as an algebraic variety. In particular, this provides a very easy way to generate such polynomials through particular parameter choices, which is very useful, for instance, for large-scale simulation studies.

Another interesting aspect of the formula derived in this paper is as a future discovery tool. The formula for Boolean nested canalyzing functions in [4] was obtained essentially through serendipity. Once the parameterization of this class was obtained it was possible to explicitly compute the number of solutions of the parametric equations for small numbers of variables. The resulting integer sequence, giving the number of Boolean nested canalyzing rules for small numbers of variables was matched to the number of Boolean unate cascade functions, for which a formula is known. It was shown in [4] that the two classes of functions are in fact identical. This is of independent interest, since Boolean unate cascade functions have been shown to lead to binary decision diagrams with smallest average path length, suggesting that they are very efficient in processing information. The formula for multistate nested canalyzing rules provides a similar opportunity. While there is no obvious match to other function classes for small fields, it is worth, in our opinion, to pursue this discovery approach further for larger fields.

8 Appendix 1

In this appendix we derive the polynomial form for the indicator functions Q_S . First, for each $r \in \mathbb{F}$ we denote the indicator function of the

singleton set $\{r\}$ by P_r , i.e., for $x_0 \in \mathbb{F}$,

$$P_r(x_0) = \begin{cases} 1 & \text{if } x_0 = r \\ 0 & \text{if } x_0 \neq r \end{cases}$$

The polynomial form of P_r is given in the following lemma.

Lemma 8.1. *For $r \in \mathbb{F}$, we have*

$$P_r(x) = (p-1) \prod_{\substack{a \in \mathbb{F} \\ a \neq r}} (x-a),$$

which has the expanded form

$$P_r(x) = (p-1) \left[x^{p-1} + rx^{p-2} + r^2x^{p-3} + \dots + r^{p-2}x + \prod_{\substack{a \in \mathbb{F} \\ a \neq r}} a \right]$$

Proof. Let $g(x) = (p-1) \prod_{\substack{a \in \mathbb{F} \\ a \neq r}} (x-a)$. We want to prove that $g(x) = P_r(x)$ for all $x \in \mathbb{F}$. Clearly $g(x_0) = 0$ if $x_0 \neq r$. It remains to prove that $g(r) = 1$. From the definition of g , it can be expanded as

$$g(x) = (p-1)(x(x-1)\dots(x-(r-1))\dots(x-(r+1))\dots(x-(p-1))).$$

Then

$$\begin{aligned} g(r) &= (p-1)r!(-1)^{p-1-r}(p-1-r)! \\ &= (p-1)(-1)^r(-1)(-2)\dots(-r) \\ &\quad (-1)^{p-1-r}(p-1-r)! \end{aligned}$$

Now, since $p-i = -i \pmod{p}$ for $i = 1, \dots, r$, we get

$$\begin{aligned} g(r) &= (p-1)(-1)^{p-1}(p-1)(p-2)\dots \\ &\quad (p-r)(p-1-r)! \\ &= (p-1)(p-1)! \\ &= (p-1)(p-1) \text{ (from Wilson's Theorem)} \\ &= 1 \end{aligned}$$

This proves the first assertion. For the second claim, from the previous formula for P_r , we see that P_r is a polynomial of degree $p-1$, so P_r can be written in the form

$$P_r(x) = (p-1)[x^{p-1} + a_{p-2}x^{p-2} + a_{p-3}x^{p-3} + \dots + a_1x + a_0],$$

where

$$a_{p-j} = (-1)^j \sum_{\substack{b_1, \dots, b_j \in \mathbb{F} \\ b_1, \dots, b_j \neq r}} b_1 \dots b_j,$$

for $j \in \{2, \dots, p-1\}$. Then

$$\begin{aligned} a_{p-j} &= \\ & (-1)^j \sum_{\substack{b_1=0 \\ b_1 \neq r}}^{p-1} \dots \sum_{\substack{b_{j-1}=0 \\ b_{j-1} \neq r}}^{p-1} \left[\sum_{\substack{b_j \in \mathbb{F} \\ b_j \neq r}} b_1 \dots b_{j-1} b_j \right] \\ &= (-1)^j \sum_{\substack{b_1=0 \\ b_1 \neq r}}^{p-1} \dots \sum_{\substack{b_{j-1}=0 \\ b_{j-1} \neq r}}^{p-1} \left[b_1 \dots b_{j-1} \sum_{\substack{b_j \in \mathbb{F} \\ b_j \neq r}} b_j \right] \\ &= (-1)^j \sum_{\substack{b_1=0 \\ b_1 \neq r}}^{p-1} \dots \sum_{\substack{b_{j-1}=0 \\ b_{j-1} \neq r}}^{p-1} \left[b_1 \dots b_{j-1} (-r) \right] \\ &= (-1)^j \sum_{\substack{b_1=0 \\ b_1 \neq r}}^{p-1} \dots \sum_{\substack{b_{j-1}=0 \\ b_{j-1} \neq r}}^{p-1} \left[b_1 \dots b_{j-2} (-r) \sum_{\substack{b_{j-1} \in \mathbb{F} \\ b_{j-1} \neq r}} b_{j-1} \right] \\ &= (-1)^j \sum_{\substack{b_1=0 \\ b_1 \neq r}}^{p-1} \dots \sum_{\substack{b_{j-1}=0 \\ b_{j-1} \neq r}}^{p-1} \left[b_1 \dots b_{j-2} (-r) (-r) \right] \\ &= (-1)^j (-r)^j \\ &= r^j. \end{aligned}$$

Finally,

$$a_0 = (-1)^{p-1} \prod_{\substack{a \in \mathbb{F} \\ a \neq r}} a = \prod_{\substack{a \in \mathbb{F} \\ a \neq r}} a.$$

Remark 8.1. Note that $p-1$ is even for $p > 2$ and $1 = -1$ for $p = 2$.

This completes the proof. \square

Finally, the polynomial form of Q_S is given in the following lemma.

Lemma 8.2. For $S \subset \mathbb{F}$, we have

$$Q_S(x) = \sum_{r \in \mathbb{F} \setminus S} P_r(x)$$

Proof. Clearly, if $x_0 \in S$, then $P_r(x_0) = 0$ for all $r \in \mathbb{F} \setminus S$. Therefore

$$Q_S(x_0) = \sum_{r \in \mathbb{F} \setminus S} P_r(x_0) = 0$$

Similarly, if $x_0 \notin S$, then $P_{x_0}(x_0) = 1$ and $P_r(x_0) = 0$ for all $r \neq x_0$ in $\mathbb{F} \setminus S$. Therefore

$$Q_S(x_0) = \sum_{r \in \mathbb{F} \setminus S} P_r(x_0) = 1.$$

This completes the proof. \square

9 Appendix 2

Here we give the proof of Theorem 4.2.

Proof. Let us first assume that the polynomial f is a nested canalizing function with canalizing input sets S_1, \dots, S_n and canalizing output values b_1, \dots, b_{n+1} . Then, by Theorem 3.1, f can be expanded as

$$f(x_1, \dots, x_n) = \sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) \prod_{i=1}^{n-j} Q_{S_i}(x_i) \right\} + b_1$$

$$= \sum_{j=0}^{n-1} \left\{ B_{n-j} \sum_{\substack{(i_1, \dots, i_{n-j}) \\ i_t \in \mathbb{F} \\ t=1, \dots, n-j}} a_{i_1}^1 \dots a_{i_{n-j}}^{n-j} x_1^{i_1} x_2^{i_2} \dots x_{n-j}^{i_{n-j}} \right\} \text{ for } j = 1, \dots, n-1, \text{ and}$$

$$+ b_1 + \sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) a_0^1 \dots a_0^{n-j} \right\},$$

where Q_{S_i} is defined as in Lemma 8.2, i.e.,

$$\begin{aligned} Q_{S_i}(x_i) &= \sum_{r \in S_i^c} P_r(x_i) \\ &= a_{p-1}^i x_i^{p-1} + a_{p-2}^i x_i^{p-2} + \dots + a_0^i. \end{aligned}$$

Now, from Lemma 8.1, we have that

$$\begin{aligned} C_{[p-1]} &= (b_{n+1} - b_n) a_{p-1}^1 \dots a_{p-1}^n \\ &= (b_{n+1} - b_n) \prod_{i=1}^n (p-1) |S_i^c|, \end{aligned}$$

and we have

$$\begin{aligned} C_{i_1 \dots i_{n-\mu}} &= \sum_{j=0}^{\mu} \left\{ B_{n-j} a_{i_1}^1 \dots a_{i_{n-\mu}}^{n-\mu} a_0^{n-\mu+1} \dots a_0^{n-j} \right\} \\ &= a_{i_1}^1 \dots a_{i_{n-\mu}}^{n-\mu} \sum_{j=0}^{\mu} \left\{ B_{n-j} a_0^{n-\mu+1} \dots a_0^{n-j} \right\} \end{aligned}$$

Now, if $j \neq n$, then

$$\begin{aligned} C_{[p-1] \setminus \{j\}}^{i_j} &= (b_{n+1} - b_n) a_{p-1}^1 \dots a_{p-1}^{j-1} a_{i_j}^j a_{p-1}^{j+1} \dots a_{p-1}^n \\ &= (b_{n+1} - b_n) a_{i_j}^j \prod_{\substack{i=1 \\ i \neq j}}^n a_{p-1}^i \end{aligned}$$

So

$$a_{i_j}^j = \left[(b_{n+1} - b_n) \prod_{\substack{i=1 \\ i \neq j}}^n a_{p-1}^i \right]^{-1} C_{[p-1] \setminus \{j\}}^{i_j}$$

$$C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} =$$

$$\sum_{j=0}^{\mu} \left\{ B_{n-j} a_{p-1}^1 \dots a_{p-1}^{n-\mu} a_0^{n-\mu+1} \dots a_0^{n-j} \right\}$$

$$= a_{p-1}^1 \dots a_{p-1}^{n-\mu} \sum_{j=0}^{\mu} \left\{ B_{n-j} a_0^{n-\mu+1} \dots a_0^{n-j} \right\}$$

$$= \left[\prod_{i=1}^{n-\mu} a_{p-1}^i \right] \sum_{j=0}^{\mu} \left\{ B_{n-j} a_0^{n-\mu+1} \dots a_0^{n-j} \right\}$$

So

$$\left[\sum_{j=0}^{\mu} \left\{ B_{n-j} a_0^{n-\mu+1} \dots a_0^{n-j} \right\} \right] =$$

$$\left[\prod_{i=1}^{n-\mu} a_{p-1}^i \right]^{-1} C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1}.$$

Now,

$$\begin{aligned} C_{i_1 \dots i_{n-\mu}} &= \prod_{j=1}^{n-\mu} a_{i_j}^j \left[\sum_{j=0}^{\mu} \left\{ B_{n-j} a_0^{n-\mu+1} \dots a_0^{n-j} \right\} \right] \\ &= \prod_{j=1}^{n-\mu} \left\{ \left[(b_{n+1} - b_n) \prod_{\substack{i=1 \\ i \neq j}}^n a_{p-1}^i \right]^{-1} \right. \\ &\quad \left. C_{[p-1] \setminus \{j\}}^{i_j} \right\} \left[\prod_{j=1}^{n-\mu} a_{p-1}^j \right]^{-1} C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} \\ &= \prod_{j=1}^{n-\mu} \left[(b_{n+1} - b_n) \prod_{i=1}^n a_{p-1}^i \right]^{-1} \\ &\quad C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} \prod_{j=1}^{n-\mu} C_{[p-1] \setminus \{j\}}^{i_j} \end{aligned}$$

$$\begin{aligned}
&= \prod_{j=1}^{n-\mu} \left[C_{[p-1]} \right]^{-1} C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} \prod_{j=1}^{n-\mu} C_{[p-1] \setminus \{j\}}^{i_j} \\
&= C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} \prod_{j=1}^{n-\mu} C_{[p-1]}^{-1} C_{[p-1] \setminus \{j\}}^{i_j}
\end{aligned}$$

Conversely, suppose Equations 4.2 - 4.7 hold for the coefficients of the polynomial $f(x_1, \dots, x_n)$ in Equation 4.8. We need to show that $f(x_1, \dots, x_n)$ is a nested canalyzing function. Let $a_{p-1}^j = (p-1) | S_j^c |$, $a_{i_j}^j = (p-1) \sum_{r \in S_j^c} r^{p-1-i_j}$, and $a_0^j = Q_{S_j}(0)$ for $j = 1, \dots, n$. Then

$$C_{[p-1]} = (b_{n+1} - b_n) \prod_{i=1}^n a_{p-1}^i,$$

and

$$C_{[p-1]}^{-1} C_{[p-1] \setminus \{j\}}^{i_j} = a_{i_j}^j [a_{p-1}^j]^{-1} \quad (9.1)$$

as well as

$$\begin{aligned}
&C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} = \\
&\left[\prod_{i=1}^{n-\mu} a_{p-1}^i \right] \sum_{j=0}^{\mu} \left\{ B_{n-j} a_0^{n-\mu+1} \dots a_0^{n-j} \right\}
\end{aligned}$$

Now from Equation 4.2,

$$C_{i_1 \dots i_{n-\mu}} = C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} \prod_{j=1}^{n-\mu} C_{[p-1]}^{-1} C_{[p-1] \setminus \{j\}}^{i_j}$$

and from Equation 9.1 we get

$$C_{i_1 \dots i_{n-\mu}} = C_{[0] \setminus \{1, \dots, n-\mu\}}^{p-1, \dots, p-1} \prod_{j=1}^{n-\mu} a_{i_j}^j [a_{p-1}^j]^{-1}.$$

Then

$$\begin{aligned}
C_{i_1 \dots i_{n-\mu}} &= \\
&\left[\prod_{i=1}^{n-\mu} a_{i_j}^i \right] \sum_{j=0}^{\mu} \left\{ B_{n-j} a_0^{n-\mu+1} \dots a_0^{n-j} \right\}
\end{aligned}$$

Now, from Equation 4.7 we get

$$C_{[0]} - b_1 = C_{[0] \setminus \{1\}}^{p-1} C_{[p-1] \setminus \{1\}}^0 C_{[p-1]}^{-1}.$$

Then

$$C_{[0]} = b_1 + \sum_{j=0}^{n-1} \left\{ B_{n-j} a_0^1 \dots a_0^{n-j} \right\}$$

$$f(x_1, \dots, x_n) =$$

$$\begin{aligned}
&\sum_{\mu=0}^{n-1} \left\{ \sum_{\substack{(i_1, \dots, i_{n-\mu}) \\ i_t \in \mathbb{F}_p \\ t=1, \dots, n-\mu}} C_{i_1 \dots i_{n-\mu}} x_1^{i_1} \dots x_{n-j}^{i_{n-j}} \right\} \\
&+ C_{[0]}.
\end{aligned}$$

Therefore

$$f(x_1, \dots, x_n) =$$

$$\begin{aligned}
&\sum_{j=0}^{n-1} \left\{ B_{n-j} \sum_{\substack{(i_1, \dots, i_{n-j}) \\ i_t \in \mathbb{F}_p \\ t=1, \dots, n-j}} a_{i_1}^1 \dots a_{i_{n-j}}^{n-j} x_1^{i_1} x_2^{i_2} \dots x_{n-j}^{i_{n-j}} \right\} \\
&+ b_1 + \sum_{j=0}^{n-1} \left\{ (b_{n-j+1} - b_{n-j}) a_0^1 \dots a_0^{n-j} \right\}.
\end{aligned}$$

Finally,

$$f(x_1, \dots, x_n) = \sum_{j=0}^{n-1} \left\{ B_{n-j} \prod_{i=1}^{n-j} Q_{S_i}(x_i) \right\} + b_1,$$

which is nested canalyzing by Theorem 3.1. This completes the proof. \square

References

- [1] J.T. Butler et al. (2005) Average path length of binary decision diagrams. *IEEE Trans. Comput.* **54**:1041–1053.
- [2] S. Harris et al. (2002) A model of transcriptional regulatory networks based on biases in the observed regulatory rules. *Complexity* 7 (4), 23-40.
- [3] Hinkelmann, F., Murrugarra, D., Jarrah, A.S., and Laubenbacher, R. (2010) A mathematical framework for agent-based models of complex biological networks. *Bull. Math. Biol.*, in press.
- [4] Abdul Salam Jarrah, Blessilda Raposa, and Reinhard Laubenbacher. Nested canalizing, unate cascade, and polynomial functions. *Physica D: Nonlinear Phenomena*, 233(2):167 – 174, 2007.
- [5] Winfried Just, Ilya Shmulevich, John Konvalina, “The number and probability of canalizing functions”, *Physica D* 197 (2004), pp. 211-221.
- [6] Stuart Kauffman, Carsten Peterson, B. Samuelsson, and Carl Troein. Random boolean network models and the yeast transcriptional network. *Proceedings of the National Academy of Sciences of the United States of America*, 100(25):14796–14799, 2003.
- [7] Stuart Kauffman, Carsten Peterson, Björn Samuelsson, and Carl Troein. Genetic networks with canalizing boolean rules are always stable. *Proceedings of the National Academy of Sciences of the United States of America*, 101(49):17102–17107, 2004.
- [8] R. Lidl and H. Niederreiter, *Finite Fields*, Cambridge University Press, New York, 1997.
- [9] R. Milo et al. (2004) Superfamilies of evolved and designed networks, *Science* **303**:1538–1542.
- [10] D. Murrugarra and R. Laubenbacher. (2011) Regulatory patterns in molecular interaction networks. Under review.
- [11] S. Nikolajewa, et al. *Boolean networks with biologically relevant rules show ordered behavior*. *BioSystems* 90 (2007) 40-47.
- [12] T. Sasao, et. al. *On the Number of Fanout-Free Functions and Unate Cascade Functions*. *IEEE trans. Comput.* 28 (1) (1979) 66-72.
- [13] Alan Veliz-Cuba, Abdul Salam Jarrah, and Reinhard Laubenbacher. Polynomial algebra of discrete models in systems biology. *Bioinformatics*, 26(13):1637–1643, 2010.
- [14] C.H. Waddington. Canalization of development and the inheritance of acquired characters. *Nature*, 150:563–565, November 1942.