QUADRATIC COVARIATION ESTIMATES IN NONSMOOTH STOCHASTIC CALCULUS

SERGIO ANGEL ALMADA MONTER AND YURI BAKHTIN

ABSTRACT. Given a Brownian Motion W, in this paper we study the asymptotic behavior, as $\varepsilon \to 0$, of the quadratic covariation between $f(\varepsilon W)$ and W in the case in which f is not smooth. Among the main features discovered is that the speed of the decay in the case $f \in C^{\alpha}$ is at least polynomial in ε and not exponential as expected. We use a recent representation as a backwardforward Itô integral of $[f(\varepsilon W), W]$ to prove an ε -dependent approximation scheme which is of independent interest. We get the result by providing estimates to this approximation. The results are then adapted and applied to generalize the results of [1], and [3] related to the Small Noise Exit from a Domian problem for the Saddle Case.

Non-smooth Itô's formula and Quadratic Variation and Large Deviation

1. INTRODUCTION

One of the central results of stochastic calculus is Itô's change of variables formula for twice differentiable transformations of semimartingales. It was realized recently that one also needs to study nonlinear maps that are not smooth enough to allow an application of the classical Itô formula. Various approaches to less regular changes of variables have been introduced, see [5], [6], [9], [11], [18], and references therein. These studies show that the key feature of the Itô formula, the quadratic covariation term, is well-defined under much weaker assumptions than those leading to the traditional formula. However, no nontrivial quantitative estimates of the arising quadratic covariation processes have appeared in the literature, to the best of our knowledge.

One area where such estimates are naturally needed is small random perturbations of dynamical systems. Often, in the course of a study of a stochastic system one has to make a simplifying change of coordinates, transforming the system locally to a simpler one. If the transformation map is C^2 , then one can apply the classical Itô calculus and easily control the Itô correction term. However, there are situations where a natural change of variables is less regular than C^2 , and in these cases there is no readily available tool that could be used to control the generalized Itô correction.

The goal of this paper is to close this gap and provide quantitative estimates on the generalized Itô correction term under nonclassical assumptions on the transformation.

Let us now be more precise. Let W be a standard 1-dimensional Wiener process on a complete probability space $(\Omega, \Sigma, \mathbf{P})$ and $\varepsilon > 0$ be a constant. If $q : \mathbb{R} \to \mathbb{R}$ is C^2 , then the classical Itô formula is (see [17, Section II.7])

$$g(\varepsilon W(t)) - g(0) = \varepsilon \int_0^t g'(\varepsilon W(s)) dW(s) + \frac{\varepsilon^2}{2} \int_0^t g''(\varepsilon W(s)) ds.$$

Introducing $f = g' \in C^1$, we can also rewrite the second term in the r.h.s. as quadratic covariation between $f(\varepsilon W)$ and εW : for $Q_{\varepsilon}(t) = [f(\varepsilon W), \varepsilon W](t)$, we have

$$Q_{\varepsilon}(t) = \varepsilon^2 \int_0^t f'(\varepsilon W(s)) ds, \quad t \ge 0.$$

In particular, for any T > 0, $\varepsilon^{-1} \sup_{t \leq T} Q_{\varepsilon}(t) \to 0$ in probability as $\varepsilon \to 0$. In this paper we show that this converges holds in the case in which f is not differentiable.

The motivation for this problem relies on small random perturbation of dynamical systems. Suppose that b is a vector field with a critical point at x^* and let Sdenote the flow generated by b:

$$\frac{d}{dt}S^t x = b(S^t x), \quad S^0 x = x.$$

It is well known (see Section 2.8 of [16]) that there is a continuous change of variables g so that locally around $g(x^*)$ the flow $g(S^tx)$ behaves like the linearized version of S. In the small random perturbation case, this combined with the traditional Itô formula imply (see, e.g. [3], [1]) that if g is at least C^2 , then the system

$$dX_{\varepsilon}(t) = b(X_{\varepsilon}(t))dt + \varepsilon dW(t), \quad X_{\varepsilon}(0) = x_0,$$

could be analyzed by working with the linear system

$$d\tilde{X}_{\varepsilon}(t) = \left(A\tilde{X}_{\varepsilon}(t) + \frac{\varepsilon^2}{2}\Phi_{\varepsilon}(X^{\varepsilon}(t))\right)dt + \varepsilon\sigma(X_{\varepsilon}(t))dW(t), \quad \tilde{X}_{\varepsilon}(0) = g(x_0),$$

where x_0 is close enough to x^* , A is the Jacobian of b at x^* , σ is at least a continuous matrix valued function, and $\varepsilon^2 \Phi_{\varepsilon}$ is the term corresponding to the quadratic covariation between $g'(X_{\varepsilon})$ and X_{ε} . There are well established cases for which g is known to be C^1 , see e.g. Hartman Theorem on Section 2.8 of [16]. In these cases, an already known C^1 formulation of Itô's formula implies a similar analogy between the non-linear and linear systems. Hence, estimates that show that in these cases the quadratic covariation term decays faster than the Itô term allow to reduce the local analysis to simpler exit problem for Ornstein-Uhlenbeck processes.

The analysis of the quadratic covariation [g'(X), X] in connection with extensions of Itô's formula for functions $g \notin C^2$ is fundamental for nonsmooth Itô calculus, see [8], [10], [11], [18], [19]. In [11], [18], [19] methods from backward stochastic calculus were used (see also the summary [20]), while in [8], [10] a local time approach was used. The basic result that has been explained in the cited literature from several points of view is that for T > 0,

(1)
$$Q_{\varepsilon}(t) = -\varepsilon \int_{0}^{t} f(\varepsilon W(s)) dW(s) - \varepsilon \int_{T-t}^{T} f(\varepsilon W(T-s)) dW(T-s) dW(T$$

where both integrals can be understood as Itô integrals w.r.t. appropriate filtrations. It is well known [17, page 389] that the integral with respect to $W(T - \cdot)$ in (1) is the time reversal of a semimartingale w.r.t. the natural filtration of $W(T - \cdot)$. Here the time reversal (with respect to T > 0) of a process X is understood as X(T - t) - X(T).

In this paper we exploit this structure by constructing an approximation scheme for Q_{ε} and using martingale techniques to show its consistency. As far as we know, this is the first attempt to use such a scheme in small noise analysis. See [4] for a related but different scheme for local time approximation in the case $\varepsilon = 1$. The text is organized as follows. In Section 2 we state our main results that include the martingale representation for the quadratic covariation, and, in Section 2.1, the results related to the application of non-smooth calculous to a particular small noise problem. In Section 3 we use the martingale representation to propose an approximation scheme that we then use to prove the key bound that the main results depend upon. The proofs of the main theorems are given in Section 4. In Section 6 proofs of auxiliary lemmas are given.

2. Main results

We are going to study $Q_{\varepsilon}(t) = [f(\varepsilon W), \varepsilon W](t)$ assuming that $f : \mathbb{R} \to \mathbb{R}$ is a bounded and uniformly Hölder or Lipschitz function, although these assumptions on f can be relaxed. It is convenient to formulate these assumptions in terms of modulus of continuity defined by:

$$\operatorname{osc}_{f}(\delta) = \sup_{|t-s| < \delta} |f(s) - f(t)|, \quad \delta > 0.$$

Throughout the text, we work with an arbitrary fixed number T > 0. We will not be explicit when including the dependency on T > 0 in the notation. We are ready to state the main results of the text.

Theorem 1. Suppose $\operatorname{osc}_f(\delta) \leq C_f \delta^{\alpha}$ for some $\alpha \in (0,1)$, $C_f > 0$, and all sufficiently small δ . Then, for every $\delta > 0$, $\gamma \in (0, \alpha)$, and $\mu \in (\gamma, \alpha)$, there are constants $\varepsilon_{\delta,\mu} > 0$ and $C_{\delta,\mu} > 0$ such that

$$\mathbf{P}\left\{\varepsilon^{-(1+\gamma)}\sup_{t\leq T}|Q_{\varepsilon}(t)|>\delta\right\}\leq C_{\delta,\mu}\varepsilon^{2(\alpha-\mu)/(1-\alpha)},\quad \varepsilon\in(0,\varepsilon_{\delta,\mu}).$$

In particular, for any $\gamma \in (0, \alpha)$,

$$\varepsilon^{-(1+\gamma)} \sup_{t \leq T} |Q_{\varepsilon}(t)| \xrightarrow{\mathbf{P}} 0, \quad \varepsilon \to 0.$$

This result is stronger than our initial claim that $\varepsilon^{-1}Q_{\varepsilon} \to 0$. Moreover, if α is close to 1, the exponent $1 + \gamma$ can be chosen to be close to 2.

The method we employ to prove this theorem produces the following estimate in the Lipschitz case where $\alpha = 1$:

Theorem 2. Suppose $\operatorname{osc}_f(\delta) \leq C_f \delta$, for some constant $C_f > 0$ and sufficiently small $\delta > 0$. Then, for every $\delta > 0$, $\gamma \in (0,1)$, and $\mu \in (\gamma,1)$, there are constants $\varepsilon_{\delta,\mu} > 0$ and $C_{\delta,\mu} > 0$ such that

$$\mathbf{P}\left\{\varepsilon^{-(1+\gamma)}\sup_{t\leq T}|Q_{\varepsilon}(t)|>\delta\right\}\leq C_{\delta,\mu}e^{-\varepsilon^{-(1-\mu)}},\quad\varepsilon\in(0,\varepsilon_{\delta,\mu}).$$

This theorem establishes that the rate of decay in probability is exponential in the Lipschitz case, which is coherent with the differentiable case in which almost sure convergence holds. For the Holder case, the method only shows a polynomial upper bound which in principle does not imply that the convergence rate can not be exponential.

The proof of Theorems 1 and 2 will be given in Section 4. An important part of the analysis is Theorem 9 given in Section 3 and in principle one can apply that result and its possible extensions to less regular functions f. The proof of Theorem 9 is in turn based on a forward-backward martingale representation of the quadratic covariation that we explain in Section 2.2. For now, we proceed to explain an application of the above results to a small noise problem studied in [1], [2], [3], [7], and [13].

2.1. Applications to the Small Noise Problem. In this section we consider the small noise scape from a saddle problem that, up to our knowledge, was first studied in [13]. The objective of the section is to establish the role that Theorems 1, and 2 have in the study of this problem. Let us start with the statement of the problem.

Consider a vector field $b : \mathbb{R}^d \to \mathbb{R}^d$, and a domain (open, bounded and convex set) $U \subset \mathbb{R}^d$ such that $0 \in U$ is the only critical point of b in the closure of U. That is, $0 \in U$ is the only $x \in \overline{U}$ such that b(x) = 0. Further, suppose that the vector field b is such that its Jacobian at 0, A = Db(0) has at least one eigenvalue with positive real part, and one eigenvalue with negative real part. Under this conditions, consider the flow S generated by b:

$$\frac{d}{dt}S^t x = b(S^t x), \quad S^0 x = x,$$

and its small noise perturbation,

$$dX_{\varepsilon}(t) = b(X_{\varepsilon})dt + \varepsilon dW(t).$$

The scape from a saddle problem is the study of the asymptotic behavior of the exit time

$$\tau_{\varepsilon}(x) = \inf \left\{ t > 0 : X_{\varepsilon}(t) \in \partial U \right\}, \quad x \in U,$$

and the exit location $X_{\varepsilon}(\tau_{\varepsilon}(x))$. The case of interest for this problem is when the initial condition for the diffusion $X_{\varepsilon}(0)$ lies in the invariant stable manifold

$$\mathcal{M}^s = \left\{ x : S^t x \to 0, \text{ as } t \to \infty \right\}.$$

The problem was first solved using a PDE approach in [13]. In that paper, it is shown that the exit time is asymptotically logarithmic in ε and that the exit location is concentrated on the intersection of ∂U and the invariant unstable manifold

$$\partial U \cap \mathcal{M}^u = \partial U \cap \{x : S^t x \to 0, \text{ as } t \to -\infty\}.$$

Later, [7] refined the result of the exit distribution in two dimensions, and further refinements were made in higher dimensions in [2].

In [3] a further generalization to the exit location was obtained, using the idea mentioned in the introduction of this paper. This result was later iterated to get the first result for a heteroclinic network, which is a more general case than the simple saddle case. The argument in [3] is as follows. It is well known (see Section 2.8 of [16]) that there is a continuous change of variables h so that locally around h(0) the flow $h(S^tx)$ behaves like the linearized version of S. For X_{ε} traditional Itô formula imply (see, e.g. [3], [1]) that if h is at least C^2 , then $\tilde{X}_{\varepsilon} = h(X_{\varepsilon})$ satisfies

$$d\tilde{X}_{\varepsilon}(t) = \left(A\tilde{X}_{\varepsilon}(t) + \frac{\varepsilon^2}{2}\Phi_{\varepsilon}(\tilde{X}_{\varepsilon}(t))\right)dt + \varepsilon\sigma(X_{\varepsilon}(t))dW(t), \quad \tilde{X}_{\varepsilon}(0) = h(X_{\varepsilon}(0)),$$

where x_0 is to 0, σ is at least a continuous matrix valued function, and $\varepsilon^2 \Phi_{\varepsilon}(\tilde{X}_{\varepsilon}(t))$ is the quadratic covariation term between the derivative of h evaluated at X_{ε} and X_{ε} itself. Under the assumption that $h \in C^2$, the above converges to 0 faster than the noise and hence it has no effect on the computation of the exit location. The limitation of this method is that the assumption $h \in C^2$ is quite restrictive. In [1] this restriction was studied by classifying systems that don't admit such a transformation h in the C^2 class. The results of [3] were extended in [1] in the two dimensional setting: the change of coordinates h transforms X_{ε} to a specific polynomial drift SDE in two dimensions which is then solved. In [1] it is also shown that this approach can not immediately be generalized to the high dimensional case. As a consequence, in this paper, we attack the high dimensional case by following the approach proposed in [3] but by allowing the transformation h to be not smooth.

We focus on a particular case to keep the exposition manageable. The novelty relies in the assumption on the smoothness for the change of coordinates, which is the main focus of the paper. The proof is a rearrangement of the main facts covered in the body of the paper, and its presented in Section 5. The theorem is stated in the spirit of Theorem 1 of [1].

Theorem 3. Suppose that A has spectrum $\lambda_1, ..., \lambda_d$ of real and simple eigenvalues such that $\lambda_1 > ... > \lambda_{\nu-1} > 0 > \lambda_{\nu} > ... > \lambda_d$ for some integer $\nu \leq d$. Also, assume that $h: U \to \mathbb{R}^d$ is a differentiable function with differentiable inverse, such that all its partial derivatives satisfy the conditions of Theorem 1 with $\alpha > 1/2$ and that $h(S^tx) = e^{At}h(x)$ in $U = (-\Delta, \Delta)^d$.

Denote $\partial U \cap \mathcal{M}^u = \{q_-, q_+\}$, and assume that $X_{\varepsilon}(0) = x_0 \in \mathcal{M}^s \cap U$. Then, there is a family of random vectors $(\phi_{\varepsilon})_{\varepsilon>0}$, a family of random variables $(\psi_{\varepsilon})_{\varepsilon>0}$, and a number

$$\beta = \begin{cases} 1, & \text{if } \nu = 2 \text{ and } -\lambda_{\nu} \ge \lambda_{1}, \\ -\frac{\lambda_{\nu}}{\lambda_{1}}, & \text{if } \nu = 2 \text{ and } -\lambda_{\nu} < \lambda_{1}, \\ 1 - \frac{\lambda_{1}}{\lambda_{2}}, & \text{if } \nu > 2 \text{ and } -\lambda_{\nu} \ge \lambda_{1} - \lambda_{2} \\ -\frac{\lambda_{\nu}}{\lambda_{1}}, & \text{if } \nu > 2 \text{ and } -\lambda_{\nu} < \lambda_{1} - \lambda_{2} \end{cases}$$

such that $X_{\varepsilon}(\tau_{\varepsilon}) = h(\Delta q_{\operatorname{sgn}\psi_{\varepsilon}}) + \varepsilon^{\beta}\phi_{\varepsilon}$, and the random vector

$$\left(\psi_{\varepsilon},\phi_{\varepsilon},\tau_{\varepsilon}-\frac{1}{\lambda_{1}}\log\varepsilon\right)$$

converges in distribution as $\varepsilon \to 0$.

Remark 4. The result is not a direct consequence of the results in this paper, since, as it will be clear at the beginning of Section 5, it requires a high dimensional version of the quadratic variation with drift. But we will see that to get this result the proof follows almost line by line the proof of the main results of this paper.

2.2. Forward-Backward Martingale Representation. The proof of Theorem 9 is based on a forward-backward martingale representation of the quadratic covariation. The focus of this section is to explain this representation as grounds to the full proof of Theorem 9 to be given in the next section. In order to do so, we need some conventions on our notation that we state as definition:

Definition 5. The time reversal of a process $X = (X(t))_{t \ge 0}$ with respect to T > 0 is defined by

$$\bar{X}(t) = X(T-t) - X(T), \quad t \in [0,T].$$

Likewise, the backward of X with respect to T > 0 is defined by

$$\hat{X}(t) = X(T-t), \quad t \in [0,T]$$

The starting point is the representation for $L_{\varepsilon} = \varepsilon^{-1}Q_{\varepsilon}$ implied by (1). For any T > 0,

(2)
$$L_{\varepsilon}(t) = -\int_0^t f(\varepsilon W(s))dW(s) - \int_{T-t}^T f(\varepsilon \hat{W}(s))d\hat{W}(s), \quad t \in [0,T].$$

We will find a convenient way to rewrite this expression using an enlargement of filtration approach. Denoting the natural filtration of a process $X = (X_t)_{t\geq 0}$ by $\mathcal{F}^X = (\mathcal{F}_t^X)_{t\geq 0}$, we note that that the integral with respect to W in (2) is an \mathcal{F}^W martingale, while the integral with respect to \hat{W} is the time reversal of the $\mathcal{F}^{\hat{W}}$ semimartingale

$$N_{\varepsilon}(t) = \int_{0}^{t} f(\varepsilon \hat{W}(s)) d\hat{W}(s).$$

Therefore, one of the terms in (2) is a martingale, while the other one has a nontrivial drift component. The following result reveals the structure of this time reversal.

Theorem 6. Let $\mathcal{G} = (\mathcal{G}_t)_{t \in [0,T]}$ be the minimum filtration such that W(T) is \mathcal{G}_0 measurable and $\mathcal{F}_t^{\hat{W}} \subset \mathcal{G}_t$. Then, \hat{W} is a \mathcal{G} semimartingale with Doob-Meyer decomposition given by

(3)
$$\hat{W}(t) = W(T) - \int_0^t \frac{\hat{W}(s)}{T-s} ds + \beta(t),$$

for some Brownian Motion β with respect to \mathcal{G} .

Moreover, if $\mathcal{H} = (\mathcal{H})_{t \in [0,T]}$ is the the minimum complete filtration such that W(T) is \mathcal{H}_0 measurable and $\mathcal{F}_t^\beta \subset \mathcal{H}_t$, then β is an \mathcal{H} Brownian Motion, \hat{W} is an \mathcal{H} semimartingale with the Doob-Meyer decomposition (3) and \hat{W} can be written as

(4)
$$\hat{W}(t) = W(T)(1 - t/T) + (T - t) \int_0^t \frac{d\beta(s)}{T - s}, \quad t \in [0, T].$$

Proof. The result follows from Theorem [17, Theorem VI.3].

Remark 7. In particular, since \hat{W} is \mathcal{H} adapted, for every function $F : \mathbb{R}^2 \to \mathbb{R}^2$ such that

$$\mathbf{E}\int_0^T F(\hat{W}(s))^2 ds < \infty,$$

the process $t \mapsto \int_{T-t}^{T} F(\hat{W}(s)) d\beta(s)$ is the time reversal of a martingale.

Using Theorem 6, we can obtain a representation for $L_{\varepsilon} = \varepsilon^{-1}Q_{\varepsilon}$. This is given in the following:

Corollary 8. Let \mathcal{H} be as in Theorem (6). Then, the process $L_{\varepsilon} = \varepsilon^{-1}Q_{\varepsilon}$ can be written as

(5)
$$L_{\varepsilon}(t) = -\int_{0}^{t} f(\varepsilon W(s)) dW(s) - \int_{T-t}^{T} f(\varepsilon \hat{W}(s)) d\beta(s) + \int_{0}^{t} f(\varepsilon W(s)) \frac{W(s)}{s} ds,$$

which is the sum of a \mathcal{F}^W martingale, a time reversal of an \mathcal{H} martingale and a bounded variation term.

Proof. This is an immediate consequence of Theorem 6 and (2) since W(t)/t is integrable on the interval [0, T] and f is bounded.

Theorem 6 is the main element we need to propose our approximation scheme, which is the main focus of Section 3.

3. SMALL NOISE ANALYSIS OF QUADRATIC COVARIATION.

In this section we study the quadratic covariation process $L_{\varepsilon} = [f(\varepsilon W), W]$. Recall the representation (5) given in Corollary 8. This will be one of the main ingredients in our proof.

Throughout this section, let $(n_{\epsilon})_{\epsilon>0}$ be integers such that $n_{\varepsilon} \nearrow \infty$ as $\varepsilon \to 0$. Let us define $(\delta_{\epsilon})_{\varepsilon>0}$ by $\delta_{\varepsilon} = T/n_{\varepsilon}$, and observe that $\delta_{\varepsilon} \searrow 0$ as $\epsilon \to 0$. The main result of this section is the following:

Theorem 9. Let $q_{\varepsilon} = 2\sqrt{\delta_{\varepsilon} |\log \delta_{\varepsilon}|}$, and let $(\gamma_{\varepsilon})_{\varepsilon>0}$ satisfy $\gamma_{\varepsilon} \to 0$ and

$$|\log \delta_{\varepsilon}| \frac{\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})}{q_{\varepsilon} \gamma_{\varepsilon}} \to 0, \quad \varepsilon \to 0.$$

Then, there are positive constants K_1, K_2, K_3 and ε_0 such that

$$\mathbf{P}\left\{\varepsilon^{-1}\sup_{t\leq T}|Q_{\varepsilon}(t)|>\gamma_{\varepsilon}\right\}\leq K_{1}\gamma_{\varepsilon}^{-1}e^{-K_{2}\gamma_{\varepsilon}^{2}/\mathrm{osc}_{f}(\varepsilon q_{\varepsilon})^{2}}+K_{3}\delta_{\varepsilon},\quad\varepsilon\in(0,\varepsilon_{0}).$$

The idea of the proof is to start with the representation (1) and use Theorem 6 to prove an approximation to each integral by a sum of increments. The result will follow once we combine the approximating sum for each integral into one. We will devote the rest of this section to developing this idea.

3.1. Approximating Processes. Let P_{ϵ} be the partition of the interval [0, T] given by points $0 = s_0 < ... < s_{n_{\epsilon}} = T$, where $s_i = i\delta_{\epsilon}$, for $i = 0, ..., n_{\epsilon}$. Also, define the backward partition \hat{P}_{ϵ} to be the partition of [0, T] given by points $0 = t_0 < ... < t_{n_{\epsilon}} = T$, where $t_i = T - s_{n_{\epsilon}-i}$.

For an arbitrary process Y and times $s, t \in [0, T]$ let $\Delta_{t,s}Y = Y(t) - Y(s)$. Then, for $t \in [0, T]$ we introduce the following notation:

(6)
$$S_{\epsilon}(t) = \int_{0}^{t} f(\varepsilon W(s)) dW(s),$$

(7)
$$\hat{S}_{\epsilon}(t) = \int_{T-t}^{T} f(\varepsilon \hat{W}(s)) d\hat{W}(s),$$

(8)
$$J_{\epsilon}(t) = \sum_{i=1}^{i(t)} f(\varepsilon W(s_{i-1})) \Delta_{s_i, s_{i-1}} W,$$

(9)
$$\hat{J}_{\epsilon}(t) = \sum_{i=1}^{i(t)} f(\varepsilon W(s_i)) \Delta_{s_i, s_{i-1}} W,$$

where i(t) is given by

$$i(t) = \min \left\{ j \in [0, n_{\varepsilon}] \cap \mathbb{Z} : s_j \ge t \right\}$$

The idea is to approximate each element S_{ε} and \hat{S}_{ε} with J_{ε} and \hat{J}_{ε} respectively, so we can approximate L_{ε} by $L_{\varepsilon,P_{\varepsilon}} = \hat{J}_{\varepsilon} - J_{\varepsilon}$. Note that since

$$f(\varepsilon W(s_i))\Delta_{s_i,s_{i-1}}W = -f(\varepsilon W(t_{n_{\varepsilon}-i}))\Delta_{t_{n_{\varepsilon}-i+1},t_{n_{\varepsilon}-i}}W,$$

after reordering the sum in (9), we can rewrite \hat{J}_{ε} as

(10)
$$\hat{J}_{\epsilon}(t) = -\sum_{i=n_{\varepsilon}-i(t)}^{n_{\varepsilon}-1} f(\varepsilon \hat{W}(t_i)) \Delta_{t_{i+1},t_i} \hat{W},$$

which is an integral sum of the Itô integral \hat{S}_{ε} . We will use Theorem 6 to justify the application of martingale techniques to prove that J_{ε} approximates S_{ε} and that \hat{J}_{ε} approximates \hat{S}_{ε} .

Once we have an approximation of L_{ε} by $L_{\varepsilon,P_{\varepsilon}}$, we notice that

(11)
$$L_{\varepsilon,P_{\varepsilon}}(t) = \sum_{i=1}^{i(t)} \Delta_{s_i,s_{i-1}} \left(f(\varepsilon W) \right) \Delta_{s_i,s_{i-1}} W$$

The differences in f in the above expression will be used to prove that $L_{\varepsilon,P_{\varepsilon}}(t)$ converges to 0 uniformly in probability and get the result.

We start with some preliminary results. The proofs will be postponed until Section 6 in order to keep the continuity of the paper. We state the next general lemma.

Lemma 10. Let $(M_{\varepsilon})_{\epsilon>0}$ be a family of martingales such that for every $\epsilon > 0$, $M_{\epsilon}(0) = 0$, the quadratic variation $\langle M_{\varepsilon} \rangle$ is absolutely continuous with respect to Lebesgue measure, and $\langle M_{\epsilon} \rangle (T) \leq r_{\epsilon}$. Then, for any $\delta > 0$,

$$\mathbf{P}\left\{\sup_{t\leq T}|M_{\epsilon}(t)|>\delta\right\}<\sqrt{\frac{8r_{\varepsilon}}{\pi\delta^2}}e^{-\delta^2/(2r_{\varepsilon})}.$$

We give a slight generalization of Levy's modulus of continuity lemma:

Lemma 11. For a Brownian motion B, define the modulus of continuity with respect to partition P_{ε} by

(12)
$$\delta_{B,\varepsilon} = \max_{i=1,\dots,n_{\varepsilon}} \sup_{s \in [s_{i-1},s_i]} |\Delta_{s,s_{i-1}}B|.$$

Then, there is a constant C > 0 independent of $\varepsilon > 0$ such that for any $\delta > 0$

$$\mathbf{P}\left\{\delta_{B,\varepsilon} > \delta\right\} \le \frac{C}{\delta\sqrt{\delta_{\varepsilon}}} e^{-\delta^2/(2\delta_{\varepsilon})}.$$

In particular, there is a $K_2 > 0$ such that

$$\mathbf{P}\left\{\delta_{B,\varepsilon} > q_{\varepsilon}\right\} \le K_2 \delta_{\varepsilon}, \quad \varepsilon > 0.$$

With these two results at hand we are ready to estimate $L_{\varepsilon,P_{\varepsilon}}$

Lemma 12. There is a positive constant K such that for any $\delta > 0$ and $\varepsilon > 0$,

$$\mathbf{P}\left\{\sup_{t\in[0,T]}|L_{\varepsilon,P_{\varepsilon}}(t)|>\delta\right\}\leq\mathbf{P}\left\{|\log\delta_{\varepsilon}|\mathrm{osc}_{f}(\varepsilon q_{\varepsilon})>\frac{q_{\varepsilon}\delta}{4T}\right\}+K\delta_{\varepsilon}.$$

Of course, the probability in the r.h.s. is either 0 or 1, and the estimate is meaningful only if the inequality in the curly brackets is violated.

Proof. Let us start with the simple inequality

(13)
$$\sup_{t \in [0,T]} L_{\varepsilon,P_{\varepsilon}}(t) \le \sum_{i=1}^{\circ} \left| \Delta_{s_i,s_{i-1}} f(\varepsilon W) \right| \left| \Delta_{s_i,s_{i-1}} W \right|,$$

n.

derived from (29). We estimate each term of the sum in the r.h.s. of (13).

From definition (12) it follows that

$$\max_{i=1,\dots,n_{\varepsilon}} |\Delta_{s_i,s_{i-1}} f(\varepsilon W)| \le \operatorname{osc}_f(\varepsilon \delta_{W,\varepsilon}).$$

Using this inequality and the definition of n_{ε} in (13), we see that

$$\sup_{t \in [0,T]} L_{\varepsilon,P_{\varepsilon}}(t) \le n_{\varepsilon} \operatorname{osc}_{f}(\varepsilon \delta_{W,\varepsilon}) \delta_{W,\varepsilon} \\ \le T \delta_{W,\varepsilon} \operatorname{osc}_{f}(\varepsilon \delta_{W,\varepsilon}) / \delta_{\varepsilon}.$$

Hence for every $\delta > 0$ the inequalities

$$\mathbf{P}\left\{\sup_{t\in[0,T]}|L_{\varepsilon,P_{\varepsilon}}|>\delta\right\}\leq\mathbf{P}\left\{\operatorname{osc}_{f}(\varepsilon\delta_{W,\varepsilon})\delta_{W,\varepsilon}>\delta_{\varepsilon}\delta/T,\ \delta_{W,\varepsilon}\leq q_{\varepsilon}\right\}+\mathbf{P}\left\{\delta_{W,\varepsilon}>q_{\varepsilon}\right\}$$
(14)
$$\leq\mathbf{P}\left\{\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})q_{\varepsilon}>\delta_{\varepsilon}\delta/T\right\}+\mathbf{P}\left\{\delta_{W,\varepsilon}>q_{\varepsilon}\right\}$$

hold. The second term in the r.h.s. of (14) can be bounded using Lemma 11, so we focus on the first term. For this notice that

$$\operatorname{osc}_f(\varepsilon q_\varepsilon) q_\varepsilon / \delta_\varepsilon = 4 |\log \delta_\varepsilon| \operatorname{osc}_f(\varepsilon q_\varepsilon) / q_\varepsilon,$$

which implies that

$$\mathbf{P}\left\{\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})q_{\varepsilon} > \delta_{\varepsilon}\delta/T\right\} \leq \mathbf{P}\left\{\left|\log \delta_{\varepsilon}|\operatorname{osc}_{f}(\varepsilon q_{\varepsilon}) > q_{\varepsilon}\delta/(4T)\right\}\right\}$$

The result follows after combining this fact with (14) and Lemma 11.

3.2. Approximation of L_{ε} by $L_{\varepsilon,P_{\varepsilon}}$. We have shown that $L_{\varepsilon,P_{\varepsilon}}$ converges to 0. In order to prove the convergence of L_{ε} we need to prove that $L_{\varepsilon,P_{\varepsilon}}$ approximates L_{ε} .

In order to do so define

$$M_{\epsilon}(t) := S_{\epsilon}(t) - J_{\epsilon}(t) + f(\varepsilon W(s_{i(t)-1}))\Delta_{s_{i(t)},t}W,$$

and

$$\hat{M}_{\epsilon}(t) := \hat{S}_{\epsilon}(t) + \hat{J}_{\epsilon}(t) + f(\varepsilon \hat{W}(t_{n_{\varepsilon}-i(t)})) \Delta_{T-t,n_{\varepsilon}-i(t)} \hat{W}.$$

Using (6), (8), and i(t), we see that the process M_{ε} can be written as

$$M_{\epsilon}(t) = \sum_{i=0}^{n_{\varepsilon}} \int_{s_{i-1} \wedge t}^{s_i \wedge t} \Delta_{s,s_{i-1}} f(\varepsilon W) dW(s).$$

Likewise, using (7), (9), (10) and the definition of the points t_i , we see that

(15)
$$\hat{M}_{\varepsilon}(t) = \sum_{i=0}^{n_{\varepsilon}-1} \int_{t_i \vee (T-t)}^{t_{i+1} \vee (T-t)} \Delta_{s,t_i} f(\varepsilon \hat{W}) d\hat{W}(s)$$
$$= \sum_{i=0}^{n_{\varepsilon}-1} \int_{t_i \vee (T-t)}^{t_{i+1} \vee (T-t)} \Delta_{s,t_i} f(\varepsilon \hat{W}) d\beta(s) - A_{\varepsilon}(t),$$

where we defined

(16)
$$A_{\varepsilon}(t) = \sum_{i=0}^{n_{\varepsilon}-1} \int_{t_i \vee (T-t)}^{t_{i+1} \vee (T-t)} \Delta_{s,t_i} f(\varepsilon \hat{W}) \frac{\hat{W}(s)}{T-s} ds.$$

Notice that M_{ε} is a \mathcal{F}^W martingale and \hat{M}_{ε} is the time reversal of a \mathcal{F}^{β} semimartingale. This is the main fact in the proof of the following Lemma: **Lemma 13.** There are positive constants K_1, K_2, K_3 and ε_0 such that for any $\delta > 0$,

$$\mathbf{P}\left\{\sup_{t\leq T}|\tilde{M}_{\epsilon}(t)|>\delta\right\}\leq (K_1/\delta)e^{-K_3\delta^2/\mathrm{osc}_f(\varepsilon q_{\varepsilon})^2}+K_2\delta_{\varepsilon},\quad \epsilon\in(0,\epsilon_0).$$

Here \tilde{M}_{ε} can be either M_{ε} or \hat{M}_{ε} .

The following lemma will be used in the proof of Lemma 13. The proof is postponed until Section 6.

Lemma 14. There are positive constants K_1, K_2, K_4 , and ε_0 such that for all $\delta > 0$,

$$\mathbf{P}\left\{\sup_{t\in(0,T)}|A_{\varepsilon}(t)|>\delta\right\}\leq (K_1/\delta)e^{-K_4\delta^2/\mathrm{osc}_f(\varepsilon q_{\varepsilon})^2}+K_2\delta_{\varepsilon},\quad \varepsilon\in(0,\varepsilon_0).$$

Proof of Lemma 13 . Let us start with the proof for M_{ε} . As we said before, the process M_{ε} is a martingale with quadratic variation $\Gamma_{\epsilon} = \langle M_{\epsilon} \rangle$ given by

(17)
$$\Gamma_{\epsilon}(t) = \sum_{i=1}^{n_{\varepsilon}} \int_{s_{i-1} \wedge t}^{s_i \wedge t} |\Delta_{s,s_{i-1}} f(\varepsilon W)|^2 ds.$$

In order to apply Lemma 10, we need to find a bound on the (random) function Γ_{ε} . In this case (12) implies that

$$\sup_{\in [s_{i-1},s_i]} |\Delta_{s,s_{i-1}} f(\varepsilon W)| \le \operatorname{osc}_f(\varepsilon \delta_{W,\varepsilon}),$$

for all $\varepsilon > 0$. Using this bound in (17) we see that

(18)
$$\Gamma_{\varepsilon}(T) \leq T \operatorname{osc}_{f}(\varepsilon \delta_{W,\varepsilon})^{2}$$

Lemma 10 implies that

$$\mathbf{P}\left\{\sup_{t\leq T}|M_{\varepsilon}(t)| > \delta\right\} \leq \mathbf{P}\left\{\sup_{t\leq T}|M_{\varepsilon}(t)| > \delta, \Gamma_{\varepsilon}(T) \leq T \operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2}\right\} \\
+ \mathbf{P}\left\{\Gamma_{\varepsilon}(T) > T \operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2}\right\} \\
(19) \qquad \leq \sqrt{8T \frac{\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2}}{\pi\delta^{2}}} e^{-\delta^{2}/(2T \operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2})} + \mathbf{P}\left\{\Gamma_{\varepsilon}(T) > T \operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2}\right\}$$

for all $\varepsilon > 0$ small enough. It remains to estimate the second probability in (19). Using (18) it easily follows that for each $\varepsilon > 0$,

$$\mathbf{P}\left\{\Gamma_{\varepsilon}(T) > T \operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2}\right\} \leq \mathbf{P}\left\{\operatorname{osc}_{f}(\varepsilon \delta_{W,\varepsilon}) > \operatorname{osc}_{f}(\varepsilon q_{\varepsilon})\right\}$$
$$\leq \mathbf{P}\left\{\delta_{W,\varepsilon} > q_{\varepsilon}\right\}.$$

Lemma 11 and (19) imply the desired estimate for M_{ε} .

To obtain the estimate on \hat{M}_{ε} , we notice that (15) and (16) imply

$$\hat{M}_{\varepsilon}(T-t) + A_{\varepsilon}(T-t) = \sum_{i=0}^{n_{\varepsilon}-1} \int_{t_i \vee t}^{t_{i+1} \vee t} \Delta_{s,t_i} f(\varepsilon \hat{W}) d\beta(s)$$
$$= \sum_{i=0}^{n_{\varepsilon}-1} \int_{t_i}^{t_{i+1}} \Delta_{s,t_i} f(\varepsilon \hat{W}) d\beta(s) - \sum_{i=0}^{n_{\varepsilon}-1} \int_{t_i \wedge t}^{t_{i+1} \wedge t} \Delta_{s,t_i} f(\varepsilon \hat{W}) d\beta(s).$$

Then, it follows that

$$\begin{split} \sup_{t \le T} \left| \hat{M}(t) - A_{\varepsilon}(t) \right| &= \sup_{t \le T} \left| \hat{M}(T - t) - A_{\varepsilon}(T - t) \right| \\ &\leq \left| \sum_{i=0}^{n_{\varepsilon} - 1} \int_{t_{i}}^{t_{i+1}} \Delta_{s,t_{i}} f(\varepsilon \hat{W}) d\beta(s) \right| + \sup_{t \le T} \left| \sum_{i=0}^{n_{\varepsilon} - 1} \int_{t_{i} \wedge t}^{t_{i+1} \wedge t} \Delta_{s,t_{i}} f(\varepsilon \hat{W}) d\beta(s) \right| \\ &\leq 2 \sup_{t \le T} \left| \sum_{i=0}^{n_{\varepsilon} - 1} \int_{t_{i} \wedge t}^{t_{i+1} \wedge t} \Delta_{s,t_{i}} f(\varepsilon \hat{W}) d\beta(s) \right|. \end{split}$$

Using this bound to proceed in the same way as we did for M_{ε} , we obtain that for any $\delta > 0$,

$$\mathbf{P}\left\{\sup_{t\leq T}\left|\hat{M}(t)-A_{\varepsilon}(T)\right|>\delta\right\}\leq (K_{1}/\delta)e^{-K_{2}\delta^{2}/\mathrm{osc}_{f}(\varepsilon q_{\varepsilon})^{2}}+K_{2}\delta_{\varepsilon},$$

for all $\varepsilon > 0$ small enough. Since

$$\mathbf{P}\left\{\sup_{t\leq T} \left|\hat{M}(t)\right| > \delta\right\} \leq \mathbf{P}\left\{\sup_{t\leq T} \left|\hat{M}(t) - A_{\varepsilon}(t)\right| > \delta/2\right\} + \mathbf{P}\left\{\sup_{t\leq T} \left|A_{\varepsilon}(t)\right| > \delta/2\right\},$$

the result follows from Lemma 14.

the result follows from Lemma 14.

A consequence of Lemma 13 is the approximation of the quadratic covariation $L_{\epsilon} = [f(\varepsilon W), W]$ by $L_{\varepsilon, P_{\varepsilon}}$, given in the following Lemma:

Lemma 15. If $(\gamma_{\varepsilon})_{\varepsilon>0}$ is such that $\gamma_{\varepsilon} \to 0$ and $\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})q_{\varepsilon}\gamma_{\varepsilon}^{-1} \to 0$ as $\varepsilon \to 0$, then there are positive constants K_1, K_2, K_5 , and ε_0 such that

$$\mathbf{P}\left\{\sup_{t\leq T}|L_{\epsilon}(t)-L_{\epsilon,P_{\epsilon}}(t)|>\gamma_{\varepsilon}\right\}\leq K_{1}\gamma_{\varepsilon}^{-1}e^{-K_{5}\gamma_{\varepsilon}^{2}/\mathrm{osc}_{f}(\varepsilon q_{\varepsilon})^{2}}+K_{2}\delta_{\varepsilon},\quad\epsilon\in(0,\epsilon_{0}).$$

Proof. Let $(\gamma_{\varepsilon})_{\varepsilon>0}$ be as in the statement of the Lemma. By the definition of M_{ε} and \hat{M}_{ε} , it follows that

(20)
$$|L_{\epsilon}(t) - L_{\epsilon,P_{\epsilon}}(t)| \leq |M_{\epsilon}| + |\tilde{M}_{\epsilon}| + |\Delta_{s_{i(t)},s_{i(t)-1}}f(\varepsilon W)\Delta_{s_{i(t)},s_{i(t)-1}}W|.$$

The result follows as a consequence of Lemmas 11 and 13. Indeed, since

$$|\Delta_{s_{i(t)},s_{i(t)-1}}f(\varepsilon W)\Delta_{s_{i(t)},s_{i(t)-1}}W| \le \operatorname{osc}_f(\varepsilon \delta_{W,\varepsilon})\delta_{W,\varepsilon},$$

Lemma 11 implies

$$\begin{aligned} \mathbf{P}\left\{|\Delta_{s_{i(t)},s_{i(t)-1}}f(\varepsilon W)\Delta_{s_{i(t)},s_{i(t)-1}}W| > \gamma_{\varepsilon}\right\} &\leq \mathbf{P}\left\{\operatorname{osc}_{f}(\varepsilon\delta_{W,\varepsilon})\delta_{W,\varepsilon} > \gamma_{\varepsilon}, \delta_{W,\varepsilon} \leq q_{\varepsilon}\right\} \\ &+ \mathbf{P}\left\{\delta_{W,\varepsilon} > q_{\varepsilon}\right\} \\ &\leq \mathbf{P}\left\{\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})q_{\varepsilon} > \gamma_{\varepsilon}\right\} + K_{2}\delta_{\varepsilon}.\end{aligned}$$

Hence, there is a $\varepsilon_0 > 0$ such that

$$\mathbf{P}\left\{|\Delta_{s_{i(t)},s_{i(t)-1}}f(\varepsilon W)\Delta_{s_{i(t)},s_{i(t)-1}}W| > \gamma_{\varepsilon}\right\} \le K_2\delta_{\varepsilon}, \quad \varepsilon \in (0,\varepsilon_0).$$

Using this bound and Lemma 13 in (20), we obtain

$$\mathbf{P}\left\{\sup_{t\leq T}|L_{\epsilon}(t)-L_{\epsilon,P_{\epsilon}}(t)|>\gamma_{\varepsilon}\right\}\leq K_{1}\gamma_{\varepsilon}^{-1}e^{-K_{5}\gamma_{\varepsilon}^{2}/\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2}}+K_{2}\delta_{\varepsilon},\quad \varepsilon\in(0,\varepsilon_{0}).$$

The proof is finished.

4. Proof of Theorems 1, 2 and 9

Proof of Theorem 9. The result is a consequence of Lemmas 12 and 15. Indeed, if $(\gamma_{\varepsilon})_{\varepsilon>0}$ is as in the statement of the Theorem, it is immediate to see that

(21)

$$\mathbf{P}\left\{\varepsilon^{-1}\sup_{t\leq T}|Q_{\varepsilon}(t)| > \gamma_{\varepsilon}\right\} \leq \mathbf{P}\left\{\sup_{t\leq T}|L_{\epsilon}(t) - L_{\epsilon,P_{\epsilon}}(t)| > \gamma_{\varepsilon}/2\right\} + \mathbf{P}\left\{\sup_{t\in[0,T]}|L_{\varepsilon,P_{\varepsilon}}(t)| > \gamma_{\varepsilon}/2\right\}.$$

The result will follow by applying Lemmas 12 and 15 to the two terms in r.h.s. of (21).

First, note that

$$\eta_{\varepsilon} = |\log \delta_{\varepsilon}| \frac{\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})}{q_{\varepsilon} \gamma_{\varepsilon}} \to 0, \quad \varepsilon \to 0,$$

implies that $\operatorname{osc}_f(\varepsilon q_\varepsilon)q_\varepsilon\gamma_\varepsilon^{-1} = 4\eta_\varepsilon\delta_\varepsilon \to 0$, as $\varepsilon \to 0$. Hence, from Lemma 15 we get that for some positive constants K'_1, K_2, K'_5 and ε'_0

(22)
$$\mathbf{P}\left\{\sup_{t\leq T}|L_{\epsilon}(t)-L_{\epsilon,P_{\epsilon}}(t)|>\gamma_{\varepsilon}/2\right\}\leq K_{1}'\gamma_{\varepsilon}^{-1}e^{-K_{5}'\gamma_{\varepsilon}^{2}/\operatorname{osc}_{f}(\varepsilon q_{\varepsilon})^{2}}+K_{2}\delta_{\varepsilon},$$

for all $\varepsilon \in (0, \varepsilon_0)$. Likewise, since $\eta_{\varepsilon} \to 0$ as $\varepsilon \to 0$, Lemma 12 implies that for some positive constants ε_1 and K,

(23)
$$\mathbf{P}\left\{\sup_{t\in[0,T]}|L_{\varepsilon,P_{\varepsilon}}(t)|>\gamma_{\varepsilon}/2\right\}\leq K\delta_{\varepsilon}, \quad \varepsilon\in(0,\varepsilon_{1}).$$

The result follows by using (22) and (23) in (21).

Proof of Theorem 1. The proof is a consequence of Theorem 9. Indeed, let us find a family $(\delta_{\varepsilon})_{\varepsilon>0}$ such that $\delta_{\varepsilon} \to 0$ and

$$\lim_{\varepsilon \to 0} |\log \delta_{\varepsilon}| \frac{\operatorname{osc}_f(\varepsilon q_{\varepsilon})}{q_{\varepsilon}} = 0.$$

Let $A(\delta_{\varepsilon}, \varepsilon) = |\log \delta_{\varepsilon}| \operatorname{osc}_{f}(\varepsilon q_{\varepsilon}) q_{\varepsilon}^{-1}$. A straightforward calculation gives

$$A(\delta_{\varepsilon},\varepsilon) \le C_f \varepsilon^{\alpha} \delta_{\varepsilon}^{(\alpha-1)/2} |\log \delta_{\varepsilon}|^{(\alpha+1)/2}.$$

Let $\mu \in (\gamma, \alpha)$ and take $\delta_{\varepsilon} = \varepsilon^{2(\alpha-\mu)/(1-\alpha)}$. Then, $A(\varepsilon^{2(\alpha-\mu)/(1-\alpha)}, \varepsilon) \leq \hat{A}(\varepsilon)$, where $\hat{A}(\varepsilon)$ is given by

$$\hat{A}(\varepsilon) = C_{\alpha,f} \varepsilon^{\mu} |\log \varepsilon|^{(\alpha+1)/2},$$

for some constant $C_{\alpha,f} > 0$ independent of $\varepsilon > 0$. So, we can use this δ_{ε} in Theorem 9 to get that

(24)
$$\mathbf{P}\left\{\varepsilon^{-1}\sup_{t\leq T}|Q_{\varepsilon}(t)| > \delta\right\} \leq K_{1}\delta^{-1}\exp\left\{-C_{0}\frac{\left(\delta\varepsilon^{-\alpha(1-\mu)/(1-\alpha)}\right)^{2}}{|\log\varepsilon|^{\alpha}}\right\} + K_{2}\varepsilon^{2(\alpha-\mu)/(1-\alpha)},$$

for all $\varepsilon > 0$ small enough and constants $K_1, K_2, C_0 > 0$ independent of $\varepsilon > 0$ and $\delta > 0$.

$$\square$$

Theorem 9 actually implies that inequality (24) remains true as long as $\hat{A}(\varepsilon)/\delta \rightarrow 0$, as $\varepsilon \rightarrow 0$. So, since $\gamma \in (0, \mu)$, we can substitute $\varepsilon^{\gamma} \delta$ for δ in (24) to get that

$$\mathbf{P}\left\{\varepsilon^{-(1+\gamma)}\sup_{t\leq T}|Q_{\varepsilon}(t)|>\delta\right\}\leq K_{1}\delta^{-1}\varepsilon^{-\gamma}\exp\left\{-C_{0}\frac{\delta^{2}\varepsilon^{2(-(\alpha-\gamma)+\alpha(\mu-\gamma))/(1-\alpha)}}{|\log\varepsilon|^{\alpha}}\right\}$$
(25)
$$+K_{2}\varepsilon^{2(\alpha-\mu)/(1-\alpha)}.$$

Since $\alpha \in (0, 1)$ and $\mu < \alpha$, we have

$$\alpha(\mu - \gamma) < \alpha(\alpha - \gamma) < \alpha - \gamma.$$

Using this fact in (25) we get that

$$\mathbf{P}\left\{\varepsilon^{-(1+\gamma)}\sup_{t\leq T}|Q_{\varepsilon}(t)|>\delta\right\}\leq K_{3}\varepsilon^{2(\alpha-\mu)/(1-\alpha)},$$

for some $K_3 > 0$, any $\delta > 0$, and all $\varepsilon > 0$ small enough. The result is proved. \Box

Proof of Theorem 2. The proof follows the same steps as the proof of Theorem 1. The first step is to to follow Theorem 9 by finding a family $(\delta_{\varepsilon})_{\varepsilon>0}$ such that $\delta_{\varepsilon} \to 0$ and

$$\lim_{\varepsilon \to 0} |\log \delta_{\varepsilon}| \varepsilon = 0$$

Given $\gamma \in (0, 1)$, we propose $\delta_{\varepsilon} = e^{-\varepsilon^{-(1-\mu)}}$, for $\mu \in (\gamma, 1)$. In this case, $|\log \delta_{\varepsilon}| \varepsilon = \varepsilon^{\mu}$, so Theorem 9 implies that

$$\mathbf{P}\left\{\varepsilon^{-1}\sup_{t\leq T}|Q_{\varepsilon}(t)|>\delta\right\}\leq K_{1}\delta^{-1}\exp\left\{-K_{4}\delta^{2}\varepsilon^{-(1+\mu)}e^{\varepsilon^{-(1-\mu)}}\right\}$$
$$+K_{2}e^{-\varepsilon^{-(1-\mu)}}.$$

As in the proof of Theorem 1, we can substitute $\delta \varepsilon^{\gamma}$ instead of δ in the last inequality. We can finish the proof by extracting the leading term in the resulting estimate.

5. Proof of Theorem 3

Using the results from [18] (see also [15], and reference therein) we observe that $Y_{\varepsilon} = h(X_{\varepsilon})$ satisfies

(26)
$$dY_{\varepsilon} = \nabla h(X_{\varepsilon}(t)) \cdot b(X_{\varepsilon}(t))dt + \varepsilon \nabla h(X_{\varepsilon}(t))dW(t) + \frac{1}{2}\mathcal{Q}_{\varepsilon}(t),$$

with initial condition $Y_{\varepsilon}(0) = h(X_{\varepsilon}(0))$, and where $\mathcal{Q}_{\varepsilon}$ is an \mathbb{R}^d -valued process with jth coordinate $\mathcal{Q}_{\varepsilon}^j$ given by

(27)
$$\mathcal{Q}_{\varepsilon}^{j}(t) = \sum_{k=1}^{d} \left[\partial_{k} h^{j}(X_{\varepsilon}), X_{\varepsilon}^{j} \right]$$
$$= \varepsilon \sum_{k=1}^{d} \left[\partial_{k} h^{j}(X_{\varepsilon}), W^{j} \right], \quad j = 1, ..., d$$

Differentiating with respect to t the identity $h(S^t x) = e^{At}h(x)$, we get that $\nabla h(x)b(x) = Ah(x)$, which combined with (26) implies

(28)
$$dY_{\varepsilon}(t) = AY_{\varepsilon}(t)dt + \varepsilon \left(\sigma(Y_{\varepsilon}(t))dW(t) + \varepsilon^{-1}\mathcal{Q}_{\varepsilon}(t)\right).$$

From this expression, to conclude the proof it is enough to show that the term $\varepsilon^{-1} \mathcal{Q}_{\varepsilon}$ in the last display converges uniformly (in an appropriate time range) towards zero in probability. The proof of this fact extends the results of this paper, but it follows the same steps (with minor modifications) as the proof of [3].

We are now going to establish what kind of convergence we need from term Q_{ε} in (28) to finish the proof, and then state the result in a separate lemma. Using the results of [2] that assert that $\frac{\tau_{\varepsilon}}{-\log \varepsilon}$ converges to a constant in probability, we can find for every v > 0, there is a large enough constant $K_v > 0$ such that

$$\mathbf{P}\left\{\tau_{\varepsilon} > -K_{\upsilon}\log\varepsilon\right\} \le \upsilon.$$

Since v is arbitrary, (27) and (28) imply that to finish the proof its enough to show that

$$\sup_{t \in [0, -K_{\upsilon} \log \varepsilon]} \left(\max_{j, k=1, \dots, d} \left[\partial_j h^k(X_{\varepsilon}), W^j \right] \right) \to 0$$

in probability as $\varepsilon \to 0$. Lemma 16 implies this result and hence finishes the proof of this theorem.

Lemma 16. Suppose $q : U \to \mathbb{R}$ is a function that satisfies the conditions of Theorem 1 with $\alpha > 1/2$. Then, for every $\Gamma > 0$ and $\delta > 0$ it follows that

$$\lim_{\varepsilon \to 0} \mathbf{P} \left\{ \sup_{t \in [0, -\Gamma \log \varepsilon]} [q(X_{\varepsilon}), W^j](t) > \delta, \tau_{\varepsilon} < -\Gamma \log \varepsilon \right\} = 0,$$

for every j = 1, ..., d.

Proof. The proof follow the exact same logic as the proof of Theorem 1 with slight modifications that we will point out. We keep the same notation as in Section 3.1 when appropriate. For instance, P_{ϵ} is a partition of the interval $[0, -\Gamma \log \varepsilon]$ given by points $0 = s_0 < \ldots < s_{n_{\epsilon}} = T_{\varepsilon} = -\Gamma \log \varepsilon$, where $s_i = i\delta_{\epsilon}$, for $i = 0, \ldots, n_{\epsilon}$. Also, define the backward partition \hat{P}_{ϵ} to be the partition of $[0, T_{\varepsilon}]$ given by points $0 = t_0 < \ldots < t_{n_{\epsilon}} = T_{\varepsilon} - s_{n_{\epsilon}-i}$. Let us fix j for the rest of the proof. Then, the idea is that the convergence

Let us fix j for the rest of the proof. Then, the idea is that the convergence towards 0 of the process $\Pi_{\varepsilon}(t) = [q(X_{\varepsilon}), W^j]$ conditioned on the sigma algebra $\mathcal{A}^j_{\varepsilon}$ generated by the history of W up to time T_{ε} except for the jth component of W, is almost identical from the main result in Theorem 1. We will show that this is the case, and then the proof will be finished due to the tower property of conditional expectations.

As mentioned before, [11], [18], [19] and [20] imply that upon fixing $\epsilon > 0$, conditioned on $\mathcal{A}_{\epsilon}^{j}$,

$$\amalg_{\varepsilon}(t) = -S_{\epsilon}(t) - \hat{S}_{\epsilon}(t),$$

where (in analogy with the notation used in Section 3.1) we defined

$$S_{\epsilon}(t) = \int_0^t f(X_{\varepsilon}(s)) dW^j(s), \text{ and } \hat{S}_{\epsilon}(t) = \int_{-\Gamma \log \varepsilon - t}^{-\Gamma \log \varepsilon} f(\hat{X}_{\varepsilon}(s)) d\hat{W}^j(s).$$

Here the time reversal is taken with respect to time $T_{\varepsilon} = -\Gamma \log \varepsilon$.

As we did before, the proof now consists on approximating the above difference by its respective sums and then show that the approximating sequence converges to 0. As expected to approximate the process Π_{ϵ} all steps will be the analogous to the ones followed in Section 3.1. In particular, \amalg_{ε} will be approximated by

(29)
$$L_{\varepsilon,P_{\varepsilon}}(t) = \sum_{i=1}^{i(t)} \Delta_{s_i,s_{i-1}} \left(q(X_{\varepsilon}) \right) \Delta_{s_i,s_{i-1}} W^j,$$

where i(t) is given by

$$i(t) = \min \left\{ j \in [0, n_{\varepsilon}] \cap \mathbb{Z} : s_j \ge t \right\}$$

To show that $L_{\varepsilon,P_{\varepsilon}}$ converges to 0, we follow line by line the proof of Lemma 12, with the only difference that n_{ε} is of order $-\delta_{\varepsilon}^{-1}\log\varepsilon$, and that the modulus of continuity of X_{ε} is now of the order $\max(\varepsilon\delta_{\varepsilon,W},\delta_{\varepsilon})$. Proceeding as described, we obtain that there is a positive constant K such that for any $\delta > 0$ and $\varepsilon > 0$, (30)

$$\mathbf{P}\left\{\sup_{t\in[0,-\Gamma\log\varepsilon]}|L_{\varepsilon,P_{\varepsilon}}(t)|>\delta\right\}\leq\mathbf{P}\left\{|\log\delta_{\varepsilon}|\operatorname{osc}_{q}(\max(\varepsilon q_{\varepsilon},q_{\varepsilon}^{2}))>\frac{q_{\varepsilon}\delta}{-4\Gamma\log\varepsilon}\right\}+K\delta_{\varepsilon}.$$

By choosing $\delta_{\varepsilon} = \varepsilon^2$, it follows that q_{ε} is of the order $-\varepsilon \log \varepsilon$, and $\max(\varepsilon q_{\varepsilon}, q_{\varepsilon}^2)$ is of the order $-\varepsilon^2(\log \varepsilon)^2$. Hence, in this case, from the last display, to ensure that $L_{\varepsilon,P_{\varepsilon}}$ converges to 0, we need that $\varepsilon^{2\alpha-1} \to 0$, as $\varepsilon \to 0$. That is, we need $\alpha > 1/2$, as stated in the statement of the theorem.

We are just left to show that the difference $L_{\varepsilon} - L_{\varepsilon,P_{\varepsilon}}$ converges to 0 under the additional conditions that δ_{ε} is of order ε^2 . In this case, the method used in Section 3.2 to proof Lemma 15 follow line by line with the appropriate modifications related to the modulus of continuity of X_{ε} , and the logarithmic grow of ε . We leave the reader to fill the details.

6. Additional Proofs

Proof of Lemma 10. For each $\epsilon > 0$, we use the representation of martingales as time changed Brownian Motion [12, Theorem 3.4.2] to see that $M_{\epsilon} = B(\langle M_{\epsilon} \rangle)$ in distribution in the space of continuous functions, for some Brownian Motion B (see [12, Theorem 3.4.2]). Therefore,

$$\mathbf{P}\left\{\sup_{t\leq T}|M_{\epsilon}(t)|>\delta\right\}\leq \mathbf{P}\left\{\sup_{t\leq r_{\epsilon}}|B(t)|>\delta\right\}.$$

Now the symmetry of B, reflection principle [12, Section 2.6], and Brownian scaling (self-similarity) imply that

$$\mathbf{P}\left\{\sup_{t\leq r_{\epsilon}}|B(t)|>\delta\right\} = \mathbf{P}\left\{\sup_{t\leq r_{\epsilon}}\max\{B(t), -B(t)\}>\delta\right\}$$
$$\leq 2\mathbf{P}\left\{\sup_{t\leq r_{\epsilon}}B(t)>\delta\right\}$$
$$\leq 4\mathbf{P}\left\{B(r_{\epsilon})>\delta\right\}$$
$$= 4\mathbf{P}\left\{\sqrt{r_{\epsilon}}B(1)>\delta\right\}.$$

The result follows by a standard Gaussian Tail estimate.

Proof of Lemma 11. Fix $\delta > 0$ and note that

(31)
$$\mathbf{P}\left\{\delta_{B,\varepsilon} > \delta\right\} \le \sum_{i=1}^{n_{\varepsilon}} \mathbf{P}\left\{\sup_{s \in (s_{i-1},s_i)} |\Delta_{s,s_{i-1}}B| > \delta\right\}.$$

We bound each of the probabilities in this sum. Since the process $\Delta_{s,s_{i-1}}B$ is equal in distribution, on the space of continuous functions, to a Brownian Motion itself up to a time shift, we can use reflection principle [12, Theorem 2.9.25] and standard Gaussian bounds to get

$$\mathbf{P}\left\{\sup_{s\in(s_{i-1},s_i)}|\Delta_{s,s_{i-1}}B|>\delta\right\}\leq 4\mathbf{P}\left\{B(\delta_{\varepsilon})>\delta\right\}\\\leq \delta^{-1}\sqrt{\frac{8\delta_{\varepsilon}}{\pi}}e^{-\delta^2/2\delta_{\varepsilon}}$$

Substituting this expression in (31) and using the fact that $n_{\varepsilon} \leq 2T/\delta_{\varepsilon}$, we see that there is a constant C > 0 independent of $\varepsilon > 0$ such that for any $\delta > 0$

$$\mathbf{P}\left\{\delta_{B,\varepsilon} > \delta\right\} \le \frac{C}{\delta\sqrt{\delta_{\varepsilon}}} e^{-\delta^2/(2\delta_{\varepsilon})}$$

as expected.

To prove the second part, use $\delta = q_{\varepsilon} = 2\sqrt{-\delta_{\varepsilon}\log\delta_{\varepsilon}}$ in the last expression to get that

$$\mathbf{P}\left\{\delta_{B,\varepsilon} > q_{\varepsilon}\right\} \leq \frac{C}{2\delta_{\varepsilon}\sqrt{-\log\delta_{\varepsilon}}}e^{2\log\delta_{\varepsilon}}$$
$$= \frac{C\delta_{\varepsilon}}{2\sqrt{-\log\delta_{\varepsilon}}}$$
$$\leq K_{2}\delta_{\varepsilon}.$$

Hence the result follows.

Proof of Lemma 14. We start with a basic inequality

$$\sup_{t \in [0,T]} |A_{\varepsilon}(t)| \leq \sum_{i=0}^{n_{\varepsilon}-1} \int_{s_i}^{s_{i+1}} |\Delta_{s,s_i} f(\varepsilon \hat{W})| \frac{|\hat{W}(s)|}{T-s} ds$$
$$\leq 2\sqrt{T} \operatorname{osc}_f(\varepsilon \delta_{W,\varepsilon}) \sup_{s \leq T} \frac{|\hat{W}(s)|}{\sqrt{T-s}}$$
$$\leq 2\sqrt{T} \operatorname{osc}_f(\varepsilon \delta_{W,\varepsilon}) \sup_{s \leq T} \frac{|W(s)|}{\sqrt{s}}.$$

It implies that

$$\begin{split} \mathbf{P}\left\{\sup_{t\in[0,T]}|A_{\varepsilon}(t)|>\delta\right\} &\leq \mathbf{P}\left\{\sup_{s\leq T}\frac{|W(s)|}{\sqrt{s}}>\frac{\delta}{2\mathrm{osc}_{f}(\varepsilon q_{\varepsilon})\sqrt{T}}\right\} + \mathbf{P}\left\{\delta_{W,\varepsilon}>q_{\varepsilon}\right\}\\ &\leq \mathbf{P}\left\{\sup_{s\leq T}\frac{|W(s)|}{\sqrt{s}}>\frac{\delta}{2\mathrm{osc}_{f}(\varepsilon q_{\varepsilon})\sqrt{T}}\right\} + K_{1}\delta_{\varepsilon}, \end{split}$$

for some constant $K_1 > 0$ independent of $\delta > 0$ and $\varepsilon > 0$. To finish the proof, we need to study the tail probability of the random variable $A = \sup_{s < T} |W(s)|/\sqrt{s}$.

In order to study the tail decay of the random variable A, note that, due to the symmetry of Brownian Motion,

$$\mathbf{P}\left\{A > \delta\right\} \le 2\mathbf{P}\left\{\sup_{t \le T} \frac{W(t)}{\sqrt{t}} > \delta\right\}.$$

So it is sufficient to focus on the tail probabilities of the random variable $N = \sup_{t \leq T} (W(t)/\sqrt{t})$, which is the supremum of a Gaussian process.

Equip the interval [0, T] with the metric ρ given by

$$\rho(s,t)^2 = \mathbf{E} \left(\frac{W(s)}{\sqrt{s}} - \frac{W(t)}{\sqrt{t}} \right)^2$$
$$= 2 \left(1 - \sqrt{\frac{s \wedge t}{s \vee t}} \right), \quad s,t \in [0,T].$$

We denote by $B_{\theta}(t) \subset [0,T]$ the ρ -ball of radius $\theta > 0$ centered at $t \in [0,T]$. Let H_{θ} be the minimum number of balls of radius θ needed in order to cover [0,T]. According to [14][Section 14, Theorem 1], if

(32)
$$\int_0^{\sigma/2} \sqrt{|\log H_\theta|} d\theta < \infty,$$

with $\sigma = \sup_{t \in [0,T]} \operatorname{var}(W(t)/\sqrt{t}) = 1$, then $\mathbf{E}N < \infty$. Then, it is standard to see [14][Corollary 2, Section 14] that there is a $\zeta_0 > \mathbf{E}N$, such that for any $\zeta > \zeta_0$

(33)
$$\mathbf{P}\left\{|N - \mathbf{E}N| > \zeta\right\} \le Ce^{-\zeta^2/2}/\zeta,$$

for some universal constant C > 0.

In our situation, if the integral in (32) is finite, this will be enough to finish the proof. Indeed, assuming (33), there is an $\varepsilon_0 > 0$ such that

$$\begin{split} \mathbf{P} \left\{ \sup_{s \leq T} \frac{|W(s)|}{\sqrt{s}} > \frac{\delta}{2 \mathrm{osc}_f(\varepsilon q_\varepsilon) \sqrt{T}} \right\} &\leq 2 \mathbf{P} \left\{ N > \frac{\delta}{2 \mathrm{osc}_f(\varepsilon q_\varepsilon) \sqrt{T}} \right\} \\ &\leq 2 \mathbf{P} \left\{ N - \mathbf{E}N > \frac{\delta}{2 \mathrm{osc}_f(\varepsilon q_\varepsilon) \sqrt{T}} - \mathbf{E}N \right\} \\ &\leq C_1 \left(\frac{\delta}{2\sqrt{T} \mathrm{osc}_f(\varepsilon q_\varepsilon)} - \mathbf{E}N \right)^{-1} \exp \left\{ -C_2 \left(\frac{\delta}{2\sqrt{T} \mathrm{osc}_f(\varepsilon q_\varepsilon)} - \mathbf{E}N \right)^2 \right\} \\ &\leq (C_1/\delta) \exp \left\{ -C_3 \frac{\delta^2}{\mathrm{osc}_f(\varepsilon q_\varepsilon)^2} \right\}, \end{split}$$

for some constants $C_1, C_2, C_3 > 0$ independent of ε and δ , and all $\varepsilon \in (0, \varepsilon_0)$. Hence we just need to show that the integral (32) is finite.

We are going to give an estimate of H_{θ} , $\theta \in (0, 1/2)$. Suppose $0 \leq s < t \leq T$, then $s \in B_{\theta}(t)$ if and only if

$$\sqrt{s} \ge \sqrt{t}(1 - \theta^2/2).$$

Therefore, if s and t belong to the same ball of radius $\theta \in (0, 1/2)$, then

$$|t-s| \le T\theta^2.$$

Hence, $H_{\theta} \leq 2/\theta^2$, and $\sqrt{|\log H_{\theta}|}$ is integrable on the interval [0, 1/2], which implies our claim.

References

- Sergio Angel Almada and Yuri Bakhtin. Normal forms approach to diffusion near hyperbolic equilibria. Nonlinearity, 24(6):1883, 2011.
- [2] Yuri Bakhtin. Exit asymptotics for small diffusion about an unstable equilibrium. Stochastic Process. Appl., 118(5):839–851, 2008.
- [3] Yuri Bakhtin. Noisy heteroclinic networks. Probability Theory and Related Fields, 150(1-2):1–42, 2011.
- [4] Blandine Bérard Bergery and Pierre Vallois. Approximation via regularization of the local time of semimartingales and Brownian motion. *Stochastic Process. Appl.*, 118(11):2058–2070, 2008.
- [5] N. Bouleau. Formules de changement de variables. Ann. Inst. H. Poincaré Probab. Statist., 20(2):133-145, 1984.
- [6] Nicolas Bouleau and Marc Yor. Sur la variation quadratique des temps locaux de certaines semimartingales. C. R. Acad. Sci. Paris Sér. I Math., 292(9):491–494, 1981.
- [7] Martin V. Day. On the exit law from saddle points. Stochastic Processes and their Applications, 60(2):287 - 311, 1995.
- [8] Nathalie Eisenbaum. Integration with respect to local time. Potential Anal., 13(4):303–328, 2000.
- [9] Nathalie Eisenbaum. On Itô's formula of Föllmer and Protter. In Séminaire de Probabilités, XXXV, volume 1755 of Lecture Notes in Math., pages 390–395. Springer, Berlin, 2001.
- [10] Nathalie Eisenbaum. Local time-space calculus for reversible semimartingales. In Séminaire de Probabilités XL, volume 1899 of Lecture Notes in Math., pages 137–146. Springer, Berlin, 2007.
- [11] Hans Föllmer, Philip Protter, and Albert N. Shiryayev. Quadratic covariation and an extension of Itô's formula. *Bernoulli*, 1(1-2):149–169, 1995.
- [12] Ioannis Karatzas and Steven E. Shreve. Brownian motion and stochastic calculus, volume 113 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1988.
- [13] Yuri Kifer. The exit problem for small random perturbations of dynamical systems with a hyperbolic fixed point. Israel J. Math., 40(1):74–96, 1981.
- [14] M. A. Lifshits. Gaussian random functions, volume 322 of Mathematics and its Applications. Kluwer Academic Publishers, Dordrecht, 1995.
- [15] S. Moret and D. Nualart. Generalization of itô's formula for smooth nondegenerate martingales. Stochastic Processes and their Applications, 91(1):115 – 149, 2001.
- [16] Lawrence Perko. Differential equations and dynamical systems, volume 7 of Texts in Applied Mathematics. Springer-Verlag, New York, third edition, 2001.
- [17] Philip E. Protter. Stochastic integration and differential equations, volume 21 of Applications of Mathematics (New York). Springer-Verlag, Berlin, second edition, 2004. Stochastic Modelling and Applied Probability.
- [18] F. Russo and P. Vallois. Itô formula for C¹-functions of semimartingales. Probab. Theory Related Fields, 104(1):27–41, 1996.
- [19] Francesco Russo and Pierre Vallois. The generalized covariation process and Itô formula. Stochastic Process. Appl., 59(1):81–104, 1995.
- [20] Francesco Russo and Pierre Vallois. Elements of stochastic calculus via regularization. In Séminaire de Probabilités XL, volume 1899 of Lecture Notes in Math., pages 147–185. Springer, Berlin, 2007.